



**Universiteit  
Leiden**  
The Netherlands

**Analecta Praehistorica Leidensia 37/38 / Schipluiden : a neolithic settlement on the Dutch North Sea coast c. 3500 CAL BC**

Kooijmans, L.P.L.; Jongste, P.; et al., ; Jongste, P.F.B.; Kooijmans, L.P.L.

**Citation**

Kooijmans, L. P. L., Jongste, P., & Et al.,. (2006). Analecta Praehistorica Leidensia 37/38 / Schipluiden : a neolithic settlement on the Dutch North Sea coast c. 3500 CAL BC, 516.  
Retrieved from <https://hdl.handle.net/1887/33080>

Version: Not Applicable (or Unknown)  
License: [Leiden University Non-exclusive license](#)  
Downloaded from: <https://hdl.handle.net/1887/33080>

**Note:** To cite this publication please use the final published version (if applicable).

# ANALECTA PRAEHISTORICA LEIDENSIA 37/38

---

PUBLICATION OF THE FACULTY OF ARCHAEOLOGY  
LEIDEN UNIVERSITY

## SCHIJPLUIDEN


A NEOLITHIC SETTLEMENT ON THE DUTCH  
NORTH SEA COAST *c.* 3500 CAL BC

EDITED BY LEENDERT P. LOUWE KOOIJMANS  
AND PETER F.B. JONGSTE



LEIDEN UNIVERSITY 2006

The publication of this volume was made possible by financial and organisational support from:



Translation by Susan Mellor

Series editors: Corrie Bakels / Hans Kamermans

Copy editors of this volume: Leendert Louwe Kooijmans / Peter Jongste

Editors of illustrations: Walter Laan and Alastair Allen, Archol BV

Copyright 2006 by the Faculty of Archaeology, Leiden

ISSN 0169-7447

ISBN-10: 90-73368-21-9

ISBN-13: 978-90-73368-21-7

Subscriptions to the series *Analecta Praehistorica Leidensia*  
and single volumes can be ordered exclusively at:

Faculty of Archaeology  
P.O. Box 9515  
NL-2300 RA Leiden  
the Netherlands

# Contents

## **Preface** IX

*Leendert Louwe Kooijmans*

## **PART I INTRODUCTION** 1

### **1 Discovery and working method** 3

*Peter Jongste*

*Leendert Louwe Kooijmans*

### **2 Stratigraphy and chronology of the site** 19

*Joanne Mol*

*Leendert Louwe Kooijmans*

*Tom Hamburg*

### **3 Features** 39

*Tom Hamburg*

*Leendert Louwe Kooijmans*

### **4 The archaeological remains: a critical spatial approach** 67

*Milco Wansleebe*

*Leendert Louwe Kooijmans*

## **PART II MAN AND MATERIALS** 89

### **5 Graves and human remains** 91

*Liesbeth Smits*

*Leendert Louwe Kooijmans*

### **6 The Schipluiden pottery** 113

*Daan Raemaekers*

*Michiel Rooke*

### **7 Flint, procurement and use** 129

*Annelou van Gijn*

*Veronique van Betuw*

*Annemieke Verbaas*

*Karsten Wentink*

### **8 Stone, procurement and use** 167

*Annelou van Gijn*

*Rob Houkes*

- 9     **Ornaments of jet, amber and bone** 195  
        *Annelou van Gijn*
- 10    **Implements of bone and antler: a Mesolithic tradition continued** 207  
        *Annelou van Gijn*
- 11    **Wooden artefacts** 225  
        *Leendert Louwe Kooijmans*  
        *Laura Kooistra*
- 12    **Fabrics of fibres and strips of bark** 253  
        *Laura Kooistra*
- 13    **Birch bark tar** 261  
        *Annelou van Gijn*  
        *Jaap Boon*
- PART III   ECOLOGY AND ECONOMY 267
- 14    **Coastal evolution of Delfland and the Schipluiden microregion in relation to Neolithic settlement** 269  
        *Joanne Mol*
- 15    **Diatoms** 285  
        *Hein de Wolf*  
        *Piet Cleveringa*
- 16    **Molluscs** 297  
        *Wim Kuijper*
- 17    **Coprolites, macroscopic analysis** 301  
        *Mark van Waijjen*  
        *Caroline Vermeeren*
- 18    **Pollen analysis and the reconstruction of the former vegetation** 305  
        *Corrie Bakels*
- 19    **Botanical remains and plant food subsistence** 317  
        *Lucy Kubiak-Martens*
- 20    **Roots, tubers and processed plant food in the local diet** 339  
        *Lucy Kubiak-Martens*
- 20a   **Analytical report on some archaeological charred residues from Schipluiden** 353  
        *Jaap Boon*
- 21    **Wood and charcoal** 363  
        *Laura Kooistra*
- 22    **Mammals** 375  
        *Jørn Zeiler*

<b>23</b>	<b>Birds</b>	<b>421</b>
	<i>Jørn Zeiler</i>	
<b>24</b>	<b>Background fauna: small mammals, amphibians and reptiles</b>	<b>443</b>
	<i>Jørn Zeiler</i>	
<b>25</b>	<b>Fish</b>	<b>449</b>
	<i>Dick Brinkhuizen</i>	
<b>26</b>	<b>Insects</b>	<b>471</b>
	<i>Tom Hakbijl</i>	
<b>PART IV SYNTHESIS</b>		<b>483</b>
<b>27</b>	<b>Schipluiden: a synthetic view</b>	<b>485</b>
	<i>Leendert Louwe Kooijmans</i>	

*Shell samples were collected from several clastic deposits in the sections and borings. They inform us about the salinity and dynamics of the environment before and shortly after the period of occupation, and in the Gantel system. Shell pottery temper was identified as mainly crushed cockle.*

### 16.1 INTRODUCTION

In the archaeological investigation special attention was paid to the occurrence of molluscs as they could have provided information on the local environment before, during and after the period of occupation and on the part played by molluscs in the occupants' diet. In actual fact, however, molluscs were encountered in only small quantities.

Shell fragments were only sporadically encountered in the settlement area, but part of the pottery was tempered with ground shells. A very small number of tiny shell fragments were found in the zoological analysis of sieve residues.

Shells were found in several deposits in the borings. The overlying Dunkirk I deposits in particular were found to contain dense concentrations of shells. The sediments of the boring samples were rinsed with water through a sieve with a mesh width of 0.25 mm. The residues were then microscopically examined.

See chapters 2 and 14 for information on the stratigraphy and the physical geography of the excavated settlement and its surroundings.

### 16.2 RESULTS

#### 16.2.1 The settlement

Virtually no molluscs were found in the actual settlement area or in the surrounding deposits. As far as the dune itself is concerned, this could be attributable to dissolution in the decalcified dune sand, in which bone had indeed also survived in only very small quantities and in a poor state of preservation. This does however not hold for the peripheral zone. The conclusion must hence be that species such as edible mussel and edible cockle, which are commonly – and often in large quantities – encountered in coastal settlements, did not feature on the occupants' menu.

At Schipluiden only 16 tiny fragments of marine mollusc species were encountered in 11 find assemblages. Grouped according to phase they are as follows:

phase 1	mussel ( <i>Mytilus edulis</i> )
phase 1-2a	mussel, peppery furrow shell ( <i>Scrobicularia plana</i> )
phase 2a	peppery furrow shell, lagoon cockle ( <i>Cerastoderma glaucum</i> )
phase 3	peppery furrow shell.
phases 1-3	mussel, edible cockle ( <i>Cerastoderma edule</i> ), peppery furrow shell

These four species and the small numbers in which they were found do not yield any information suggesting changes in the local environment in the course of the investigated period. They do probably not represent the remains of consumed molluscs. The fragments have thin walls and appear to derive from small specimens. They may have been brought to the site by humans (whether or not deliberately) or deposited there during high tides. The encountered species may well have lived in a marine environment with a reduced saline content (brackish).

One of the samples (phases 1-2a) contained a few fragments of the shells of land snails: an amber snail and the shell of the beautiful snail. Both species may have occurred naturally at the site; they are common species characteristic of a wide range of open terrestrial environments, including high-lying salt marshes and damp dunes.

There will undoubtedly have been other snail and mussel species both on land and in the water, though the mollusc fauna may have been poor due to extreme conditions, for example extreme temperatures or saline contents.

#### 16.2.2 Borings in the surrounding area

In the boring research that was conducted in the site's surroundings with the aim of reconstructing the former landscape (see chapter 14) a few small shell-containing samples were collected of several successive deposits (table 16.1). The majority had a volume of only 10 cm<sup>3</sup>.

The plant and animal remains contained in these very small samples (table 16.1) yielded the following environmental evidence.

The Early Holocene clay was deposited in a coastal area under saline and brackish conditions. The area was shielded from the open sea and was a kind of tidal-flat area or a lagoon.

boring	68	68	73	173	174	68	181	151	181
depth in cm below surface	645	460	370 - 377	460 - 470	470	317 - 320	370 - 375	413 - 424	260 - 270
sediment	sandy cl.	sandy cl.	wd/sand	cl. sand	cl. sand	sandy cl.	cl. sand	clay	clay
Unit	Earl. Holo.	R-Z	R-Z	R-Z	R-Z	40/19	40/20	2	2
volume in cm <sup>3</sup>	10	10	5	10	10	10	10	100	50
molluscs									
<i>Barnea candida</i>	—	—	1	—	—	—	—	—	—
<i>Cerastoderma edule</i>	—	1	1	x	—	—	—	x	—
<i>Mytilus edulis</i>	x	x	x	x	x	1	1	—	x
<i>Littorina saxatilis</i>	x	—	—	—	—	—	—	—	—
<i>Macoma balthica</i>	—	—	1 doublet	—	—	—	—	x	x
<i>Scrobicularia plana</i>	—	1 doublet	—	x, 1 doublet	1 doublet	1 doublet	—	—	1
<i>Spisula subtruncata</i>	—	1	—	—	—	—	—	—	—
<i>Hydrobia ulvae</i>	xx	1	x	x	x	x	—	x	x
<i>Hydrobia ventrosa</i>	xx	—	—	—	—	—	—	xx	—
other animal remains									
foraminifers	—	—	x	x	x	—	—	xxx	xx
ostracods	—	—	x	xx	x	x	—	xxx	xx
<i>Nereis</i> sp. (a ragworm) jaws	x	—	—	1	—	1	—	x	—
<i>Echinocardium cordatum</i> spines	—	x	x	x	x	x	—	—	x
sea-mats	—	—	1	—	—	—	—	—	—
sponges spicula	—	—	—	—	—	—	—	x	—
plant remains (seeds)									
<i>Agrostis</i> sp.	—	—	—	—	—	—	—	—	1
<i>Aster tripolium</i>	—	—	—	—	—	—	—	—	1
<i>Atriplex</i> sp.	—	—	—	—	—	—	—	—	1
<i>Bolboschoenus maritimus</i>	—	—	—	—	—	—	1	—	—
<i>Carex</i> sp.	—	—	—	—	—	—	—	x	1
<i>Chenopodium glaucum/rubrum</i>	—	—	—	—	—	—	—	—	2
<i>Mentha</i> sp.	—	—	—	—	—	—	—	4	—
<i>Suaeda maritima</i>	—	1	—	—	—	—	16	—	—
<i>Scirpus</i> sp.	—	1	—	—	—	—	—	—	—
charcoal	—	x	—	—	—	—	—	—	—
x = 2 - 20      cl = clay, clayey xx = 20 - 50      wd = wood xxx = hundreds      R-Z = Rijswijk-Zoetermeer sands									

Table 16.1 Identifications of molluscs from bore hole samples.

The Rijswijk-Zoetermeer sands originated in a saline (marine) environment (tidal-flat area) that was connected to the open sea. One sample (boring 73) contained a piece of wood with drill holes and a white paddick shell.

Unit 40/Unit 19 originated in a saline tidal-flat area.

Unit 2 was deposited in a brackish (or saline) coastal area: a calm tidal-flat area or a lagoon.

Unit 0 (Dunkirk) was formed in the Gantel system (see section 2.2.3). Molluscs characteristic of marine, brackish, freshwater and terrestrial environments were all represented. The total picture is indicative of sedimentation in an estuary:

a transitional area between land and sea where freshwater flowed into the sea; see also section 16.2.4.

### 16.2.3 Pottery temper

Some of the prehistoric pottery was tempered with shells fragments (chapter 6). Six sherds from phase 2a were microscopically examined to see whether any of the employed species could be identified. The shell remains were burned and visible in the clay matrix as a fragment or an impression. The small dimensions (at most a few mm<sup>2</sup>) and damage made it very difficult to determine the species,



but some information was nevertheless obtained. The shells were found to derive from cockle (edible or lagoon cockle), a mudsnail (*Hydrobia ulvae*/*Hydrobia ventrosa*) and an unknown bivalve.

In addition, some foraminifers, large diatoms, fine sand grains and fine plant matter (carbonised) were recorded. One of the sherds contained a carbonised rough club-rush seed.

Both the shells and the clay indicate that the pottery was made from material deposited in a saline or brackish environment. The latter option is the most likely. And both the shells and the clay were probably easily accessible in the site's surroundings. The clay will have been formed in the period preceding the occupation phase concerned (2a). The most likely source of the clay is Unit 19.

#### 16.2.4 Unit 0 (*Dunkirk I* deposits)

A large sample was taken from the base of the Dunkirk I deposit, which was laid down in the (estuarine) area of the tidal inlet/river Gantel and covered the dune. It dates from approximately 500 to 0 cal BC. Long grooves were observed at the base of this deposit (see section 2.2.3). A sample of one litre of sand mixed with layers of clay was taken from the fill of one of those grooves. This sample was studied for the presence of molluscs. In addition, another 5 litres of sand was studied to see whether it would yield comparable evidence. That was indeed found to be the case.

The molluscs encountered in the sample (table 16.2) proved to derive from different environments. Seven of the represented species favour a marine environment and three prefer brackish conditions. Some marine species, such as the mussel, Baltic tellin, peppery furrow shell, mudsnail and periwinkle, can tolerate reduced saline contents and are hence encountered in brackish environments, too. The represented range of species is indicative not of open saline water (the North Sea) but of a calm saline, tidal-flat kind of coastal area. The assemblage includes both poorly preserved and well-preserved shells.

The freshwater shells were all well preserved. They represent a fairly narrow faunal range, which nevertheless includes some remarkable species, in particular the swollen spire snail – a mollusc favouring freshwater tidal areas that tolerates slightly elevated saline contents. The dwarf pond snail and the ram's-horn tolerate desiccation of their biotope. They occur in shallow ponds surrounded by vegetation and at the edges of larger areas of water. The marsh snail can also survive in such environments. Finally, two terrestrial species were also encountered. Both species are encountered on damp banks.

As far as the reconstruction of the former environment is concerned there are two possibilities. The deposit may have been formed in a coastal area where freshwater flowed into the sea. In such an area, a mixture of saline and freshwater will have led to the formation of a brackish environment in

Species characteristic of a marine environment (poorly to well-preserved)

<i>Cerastoderma edule</i>	some valves and fragments
<i>Cerastoderma glaucum</i>	some valves and fragments
<i>Mytilus edulis</i>	several dozen doublets and valves and some fragments
<i>Spisula subtruncata</i>	some valves and fragments
<i>Macoma balthica</i>	some doublets, valves and fragments
<i>Scrobicularia plana</i>	1 valve and some fragments
<i>Hydrobia ventrosa</i>	several dozen
<i>Hydrobia cf acuta</i>	a few
<i>Hydrobia ulvae</i>	a few
<i>Littorina</i> sp.	1

Species characteristic of a freshwater environment (well-preserved)

<i>Mercuria confusa</i>	11
<i>Valvata piscinalis</i>	1
<i>Stagnicola palustris</i>	25
<i>Galba truncatula</i>	5
<i>Radix ovata</i>	2
<i>Anisus leucostoma</i>	1

Species characteristic of a terrestrial environment (well-preserved)

<i>Oxyloma</i> sp.	5
<i>Vallonia pulchella</i>	1

barnacle, plates	some
<i>Echinocardium cordatum</i>	some fragments, dozens of spines
ostracods, shells	dozens
foraminifers	
plant remains	small quantities

Table 16.2 Molluscs from the lower part of Unit 0. The sample, consisting of one litre of sand mixed with thin layers of clay, was sieved through a 0.25-mm mesh width.

which all of the encountered molluscs could have lived. The shells will then have been washed some distance away from their original positions by currents. The second possibility is an area with alternating periods of strong and less strong marine influence. In that case the freshwater species will have lived here in periods with strong discharge of freshwater and the marine species will have colonised the area in periods of pronounced marine influence. The brackish sediments will have been laid down in the transitional periods.

The majority of the other represented animal species favour a saline environment. The seeds contained in the same sample derive from plants that grow on the banks of areas of freshwater (some also in brackish water) and in swamps. The few aquatic plants encountered are characteristic of freshwater and brackish water. These results agree well with those of the mollusc analysis.

## 16.3 CONCLUSION

All the investigated deposits were laid down in a marine coastal area shielded from the open sea. In some periods the environment was saline due to a good connection with the open sea (Rijswijk-Zoetermeer sands), in others it was characterised by a reduced saline content (brackish), represented by the Early Holocene clay and Unit 2. During the deposition of the Dunkirk I sediment by the Gantel the area was an estuary. The environments in this coastal area ranged from tidal gullies in open connection with the sea to calm tidal flats and salt marshes.

Deposits from the period of the excavated settlement on the dune contained few mollusc remains. The small

quantities encountered point to a calm saline or brackish environment.

W.J. Kuijper  
Faculty of Archaeology  
Leiden University  
PO Box 9515  
2300 RA Leiden  
The Netherlands  
w.j.kuijper@arch.leidenuniv.nl

scientific	English	Dutch
<i>Mollusca</i>	molluscs/shells	mollusken, weekdieren, schelpen
<i>Anisus leucostoma</i>	button ram's-horn	geronde schijfhoren
<i>Barnea candida</i>	white piddock	witte boormossel
<i>Cerastoderma edule</i>	edible cockle	gewone kokkel
<i>Cerastoderma glaucum</i>	lagoon cockle	brakwaterkokkel
<i>Galba truncatula</i>	dwarf pond snail	leverbotlak
<i>Hydrobia cf. acuta</i>	mudsnail	vergeten brakwaterhoren
<i>Hydrobia ventrosa</i>	spire mudsnail	opgezwollen brakwaterhoren
<i>Hydrobia ulvae</i>	mudsnail	wadslakje
<i>Littorina sp.</i>	periwinkle	alikuik
<i>Macoma balthica</i>	Baltic tellin	nonnetje
<i>Mercuria confusa</i>	swollen spire snail	getijdenslak
<i>Mytilus edulis</i>	edible mussel	gewone mossel
<i>Oxyloma sp.</i>	amber snail	barnsteenslak
<i>Radix ovata</i>	common pond snail	ovale poelslak
<i>Scrobicularia plana</i>	peppery furrow shell	platte slijkgaper
<i>Spisula subtruncata</i>	cut trough shell	halfgeknotte strandschelp
<i>Stagnicola palustris</i>	marsh pond snail	moeraspoelslak
<i>Vallonia pulchella</i>	smooth grass snail	fraaie jachthorenslak
<i>Valvata piscinalis</i>	common valve snail	vijverpluimdrager

Table 16.3 Glossary of the scientific, English and Dutch names of the species mentioned in the text.