



Universiteit
Leiden
The Netherlands

Quenched large deviation principle for words in a letter sequence

Birkner, M.G.; Greven, A.; Hollander, W.T.F. den

Citation

Birkner, M. G., Greven, A., & Hollander, W. T. F. den. (2010). Quenched large deviation principle for words in a letter sequence. *Probability Theory And Related Fields*, 148(3-4), 403-456. Retrieved from <https://hdl.handle.net/1887/59868>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/59868>

Note: To cite this publication please use the final published version (if applicable).

Quenched large deviation principle for words in a letter sequence

Matthias Birkner ¹
 Andreas Greven ²
 Frank den Hollander ^{3 4}

13th May 2009

Abstract

When we cut an i.i.d. sequence of letters into words according to an independent renewal process, we obtain an i.i.d. sequence of words. In the *annealed* large deviation principle (LDP) for the empirical process of words, the rate function is the specific relative entropy of the observed law of words w.r.t. the reference law of words. In the present paper we consider the *quenched* LDP, i.e., we condition on a typical letter sequence. We focus on the case where the renewal process has an *algebraic* tail. The rate function turns out to be a sum of two terms, one being the annealed rate function, the other being proportional to the specific relative entropy of the observed law of letters w.r.t. the reference law of letters, with the former being obtained by concatenating the words and randomising the location of the origin. The proportionality constant equals the tail exponent of the renewal process. Earlier work by Birkner considered the case where the renewal process has an exponential tail, in which case the rate function turns out to be the first term on the set where the second term vanishes and to be infinite elsewhere. In a companion paper the annealed and the quenched LDP are applied to the collision local time of transient random walks, and the existence of an intermediate phase for a class of interacting stochastic systems is established.

Key words: Letters and words, renewal process, empirical process, annealed vs. quenched, large deviation principle, rate function, specific relative entropy.

MSC 2000: 60F10, 60G10.

Acknowledgement: This work was supported in part by DFG and NWO through the Dutch-German Bilateral Research Group “Mathematics of Random Spatial Models from Physics and Biology”. MB and AG are grateful for hospitality at EURANDOM. We also thank an anonymous referee for her/his careful reading and helpful comments.

¹Department Biologie II, Abteilung Evolutionsbiologie, University of Munich (LMU), Grosshaderner Str. 2, 82152 Planegg-Martinsried, Germany

²Mathematisches Institut, Universität Erlangen-Nürnberg, Bismarckstrasse 1 $\frac{1}{2}$, 91054 Erlangen, Germany

³Mathematical Institute, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands

⁴EURANDOM, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

1 Introduction and main results

1.1 Problem setting

Let E be a finite set of *letters*. Let $\tilde{E} = \cup_{n \in \mathbb{N}} E^n$ be the set of finite *words* drawn from E . Both E and \tilde{E} are Polish spaces under the discrete topology. Let $\mathcal{P}(E^{\mathbb{N}})$ and $\mathcal{P}(\tilde{E}^{\mathbb{N}})$ denote the set of probability measures on sequences drawn from E , respectively, \tilde{E} , equipped with the topology of weak convergence. Write θ and $\tilde{\theta}$ for the left-shift acting on $E^{\mathbb{N}}$, respectively, $\tilde{E}^{\mathbb{N}}$. Write $\mathcal{P}^{\text{inv}}(E^{\mathbb{N}})$, $\mathcal{P}^{\text{erg}}(E^{\mathbb{N}})$ and $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, $\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})$ for the set of probability measures that are invariant and ergodic under θ , respectively, $\tilde{\theta}$.

For $\nu \in \mathcal{P}(E)$, let $X = (X_i)_{i \in \mathbb{N}}$ be i.i.d. with law ν . Without loss of generality we will assume that $\text{supp}(\nu) = E$ (otherwise we replace E by $\text{supp}(\nu)$). For $\rho \in \mathcal{P}(\mathbb{N})$, let $\tau = (\tau_i)_{i \in \mathbb{N}}$ be i.i.d. with law ρ having infinite support and satisfying the *algebraic tail property*

$$\lim_{\substack{n \rightarrow \infty \\ \rho(n) > 0}} \frac{\log \rho(n)}{\log n} =: -\alpha, \quad \alpha \in (1, \infty). \quad (1.1)$$

(No regularity assumption will be necessary for $\text{supp}(\rho)$.) Assume that X and τ are independent and write \mathbb{P} to denote their joint law. Cut words out of X according to τ , i.e., put (see Figure 1)

$$T_0 := 0 \quad \text{and} \quad T_i := T_{i-1} + \tau_i, \quad i \in \mathbb{N}, \quad (1.2)$$

and let

$$Y^{(i)} := (X_{T_{i-1}+1}, X_{T_{i-1}+2}, \dots, X_{T_i}), \quad i \in \mathbb{N}. \quad (1.3)$$

Then, under the law \mathbb{P} , $Y = (Y^{(i)})_{i \in \mathbb{N}}$ is an i.i.d. sequence of words with marginal law $q_{\rho, \nu}$ on \tilde{E} given by

$$q_{\rho, \nu}((x_1, \dots, x_n)) := \mathbb{P}(Y^{(1)} = (x_1, \dots, x_n)) = \rho(n) \nu(x_1) \cdots \nu(x_n), \quad (1.4)$$

$n \in \mathbb{N}, x_1, \dots, x_n \in E.$




Figure 1: Cutting words from a letter sequence according to a renewal process.

For $N \in \mathbb{N}$, let $(Y^{(1)}, \dots, Y^{(N)})^{\text{per}}$ stand for the periodic extension of $(Y^{(1)}, \dots, Y^{(N)})$ to an element of $\tilde{E}^{\mathbb{N}}$, and define

$$R_N := \frac{1}{N} \sum_{i=0}^{N-1} \delta_{\tilde{\theta}^i(Y^{(1)}, \dots, Y^{(N)})^{\text{per}}} \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}), \quad (1.5)$$

the *empirical process of N -tuples of words*. By the ergodic theorem, we have

$$\text{w-} \lim_{N \rightarrow \infty} R_N = q_{\rho, \nu}^{\otimes \mathbb{N}} \quad \mathbb{P}\text{-a.s.}, \quad (1.6)$$

with $\text{w-} \lim$ denoting the weak limit. The following large deviation principle (LDP) is standard (see e.g. Dembo and Zeitouni [5], Corollaries 6.5.15 and 6.5.17). For $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ let

$$H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) := \lim_{N \rightarrow \infty} \frac{1}{N} h \left(Q|_{\mathcal{F}_N} | (q_{\rho, \nu}^{\otimes \mathbb{N}})|_{\mathcal{F}_N} \right) \in [0, \infty] \quad (1.7)$$

be the *specific relative entropy* of Q w.r.t. $q_{\rho,\nu}^{\otimes N}$, where $\mathcal{F}_N = \sigma(Y^{(1)}, \dots, Y^{(N)})$ is the sigma-algebra generated by the first N words, $Q|_{\mathcal{F}_N}$ is the restriction of Q to \mathcal{F}_N , and $h(\cdot | \cdot)$ denotes relative entropy. (For general properties of entropy, see Walters [13], Chapter 4.)

Theorem 1.1. [Annealed LDP] *The family of probability distributions $\mathbb{P}(R_N \in \cdot)$, $N \in \mathbb{N}$, satisfies the LDP on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ with rate N and with rate function $I^{\text{ann}}: \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) \rightarrow [0, \infty]$ given by*

$$I^{\text{ann}}(Q) = H(Q | q_{\rho,\nu}^{\otimes N}). \quad (1.8)$$

This rate function is lower semi-continuous, has compact level sets, has a unique zero at $Q = q_{\rho,\nu}^{\otimes N}$, and is affine.

The LDP for R_N arises from the LDP for N -tuples via a projective limit theorem. The ratio under the limit in (1.7) is the rate function for N -tuples according to Sanov's theorem (see e.g. den Hollander [8], Section II.5), and is non-decreasing in N .

1.2 Main theorems

Our aim in the present paper is to derive the LDP for $\mathbb{P}(R_N \in \cdot | X)$, $N \in \mathbb{N}$. To state our result, we need some more notation.

Let $\kappa: \tilde{E}^{\mathbb{N}} \rightarrow E^{\mathbb{N}}$ denote the *concatenation map* that glues a sequence of words into a sequence of letters. For $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ such that

$$m_Q := \mathbb{E}_Q[\tau_1] < \infty, \quad (1.9)$$

define $\Psi_Q \in \mathcal{P}^{\text{inv}}(E^{\mathbb{N}})$ as

$$\Psi_Q(\cdot) := \frac{1}{m_Q} \mathbb{E}_Q \left[\sum_{k=0}^{\tau_1-1} \delta_{\theta^k \kappa(Y)}(\cdot) \right]. \quad (1.10)$$

Think of Ψ_Q as the *shift-invariant version of the concatenation of Y under the law Q obtained after randomising the location of the origin.*

For $\text{tr} \in \mathbb{N}$, let $[\cdot]_{\text{tr}}: \tilde{E} \rightarrow [\tilde{E}]_{\text{tr}} := \cup_{n=1}^{\text{tr}} E^n$ denote the *word length truncation* map defined by

$$y = (x_1, \dots, x_n) \mapsto [y]_{\text{tr}} := (x_1, \dots, x_{n \wedge \text{tr}}), \quad n \in \mathbb{N}, x_1, \dots, x_n \in E. \quad (1.11)$$

Extend this to a map from $\tilde{E}^{\mathbb{N}}$ to $[\tilde{E}]_{\text{tr}}^{\mathbb{N}}$ via

$$[(y^{(1)}, y^{(2)}, \dots)]_{\text{tr}} := ([y^{(1)}]_{\text{tr}}, [y^{(2)}]_{\text{tr}}, \dots) \quad (1.12)$$

and to a map from $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv}}([\tilde{E}]_{\text{tr}}^{\mathbb{N}})$ via

$$[Q]_{\text{tr}}(A) := Q(\{z \in \tilde{E}^{\mathbb{N}}: [z]_{\text{tr}} \in A\}), \quad A \subset [\tilde{E}]_{\text{tr}}^{\mathbb{N}} \text{ measurable}. \quad (1.13)$$

Note that if $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, then $[Q]_{\text{tr}}$ is an element of the set

$$\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}}) = \{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}): m_Q < \infty\}. \quad (1.14)$$

Theorem 1.2. [Quenched LDP] *Assume (1.1). Then, for $\nu^{\otimes N}$ -a.s. all X , the family of (regular) conditional probability distributions $\mathbb{P}(R_N \in \cdot | X)$, $N \in \mathbb{N}$, satisfies the LDP on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ with rate N and with deterministic rate function $I^{\text{que}}: \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) \rightarrow [0, \infty]$ given by*

$$I^{\text{que}}(Q) := \begin{cases} I^{\text{fin}}(Q), & \text{if } Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}}), \\ \lim_{\text{tr} \rightarrow \infty} I^{\text{fin}}([Q]_{\text{tr}}), & \text{otherwise,} \end{cases} \quad (1.15)$$

where

$$I^{\text{fin}}(Q) := H(Q | q_{\rho,\nu}^{\otimes N}) + (\alpha - 1) m_Q H(\Psi_Q | \nu^{\otimes N}). \quad (1.16)$$

Theorem 1.3. *The rate function I^{que} is lower semi-continuous, has compact level sets, has a unique zero at $Q = q_{\rho, \nu}^{\otimes \mathbb{N}}$, and is affine. Moreover, it is equal to the lower semi-continuous extension of I^{fin} from $\mathcal{P}^{\text{inv, fin}}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$.*

Theorem 1.2 will be proved in Sections 3–5, Theorem 1.3 in Section 6.

A remarkable aspect of (1.16) in relation to (1.8) is that it *quantifies* the difference between the quenched and the annealed rate function. Note the appearance of the tail exponent α . We have not been able to find a simple formula for $I^{\text{que}}(Q)$ when $m_Q = \infty$. In Appendix A we will show that the annealed and the quenched rate function are continuous under truncation of word lengths, i.e.,

$$I^{\text{ann}}(Q) = \lim_{\text{tr} \rightarrow \infty} I^{\text{ann}}([Q]_{\text{tr}}), \quad I^{\text{que}}(Q) = \lim_{\text{tr} \rightarrow \infty} I^{\text{que}}([Q]_{\text{tr}}), \quad Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}). \quad (1.17)$$

Theorem 1.2 is an extension of Birkner [2], Theorem 1. In that paper, the quenched LDP is derived under the assumption that the law ρ satisfies the *exponential tail property*

$$\exists C < \infty, \lambda > 0: \quad \rho(n) \leq C e^{-\lambda n} \quad \forall n \in \mathbb{N} \quad (1.18)$$

(which includes the case where $\text{supp}(\rho)$ is finite). The rate function governing the LDP is given by

$$I^{\text{que}}(Q) := \begin{cases} H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}), & \text{if } Q \in \mathcal{R}_\nu, \\ \infty, & \text{if } Q \notin \mathcal{R}_\nu, \end{cases} \quad (1.19)$$

where

$$\mathcal{R}_\nu := \left\{ Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : \text{w-}\lim_{L \rightarrow \infty} \frac{1}{L} \sum_{k=0}^{L-1} \delta_{\theta^k \kappa(Y)} = \nu^{\otimes \mathbb{N}} \text{ } Q \text{ - a.s.} \right\}. \quad (1.20)$$

Think of \mathcal{R}_ν as the set of those Q 's for which *the concatenation of words has the same statistical properties as the letter sequence X* . This set is not closed in the weak topology: its closure is $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$.

We can include the cases where ρ satisfies (1.1) with $\alpha = 1$ or $\alpha = \infty$.

Theorem 1.4. (a) *If $\alpha = 1$, then the quenched LDP holds with $I^{\text{que}} = I^{\text{ann}}$ given by (1.8).*
(b) *If $\alpha = \infty$, then the quenched LDP holds with rate function*

$$I^{\text{que}}(Q) = \begin{cases} H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) & \text{if } \lim_{\text{tr} \rightarrow \infty} m_{[Q]_{\text{tr}}} H(\Psi_{[Q]_{\text{tr}}} \mid \nu^{\otimes \mathbb{N}}) = 0, \\ \infty & \text{otherwise.} \end{cases} \quad (1.21)$$

Theorem 1.4 will be proved in Section 7. Part (a) says that the quenched and the annealed rate function are identical when $\alpha = 1$. Part (b) says that (1.19) can be viewed as the limiting case of (1.16) as $\alpha \rightarrow \infty$. Indeed, it was shown in Birkner [2], Lemma 2, that on $\mathcal{P}^{\text{inv, fin}}(\tilde{E}^{\mathbb{N}})$:

$$\Psi_Q = \nu^{\otimes \mathbb{N}} \text{ if and only if } Q \in \mathcal{R}_\nu. \quad (1.22)$$

Hence, (1.21) and (1.19) agree on $\mathcal{P}^{\text{inv, fin}}(\tilde{E}^{\mathbb{N}})$, and the rate function (1.21) is the lower semicontinuous extension of (1.19) to $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$. By Birkner [2], Lemma 7, the expressions in (1.21) and (1.19) are identical if ρ has exponentially decaying tails. In this sense, Part (b) generalises the result in Birkner [2], Theorem 1, to arbitrary ρ with a tail that decays faster than algebraic.

Let $\pi_1: \tilde{E}^{\mathbb{N}} \rightarrow \tilde{E}$ be the projection onto the first word, and let $\mathcal{P}(\tilde{E})$ be the set of probability measures on \tilde{E} . An application of the contraction principle to Theorem 1.2 yields the following.

Corollary 1.5. *Under the assumptions of Theorem 1.2, for $\nu^{\otimes \mathbb{N}}$ -a.s. all X , the family of (regular) conditional probability distributions $\mathbb{P}(\pi_1 R_N \in \cdot | X)$, $N \in \mathbb{N}$, satisfies the LDP on $\mathcal{P}(\tilde{E})$ with rate N and with deterministic rate function $I_1^{\text{que}}: \mathcal{P}(\tilde{E}) \rightarrow [0, \infty]$ given by*

$$I_1^{\text{que}}(q) := \inf \{ I^{\text{que}}(Q) : Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}), \pi_1 Q = q \}. \quad (1.23)$$

This rate function is lower semi-continuous, has compact levels sets, has a unique zero at $q = q_{\rho, \nu}$, and is convex.

Corollary 1.5 shows that the rate function in Birkner [1], Theorem 6, must be replaced by (1.23). It does not appear possible to evaluate the infimum in (1.23) explicitly in general. For a $q \in \mathcal{P}(\tilde{E})$ with finite mean length and $\Psi_{q^{\otimes \mathbb{N}}} = \nu^{\otimes \mathbb{N}}$, we have $I_1^{\text{que}}(q) = h(q | q_{\rho, \nu})$.

By taking projective limits, it is possible to extend Theorems 1.2–1.3 to more general letter spaces. See, e.g., Deuschel and Stroock [6], Section 4.4, or Dembo and Zeitouni [5], Section 6.5, for background on (specific) relative entropy in general spaces. The following corollary will be proved in Section 8.

Corollary 1.6. *The quenched LDP also holds when E is a Polish space, with the same rate function as in (1.15–1.16).*

In the companion paper [3] the annealed and quenched LDP are applied to the collision local time of transient random walks, and the existence of an intermediate phase for a class of interacting stochastic systems is established.

1.3 Heuristic explanation of main theorems

To explain the background of Theorem 1.2, we begin by recalling a few properties of entropy. Let $H(Q)$ denote the *specific entropy* of $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ defined by

$$H(Q) := \lim_{N \rightarrow \infty} \frac{1}{N} h(Q|_{\mathcal{F}_N}) \in [0, \infty], \quad (1.24)$$

where $h(\cdot)$ denotes entropy. The sequence under the limit in (1.24) is non-increasing in N . Since $q_{\rho, \nu}^{\otimes \mathbb{N}}$ is a product measure, we have the identity (recall (1.2–1.4))

$$\begin{aligned} H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) &= -H(Q) - \mathbb{E}_Q[\log q_{\rho, \nu}(Y_1)] \\ &= -H(Q) - \mathbb{E}_Q[\log \rho(\tau_1)] - m_Q \mathbb{E}_{\Psi_Q}[\log \nu(X_1)]. \end{aligned} \quad (1.25)$$

Similarly,

$$H(\Psi_Q | \nu^{\otimes \mathbb{N}}) = -H(\Psi_Q) - \mathbb{E}_{\Psi_Q}[\log \nu(X_1)]. \quad (1.26)$$

Below, for a discrete random variable Z with a law Q on a state space \mathcal{Z} we will write $Q(Z)$ for the random variable $f(Z)$ with $f(z) = Q(Z = z)$, $z \in \mathcal{Z}$. Abbreviate

$$K^{(N)} := \kappa(Y^{(1)}, \dots, Y^{(N)}) \quad \text{and} \quad K^{(\infty)} := \kappa(Y). \quad (1.27)$$

In analogy with (1.14), define

$$\mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}}) := \left\{ Q \in \mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}}) : m_Q < \infty \right\}. \quad (1.28)$$

Lemma 1.7. [Birkner [2], Lemmas 3 and 4]

Suppose that $Q \in \mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$ and $H(Q) < \infty$. Then, Q -a.s.,

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log Q(K^{(N)}) &= -m_Q H(\Psi_Q), \\ \lim_{N \rightarrow \infty} \frac{1}{N} \log Q(\tau_1, \dots, \tau_N \mid K^{(N)}) &=: -H_{\tau|K}(Q), \\ \lim_{N \rightarrow \infty} \frac{1}{N} \log Q(Y^{(1)}, \dots, Y^{(N)}) &= -H(Q), \end{aligned} \tag{1.29}$$

with

$$m_Q H(\Psi_Q) + H_{\tau|K}(Q) = H(Q). \tag{1.30}$$

Equation (1.30), which follows from (1.29) and the identity

$$Q(K^{(N)})Q(\tau_1, \dots, \tau_N \mid K^{(N)}) = Q(Y^{(1)}, \dots, Y^{(N)}), \tag{1.31}$$

identifies $H_{\tau|K}(Q)$. Think of $H_{\tau|K}(Q)$ as the *conditional specific entropy of word lengths under the law Q given the concatenation*. Combining (1.25–1.26) and (1.30), we have

$$H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) = m_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}}) - H_{\tau|K}(Q) - \mathbb{E}_Q[\log \rho(\tau_1)]. \tag{1.32}$$

The term $-H_{\tau|K}(Q) - \mathbb{E}_Q[\log \rho(\tau_1)]$ in (1.32) can be interpreted as the *conditional specific relative entropy of word lengths under the law Q w.r.t. $\rho^{\otimes \mathbb{N}}$ given the concatenation*.

Note that $m_Q < \infty$ and $H(Q) < \infty$ imply that $H(\Psi_Q) < \infty$, as can be seen from (1.30). Also note that $-\mathbb{E}_{\Psi_Q}[\log \nu(X_1)] < \infty$ because E is finite, and $-\mathbb{E}_Q[\log \rho(\tau_1)] < \infty$ because of (1.1) and $m_Q < \infty$, implying that (1.25–1.26) are proper.

We are now ready to give a heuristic explanation of Theorem 1.2. Let

$$R_{j_1, \dots, j_N}^N(X), \quad 0 < j_1 < \dots < j_N < \infty, \tag{1.33}$$

denote the empirical process of N -tuples of words when X is cut at the points j_1, \dots, j_N (i.e., when $T_i = j_i$ for $i = 1, \dots, N$; see (3.16–3.17) for a precise definition). Fix $Q \in \mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$. The probability $\mathbb{P}(R_N \approx Q \mid X)$ is a sum over all N -tuples j_1, \dots, j_N such that $R_{j_1, \dots, j_N}^N(X) \approx Q$, weighted by $\prod_{i=1}^N \rho(j_i - j_{i-1})$ (with $j_0 = 0$). The fact that $R_{j_1, \dots, j_N}^N(X) \approx Q$ has three consequences:

- (1) The j_1, \dots, j_N must cut $\approx N$ substrings out of X of total length $\approx Nm_Q$ that look like the concatenation of words that are Q -typical, i.e., that look as if generated by Ψ_Q (possibly with gaps in between). This means that most of the cut-points must hit atypical pieces of X . We expect to have to shift X by $\approx \exp[Nm_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}})]$ in order to find the first contiguous substring of length Nm_Q whose empirical shifts lie in a small neighbourhood of Ψ_Q . By (1.1), the probability for the single increment $j_1 - j_0$ to have the size of this shift is $\approx \exp[-N\alpha m_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}})]$.
- (2) The combinatorial factor $\exp[NH_{\tau|K}(Q)]$ counts how many “local perturbations” of j_1, \dots, j_N preserve the property that $R_{j_1, \dots, j_N}^N(X) \approx Q$.
- (3) The statistics of the increments $j_1 - j_0, \dots, j_N - j_{N-1}$ must be close to the distribution of word lengths under Q . Hence, the weight factor $\prod_{i=1}^N \rho(j_i - j_{i-1})$ must be $\approx \exp[N\mathbb{E}_Q[\log \rho(\tau_1)]]$ (at least, for Q -typical pieces).

The contributions from (1)–(3), together with the identity in (1.32), explain the formula in (1.16) on $\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$. Considerable work is needed to extend (1)–(3) from $\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$. This is explained in Section 3.5.

In (1), instead of having a single large increment preceding a single contiguous substring of length Nm_Q , it is possible to have several large increments preceding several contiguous substrings, which together have length Nm_Q . The latter gives rise to the same contribution, and so there is some entropy associated with the choice of the large increments. Lemma 2.1 in Section 2.1 is needed to control this entropy, and shows that it is negligible.

1.4 Outline

Section 2 collects some preparatory facts that are needed for the proofs of the main theorems, including a lemma that controls the entropy associated with the locations of the large increments in the renewal process. In Section 3 and 4 we prove the large deviation upper, respectively, lower bound. The proof of the former is long (taking up about half of the paper) and requires a somewhat lengthy construction with combinatorial, functional analytic and ergodic theoretic ingredients. In particular, extending the lower bound from ergodic to non-ergodic probability measures is technically involved. The proofs of Theorems 1.2–1.4 are in Sections 5–7, that of Corollary 1.6 is in Section 8. Appendix A contains a proof that the annealed and the quenched rate function are continuous under the truncation of the word length approximation.

2 Preparatory facts

Section 2.1 proves a core lemma that is needed to control the entropy of large increments in the renewal process. Section 2.2 shows that the tail property of ρ is preserved under convolutions.

2.1 A core lemma

As announced at the end of Section 1.3, we need to account for the entropy that is associated with the locations of the large increments in the renewal process. This requires the following combinatorial lemma.

Lemma 2.1. *Let $\omega = (\omega_l)_{l \in \mathbb{N}}$ be i.i.d. with $\mathbb{P}(\omega_1 = 1) = 1 - \mathbb{P}(\omega_1 = 0) = p \in (0, 1)$, and let $\alpha \in (1, \infty)$. For $N \in \mathbb{N}$, let*

$$S_N(\omega) := \sum_{\substack{0 < j_1 < \dots < j_N < \infty \\ \omega_{j_1} = \dots = \omega_{j_N} = 1}} \prod_{i=1}^N (j_i - j_{i-1})^{-\alpha} \quad (j_0 = 0) \quad (2.1)$$

and put

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log S_N(\omega) =: -\phi(\alpha, p) \quad \omega - a.s. \quad (2.2)$$

(the limit being ω -a.s. constant by tail triviality). Then

$$\lim_{p \downarrow 0} \frac{\phi(\alpha, p)}{\alpha \log(1/p)} = 1. \quad (2.3)$$

Proof. Let $\tau_N := \min\{l \in \mathbb{N} : \omega_l = \omega_{l+1} = \dots = \omega_{l+N-1}\}$. In (2.1), choosing $j_1 = \tau_N$ and $j_i = j_{i-1} + 1$ for $i = 2, \dots, N$, we see that $S_N(\omega) \geq \tau_N^{-\alpha}$. Since

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log \tau_N \rightarrow \log(1/p) \quad \omega - a.s., \quad (2.4)$$

we have

$$\phi(\alpha, p) \leq \alpha \log(1/p) \quad \forall p \in (0, 1). \quad (2.5)$$

To show that this bound is sharp in the limit as $p \downarrow 0$, we estimate fractional moments of $S_N(\omega)$.

For any $\beta \in (1/\alpha, 1]$, using that $(u+v)^\beta \leq u^\beta + v^\beta$, $u, v \geq 0$, we get

$$\begin{aligned} \mathbb{E} \left[S_N(\omega)^\beta \right] &\leq \sum_{0 < j_1 < \dots < j_N < \infty} \mathbb{E} \left[\mathbb{1}_{\{\omega_{j_1} = \dots = \omega_{j_N} = 1\}} \prod_{i=1}^N (j_i - j_{i-1})^{-\alpha\beta} \right] \\ &= \sum_{0 < j_1 < \dots < j_N < \infty} p^N \prod_{i=1}^N (j_i - j_{i-1})^{-\alpha\beta} \\ &= [p \zeta(\alpha\beta)]^N, \end{aligned} \quad (2.6)$$

where $\zeta(s) = \sum_{n \in \mathbb{N}} n^{-s}$, $s > 1$, is Riemann's ζ -function. Hence, for any $\varepsilon > 0$, Markov's inequality yields

$$\begin{aligned} \mathbb{P} \left(\frac{1}{N} \log S_N(\omega) \geq \frac{1}{\beta} [\log p + \log \zeta(\alpha\beta) + \varepsilon] \right) \\ = \mathbb{P} \left(S_N(\omega)^\beta \geq e^{\varepsilon N} [p \zeta(\alpha\beta)]^N \right) \leq e^{-\varepsilon N} [p \zeta(\alpha\beta)]^{-N} \mathbb{E} \left[S_N(\omega)^\beta \right] \leq e^{-\varepsilon N}. \end{aligned} \quad (2.7)$$

Thus, by the first Borel-Cantelli Lemma,

$$-\phi(\alpha, p) = \limsup_{N \rightarrow \infty} \frac{1}{N} \log S_N(\omega) \leq \frac{1}{\beta} [\log p + \log \zeta(\alpha\beta)] \quad \text{a.s.} \quad (2.8)$$

Now let $p \downarrow 0$, followed by $\beta \downarrow 1/\alpha$ to obtain the claim. \square

Remark 2.2. Note that $\mathbb{E}[S_N(\omega)] = (p\zeta(\alpha))^N$, while typically $S_N(\omega) \approx p^{\alpha N}$. In the above computation, this is verified by bounding suitable non-integer moments of $S_N(\omega)/p^{\alpha N}$. Estimating non-integer moments in situations when the mean is inconclusive is a useful technique in a variety of different probabilistic contexts. See, e.g., Holley and Liggett [9] and Toninelli [12]. The proof of Lemma 2.1 above is similar to that of Toninelli [12], Theorem 2.1.

2.2 Convolution preserves polynomial tail

The following lemma will be needed in Sections 3.3 and 3.5. For $m \in \mathbb{N}$, let ρ^{*m} denote the m -fold convolution of ρ .

Lemma 2.3. *Suppose that ρ satisfies $\rho(n) \leq C_\rho n^{-\alpha}$, $n \in \mathbb{N}$, for some $C_\rho < \infty$. Then*

$$\rho^{*m}(n) \leq (C_\rho \vee 1) m^{\alpha+1} n^{-\alpha} \quad \forall m, n \in \mathbb{N}. \quad (2.9)$$

Proof. If $n \leq m$, then the right-hand side of (2.9) is ≥ 1 . So, let us assume that $n > m$. Then

$$\begin{aligned} \rho^{*m}(n) &= \sum_{\substack{x_1, \dots, x_m \geq 1 \\ x_1 + \dots + x_m = n}} \prod_{i=1}^m \rho(x_i) \leq \sum_{j=1}^m \sum_{\substack{x_1, \dots, x_m \geq 1 \\ x_1 + \dots + x_m = n \\ x_j = x_1 \vee \dots \vee x_m}} \rho(x_j) \prod_{i \neq j} \rho(x_i) \\ &\leq m C_\rho [n/m]^{-\alpha} \sum_{x_1, \dots, x_{m-1} \geq 1} \prod_{i=1}^{m-1} \rho(x_i) \\ &= m C_\rho [n/m]^{-\alpha} \leq C_\rho m^{\alpha+1} n^{-\alpha}. \end{aligned} \quad (2.10)$$

\square

3 Upper bound

The following upper bound will be used in Section 5 to derive the upper bound in the definition of the LDP.

Proposition 3.1. *For any $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ and any $\varepsilon > 0$, there is an open neighbourhood $\mathcal{O}(Q) \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ of Q such that*

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{O}(Q) \mid X) \leq -I^{\text{fin}}(Q) + \varepsilon \quad X - a.s. \quad (3.1)$$

We remark that since $|E| < \infty$ we automatically have $I^{\text{fin}}(Q) \in [0, \infty)$ for all $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$, so the right-hand side of (3.1) is finite.

Proof. It suffices to consider the case $\Psi_Q \neq \nu^{\otimes \mathbb{N}}$. The case $\Psi_Q = \nu^{\otimes \mathbb{N}}$, for which $I^{\text{fin}}(Q) = H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}})$ as is seen from (1.16), is contained in the upper bound in Birkner [2], Lemma 8. Alternatively, by lower semicontinuity of $Q' \mapsto H(Q' \mid q_{\rho, \nu}^{\otimes \mathbb{N}})$, there is a neighbourhood $\mathcal{O}(Q)$ such that

$$\inf_{Q' \in \overline{\mathcal{O}(Q)}} H(Q' \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) \geq H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) - \varepsilon = I^{\text{fin}}(Q) - \varepsilon, \quad (3.2)$$

where $\overline{\mathcal{O}(Q)}$ denotes the closure of $\mathcal{O}(Q)$ (in the weak topology), and we can use the annealed bound.

In Sections 3.1–3.5 we first prove Proposition 3.1 under the assumption that there exist $\alpha \in (1, \infty)$, $C_\rho < \infty$ such that

$$\rho(n) \leq C_\rho n^{-\alpha}, \quad n \in \mathbb{N}, \quad (3.3)$$

which is needed in Lemma 2.3. In Section 3.6 we show that this can be replaced by (1.1). In Sections 3.1–3.4, we first consider $Q \in \mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$ (recall (1.28)). Here, we turn the heuristics from Section 1.3 into a rigorous proof. In Section 3.5 we remove the ergodicity restriction. The proof is long and technical (taking up more than half of the paper).

3.1 Step 1: Consequences of ergodicity

We will use the ergodic theorem to construct specific neighborhoods of $Q \in \mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$ that are well adapted to formalize the strategy of proof outlined in our heuristic explanation of the main theorem in Section 1.3.

Fix $\varepsilon_1, \delta_1 > 0$. By the ergodicity of Q and Lemma 1.7, the event (recall (1.9) and (1.27))

$$\begin{aligned} & \left\{ \frac{1}{M} |K^{(M)}| \in m_Q + [-\varepsilon_1, \varepsilon_1] \right\} \\ & \cap \left\{ -\frac{1}{M} \log Q(K^{(M)}) \in m_Q H(\Psi_Q) + [-\varepsilon_1, \varepsilon_1] \right\} \\ & \cap \left\{ -\frac{1}{M} \log Q(Y^{(1)}, \dots, Y^{(M)}) \in H(Q) + [-\varepsilon_1, \varepsilon_1] \right\} \\ & \cap \left\{ \frac{1}{M} \sum_{k=1}^{|K^{(M)}|} \log \nu((K^{(M)})_k) \in m_Q \mathbb{E}_{\Psi_Q} [\log \nu(X_1)] + [-\varepsilon_1, \varepsilon_1] \right\} \\ & \cap \left\{ \frac{1}{M} \sum_{i=1}^M \log \rho(\tau_i) \in \mathbb{E}_Q [\log \rho(\tau_1)] + [-\varepsilon_1, \varepsilon_1] \right\} \end{aligned} \quad (3.4)$$

has Q -probability at least $1 - \delta_1/4$ for M large enough (depending on Q), where $|K^{(M)}|$ is the length of the string of letters $K^{(M)}$. Hence, there is a finite number A of sentences of length M , denoted by

$$(z_a)_{a=1,\dots,A} \text{ with } z_a := (y^{(a,1)}, \dots, y^{(a,M)}) \in \tilde{E}^M, \quad (3.5)$$

such that for $a = 1, \dots, A$,

$$\begin{aligned} |\kappa(z_a)| &\in [M(m_Q - \varepsilon_1), M(m_Q + \varepsilon_1)], \\ Q(K^{(M)} = \kappa(z_a)) &\in [\exp[-M(m_Q H(\Psi_Q) + \varepsilon_1)], \exp[-M(m_Q H(\Psi_Q) - \varepsilon_1)]], \\ Q((Y^{(1)}, \dots, Y^{(M)}) = z_a) &\in [\exp[-M(H(Q) + \varepsilon_1)], \exp[-M(H(Q) - \varepsilon_1)]], \\ \sum_{k=1}^{|\kappa(z_a)|} \log \nu((\kappa(z_a))_k) &\in [M(m_Q \mathbb{E}_{\Psi_Q}[\log \nu(X_1)] - \varepsilon_1), M(m_Q \mathbb{E}_{\Psi_Q}[\log \nu(X_1)] + \varepsilon_1)], \\ \sum_{i=1}^M \log \rho(|y^{(a,i)}|) &\in [M(\mathbb{E}_Q[\log \rho(\tau_1)] - \varepsilon_1), M(\mathbb{E}_Q[\log \rho(\tau_1)] + \varepsilon_1)], \end{aligned} \quad (3.6)$$

and

$$\sum_{a=1}^A Q((Y^{(1)}, \dots, Y^{(M)}) = z_a) \geq 1 - \frac{\delta_1}{2}. \quad (3.7)$$

Note that (3.7) and the third line of (3.6) imply that

$$A \in \left[\left(1 - \frac{\delta_1}{2}\right) \exp [M(H(Q) - \varepsilon_1)], \exp [M(H(Q) + \varepsilon_1)] \right]. \quad (3.8)$$

Abbreviate

$$\mathcal{A} := \{z_a, a = 1, \dots, A\}. \quad (3.9)$$

Let

$$\mathcal{B} := \{\zeta^{(b)}, b = 1, \dots, B\} = \{\kappa(z_a), a = 1, \dots, A\} \quad (3.10)$$

be the set of strings of letters arising from concatenations of the individual z_a 's, and let

$$I_b := \{1 \leq a \leq A: \kappa(z_a) = \zeta^{(b)}\}, \quad b = 1, \dots, B, \quad (3.11)$$

so that $|I_b|$ is the number of sentences in \mathcal{A} giving a particular string in \mathcal{B} . By the second line of (3.6), we can bound B as

$$B \leq \exp [M(m_Q H(\Psi_Q) + \varepsilon_1)], \quad (3.12)$$

because $\sum_{b=1}^B Q(K^{(M)} = \zeta^{(b)}) \leq 1$ and each summand is at least $\exp[-M(m_Q H(\Psi_Q) + \varepsilon_1)]$. Furthermore, we have

$$|I_b| \leq \exp [M(H_{\tau|K}(Q) + 2\varepsilon_1)], \quad b = 1, \dots, B, \quad (3.13)$$

since

$$\begin{aligned} \exp [-M(m_Q H(\Psi_Q) - \varepsilon_1)] &\geq Q(\kappa(Y^{(1)}, \dots, Y^{(M)}) = \zeta^{(b)}) \\ &\geq \sum_{a \in I_b} Q((Y^{(1)}, \dots, Y^{(M)}) = z_a) \geq |I_b| \exp [-M(H(Q) + \varepsilon_1)], \end{aligned} \quad (3.14)$$

and $H(Q) - m_Q H(\Psi_Q) = H_{\tau|K}(Q)$ by (1.30).

3.2 Step 2: Good sentences in open neighbourhoods

Define the following open neighbourhood of Q (recall (3.9))

$$\mathcal{O} := \left\{ Q' \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : Q'_{|\mathcal{F}_M}(\mathcal{A}) > 1 - \delta_1 \right\}. \quad (3.15)$$

Here, $Q(z)$ is shorthand for $Q((Y^{(1)}, \dots, Y^{(M)}) = z)$. For $x \in E^{\mathbb{N}}$ and for a vector of cut-points $(j_1, \dots, j_N) \in \mathbb{N}^N$ with $0 < j_1 < \dots < j_N < \infty$ and $N > M$, let

$$\xi_N := (\xi^{(i)})_{i=1, \dots, N} = (x|_{(0, j_1]}, x|_{(j_1, j_2]}, \dots, x|_{(j_{N-1}, j_N]}) \in \tilde{E}^N \quad (3.16)$$

(with $(0, j_1]$ shorthand notation for $(0, j_1] \cap \mathbb{N}$, etc.) be the sequence of words obtained by cutting x at the positions j_i , and let

$$R_{j_1, \dots, j_N}^N(x) := \frac{1}{N} \sum_{i=0}^{N-1} \delta_{\tilde{\theta}^i(\xi_N)^{\text{per}}} \quad (3.17)$$

be the corresponding empirical process. By (3.15),

$$\begin{aligned} R_{j_1, \dots, j_N}^N(x) \in \mathcal{O} &\implies \\ \#\left\{ 1 \leq i \leq N - M : (x|_{(j_{i-1}, j_i]}, \dots, x|_{(j_{i+M-1}, j_{i+M}]}) \in \mathcal{A} \right\} &\geq N(1 - \delta_1) - M. \end{aligned} \quad (3.18)$$

Note that (3.18) implies that the sentence ξ_N contains at least

$$C := \lfloor (1 - \delta_1)N/M \rfloor - 1 \quad (3.19)$$

disjoint subsentences from the set \mathcal{A} , i.e., there are $1 \leq i_1, \dots, i_C \leq N - M$ with $i_c - i_{c-1} \geq M$ for $c = 1, \dots, C$ such that

$$(\xi^{(i_c)}, \xi^{(i_c+1)}, \dots, \xi^{(i_c+M-1)}) \in \mathcal{A} \quad (3.20)$$

(we implicitly assume that N is large enough so that $C > 1$). Indeed, we can e.g. construct the i_c 's iteratively as

$$\begin{aligned} i_0 &= -M, \\ i_c &= \min \left\{ k \geq i_{c-1} + M : \text{a sentence from } \mathcal{A} \text{ starts at position } k \text{ in } \xi_N \right\}, \\ c &= 1, \dots, C, \end{aligned} \quad (3.21)$$

and we can continue the iteration as long as $cM + \delta_1 N \leq N$. But (3.20) in turn implies that the j_{i_c} 's cut out of x at least C disjoint subwords from \mathcal{B} , i.e.,

$$x|_{(j_{i_c}, j_{i_c+M})} \in \mathcal{B}, \quad c = 1, \dots, C. \quad (3.22)$$

3.3 Step 3: Estimate of the large deviation probability

Using Steps 1 and 2, we estimate (recall (3.15))

$$\mathbb{P}(R_N \in \mathcal{O} | X) = \sum_{0 < j_1 < \dots < j_N < \infty} \mathbb{1}_{\mathcal{O}}(R_{j_1, \dots, j_N}^N(X)) \prod_{i=1}^N \rho(j_i - j_{i-1}) \quad (3.23)$$

from above as follows. Fix a vector of cut-points (j_1, \dots, j_N) giving rise to a non-zero contribution in the right-hand side of (3.23). We think of this vector as describing a particular way of cutting X




Figure 2: Looking for good subsentences and filling subsentences (see below (3.25)).

into a sentence of N words. By (3.22), at least C (recall 3.19) of the j_c 's must be cut-points where a word from \mathcal{B} is written on X , and these C subwords must be disjoint. As words in \mathcal{B} arise from concatenations of sentences from \mathcal{A} , this means we can find

$$\ell_1 < \dots < \ell_C, \quad \{\ell_1, \dots, \ell_C\} \subset \{0, j_1, \dots, j_N\} \quad \text{and} \quad \zeta_1, \dots, \zeta_C \in \mathcal{A} \quad (3.24)$$

such that

$$X|_{(\ell_c, \ell_c + |\kappa(\zeta_c)|]} = \kappa(\zeta_c) =: \eta^{(c)} \in \mathcal{B} \quad \text{and} \quad \ell_c \geq \ell_{c-1} + |\kappa(\zeta_{c-1})|, \quad c = 1, \dots, C-1. \quad (3.25)$$

We call ζ_1, \dots, ζ_C the *good* subsentences.

Note that once we fix the ℓ_c 's and the ζ_c 's, this determines $C+1$ *filling* subsentences (some of which may be empty) consisting of the words between the good subsentences. See Figure 2 for an illustration. In particular, this determines numbers $m_1, \dots, m_{C+1} \in \mathbb{N}$ such that $m_1 + \dots + m_{C+1} = N - CM$, where m_c is the number of words we cut between the $(c-1)$ -st and the c -th good subsentence (and m_{C+1} is the number of words after the C -th good subsentence).

Next, let us fix good $\ell_1 < \dots < \ell_C$ and $\eta^{(1)}, \dots, \eta^{(C)} \in \mathcal{B}$, satisfying

$$X|_{(\ell_c, \ell_c + |\eta^{(c)}|]} = \eta^{(c)}, \quad \ell_c \geq \ell_{c-1} + |\eta^{(c-1)}|, \quad c = 1, \dots, C. \quad (3.26)$$

To estimate how many different choices of (j_1, \dots, j_N) may lead to this particular $((\ell_c), (\eta^{(c)}))$, we proceed as follows. There are at most

$$(2M\varepsilon_1)^C \exp [M(H_{\tau|K}(Q) + 2\varepsilon_1)]^C \leq \exp [N(H_{\tau|K}(Q) + \delta_2)] \quad (3.27)$$

possible choices for the word lengths inside these good subsentences. Indeed, by the first line of (3.6), at most $2M\varepsilon_1$ different elements of \mathcal{B} can start at any given position ℓ_c and, by (3.13), each of them can be cut in at most $\exp [M(H_{\tau|K}(Q) + 2\varepsilon_1)]$ different ways to obtain an element of \mathcal{A} . In (3.27), $\delta_2 = \delta_2(\varepsilon_1, \delta_1, M)$ can be made arbitrarily small by choosing M large and ε_1, δ_1 small. Furthermore, there are at most

$$\binom{N - C(M-1)}{C} \leq \exp[\delta_3 N] \quad (3.28)$$

possible choices of the m_c 's, where $\delta_3 = \delta_3(\delta_1, M)$ can be made arbitrarily small by choosing M large and δ_1 small.

Next, we estimate the value of $\prod_{i=1}^N \rho(j_i - j_{i-1})$ for any (j_1, \dots, j_N) leading to the given $((\ell_c), (\eta^{(c)}))$. In view of the fifth line of (3.6), we have

$$\begin{aligned} & \prod_{i=1}^N \rho(j_i - j_{i-1}) \mathbb{1}_{\{\text{the } i\text{-th word falls inside the } C \text{ good subtences}\}} \\ & \leq \exp [CM(\mathbb{E}_Q[\log \rho(\tau_1)] + \varepsilon_1)] \\ & \leq \exp [N(\mathbb{E}_Q[\log \rho(\tau_1)] + \delta_4)], \end{aligned} \tag{3.29}$$

where $\delta_4 = \delta_4(\varepsilon_1, \delta_1, M)$ can be made arbitrarily small by choosing M large and ε_1, δ_1 small. The filling subtences have to exactly fill up the gaps between the good subtences and so, for a given choice of $(\ell_c), (\eta^{(c)})$ and (m_c) , the contribution to $\prod_{i=1}^N \rho(j_i - j_{i-1})$ from the filling subtences is $\prod_{c=1}^C \rho^{*m_c}(\ell_c - \ell_{c-1} - |\eta^{(c-1)}|)$ (the term for $c = 1$ is to be interpreted as $\rho^{*m_1}(\ell_1)$, and ρ^{*0} as δ_0). By Lemma 2.3, using (3.3),

$$\begin{aligned} & \prod_{c=1}^C \rho^{*m_c}(\ell_c - \ell_{c-1} - |\eta^{(c-1)}|) \\ & \leq (C_\rho \vee 1)^C \left(\prod_{c=1}^C m_c^{\alpha+1} \right) \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\eta^{(c-1)}|) \vee 1)^{-\alpha} \\ & \leq (C_\rho \vee 1)^C \left(\frac{N - CM}{C} \right)^{(\alpha+1)C} \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\eta^{(c-1)}|) \vee 1)^{-\alpha} \\ & \leq \exp[N\delta_5] \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\eta^{(c-1)}|) \vee 1)^{-\alpha}, \end{aligned} \tag{3.30}$$

where $\delta_5 = \delta(\delta_1, M)$ can be made arbitrarily small by choosing M large and δ_1 small. For the second inequality, we have used the fact that the product $\prod_{c=1}^C m_c^{\alpha+1}$ is maximal when all factors are equal.

Combining (3.23–3.30), we obtain

$$\begin{aligned} \mathbb{P}(R_N \in \mathcal{O} \mid X) & \leq \exp \left[N \left(H_{\tau|K}(Q) + \mathbb{E}_Q[\log \rho(\tau_1)] + \delta_2 + \delta_3 + \delta_4 + \delta_5 \right) \right] \\ & \quad \times \sum_{(\ell_c), (\eta^{(c)}) \text{ good}} \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\eta^{(c-1)}|) \vee 1)^{-\alpha}. \end{aligned} \tag{3.31}$$

Combining (3.31) with Lemma 3.2 below, and recalling the identity in (1.32), we obtain the result in Proposition 3.1 for ρ satisfying (3.3), with \mathcal{O} defined in (3.15) and $\varepsilon = \delta_2 + \delta_3 + \delta_4 + \delta_5 + \delta_6$. Note that ε can be made arbitrarily small by choosing ε_1, δ_1 small and M large.

3.4 Step 4: Cost of finding good sentences

Lemma 3.2. *For $\varepsilon_1, \delta_1 > 0$ and $M \in \mathbb{N}$,*

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \frac{1}{N} \log \left[\sum_{(\ell_c), (\eta^{(c)}) \text{ good}} \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\eta^{(c-1)}|) \vee 1)^{-\alpha} \right] \\ & \leq -\alpha m_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}}) + \delta_6 \quad \text{a.s.}, \end{aligned} \tag{3.32}$$

where $\delta_6 = \delta(\varepsilon_1, \delta_1, M)$ can be made arbitrarily small by choosing M large and ε_1, δ_1 small.

Proof. Note that, by the fourth line of (3.6), for any $\eta \in \mathcal{B}$ (recall (3.10)) and $k \in \mathbb{N}$,

$$\mathbb{P}(\eta \text{ starts at position } k \text{ in } X) \leq \exp [M(m_Q \mathbb{E}_{\Psi_Q} [\log \nu(X_1)] + \varepsilon_1)]. \quad (3.33)$$

Combining this with (3.12), we get

$$\begin{aligned} & \mathbb{P}(\text{some element of } \mathcal{B} \text{ starts at position } k \text{ in } X) \\ & \leq \exp [M(m_Q \mathbb{E}_{\Psi_Q} [\log \nu(X_1)] + \varepsilon_1)] \times \exp [M(m_Q H(\Psi_Q) + \varepsilon_1)] \\ & = \exp [-M(m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - 2\varepsilon_1)], \end{aligned} \quad (3.34)$$

where we use (1.26).

Next, we *coarse-grain* the sequence X into blocks of length

$$L := \lfloor M(m_Q - \varepsilon_1) \rfloor, \quad (3.35)$$

and compare the coarse-grained sequence with a *low-density Bernoulli sequence*. To this end, define a $\{0, 1\}$ -valued sequence $(A_l)_{l \in \mathbb{N}}$ inductively as follows. Put $A_0 := 0$, and, for $l \in \mathbb{N}$ given that A_0, A_1, \dots, A_{l-1} have been assigned values, define A_l by distinguishing the following two cases:

(1) If $A_{l-1} = 0$, then

$$A_l := \begin{cases} 1, & \text{if in } X \text{ there is a word } \eta \in \mathcal{B} \text{ starting in } ((l-1)L, lL], \\ 0, & \text{otherwise.} \end{cases} \quad (3.36)$$

(2) If $A_{l-1} = 1$, then

$$A_l := \begin{cases} 1, & \text{if in } X \text{ there are words } \eta, \eta' \in \mathcal{B} \text{ starting in } ((l-2)L, (l-1)L], \\ & \text{respectively, } ((l-1)L, lL] \text{ and occurring disjointly,} \\ 0, & \text{otherwise.} \end{cases} \quad (3.37)$$

Put

$$p := L \exp [-M(m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - 2\varepsilon_1)]. \quad (3.38)$$

Then we claim

$$\mathbb{P}(A_1 = a_1, \dots, A_n = a_n) \leq p^{a_1 + \dots + a_n}, \quad n \in \mathbb{N}, a_1, \dots, a_n \in \{0, 1\}. \quad (3.39)$$

In order to verify (3.39), fix $a_1, \dots, a_n \in \{0, 1\}$ with $a_1 + \dots + a_n = m$. By construction, for the event in the left-hand side of (3.39) to occur there must be m non-overlapping elements of \mathcal{B} at certain positions in X . By (3.34), the occurrence of any m fixed starting positions has probability at most

$$\exp [-mM(m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - 2\varepsilon_1)], \quad (3.40)$$

while the choice of the a_l 's dictates that there are at most L^m possibilities for the starting points of the m words.

By (3.39), we can couple the sequence $(A_l)_{l \in \mathbb{N}}$ with an i.i.d. Bernoulli(p)-sequence $(\omega_l)_{l \in \mathbb{N}}$ such that

$$A_l \leq \omega_l \quad \forall l \in \mathbb{N} \quad \text{a.s.} \quad (3.41)$$

(Note that (3.39) guarantees the existence of such a coupling for any fixed n . In order to extend this existence to the infinite sequence, observe that the set of functions depending on finitely many

coordinates is dense in the set of continuous increasing functions on $\{0, 1\}^{\mathbb{N}}$, and use the results in Strassen [11].)

Each admissible choice of ℓ_1, \dots, ℓ_C in (3.32) leads to a C -tuple $i_1 < \dots < i_C$ such that $A_{i_1} = \dots = A_{i_C} = 1$ (since it cuts out non-overlapping words, which is compatible with (3.36–3.37)), and for any such (i_1, \dots, i_C) there are at most L^C different admissible choices of the ℓ_c 's. Thus, we have

$$\sum_{(\ell_c), (\eta^{(c)}) \text{ good}} \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\eta^{(c-1)}|) \vee 1)^{-\alpha} \leq L^C L^{-\alpha} \sum_{\substack{0 < i_1 < \dots < i_C < \infty \\ A_{i_1} = \dots = A_{i_C} = 1}} \prod_{c=1}^C (i_c - i_{c-1})^{-\alpha}. \quad (3.42)$$

Using (3.19) and recalling the definition of $\phi(\alpha, p)$ in (2.2), we have

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log [\text{r.h.s. (3.42)}] \leq \frac{1 - \delta_1}{M} \left(\log(Mm_Q) - \phi(\alpha, p) \right) \quad (\omega, A) - a.s. \quad (3.43)$$

From (3.38) we know that $\log(1/p) \sim M(m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - 2\varepsilon_1)$ as $M \rightarrow \infty$ and so, by Lemma 2.1, we have

$$\text{r.h.s. (3.43)} \leq -(1 - \varepsilon_2)\alpha(m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - 2\varepsilon_1) \quad (3.44)$$

for any $\varepsilon_2 \in (0, 1)$, provided M is large enough. This completes the proof of Lemma 3.2, and hence of Proposition 3.1 for $Q \in \mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}})$. \square

3.5 Step 5: Removing the assumption of ergodicity

Sections 3.1–3.4 contain the main ideas behind the proof of Proposition 3.1. In the present section we extend the bound from $\mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv, fin}}(\tilde{E}^{\mathbb{N}})$. This requires setting up a variant of the argument in Sections 3.1–3.4 in which the ergodic components of Q are ‘‘approximated with a common length scale on the letter level’’. This turns out to be technically involved and to fall apart into 6 substeps.

Let $Q \in \mathcal{P}^{\text{inv, fin}}(\tilde{E}^{\mathbb{N}})$ have a non-trivial ergodic decomposition

$$Q = \int_{\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})} Q' W_Q(dQ'), \quad (3.45)$$

where W_Q is a probability measure on $\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})$ (Georgii [7], Proposition 7.22). We may assume w.l.o.g. that $H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) < \infty$, otherwise we can simply employ the annealed bound. Thus, W_Q is in fact supported on $\mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}}) \cap \{Q' : H(Q' | q_{\rho, \nu}^{\otimes \mathbb{N}}) < \infty\}$.

Fix $\varepsilon > 0$. In the following steps, we will construct an open neighbourhood $\mathcal{O}(Q) \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ of Q satisfying (3.1) (for technical reasons with ε replaced by some $\varepsilon' = \varepsilon'(\varepsilon)$ that becomes arbitrarily small as $\varepsilon \downarrow 0$).

3.5.1 Preliminaries

Observing that

$$m_Q = \int_{\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})} m_{Q'} W_Q(dQ') < \infty, \quad H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) = \int_{\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})} H(Q' | q_{\rho, \nu}^{\otimes \mathbb{N}}) W_Q(dQ') < \infty, \quad (3.46)$$

we can find $K_0, K_1, m^* > 0$ and a compact set

$$\mathcal{C} \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) \cap \text{supp}(W_Q) \cap \{Q : H(\cdot | q_{\rho, \nu}^{\otimes \mathbb{N}}) \leq K_0\} \quad (3.47)$$

such that

$$\sup\{H(\Psi_P | \nu^{\otimes \mathbb{N}}) : P \in \mathcal{C}\} \leq K_1, \quad (3.48)$$

$$\sup\{m_P : P \in \mathcal{C}\} \leq m^*, \quad (3.49)$$

$$\text{the family } \{\mathcal{L}_P(\tau_1) : P \in \mathcal{C}\} \text{ is uniformly integrable,} \quad (3.50)$$

$$W_Q(\mathcal{C}) \geq 1 - \varepsilon/2, \quad (3.51)$$

$$\int_{\mathcal{C}} H(Q' | q_{\rho, \nu}^{\otimes \mathbb{N}}) W_Q(dQ') \geq H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) - \varepsilon/2, \quad (3.52)$$

$$\int_{\mathcal{C}} m_{Q'} H(\Psi_{Q'} | \nu^{\otimes \mathbb{N}}) W_Q(dQ') \geq m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - \varepsilon/2. \quad (3.53)$$

In order to check (3.50), observe that $\mathbb{E}_Q[\tau_1] < \infty$ implies that there is a sequence (c_n) with $\lim_{n \rightarrow \infty} c_n = \infty$ such that

$$\mathbb{E}_Q[\tau_1 \mathbb{1}_{\{\tau_1 \geq c_n\}}] \leq \frac{6}{\pi^2 n^3} \frac{\varepsilon}{6}, \quad n \in \mathbb{N}. \quad (3.54)$$

Put

$$\hat{A}_n := \{Q' \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : \mathbb{E}_{Q'}[\tau_1 \mathbb{1}_{\{\tau_1 \geq c_n\}}] > 1/n\} \quad (3.55)$$

and $A := \bigcap_{n \in \mathbb{N}} (\hat{A}_n)^c$. Each \hat{A}_n is open, hence A is closed, and by the Markov inequality we have

$$W_Q(\{Q' : \mathbb{E}_{Q'}[\tau_1 \mathbb{1}_{\{\tau_1 \geq c_n\}}] > 1/n\}) \leq n \mathbb{E}_Q[\tau_1 \mathbb{1}_{\{\tau_1 \geq c_n\}}] \leq \frac{6}{\pi^2 n^2} \frac{\varepsilon}{6}. \quad (3.56)$$

Thus,

$$W_Q(A^c) = W_Q(\bigcup_{n \in \mathbb{N}} \hat{A}_n) \leq \frac{\varepsilon}{6} \sum_{n \in \mathbb{N}} \frac{6}{\pi^2 n^2} = \frac{\varepsilon}{6}. \quad (3.57)$$

This implies that the mapping

$$Q' \mapsto m_{Q'} H(\Psi_{Q'} | \nu^{\otimes \mathbb{N}}) \quad \text{is lower semicontinuous on } \mathcal{C}. \quad (3.58)$$

Indeed, if $w - \lim_{n \rightarrow \infty} Q'_n = Q''$ and $(Q'_n) \subset \mathcal{C}$, then $\lim_{n \rightarrow \infty} \mathbb{E}_{Q'_n}[\tau_1] = \lim_{n \rightarrow \infty} m_{Q'_n} = m_{Q''} = \mathbb{E}_{Q''}[\tau_1]$ and $w - \lim_{n \rightarrow \infty} \Psi_{Q'_n} = \Psi_{Q''}$ by uniform integrability (see Birkner [2], Remark 7).

Furthermore, we can find $N_0, L_0 \in \mathbb{N}$ with $L_0 \leq N_0$ and a finite set $\tilde{W} \subset \tilde{E}^{N_0}$ such that the following holds. Let

$$W := \left\{ \pi_{L_0}(\theta^i \kappa(\zeta)) : \zeta = (\zeta^{(1)}, \dots, \zeta^{(N_0)}) \in \tilde{W}, 0 \leq i < |\zeta^{(1)}| \right\} \quad (3.59)$$

be the set of words of length L_0 obtained by concatenating sentences from \tilde{W} , possibly shifting the ‘‘origin’’ inside the first word and restricting to the first L_0 letters. Then, denoting by \mathcal{D} the set of all $P \in \mathcal{P}^{\text{inv, fin}}(\tilde{E}^{\mathbb{N}}) \cap \mathcal{C}$ that satisfy

$$\sum_{\zeta \in \tilde{W}} P(\zeta) \geq 1 - \frac{\varepsilon}{3c_{\lceil 3/\varepsilon \rceil}}, \quad \forall \xi \in W : \Psi_P(\xi) \leq \frac{1 + \varepsilon/2}{m_P} \mathbb{E}_P \left[\mathbb{1}_{\tilde{W}}(\pi_{N_0} Y) \sum_{i=0}^{\tau_1 - 1} \mathbb{1}_{\{\xi\}}(\pi_{L_0} \theta^i \kappa(Y)) \right] \quad (3.60)$$

$$H(P | q_{\rho, \nu}^{\otimes \mathbb{N}}) + \varepsilon/4 \geq \frac{1}{N_0} \sum_{\zeta \in \tilde{W}} P(\zeta) \log \frac{P(\zeta)}{q_{\rho, \nu}^{\otimes N_0}(\zeta)} \geq H(P | q_{\rho, \nu}^{\otimes \mathbb{N}}) - \varepsilon/4, \quad (3.61)$$

$$m_P H(\Psi_P | \nu^{\otimes \mathbb{N}}) + \varepsilon/4 \geq \frac{m_P}{L_0} \sum_{w \in W} \Psi_P(w) \log \frac{\Psi_P(w)}{\nu^{\otimes L_0}(w)} \geq m_P H(\Psi_P | \nu^{\otimes \mathbb{N}}) - \varepsilon/4, \quad (3.62)$$

we can choose N_0 , L_0 and \widetilde{W} so large that the following inequalities hold:

$$W_Q(\mathcal{D}) \geq 1 - 3\varepsilon/4, \quad (3.63)$$

$$\int_{\mathcal{D}} H(P | q_{\rho, \nu}^{\otimes \mathbb{N}}) W_Q(dP) \geq H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) - 3\varepsilon/4, \quad (3.64)$$

$$\int_{\mathcal{D}} m_P H(\Psi_P | \nu^{\otimes \mathbb{N}}) W_Q(dP) \geq m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - 3\varepsilon/4. \quad (3.65)$$

We may choose the set \widetilde{W} in such a way that

$$\delta_{\widetilde{W}} := \min\{q_{\rho, \nu}^{\otimes N_0}(\zeta) : \zeta \in \widetilde{W}\} \cdot \frac{\min\{\nu^{\otimes L_0}(\xi) : \xi \in W\}}{\max\{|\zeta^{(1)}| : \zeta \in \widetilde{W}\}} \cdot \frac{1}{|\widetilde{W}|} > 0. \quad (3.66)$$

3.5.2 Approximating with a given length scale on the letter level

For $P \in \mathcal{P}^{\text{inv, fin}}(\widetilde{E}^{\mathbb{N}})$, we put

$$\delta_{P, \widetilde{W}} := \delta_{\widetilde{W}} \cdot \left(\min\{P(\zeta) : \zeta \in \widetilde{W}, P(\zeta) > 0\} \wedge \min\{\Psi_P(\xi) : \xi \in W, \Psi_P(\xi) > 0\} \right). \quad (3.67)$$

For $\delta > 0$ and $L \in \mathbb{N}$, we say that $P \in \mathcal{P}^{\text{inv, fin}}(\widetilde{E}^{\mathbb{N}})$ can be (δ, L) -approximated if there exists a finite subset $\mathcal{A}_P \subset \widetilde{E}^{\lceil L/m_P \rceil}$ of “ P -typical” sentences, each consisting of $\approx L/m_P$ words (we assume that $L > N_0 m_P$), such that

$$P|_{\mathcal{A}_P^{\lceil L/m_P \rceil}}(\mathcal{A}_P) > 1 - \frac{1}{2} \delta \cdot \delta_{P, \widetilde{W}} \quad (3.68)$$

and, for all $z = (y^{(1)}, \dots, y^{\lceil L/m_P \rceil}) \in \mathcal{A}_P$,

$$\begin{aligned} P(z) &\in \left[\exp[-\lceil L/m_P \rceil(H(Q) + \delta)], \exp[-\lceil L/m_P \rceil(H(Q) - \delta)] \right], \\ |\kappa(z)| &\in [L(1 - \delta), L(1 + \delta)], \\ P(K^{\lceil L/m_P \rceil} = z) &\in \left[\exp[-L(H(\Psi_Q) + \delta)], \exp[-L(H(\Psi_Q) - \delta)] \right], \\ \sum_{k=1}^{|\kappa(z)|} \log \nu(\kappa(z)_k) &\in [L(1 - \delta), L(1 + \delta)] \mathbb{E}_{\Psi_P}[\log \nu(X_1)], \\ \sum_{i=1}^{\lceil L/m_P \rceil} \log \rho(|y^{(i)}|) &\in [(L/m_P)(1 - \delta), (L/m_P)(1 + \delta)] \mathbb{E}_P[\log \rho(\tau_1)], \\ |\{z' \in \mathcal{A}_P : \kappa(z) = \kappa(z')\}| &\leq \exp[(L/m_P)(H_{\tau|K}(P) + \delta)]. \end{aligned} \quad (3.69)$$

By the third and the fourth line of (3.69) we have, using (1.26),

$$\mathbb{P}(X \text{ starts with some element of } \kappa(\mathcal{A}_P)) \leq \exp\left[-L(1 - 2\delta)H(\Psi_Q | \nu^{\otimes \mathbb{N}})\right]. \quad (3.70)$$

For P that can be (δ, L) -approximated, define an open neighbourhood of P via

$$\mathcal{U}_{(\delta, L)}(P) := \left\{ P' \in \mathcal{P}^{\text{inv}}(\widetilde{E}^{\mathbb{N}}) : \frac{P'(z)}{P(z)} \in (1 - \delta \cdot \delta_{P, \widetilde{W}}, 1 + \delta \cdot \delta_{P, \widetilde{W}}) \quad \forall z \in \mathcal{A}_P \right\}, \quad (3.71)$$

where $\mathcal{A}_P = \mathcal{A}_P(\delta, L)$ is the set from (3.68–3.69). By the results of Section 3.1 and the above, for given $P \in \mathcal{P}^{\text{erg, fin}}(\widetilde{E}^{\mathbb{N}}) \cap \mathcal{C}$ and $\delta_0 > 0$ there exist $\delta' \in (0, \delta_0)$ and L' such that

$$\forall L'' \geq L' : P \text{ can be } (\delta', L'')\text{-approximated.} \quad (3.72)$$

Assume that a given $P \in \mathcal{D}$ can be (δ, L) -approximated for some L such that $\lceil L/m_P \rceil \geq N_0$. We claim that then for any $P' \in \mathcal{D} \cap \mathcal{U}_{(\delta, L)}(P)$,

$$P'(\widetilde{E}^{\lceil L/m_P \rceil} \setminus \mathcal{A}_P) \leq 2\delta \cdot \delta_{P, \widetilde{W}}, \quad (3.73)$$

$$\forall \zeta \in \widetilde{W}: P'(\zeta) \leq \begin{cases} (1 + 3\delta)P(\zeta) & \text{if } P(\zeta) > 0, \\ 2\delta \cdot \delta_{P, \widetilde{W}} (\leq 2\delta \cdot \min\{q_{\rho, \nu}^{\otimes N_0}(\zeta'): \zeta' \in \widetilde{W}\}) & \text{otherwise,} \end{cases} \quad (3.74)$$

$$\forall \xi \in W: m_{P'}\Psi_{P'}(\xi) \leq \begin{cases} (1 + \varepsilon/2)(1 + 3\delta)m_P\Psi_P(\xi) & \text{if } \Psi_P(\xi) > 0, \\ (1 + \varepsilon/2)2\delta \min\{\nu^{\otimes L_0}(\xi'): \xi' \in W\} & \text{otherwise,} \end{cases} \quad (3.75)$$

$$m_{P'} \geq (1 - 3\delta)(m_P - \varepsilon) (\geq (1 - 3\delta - \varepsilon)m_P). \quad (3.76)$$

(3.73) follows from (3.68) and (3.71). To verify (3.74), note that, for $\zeta \in \widetilde{W}$,

$$\begin{aligned} P'(\zeta) &\leq \sum_{z \in \mathcal{A}_P: \pi_{N_0}(z) = \zeta} P'(z) + \sum_{z \in \widetilde{E}^{\lceil L/m_P \rceil} \setminus \mathcal{A}_P: \pi_{N_0}(z) = \zeta} P'(z) \\ &\leq (1 + \delta) \sum_{z \in \mathcal{A}_P: \pi_{N_0}(z) = \zeta} P(z) + P'(\widetilde{E}^{\lceil L/m_P \rceil} \setminus \mathcal{A}_P) \end{aligned} \quad (3.77)$$

and use (3.73) on the last term in the second line, observing that $\delta_{P, \widetilde{W}} \leq P(\zeta)$ whenever $\zeta \in \widetilde{W}$ and $P(\zeta) > 0$. To verify (3.75), observe that, for $\xi \in W$ (recall the definition of $\Psi_{P'}$ from (1.10)), using (3.60),

$$\begin{aligned} (1 + \varepsilon/2)^{-1} m_{P'}\Psi_{P'}(\xi) &\leq \sum_{\zeta \in \widetilde{W}} P'(\zeta) \sum_{i=0}^{|\zeta^{(1)}|-1} \mathbb{1}_{\{\xi\}}(\pi_{L_0}(\theta^i \kappa(\zeta))) \\ &\leq (1 + \delta)m_P\Psi_P(\xi) + \sum_{\zeta \in \widetilde{W}: P(\zeta)=0} |\zeta^{(1)}|P'(\zeta) \end{aligned} \quad (3.78)$$

and that the sum in the second line above is bounded by $|\widetilde{W}| \cdot \max\{|\zeta^{(1)}| : \zeta \in \widetilde{W}\} \cdot 2\delta \cdot \delta_{P, \widetilde{W}}$, which is not more than $2\delta m_P\Psi_P(\xi)$ if $\Psi_P(\xi) > 0$ and not more than $2\delta \min\{\nu^{\otimes L_0}(\xi'): \xi' \in W\}$ otherwise. Lastly, to verify (3.76), note that

$$P'(\zeta) \geq (1 - 3\delta)P(\zeta) \quad \forall \zeta \in \widetilde{W} \quad (3.79)$$

(which can be proved in the same way as (3.74)), so that

$$m_{P'} = \sum_{y \in \widetilde{E}} |y|P'(y) \geq \sum_{\zeta \in \widetilde{W}} |\zeta^{(1)}|P'(\zeta) \geq (1 - 3\delta) \sum_{\zeta \in \widetilde{W}} |\zeta^{(1)}|P(\zeta). \quad (3.80)$$

Furthermore,

$$m_P \leq \sum_{\zeta \in \widetilde{W}} |\zeta^{(1)}|P(\zeta) + c_{\lceil 3/\varepsilon \rceil} P(\widetilde{E}^{N_0} \setminus \widetilde{W}) + \sum_{y \in \widetilde{E}: |y| > c_{\lceil 3/\varepsilon \rceil}} |y|P(y). \quad (3.81)$$

Observing that the second and the third term on the right-hand side are each at most $\varepsilon/3$, we find that (3.80–3.81) imply (3.76).

Finally, observe that (3.74–3.76) imply that there exists $\delta_0 (= \delta_0(\varepsilon)) > 0$ with the following property: For any $P, P' \in \mathcal{D}$ such that P can be (δ, L) -approximated for some L with $\lceil L/m_P \rceil \geq N_0$ and $\delta \leq \delta_0$ and $P' \in \mathcal{U}_{(\delta, L)}(P)$, we have

$$H(P' | q_{\rho, \nu}^{\otimes N}) \leq (1 + \varepsilon) \left(H(P | q_{\rho, \nu}^{\otimes N}) + \varepsilon \right) \quad \text{and} \quad (3.82)$$

$$m_{P'} H(\Psi_{P'} | \nu^{\otimes N}) \leq (1 + \varepsilon) \left(m_P H(\Psi_P | \nu^{\otimes N}) + \varepsilon \right). \quad (3.83)$$

Here, (3.82) follows from the observation

$$\begin{aligned}
& H(P' | q_{\rho, \nu}^{\otimes \mathbb{N}}) - \frac{\varepsilon}{4} \\
& \leq \frac{1}{N_0} \sum_{\zeta \in \widetilde{W}} P'(\zeta) \log \frac{P'(\zeta)}{q_{\rho, \nu}^{\otimes N_0}(\zeta)} \\
& \leq \frac{1+3\delta}{N_0} \sum_{\zeta \in \widetilde{W}} P(\zeta) \log \frac{(1+3\delta)P(\zeta)}{q_{\rho, \nu}^{\otimes N_0}(\zeta)} + \frac{1}{N_0} \sum_{\zeta \in \widetilde{W}: P(\zeta)=0} P'(\zeta) \log \frac{\min\{q_{\rho, \nu}^{\otimes N_0}(\zeta') : \zeta' \in \widetilde{W}\}}{q_{\rho, \nu}^{\otimes N_0}(\zeta)} \\
& \leq (1+3\delta) \left(H(P | q_{\rho, \nu}^{\otimes \mathbb{N}}) + \frac{\varepsilon}{4} \right) + \frac{1+3\delta}{N_0} \log(1+3\delta).
\end{aligned} \tag{3.84}$$

Similarly, observing that

$$\begin{aligned}
& m_{P'} \sum_{\xi \in W} \Psi_{P'}(\xi) \log \frac{m_{P'} \Psi_{P'}(\xi)}{m_{P'} \nu^{\otimes L_0}(\xi)} \\
& \leq \left(1 + \frac{\varepsilon}{2}\right) (1+3\delta) m_P \sum_{\xi \in W} \Psi_P(\xi) \log \frac{(1+\varepsilon/2)(1+3\delta)m_P \Psi_P(\xi)}{(1-3\delta-\varepsilon)m_P \nu^{\otimes L_0}(\xi)} \\
& \quad + m_{P'} \sum_{\xi \in W: \Psi_P(\xi)=0} \Psi_{P'}(\xi) \log \frac{(1+\varepsilon/2)2\delta \min\{\nu^{\otimes L_0}(\xi') : \xi' \in W\}}{\nu^{\otimes L_0}(\xi)} \\
& \leq \left(1 + \frac{\varepsilon}{2}\right) (1+3\delta) L_0 m_P H(\Psi_P | \nu^{\otimes \mathbb{N}}) + \varepsilon/2 + m^* \log \frac{(1+3\delta)(1+\varepsilon/2)}{1-3\delta-\varepsilon}
\end{aligned} \tag{3.85}$$

we obtain (3.83) in view of (3.62).

3.5.3 Approximating the ergodic decomposition

In the previous subsection, we have approximated a given $P \in \mathcal{P}^{\text{erg,fin}}(\widetilde{E}^{\mathbb{N}})$, i.e., we have constructed a certain neighbourhood of P w.r.t. the weak topology, which requires only conditions on the frequencies of sentences whose concatenations are $\approx L$ letters long. While the required L will in general vary with P , we now want to construct a compact $\mathcal{C}' \subset \mathcal{C}$ such that $W_Q(\mathcal{C}')$ is still close to 1 and all $P \in \mathcal{C}'$ can be approximated on the same scale L (on the letter level). To this end, let

$$\mathcal{D}_{\varepsilon', L'} := \{P \in \mathcal{D} : P \text{ can be } (\varepsilon', L')\text{-approximated}\}. \tag{3.86}$$

By (3.72), we have

$$\bigcup_{\substack{\varepsilon' \in (0, \varepsilon/2) \\ L' \in \mathbb{N}}} \mathcal{D}_{\varepsilon', L'} = \mathcal{P}^{\text{erg,fin}}(\widetilde{E}^{\mathbb{N}}) \cap \mathcal{C}, \tag{3.87}$$

so, in view of (3.51–3.53), we can choose

$$0 < \varepsilon_1 < \frac{\varepsilon}{2m^*(1 \vee K_1)} \wedge \frac{\delta_0}{2} \tag{3.88}$$

and $L \in \mathbb{N}$ such that

$$W_Q(\mathcal{D}_{\varepsilon_1, L}) \geq 1 - \varepsilon, \tag{3.89}$$

$$\int_{\mathcal{D}_{\varepsilon_1, L}} H(Q' | q_{\rho, \nu}^{\otimes \mathbb{N}}) W_Q(dQ') \geq H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) - \varepsilon, \tag{3.90}$$

$$\int_{\mathcal{D}_{\varepsilon_1, L}} m_{Q'} H(\Psi_{Q'} | \nu^{\otimes \mathbb{N}}) W_Q(dQ') \geq m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - \varepsilon. \tag{3.91}$$

For $P \in \mathcal{D}_{\varepsilon_1, L}$, let

$$\mathcal{U}'(P) := \left\{ P' \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : \frac{P'(z)}{P(z)} \in \left(1 - \frac{\varepsilon_1}{2} \delta_{P, \tilde{W}}, 1 + \frac{\varepsilon_1}{2} \delta_{P, \tilde{W}} \right) \quad \forall z \in \mathcal{A}_P \right\}, \quad (3.92)$$

where \mathcal{A}_P is the set from (3.68–3.69) that appears in the definition of $\mathcal{U}_{(\varepsilon_1, L)}(P)$ and $\delta_{P, \tilde{W}}$ is defined in (3.67). Note that $\mathcal{U}'(P) \subset \mathcal{U}_{(\varepsilon_1, L)}(P)$. Indeed, $\inf_{P \in \mathcal{D}_{\varepsilon_1, L}} \text{dist}(\mathcal{U}'(P), \mathcal{U}_{(\varepsilon_1, L)}(P)^c) > 0$ if we metrize the weak topology. Consequently,

$$\mathcal{C}' := \mathcal{C} \cap \overline{\cup_{P \in \mathcal{D}_{\varepsilon_1, L}} \mathcal{U}'(P)} \quad (\supset \mathcal{D}_{\varepsilon_1, L}) \quad (3.93)$$

is compact and satisfies $W_Q(\mathcal{C}') \geq 1 - \varepsilon$, and

$$\mathcal{C}' \subset \bigcup_{P \in \mathcal{D}_{\varepsilon_1, L}} \mathcal{U}_{(\varepsilon_1, L)}(P) \quad (3.94)$$

is an open cover. By compactness there exist $R \in \mathbb{N}$ and (pairwise different) $Q_1, \dots, Q_R \in \mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}}) \cap \mathcal{C}'$ such that

$$\mathcal{U}_{(\varepsilon_1, L)}(Q_1) \cup \dots \cup \mathcal{U}_{(\varepsilon_1, L)}(Q_R) \supset \mathcal{C}', \quad (3.95)$$

where $\mathcal{U}_{(\varepsilon_1, L)}(Q_r)$ is of the type (3.71) with a set $\mathcal{A}_r \subset \tilde{E}^{M_r}$ satisfying (3.68–3.69) with P replaced by Q_r , and $M_r = \lceil L/m_{Q_r} \rceil$.

For $z \in \cup_{n \in \mathbb{N}} \tilde{E}^n$ consider the probability measure on $[0, 1]$ given by $\mu_{Q, z}(B) := W_Q(\{Q' \in \mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}}) : Q'(z) \in B\})$, $B \subset [0, 1]$ measurable. Observing that

$$\bigcup_{r=1}^R \bigcup_{z \in \mathcal{A}_r} \{u \in [0, 1] : u \text{ is an atom of } \mu_{Q, z}\} \quad (3.96)$$

is at most countable, we can find $\varepsilon_2 \in [\varepsilon_1, \varepsilon_1 + \varepsilon_2^2)$ (note that still $\varepsilon_2 < 2\varepsilon_1$) and $\tilde{\delta} > 0$ such that

$$W_Q \left(\left\{ Q' \in \mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}}) : \begin{array}{l} Q'(z)/Q_r(z) \in [1 - (\varepsilon_2 + \tilde{\delta})\delta_{Q_r, \tilde{W}}, 1 - (\varepsilon_2 - \tilde{\delta})\delta_{Q_r, \tilde{W}}] \text{ or} \\ Q'(z)/Q_r(z) \in [1 + (\varepsilon_2 - \tilde{\delta})\delta_{Q_r, \tilde{W}}, 1 + (\varepsilon_2 + \tilde{\delta})\delta_{Q_r, \tilde{W}}] \\ \text{for some } r \in \{1, \dots, R\} \text{ and } z \in \mathcal{A}_r \end{array} \right\} \right) \leq \frac{\varepsilon}{1 \vee K_0 \vee m^* K_1}. \quad (3.97)$$

Define “disjointified” versions of the $\mathcal{U}_{(\varepsilon, L)}(Q_r)$ as follows. For $r = 1, \dots, R$, put iteratively

$$\tilde{\mathcal{U}}_r := \left\{ Q' \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : \begin{array}{l} Q'(z) \in Q_r(z)[1 - \varepsilon_2 \delta_{Q_r, \tilde{W}}, 1 + \varepsilon_2 \delta_{Q_r, \tilde{W}}] \text{ for all } z \in \mathcal{A}_r \\ \text{and for each } r' < r \text{ there is } z' \in \mathcal{A}_{r'} \text{ such that} \\ Q'(z') \notin Q_{r'}(z')[1 - (\varepsilon_2 + \tilde{\delta})\delta_{Q_{r'}, \tilde{W}}, 1 + (\varepsilon_2 + \tilde{\delta})\delta_{Q_{r'}, \tilde{W}}] \end{array} \right\}. \quad (3.98)$$

It may happen that some of the $\tilde{\mathcal{U}}_r$ are empty or satisfy $W_Q(\tilde{\mathcal{U}}_r) = 0$. We then (silently) remove these and re-number the remaining ones. Note that each $\tilde{\mathcal{U}}_r$ is an open subset of $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ and

$$W_Q(\cup_{r=1}^R \tilde{\mathcal{U}}_r) = \sum_{r=1}^R W_Q(\tilde{\mathcal{U}}_r) \geq 1 - 2\varepsilon, \quad (3.99)$$

since $W_Q(\mathcal{C}' \setminus \cup_{r=1}^R \tilde{\mathcal{U}}_r) \leq \varepsilon$.

For $r = 1, \dots, R$, we have, using (3.82–3.83) and the choice of ε_2 ($\leq 2\varepsilon_1 \leq \delta_0$),

$$W_Q(\tilde{\mathcal{U}}_r \cap \mathcal{D}) \left(H(Q_r | q_{\rho, \nu}^{\otimes N}) + \varepsilon \right) \geq \frac{1}{1 + \varepsilon} \int_{\tilde{\mathcal{U}}_r \cap \mathcal{D}} H(Q' | q_{\rho, \nu}^{\otimes N}) W_Q(dQ'), \quad (3.100)$$

$$W_Q(\tilde{\mathcal{U}}_r \cap \mathcal{D}) \left(m_{Q_r} H(\Psi_{Q_r} | \nu^{\otimes N}) + \varepsilon \right) \geq \frac{1}{1 + \varepsilon} \int_{\tilde{\mathcal{U}}_r \cap \mathcal{D}} m_{Q'} H(\Psi_{Q'} | \nu^{\otimes N}) W_Q(dQ'), \quad (3.101)$$

so that altogether, using (3.90–3.91),

$$\begin{aligned} \sum_{r=1}^R W_Q(\tilde{\mathcal{U}}_r) \left\{ H(Q_r | q_{\rho, \nu}^{\otimes N}) + (\alpha - 1) m_{Q_r} H(\Psi_{Q_r} | \nu^{\otimes N}) \right\} \\ \geq \frac{1}{1 + \varepsilon} \left(H(Q | q_{\rho, \nu}^{\otimes N}) + (\alpha - 1) m_Q H(\Psi_Q | \nu^{\otimes N}) \right) - 2\alpha\varepsilon. \end{aligned} \quad (3.102)$$

3.5.4 More layers: long sentences with the right pattern frequencies

For $z \in \cup_{n \in \mathbb{N}} \tilde{E}^n$ and $\xi = (\xi^{(1)}, \dots, \xi^{(\tilde{M})}) \in \tilde{E}^{\tilde{M}}$ (with $\tilde{M} > |z|$), let

$$\text{freq}_z(\xi) = \frac{1}{\tilde{M}} \left| \{1 \leq i \leq \tilde{M} - |z| + 1 : (\xi^{(i)}, \dots, \xi^{(i+|z|-1)}) = z\} \right| \quad (3.103)$$

be the empirical frequency of z in ξ . Note that, for any $P \in \mathcal{P}^{\text{erg, fin}}(\tilde{E}^{\mathbb{N}})$, $z \in \cup_{n \in \mathbb{N}} \tilde{E}^n$ and $\varepsilon' > 0$, we have

$$\lim_{\tilde{M} \rightarrow \infty} P \left(\{ \xi \in \tilde{E}^{\tilde{M}} : \text{freq}_z(\xi) \in P(z)(1 - \varepsilon', 1 + \varepsilon') \} \right) = 1 \quad (3.104)$$

and

$$\lim_{\tilde{M} \rightarrow \infty} P \left(\{ \xi \in \tilde{E}^{\tilde{M}} : |\kappa(\xi)| \in M(m_P - \varepsilon', m_P + \varepsilon') \} \right) = 1. \quad (3.105)$$

For $\tilde{M} \in \mathbb{N}$ and $r \in \{1, \dots, R\}$, put

$$V_{r, \tilde{M}} := \left\{ \xi \in \tilde{E}^{\tilde{M}} : \begin{array}{l} |\kappa(\xi)| \in \tilde{M}(m_{Q_r} - \varepsilon_2, m_{Q_r} + \varepsilon_2), \\ \text{freq}_z(\xi) \in Q_r(z)(1 - \varepsilon_2 \delta_{Q_r, \tilde{W}}, 1 + \varepsilon_2 \delta_{Q_r, \tilde{W}}) \text{ for all } z \in \mathcal{A}_r, \\ \text{and for each } r' < r \text{ there is a } z' \in \mathcal{A}_{r'} \text{ such that} \\ \text{freq}_{z'}(\xi) \notin Q_{r'}(z') [1 - (\varepsilon_2 + \tilde{\delta}) \delta_{Q_{r'}, \tilde{W}}, 1 + (\varepsilon_2 + \tilde{\delta}) \delta_{Q_{r'}, \tilde{W}}] \end{array} \right\}. \quad (3.106)$$

Note that when $|E| < \infty$, also $|V_{r, \tilde{M}}| < \infty$. Furthermore, $V_{r, \tilde{M}} \cap V_{r', \tilde{M}} = \emptyset$ for $r \neq r'$. For $\xi \in V_{r, \tilde{M}}$, we have

$$\left| \left\{ 1 \leq i \leq \tilde{M} - M_r + 1 : (\xi^{(i)}, \xi^{(i+1)}, \dots, \xi^{(i+M_r-1)}) \in \mathcal{A}_r \right\} \right| \geq \tilde{M}(1 - 2\varepsilon_2), \quad (3.107)$$

in particular, there are at least $K_r := \lfloor \tilde{M}(1 - 3\varepsilon_2)/M_r \rfloor$ elements $z_1, \dots, z_{K_r} \in \mathcal{A}_r$ (not necessarily distinct) appearing in this order as disjoint subwords of ξ . The z_k 's can for example be constructed in a “greedy” way, parsing ξ from left to right as in Section 3.2 (see, in particular, (3.21)). This implies, in particular, that

$$\begin{aligned} \prod_{i=1}^{\tilde{M}} \rho(|\xi^{(i)}|) &\leq \prod_{k=1}^{K_r} \prod_{w \text{ in } z_k} \rho(|w|) \leq \left(\exp \left[(1 - \varepsilon_2) \tilde{M}_r \mathbb{E}_{Q_r} [\log \rho(\tau_1)] \right] \right)^{K_r} \\ &\leq \exp \left[(1 - 4\varepsilon_2) \tilde{M} \mathbb{E}_{Q_r} [\log \rho(\tau_1)] \right] \leq \exp \left[\tilde{M} \mathbb{E}_{Q_r} [\log \rho(\tau_1)] + \tilde{M} \varepsilon'_\rho \right] \end{aligned} \quad (3.108)$$

if \widetilde{M} is large enough, where $c'_\rho := \sup_{k \in \text{supp}(\rho)} \{-\log(\rho(k))/k\} (< \infty)$ and we use that $\varepsilon_2 m^* \leq \varepsilon$ by definition. Furthermore, for each $r \in \{1, \dots, R\}$ and $\eta \in V_{r, \widetilde{M}}$, we have

$$|\{\zeta \in V_{r, \widetilde{M}}: \kappa(\zeta) = \kappa(\eta)\}| \leq \exp \left[\widetilde{M}(H_{\tau|K}(Q_r) + \delta_1) \right], \quad (3.109)$$

where δ_1 can be made arbitrarily small by choosing ε small. (Note that the quantity on the left-hand side is the number of ways in which $\kappa(\eta)$ can be “re-cut” to obtain another element of $V_{r, \widetilde{M}}$.) In order to check (3.109), we note that any $\zeta \in V_{r, \widetilde{M}}$ must contain at least K_r disjoint subsentences from \mathcal{A}_r , and each $z \in \mathcal{A}_r \subset \widetilde{E}^{M_r}$ satisfies $|\kappa(z)| \geq L$. Hence there are at most

$$\binom{\widetilde{M}(m_{Q_r} + \varepsilon_2) - K_r(L - 1)}{K_r} \leq 2^{4\varepsilon_2 \widetilde{M} m_{Q_r}} \leq 2^{4\varepsilon_2 m^* \widetilde{M}} \quad (3.110)$$

choices for the positions in the letter sequence $\kappa(\eta)$ where the concatenations of the disjoint subsentences from \mathcal{A}_r can begin, and there are at most

$$\binom{\widetilde{M} - K_r(M_r - 1)}{K_r} \leq 2^{3\varepsilon_2 \widetilde{M}} \quad (3.111)$$

choices for the positions in the word sequence ζ where the subsentences from \mathcal{A}_r can begin. By construction (recall the last line of (3.69)), each $z \in \mathcal{A}_r$ can be “re-cut” in not more than $\exp[(L/m_{Q_r})(H_{\tau|K}(Q_r) + \varepsilon_2)]$ many ways. Combining these observations with the fact that

$$\left(\exp \left[(L/m_{Q_r})(H_{\tau|K}(Q_r) + \varepsilon_2) \right] \right)^{K_r} \leq \exp \left[\frac{\widetilde{M}}{M_r} M_r (H_{\tau|K}(Q_r) + \varepsilon_2) \right], \quad (3.112)$$

we get (3.109) with $\delta_1 := \varepsilon_2 + 3\varepsilon_2 \log 2 + 4\varepsilon_2 m^* \log 2$.

We see from (3.104–3.105) and the definitions of \widetilde{U}_r and $V_{r, \widetilde{M}}$ that, for any $\varepsilon' > 0$

$$\bigcup_{\widetilde{M} \in \mathbb{N}} \left\{ P \in \widetilde{U}_r: P(V_{r, \widetilde{M}}) > 1 - \varepsilon' \right\} = \widetilde{U}_r. \quad (3.113)$$

Put $\varepsilon_3 := \varepsilon_2 \min_{r=1, \dots, R} W_Q(\widetilde{U}_r) (\leq \varepsilon_2)$. We can choose \widetilde{M} so large that

$$W_Q \left(\left\{ P \in \widetilde{U}_r: P(V_{r, \widetilde{M}}) > 1 - \frac{\varepsilon_3}{4} \right\} \right) > W_Q(\widetilde{U}_r) \left(1 - \frac{\varepsilon_2}{2} \right), \quad r = 1, \dots, R. \quad (3.114)$$

For $M' > \widetilde{M}$ and $r = 1, \dots, R$, put

$$W_{r, M'} := \left\{ \zeta \in \widetilde{E}^{M'}: \text{freq}_{V_{r, \widetilde{M}}}(\zeta) > 1 - \varepsilon_3/2 \right\}. \quad (3.115)$$

Note that for $r \neq r'$ (because $V_{r, \widetilde{M}} \cap V_{r', \widetilde{M}} = \emptyset$) there cannot be much overlap between $\zeta \in W_{r, M'}$ and $\eta \in W_{r', M'}$:

$$\max\{k: k\text{-suffix of } \zeta = k\text{-prefix of } \eta\} \leq \varepsilon_3 M' \quad (3.116)$$

(here, the k -prefix of $\eta \in \widetilde{E}^n$, $k < n$, consists of the first k words, the k -suffix of the last k words). To see this, note that any subsequence of length k of ζ must contain at least $(k - \varepsilon_3 M'/2)_+$ positions where a sentence from $V_{r, \widetilde{M}}$ starts, and any subsequence of length k of η must contain at least $(k - \varepsilon_3 M'/2)_+$ positions where a sentence from $V_{r', \widetilde{M}}$ starts, so any k appearing in (3.116) must satisfy $2(k - \varepsilon_3 M'/2)_+ \leq k$, which enforces $k \leq \varepsilon_3 M'$.

Observe that (3.115) implies that we may choose M' so large that for $r = 1, \dots, R$,

$$\text{each } \zeta \in W_{r, M'} \text{ contains at least } (1 - \varepsilon_3) \frac{M'}{M} \text{ disjoint subsentences from } V_{r, \widetilde{M}}. \quad (3.117)$$

For $P \in \mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$ with $P(V_{r,\tilde{M}}) > 1 - \varepsilon_3/3$ we have

$$\lim_{M' \rightarrow \infty} P(W_{r,M'}) = 1, \quad (3.118)$$

and hence

$$\bigcup_{M' > \tilde{M}} \left\{ P \in \tilde{\mathcal{U}}_r : P(W_{r,M'}) > 1 - \varepsilon_2 \right\} \supset \left\{ P \in \tilde{\mathcal{U}}_r : P(V_{r,\tilde{M}}) > 1 - \varepsilon_3/3 \right\}, \quad (3.119)$$

and so we can choose M' so large that

$$W_Q \left(\left\{ P \in \tilde{\mathcal{U}}_r : P(W_{r,M'}) > 1 - \varepsilon_2 \right\} \right) > W_Q(\tilde{\mathcal{U}}_r)(1 - \varepsilon_2), \quad r = 1, \dots, R. \quad (3.120)$$

Now define

$$\mathcal{O}(Q) := \left\{ Q' \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : Q'(W_{r,M'}) > W_Q(\tilde{\mathcal{U}}_r)(1 - 2\varepsilon_2), r = 1, \dots, R \right\}. \quad (3.121)$$

Note that $\mathcal{O}(Q)$ is open in the weak topology on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, since it is defined in terms of requirements on certain finite marginals of Q' , and that for $r = 1, \dots, R$,

$$Q(W_{r,M'}) = \int_{\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})} Q'(W_{r,M'}) W_Q(dQ') \geq \int_{\tilde{\mathcal{U}}_r} Q'(W_{r,M'}) W_Q(dQ') \geq (1 - \varepsilon_2)^2 W_Q(\tilde{\mathcal{U}}_r) \quad (3.122)$$

by (3.120), so that in fact $Q \in \mathcal{O}(Q)$.

3.5.5 Estimating the large deviation probability: good loops and filling loops

Consider a choice of “cut-points” $j_1 < \dots < j_N$ as appearing in the sum in (3.23). Note that, by the definition of $\mathcal{O}(Q)$ (recall (3.16–3.17)),

$$R_{j_1, \dots, j_N}^N(X) \in \mathcal{O}(Q) \quad (3.123)$$

enforces

$$\left| \left\{ 1 \leq i \leq N - M' : (X|_{(j_{i-1}, j_i]}, \dots, X|_{(j_{i+M'-1}, j_{i+M'}]}) \in W_{r,M'} \right\} \right| \geq N W_Q(\tilde{\mathcal{U}}_r)(1 - 3\varepsilon_2), \quad r = 1, \dots, R, \quad (3.124)$$

when N is large enough. This fact, together with (3.116), enables us to pick at least

$$J := \sum_{r=1}^R \lceil (1 - 4\varepsilon_2)N/M' \rceil W_Q(\tilde{\mathcal{U}}_r) \quad (3.125)$$

subsentences ζ_1, \dots, ζ_J occurring as disjoint subsentences in this order on ξ_N such that

$$\left| \left\{ 1 \leq j \leq J : \zeta_j \in W_{r,M'} \right\} \right| > (1 - 4\varepsilon_2) W_Q(\tilde{\mathcal{U}}_r) \frac{N}{M'}, \quad r = 1, \dots, R, \quad (3.126)$$

where we note that $J \geq (1 - 4\varepsilon_2)(1 - 2\varepsilon)(N/M') (\geq (1 - 8\varepsilon)(N/M'))$ by (3.99). Indeed, we can for example construct these ζ_j 's iteratively in a “greedy” way, parsing through ξ_N from left to right and always picking the next possible subsentence from one of the R types whose count does not yet exceed $(1 - 4\varepsilon_2) W_Q(\tilde{\mathcal{U}}_r) (N/M')$, as follows. Let $k_{s,r}$ be total number of subsentences of type r we have chosen after the s -th step ($k_{0,1} = \dots = k_{0,R} = 0$). If in the s -th step we have picked $\zeta_s = (\xi_N^{(p)}, \dots, \xi_N^{(p+M'-1)})$ at position p , then let

$$p' := \min \left\{ i \geq p + M' : \text{at position } i \text{ in } \xi_N \text{ starts a sentence from } W_{u,M'} \text{ for some } u \in U_s \right\}, \quad (3.127)$$

where $U_s := \{r: k_{r,s} < (1 - 4\varepsilon_2)W_Q(\tilde{\mathcal{U}}_r)(N/M')\}$, pick the next subsentence ζ_{s+1} starting at position p' (say, of type u) and increase the corresponding $k_{s+1,u}$. Repeat this until $k_{s,r} \geq (1 - 4\varepsilon_2)W_Q(\tilde{\mathcal{U}}_r)(N/M')$ for $r = 1, \dots, R$.

In order to verify that this algorithm does not get stuck, let $\text{rem}(s, r)$ be the ‘‘remaining’’ number of positions (to the right of the position where the word was picked in the s -th step) where a subsentence from $W_{r,M'}$ begins on ξ_N . By (3.124), we have

$$\text{rem}(0, r) \geq NW_Q(\tilde{\mathcal{U}}_r)(1 - 3\varepsilon_2). \quad (3.128)$$

If in the s -th step a subsentence of type r is picked, then we have $\text{rem}(s+1, r) \geq \text{rem}(s, r) - M'$, and for $r' \neq r$ we have $\text{rem}(s+1, r') \geq \text{rem}(s, r') - \varepsilon_3 M'$ by (3.116). Thus,

$$\begin{aligned} \text{rem}(s, r) &\geq \text{rem}(0, r) - k_{s,r}M' - (s - k_{s,r})\varepsilon_3 M' \\ &= \text{rem}(0, r) - k_{s,r}(1 - \varepsilon_3)M' - s\varepsilon_3 M', \end{aligned} \quad (3.129)$$

which is > 0 as long as $k_{s,r} < (1 - 4\varepsilon_2)W_Q(\tilde{\mathcal{U}}_r)(N/M')$ and $s < J$.

A. Combinatorial consequences. By (3.117) and (3.126), $R_{j_1, \dots, j_N}^N(X) \in \mathcal{O}(Q)$ implies that ξ_N contains at least

$$C := \sum_{r=1}^R \left[(1 - 4\varepsilon_2)W_Q(\tilde{\mathcal{U}}_r) \frac{N}{M'} \right] \left[(1 - \varepsilon_2) \frac{M'}{M} \right] \quad \left(\geq (1 - 5\varepsilon_2)(1 - 2\varepsilon) \frac{N}{M} \right) \quad (3.130)$$

disjoint subsentences η_1, \dots, η_C (appearing in this order in ξ_N) such that at least

$$\frac{N}{M}(1 - 6\varepsilon_2)W_Q(\tilde{\mathcal{U}}_r) \text{ of the } \eta_c \text{'s are from } V_{r, \tilde{M}}, \quad r = 1, \dots, R. \quad (3.131)$$

Let k_1, \dots, k_C ($k_{c+1} \geq k_c + \tilde{M}$, $1 \leq c < C$) be the indices where the disjoint subsentences η_c start in ξ_N , i.e.,

$$\eta_c = \left(\xi_N^{(k_c)}, \xi_N^{(k_c+1)}, \dots, \xi_N^{(k_c + \tilde{M} - 1)} \right) \in V_{r_c, \tilde{M}}, \quad i = c, \dots, C, \quad (3.132)$$

and the r_c 's must respect the frequencies dictated by the $W_Q(\tilde{\mathcal{U}}_r)$'s as in (3.131). Thus, each choice (j_1, \dots, j_N) yielding a non-zero summand in (3.23) leads to a triple

$$(\ell_1, \dots, \ell_C), (r_1, \dots, r_C), (\bar{\eta}_1, \dots, \bar{\eta}_C) \quad (3.133)$$

such that $\bar{\eta}_c \in \kappa(V_{r_c, \tilde{M}})$, $\ell_{c+1} \geq \ell_c + |\bar{\eta}_c|$, the r_c 's respect the frequencies as in (3.131), and

$$\text{the word } \bar{\eta}_c \text{ starts at position } \ell_c \text{ in } X \text{ for } c = 1, \dots, C. \quad (3.134)$$

As in Section 3.3, we call such triples *good*, the loops inside the subsentences η_i *good* loops, the others *filling* loops.

Fix a good triple for the moment. In order to count how many choices of $j_1 < \dots < j_N$ can lead to this particular triple and to estimate their contribution, observe the following:

1. There are at most

$$\binom{N - C(\tilde{M} - 1)}{C} \leq \exp(\delta'_1 N) \quad (3.135)$$

choices for the $k_1 < \dots < k_C$, where δ'_1 can be made arbitrarily small by choosing ε small and \tilde{M} large.

2. Once the k_c 's are fixed, by (3.109) and (3.131) there are at most

$$\begin{aligned} & \prod_{r=1}^R \left(\exp \left[\widetilde{M} (H_{\tau|K}(Q_r) + \delta_1) \right] \right)^{\frac{N}{M} W_Q(\widetilde{\mathcal{U}}_r)} \\ &= \exp \left[N \sum_{r=1}^R W_Q(\widetilde{\mathcal{U}}_r) (H_{\tau|K}(Q_r) + \delta_1) \right] \end{aligned} \quad (3.136)$$

choices for the good loops and, by (3.108), for each choice of the good loops the product of the $\rho(j_k - j_{k-1})$'s inside the good loops is at most

$$\begin{aligned} & \prod_{r=1}^R \left(\exp \left[\widetilde{M} \mathbb{E}_{Q_r}[\log \rho(\tau_1)] \right] + \widetilde{M} c'_\rho \varepsilon \right)^{\frac{N}{M} W_Q(\widetilde{\mathcal{U}}_r)} \\ & \leq \exp \left[N c'_\rho \varepsilon + N \sum_{r=1}^R W_Q(\widetilde{\mathcal{U}}_r) \mathbb{E}_{Q_r}[\log \rho(\tau_1)] \right]. \end{aligned} \quad (3.137)$$

3. For each choice of the k_c 's, the contribution of the filling loops to the weight is

$$\begin{aligned} & \rho^{*(k_1-1)}(\ell_1 - 1) \prod_{c=1}^{C-1} \rho^{*(k_{c+1}-k_c-\widetilde{M})}(\ell_{c+1} - \ell_c - |\overline{\eta}_c|) \\ & \leq (C_\rho \vee 1)^C k_1^{\alpha+1} \prod_{c=1}^{C-1} (k_{c+1} - k_c - \widetilde{M})^{\alpha+1} \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\overline{\eta}_{c-1}|) \vee 1)^{-\alpha} \\ & \leq (C_\rho \vee 1)^C \left(\frac{N - C\widetilde{M}}{C} \right)^{(\alpha+1)C} \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\overline{\eta}_{c-1}|) \vee 1)^{-\alpha} \\ & \leq e^{\delta_2 N} \prod_{c=1}^C ((\ell_c - \ell_{c-1} - |\overline{\eta}_{c-1}|) \vee 1)^{-\alpha}, \end{aligned} \quad (3.138)$$

where δ_2 can be made arbitrarily small by choosing ε small and \widetilde{M} large (and we interpret $\ell_0 = 0$, $|\overline{\eta}_0| = 0$). Here, we have used Lemma 2.3 in the first inequality, as well as the fact that the product $\prod_{c=1}^{C-1} (k_{c+1} - k_c - \widetilde{M})$ is maximal when all factors are equal in the second inequality.

Combining (3.135–3.138), we see that

$$\begin{aligned} & \mathbb{P}(R_N \in \mathcal{O}(Q) | X) \\ & \leq e^{(\delta_1' + \delta_2' + \delta_1 + \varepsilon c'_\rho)N} \exp \left[N \sum_{r=1}^R W_Q(\widetilde{\mathcal{U}}_r) (H_{\tau|K}(Q_r) + \mathbb{E}_{Q_r}[\log \rho(\tau_1)]) \right] \\ & \quad \times \sum_{\substack{(\ell_i), (r_i), (\overline{\eta}_i) \\ \text{good}}}^C \prod_{i=1}^C ((\ell_i - \ell_{i-1} - |\overline{\eta}_{i-1}|) \vee 1)^{-\alpha}. \end{aligned} \quad (3.139)$$

We claim that X -a.s.

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \frac{1}{N} \log \sum_{\substack{(\ell_i), (r_i), (\overline{\eta}_i) \\ \text{good}}}^C \prod_{i=1}^C ((\ell_i - \ell_{i-1} - |\overline{\eta}_{i-1}|) \vee 1)^{-\alpha} \\ & \leq \delta_2 - \alpha \sum_{r=1}^R W_Q(\widetilde{\mathcal{U}}_r) m_{Q_r} H(\Psi_{Q_r} | \nu^{\otimes N}), \end{aligned} \quad (3.140)$$

where δ_2 can be made arbitrarily small by choosing ε small and L large. A proof of this is given below. Observe next that (3.139–3.140) (recall also (1.32)) yield that X -a.s. (with $\delta := \delta'_1 + \delta'_2 + \delta_1 + \delta_2 + \varepsilon c'_\rho$)

$$\begin{aligned}
& \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{O}(Q) | X) \\
& \leq \delta - \sum_{r=1}^R W_Q(\tilde{\mathcal{U}}_r) \left(H(Q_r | q_{\rho, \nu}^{\otimes N}) + (\alpha - 1) m_{Q_r} H(\Psi_{Q_r} | \nu^{\otimes N}) \right) \\
& \leq \delta + 2\alpha\varepsilon - \frac{1}{1 + \varepsilon} \int_{\mathcal{P}^{\text{erg}}(\tilde{E}^N)} H(Q' | q_{\rho, \nu}^{\otimes N}) + (\alpha - 1) m_{Q'} H(\Psi_{Q'} | \nu^{\otimes N}) W_Q(dQ') \\
& = -\frac{1}{1 + \varepsilon} I^{\text{fin}}(Q) + \delta + 2\alpha\varepsilon
\end{aligned} \tag{3.141}$$

(use (3.102) for the second inequality, and see (6.3) for the last equality), which completes the proof.

B. Coarse-graining X with R colours. It remains to verify (3.140), for which we employ a coarse-graining scheme similar to the one used in Section 3.4 (with block lengths $\lceil (1 - \varepsilon_2)L \rceil$, etc.) To ease notation, we silently replace L by $(1 - \varepsilon_2)L$ in the following. Split X into blocks of L consecutive letters, define a $\{0, 1\}$ -valued array $A_{i,r}$, $i \in \mathbb{N}$, $r \in \{1, \dots, R\}$ as in Section 3.4 inductively: For each r , put $A_{0,r} := 0$ and, given that $A_{0,r}, A_{1,r}, \dots, A_{l-1,r}$ have been assigned values, define A_l as follows:

(1) If $A_{l-1,r} = 0$, then

$$A_{l,r} := \begin{cases} 1, & \text{if in } X \text{ there is a word from } \kappa(\mathcal{A}_r) \text{ starting in } ((l-1)L, lL], \\ 0, & \text{otherwise.} \end{cases} \tag{3.142}$$

(2) If $A_{l-1,r} = 1$, then

$$A_l := \begin{cases} 1, & \text{if in } X \text{ there are two words from } \kappa(\mathcal{A}_r) \text{ starting in } ((l-2)L, (l-1)L], \\ & \text{respectively, } ((l-1)L, lL] \text{ and occurring disjointly,} \\ 0, & \text{otherwise.} \end{cases} \tag{3.143}$$

Put

$$p_r := L \exp \left(- (1 - 2\varepsilon_2) L H(\Psi_{Q_r} | \nu^{\otimes N}) \right). \tag{3.144}$$

Arguing as in Section 3.4, we can couple the $(A_{i,r})_{i \in \mathbb{N}, 1 \leq r \leq R}$ with an array $\bar{\omega} = (\omega_{i,r})_{i \in \mathbb{N}, 1 \leq r \leq R}$ such that $A_{i,r} \leq \omega_{i,r}$ and the sequence $((\omega_{i,1}, \dots, \omega_{i,R}))_{i \in \mathbb{N}}$ is i.i.d. with $\mathbb{P}(\omega_{i,r} = 1) = p_r$. In particular, for each r , $(\omega_{i,r})_{i \in \mathbb{N}}$ is a Bernoulli(p_r)-sequence. There may (and certainly will be if Ψ_{Q_r} and $\Psi_{Q_{r'}}$ are similar) an arbitrary dependence between the $\omega_{i,1}, \dots, \omega_{i,R}$ for fixed i , but this will be harmless in the low-density limit we are interested in.

For $r \in \{1, \dots, R\}$, put $d_r := W_Q(\tilde{\mathcal{U}}_r)(1 - 6\varepsilon_2)$, $D_r := \lceil (1 - \varepsilon_2) \widetilde{M} m_{Q_r} / L \rceil$. If $\eta_c \in V_{r_c, \widetilde{M}}$, then

$$|\kappa(\eta_c)| \in \widetilde{M} m_{Q_{r_c}} (1 - \varepsilon_2, 1 + \varepsilon_2), \tag{3.145}$$

so $\kappa(\eta_c)$ covers at least D_{r_c} consecutive L -blocks of the coarse-graining. Furthermore, as η_c in turn contains at least $D_{r_c}(1 - 3\varepsilon_2)$ disjoint subsentences from \mathcal{A}_{r_c} , we see that at least $D_{r_c}(1 - 3\varepsilon_2)$ of

these blocks must have $A_{k,r_c} = 1$. Thus, for fixed X , we read off from each good triple $(\ell_c), (r_c), (\bar{\eta}_c)$ numbers $m_1 < \dots < m_C$ such that

$$\begin{aligned} m_{c+1} &\geq m_c + D_{r_c}, \quad c = 1, \dots, C-1, \\ |\{m_c \leq k < m_c + D_{r_c} : A_{k,r_c} = 1\}| &\geq D_{r_c}(1 - 3\varepsilon_2), \quad c = 1, \dots, C, \\ |\{1 \leq c \leq C : r_c = r\}| &\geq d_r C, \quad r = 1, \dots, R. \end{aligned} \quad (3.146)$$

where m_c is the index of the L -block that contains ℓ_c . Furthermore, note that for a given ‘‘coarse-graining’’ (m_c) and (r_c) satisfying (3.146), there are at most

$$L^C \left(2\varepsilon_2 \widetilde{M} \max_{r=1, \dots, R} m_{Q_r} \right)^C \leq \exp(\delta_3 N) \quad (3.147)$$

choices for ℓ_c and $\bar{\eta}_c$ that lead to a good triple $(\ell_c), (r_c), (\bar{\eta}_c)$ with this particular coarse-graining. Indeed, for each $c = 1, \dots, C$ there are at most L choices for ℓ_c and, since each $\eta \in V_{r_c, \widetilde{M}}$ satisfies

$$|\kappa(\eta)| \in \widetilde{M} m_{Q_{r_c}} (1 - \varepsilon_2, 1 + \varepsilon_2), \quad (3.148)$$

there are at most $2\varepsilon_2 \widetilde{M} m_{Q_{r_c}}$ choices for $\bar{\eta}_c$ (note that once ℓ_c is fixed as a ‘‘starting point’’ for a word on X , choosing $\bar{\eta}_c$ in fact amounts to choosing an ‘‘endpoint’’). Note that δ_3 can be made arbitrarily small by choosing ε small and \widetilde{M} large. Finally, (3.147) and Lemma 3.3 yield (3.140). Indeed, since

$$\begin{aligned} \limsup_{N \rightarrow \infty} \frac{C}{N} &\leq \frac{1}{\widetilde{M}}, \\ \sum_{r=1}^R d_r D_r \log p_r &\leq -(1 - 8\varepsilon_2) \sum_{r=1}^R W_Q(\widetilde{U}_r) \frac{\widetilde{M} m_{Q_r}}{L} \left(LH(\Psi_{Q_r} | \nu^{\otimes N}) - \log L \right) \\ &\leq -\widetilde{M} \sum_{r=1}^R W_Q(\widetilde{U}_r) m_{Q_r} H(\Psi_{Q_r} | \nu^{\otimes N}) + \left(8\varepsilon_2 m^* K_1 + \frac{\log L}{L} \right) \widetilde{M}, \end{aligned} \quad (3.149)$$

by choosing ε small (note that $\varepsilon_2 m^* K_1 \leq \varepsilon$), L and \widetilde{M} large, and γ sufficiently close to $1/\alpha$, the right-hand side of (3.154) is smaller than the right-hand side of (3.140).

3.5.6 A multicolour version of the core lemma

The following is an extension of Lemma 2.1. Let $R \in \mathbb{N}$, $\bar{\omega}_i = (\omega_{i,1}, \dots, \omega_{i,R}) \in \{0, 1\}^R$, and assume that $(\bar{\omega}_i)_{i \in \mathbb{N}}$ is i.i.d. with

$$\mathbb{P}(\omega_{i,r} = 1) = p_r, \quad i \in \mathbb{N}, \quad r = 1, \dots, R. \quad (3.151)$$

Note that there may be an arbitrary dependence between the $\omega_{i,r}$'s for fixed i . This will be harmless in the limit we are interested in below.

Lemma 3.3. *Let $\alpha \in (1, \infty)$, $\varepsilon > 0$, $(d_1, \dots, d_R) \in [0, 1]^R$ with $\sum_{r=1}^R d_r \leq 1$, $D_1, \dots, D_R \in \mathbb{N}$, $C \in \mathbb{N}$, put*

$$S_C(\bar{\omega}) := \sum_{\substack{m_1, \dots, m_C \\ r_1, \dots, r_C}}^* \prod_{i=1}^C (m_i - m_{i-1} - D_{r_{i-1}})^{-\alpha}, \quad (3.152)$$

where the sum \sum^* extends over all pairs of C -tuples $m_0 := 0 < m_1 < \dots < m_C$ from \mathbb{N}^C and $(r_1, \dots, r_C) \in \{1, \dots, R\}^C$ satisfying the constraints

$$\begin{aligned} m_{i+1} &\geq m_i + D_{r_i}, \\ |\{1 \leq i \leq C : r_i = r\}| &\geq d_r C, \quad r = 1, \dots, R, \\ |\{m_i \leq k < m_i + D_{r_i} : \omega_{k,r_i} = 1\}| &\geq D_{r_i} (1 - \varepsilon), \quad i = 1, \dots, C. \end{aligned} \quad (3.153)$$

Then $\bar{\omega}$ -a.s.

$$\begin{aligned} & \limsup_{C \rightarrow \infty} \frac{1}{C} \log S_C(\bar{\omega}) \\ & \leq \inf_{\gamma \in (1/a, 1)} \left\{ \frac{1}{\gamma} \left(\log \zeta(a\gamma) + h(\underline{d}) + d_0 \log R + (\log 2) \sum_{r=1}^R d_r D_r + (1 - \varepsilon) \sum_{r=1}^R d_r D_r \log p_r \right) \right\}, \end{aligned} \quad (3.154)$$

where $h(\underline{d}) := -\sum_{r=0}^R d_r \log d_r$ (with $d_0 := 1 - d_1 - \dots - d_R$) is the entropy of \underline{d} .

Proof. The proof is a variation on the proof of Lemma 2.1. We again estimate fractional moments. For $\gamma \in (1/a, 1)$, we have

$$\begin{aligned} & \mathbb{E}[(S_C)^\gamma] \\ & \leq \sum'_{r_1, \dots, r_C} \sum_{\substack{m_1, \dots, m_C \\ m_{i+1} \geq m_i + D_{r_i} \forall i}} \mathbb{P} \left(\bigcap_{i=1}^C \left\{ |\{k \in [m_i, m_i + D_{r_i} - 1] : \omega_{k, r_i} = 1\}| \geq (1 - \varepsilon) D_{r_i} \right\} \right) \\ & \quad \times \prod_{i=1}^C (m_i - m_{i-1} - D_{r_{i-1}})^{-\alpha\gamma}, \end{aligned} \quad (3.155)$$

where the sum \sum' extends over all (r_1, \dots, r_C) satisfying the constraint in the second line of (3.153). Noting that

$$\begin{aligned} \mathbb{P} \left(|\{k \in [m_i, m_i + D_{r_i} - 1] : \omega_{k, r_i} = 1\}| \geq (1 - \varepsilon) D_{r_i} \right) &= \sum_{m=(1-\varepsilon)D_{r_i}}^{D_{r_i}} \binom{D_{r_i}}{m} p_r^m (1 - p_r)^{D_{r_i} - m} \\ &\leq p_r^{(1-\varepsilon)D_{r_i}} 2^{D_{r_i}} \end{aligned}$$

and

$$\begin{aligned} & |\{(r_1, \dots, r_C) \in \{1, \dots, R\}^C : \text{at least } d_r C \text{ of the } r_i = r, r = 1, \dots, R\}| \\ & \leq R^{d_0 C} \binom{C}{d_0 C \ d_1 C \ \dots \ d_R C} = \exp \left[C(d_0 \log R + h(\underline{d}) + o(1)) \right], \end{aligned}$$

we see from (3.155) that

$$\begin{aligned} \mathbb{E}[(S_C)^\gamma] &\leq \exp \left[C(d_0 \log R + h(\underline{d}) + o(1)) \right] \times \prod_{r=1}^R (2p_r^{(1-\varepsilon)})^{d_r C D_r} \\ &\quad \times \sum_{\substack{m_1, \dots, m_C \\ m_{i+1} \geq m_i + D_{r_i} \forall i}} \prod_{i=1}^C (m_i - m_{i-1} - D_{r_{i-1}})^{-\alpha\gamma} \\ &= \exp C \left[d_0 \log R + h(\underline{d}) + \log \zeta(a\gamma) + \sum_{r=1}^R d_r D_r \log 2 + (1 - \varepsilon) \sum_{r=1}^R d_r D_r \log p_r \right], \end{aligned} \quad (3.156)$$

which yields (3.154) as in the proof of Lemma 2.1. \square

3.6 Step 6: Weakening the tail assumption

We finally show how to go from (3.3) to (1.1). Suppose that ρ satisfies (1.1) with a certain $\alpha \in (1, \infty)$. Then, for any $\alpha' \in (1, \alpha)$, there is a $C_\rho(\alpha')$ such that (3.3) holds for this α' . Hence,

as shown in Sections 3.1–3.4, for any $\varepsilon > 0$ we can find a neighbourhood $\mathcal{O}(Q) \subset \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ of Q such that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{O}(Q) \mid X) \leq -H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) - (\alpha' - 1) m_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}}) + \frac{\varepsilon}{2} \quad X - a.s. \quad (3.157)$$

The right-hand side is $\leq -I^{\text{fin}}(Q) + \varepsilon$ for α' sufficiently close to α , so that we again get (3.1). \square

4 Lower bound

The following lower bound will be used in Section 5 to derive the lower bound in the definition of the LDP.

Proposition 4.1. *For any $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ and any open neighbourhood $\mathcal{U}(Q) \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ of Q ,*

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{U}(Q) \mid X) \geq -I^{\text{fin}}(Q) \quad X - a.s. \quad (4.1)$$

Proof. Suppose first that $Q \in \mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$. Then, informally, our strategy runs as follows. In X , look for the first string of length $\approx Nm_Q$ that looks typical for Ψ_Q . Make the first jump long enough so as to land at the start of this string. Make the remaining $N - 1$ jumps typical for Q . The probability of this strategy on the exponential scale is the conditional specific relative entropy of word lengths under Q w.r.t. $\rho^{\otimes \mathbb{N}}$ given the concatenation, i.e., $\approx \exp[N(H_{\tau|K}(Q) + \mathbb{E}_Q[\log \rho(\tau_1)])]$, times the probability of the first long jump. In order to find a suitable string, we have to skip ahead in X a distance $\approx \exp[Nm_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}})]$. By (1.1), the probability of the first jump is therefore $\approx \exp[-N\alpha m_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}})]$. In view of (1.16) and (1.32), this yields the claim. In the actual proof, it turns out to be technically simpler to employ a slightly different strategy, which has the same asymptotic cost, where we look not only for one contiguous piece of “ Ψ_Q -typical” letters but for a sequence of $\lceil N/M \rceil$ pieces, each of length $\approx Mm_Q$. Then we let $N \rightarrow \infty$, followed by $M \rightarrow \infty$.

More formally, we choose for $\mathcal{O}(Q)$ an open neighborhood $\mathcal{O}' \subset \mathcal{O}$ of the type introduced in Section 3.2, and we estimate $\mathbb{P}(R_N \in \mathcal{O}' \mid X)$ from below by using (3.17–3.20).

Assume first that Q is ergodic. We can then assume that the neighbourhood \mathcal{U} is given by

$$\mathcal{U} = \{Q' \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : (\pi_{L_u} Q')(\zeta_u) \in (a_u, b_u), \quad u = 1, \dots, U\} \quad (4.2)$$

for some $U \in \mathbb{N}$, $L_1, \dots, L_U \in \mathbb{N}$, $0 \leq a_u < b_u \leq 1$ and $\zeta_u \in \tilde{E}^{L_u}$, $u = 1, \dots, U$. As in Section 3.1, by ergodicity of Q we can find for each $\varepsilon > 0$ a sufficiently large $M \in \mathbb{N}$ and a set $\mathcal{A} = \{z_1, \dots, z_A\} \subset \tilde{E}^M$ of “ Q -typical sentences” satisfying (3.6–3.7) (with $\varepsilon_1 = \delta_1 = \varepsilon$, say), and additionally

$$\frac{1}{M} \left| \{0 \leq j \leq M - L_i : \pi_{L_u}(\tilde{\theta}^j z_a) = \zeta_u\} \right| \in (a_u, b_u), \quad a = 1, \dots, A, \quad u = 1, \dots, U. \quad (4.3)$$

Let $\mathcal{B} := \kappa(\mathcal{A})$. Then from (3.6–3.7) we have that, for each $b \in \mathcal{B}$,

$$|I_b| = |\{z \in \mathcal{A} : \kappa(z) = b\}| \geq \exp[M(H_{\tau|K}(Q) - 2\varepsilon)], \quad (4.4)$$

and

$$\mathbb{P}(X \text{ begins with some element of } \mathcal{B}) \geq \exp[-Mm_Q(H(\Psi_Q \mid \nu^{\otimes \mathbb{N}}) + 2\varepsilon)]. \quad (4.5)$$

Let

$$\begin{aligned} \sigma_1^{(M)} &:= \min\{i : \theta^i X \text{ begins with some element of } \mathcal{B}\}, \\ \sigma_l^{(M)} &:= \min\{i > \sigma_{l-1}^{(M)} + M(m_Q + \varepsilon) : \theta^i X \text{ begins with some element of } \mathcal{B}\}, \quad l = 2, 3, \dots \end{aligned} \quad (4.6)$$

Restricting the sum in (3.23) over $0 < j_1 < \dots < j_N < \infty$ such that $j_1 = \sigma_1^{(M)}$, $j_2 - j_1, \dots, j_M - j_{M-1}$ are the word lengths corresponding to the z_a 's compatible with $\pi_{Mm_Q}(\theta^{\tau_1} X)$, $j_{M+1} = \sigma_2^{(M)}$, etc., we see that

$$\frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{U} \mid X) \geq H_{\tau|K}(Q) + \mathbb{E}_Q[\log \rho(\tau_1)] - 3\varepsilon - \alpha \frac{1}{N} \sum_{l=1}^{\lfloor N/M \rfloor} \log(\sigma_l^{(M)} - \sigma_{l-1}^{(M)}) \quad (4.7)$$

for N sufficiently large. Hence X -a.s.

$$\begin{aligned} \liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{U} \mid X) &\geq H_{\tau|K}(Q) + \mathbb{E}_Q[\log \rho(\tau_1)] - 3\varepsilon - \alpha \frac{1}{M} \mathbb{E}[\log \sigma_1^{(M)}] \\ &\geq H_{\tau|K}(Q) + \mathbb{E}_Q[\log \rho(\tau_1)] - \alpha m_Q (H(\Psi_Q \mid \nu^{\otimes \mathbb{N}}) - 6\varepsilon) \\ &= -I^{\text{fin}}(Q) - 6\varepsilon, \end{aligned} \quad (4.8)$$

where we have used (4.5) in the second inequality. Now let $\varepsilon \downarrow 0$.

It remains to remove the restriction of ergodicity of Q , analogously to the proof of Birkner [2], Proposition 2. To that end, assume that $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ admits a non-trivial ergodic decomposition. Then, for each $\varepsilon > 0$, we can find $Q_1, \dots, Q_R \in \mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$, $\lambda_1, \dots, \lambda_R \in (0, 1)$, $\sum_{r=1}^R \lambda_r = 1$ such that $\lambda_1 Q_1 + \dots + \lambda_R Q_R \in \mathcal{U}$ and

$$\sum_{i=1}^R \lambda_r I^{\text{fin}}(Q_r) \leq I^{\text{fin}}(Q) + \varepsilon \quad (4.9)$$

(for details see Birkner [2], p. 723; employ the fact that both terms in I^{fin} are affine). For each $r = 1, \dots, R$, pick a small neighbourhood \mathcal{U}_r of Q_r such that

$$Q'_r \in \mathcal{U}_r, r = 1, \dots, R \implies \sum_{i=1}^R \lambda_r Q'_r \in \mathcal{U}. \quad (4.10)$$

Using the above strategy for Q_1 for $\lambda_1 N$ loops, then the strategy for Q_2 for $\lambda_2 N$ loops, etc., we see that

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{U} \mid X) \geq - \sum_{i=1}^R \lambda_r I^{\text{fin}}(Q_r) - 6\varepsilon \geq -I^{\text{fin}}(Q) - 7\varepsilon. \quad (4.11)$$

□

5 Proof of Theorem 1.2

Proof. The proof comes in 3 steps. We first prove that, for each word length truncation level $\text{tr} \in \mathbb{N}$, the family $\mathbb{P}([R_N]_{\text{tr}} \in \cdot \mid X)$, $N \in \mathbb{N}$, X -a.s. satisfies an LDP on

$$\mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) = \{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : Q(|Y^{(1)}| \leq \text{tr}) = 1\} \quad (5.1)$$

(recall (1.11–1.13)) with a deterministic rate function $I^{\text{fin}}([Q]_{\text{tr}})$ (this is essentially the content of Propositions 4.1 and 3.1). Note that $[Q]_{\text{tr}} = Q$ for $Q \in \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, and that $\mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ is a closed subset of $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, in particular, a Polish space under the relative topology (which is again the weak topology). After we have given the proof for fixed tr , we let $\text{tr} \rightarrow \infty$ and use a projective limit argument to complete the proof of Theorem 1.2.

1. Fix a truncation level $\text{tr} \in \mathbb{N}$. Propositions 4.1 and 3.1 combine to yield the LDP on $\mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ in the following standard manner. Note that any $Q \in \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ satisfies $m_Q < \infty$.

1a. Let $\mathcal{O} \subset \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ be *open*. Then, for any $Q \in \mathcal{O}$, there is an open neighbourhood $\mathcal{O}(Q) \subset \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ of Q such that $\mathcal{O}(Q) \subset \mathcal{O}$. The latter inclusion, together with Proposition 4.1, yields

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}([R_N]_{\text{tr}} \in \mathcal{O} \mid X) \geq -I^{\text{fin}}(Q) \quad X - a.s. \quad (5.2)$$

Optimising over $Q \in \mathcal{O}$, we get

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}([R_N]_{\text{tr}} \in \mathcal{O} \mid X) \geq - \inf_{Q \in \mathcal{O}} I^{\text{fin}}(Q) \quad X - a.s. \quad (5.3)$$

Here, note that, since $\mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ is Polish, it suffices to optimise over a countable set generating the weak topology, allowing us to transfer the X -a.s. limit from points to sets (see, e.g., Comets [4], Section III).

1b. Let $\mathcal{K} \subset \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ be *compact*. Then there exist $M \in \mathbb{N}$, $Q_1, \dots, Q_M \in \mathcal{K}$ and open neighbourhoods $\mathcal{O}(Q_1), \dots, \mathcal{O}(Q_M) \subset \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ such that $\mathcal{K} \subset \cup_{m=1}^M \mathcal{O}(Q_m)$. The latter inclusion, together with Proposition 3.1, yields

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}([R_N]_{\text{tr}} \in \mathcal{K} \mid X) \leq - \inf_{1 \leq m \leq M} I^{\text{fin}}(Q_m) + \varepsilon \quad X - a.s. \quad \forall \varepsilon > 0. \quad (5.4)$$

Extending the infimum to $Q \in \mathcal{K}$ and letting $\varepsilon \downarrow 0$ afterwards, we obtain

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}([R_N]_{\text{tr}} \in \mathcal{K} \mid X) \leq - \inf_{Q \in \mathcal{K}} I^{\text{fin}}(Q) \quad X - a.s. \quad (5.5)$$

1c. Let $\mathcal{C} \subset \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ be *closed*. Because $Q \mapsto H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}})$ has compact level sets, for any $M < \infty$ the set $\mathcal{K}_M = \mathcal{C} \cap \{Q \in \mathcal{P}_{\text{tr}}^{\text{inv}}(\tilde{E}^{\mathbb{N}}) : H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) \leq M\}$ is compact. Hence, doing *annealing* on X and using (5.5), we get

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}([R_N]_{\text{tr}} \in \mathcal{C} \mid X) \leq \max \left\{ -M, - \inf_{Q \in \mathcal{K}_M} I^{\text{fin}}(Q) \right\} \quad X - a.s. \quad (5.6)$$

Extending the infimum to $Q \in \mathcal{C}$ and letting $M \rightarrow \infty$ afterwards, we arrive at

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}([R_N]_{\text{tr}} \in \mathcal{C} \mid X) \leq - \inf_{Q \in \mathcal{C}} I^{\text{fin}}(Q) \quad X - a.s. \quad (5.7)$$

Equations (5.3) and (5.7) complete the proof of the conditional LDP for $[R_N]_{\text{tr}}$.

2. It remains to remove the truncation of word lengths. We know from Step 1 that, for every $\text{tr} \in \mathbb{N}$, the family $\mathbb{P}([R_N]_{\text{tr}} \in \cdot \mid X)$, $N \in \mathbb{N}$, satisfies the LDP on $\mathcal{P}^{\text{inv}}([\tilde{E}]_{\text{tr}}^{\mathbb{N}})$ with rate function I^{fin} . Consequently, by the Dawson-Gärtner projective limit theorem (see Dembo and Zeitouni [5], Theorem 4.6.1), the family $\mathbb{P}(R_N \in \cdot \mid X)$, $N \in \mathbb{N}$, satisfies the LDP on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ with rate function

$$I^{\text{que}}(Q) = \sup_{\text{tr} \in \mathbb{N}} I^{\text{fin}}([Q]_{\text{tr}}), \quad Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}). \quad (5.8)$$

The sup may be replaced by a limsup because the truncation may start at any level. For $Q \in \mathcal{P}^{\text{inv, fin}}(\tilde{E}^{\mathbb{N}})$, we have $\lim_{\text{tr} \rightarrow \infty} I^{\text{fin}}([Q]_{\text{tr}}) = I^{\text{fin}}(Q)$ by Lemma A.1, and so we get the claim if we can show that limsup can be replaced by a limit, which is done in Step 3. Note that I^{que} inherits from I^{fin} the properties qualifying it to be a rate function: this is part of the projective limit theorem. For I^{fin} these properties are proved in Section 6.

3. Since I^{que} is lower semi-continuous, it is equal to its lower semi-continuous regularisation

$$\tilde{I}^{\text{que}}(Q) := \sup_{\mathcal{O}(Q)} \inf_{Q' \in \mathcal{O}(Q)} I^{\text{que}}(Q'), \quad (5.9)$$

where the supremum runs over the open neighborhoods of Q . For each $\text{tr} \in \mathbb{N}$, $[Q]_{\text{tr}} \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$, while $w - \lim_{\text{tr} \rightarrow \infty} [Q]_{\text{tr}} = Q$. So, in particular,

$$I^{\text{que}}(Q) = \tilde{I}^{\text{que}}(Q) \leq \sup_n \inf_{\text{tr} \geq n} I^{\text{fin}}([Q]_{\text{tr}}) = \liminf_{\text{tr} \rightarrow \infty} I^{\text{fin}}([Q]_{\text{tr}}), \quad (5.10)$$

implying that in fact

$$I^{\text{que}}(Q) = \lim_{\text{tr} \rightarrow \infty} I^{\text{fin}}([Q]_{\text{tr}}), \quad Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}). \quad (5.11)$$

□

Lemma A.1 in Appendix A, together with (5.11), shows that $I^{\text{que}}(Q) = I^{\text{fin}}(Q)$ for $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$, as claimed in the first line of (1.15).

6 Proof of Theorem 1.3

Proof. The proof comes in 5 steps.

1. Every $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ can be decomposed as

$$Q = \int_{\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})} Q' W_Q(dQ') \quad (6.1)$$

for some unique probability measure W_Q on $\mathcal{P}^{\text{erg}}(\tilde{E}^{\mathbb{N}})$ (Georgii [7], Proposition 7.22). If $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$, then W_Q is concentrated on $\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})$ and so, by (1.9–1.10),

$$m_Q = \int_{\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})} m_{Q'} W_Q(dQ'), \quad \Psi_Q = \int_{\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})} \frac{m_{Q'}}{m_Q} \Psi_{Q'} W_Q(dQ'). \quad (6.2)$$

Since $Q \mapsto H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}})$ and $\Psi \mapsto H(\Psi \mid \nu^{\otimes \mathbb{N}})$ are affine (see e.g. Deuschel and Stroock [6], Example 4.4.41), it follows from (1.16) and (6.1–6.2) that

$$I^{\text{fin}}(Q) = \int_{\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})} I^{\text{fin}}(Q') W_Q(dQ'). \quad (6.3)$$

Since $Q \mapsto W_Q$ is affine, (6.3) shows that I^{fin} is affine on $\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$.

2. Let $(Q_n)_{n \in \mathbb{N}} \subset \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ be such that $w - \lim_{n \rightarrow \infty} Q_n = Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$. By Proposition 3.1, for any $\varepsilon > 0$ we can find an open neighbourhood $\mathcal{O}(Q) \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ of Q such that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{O}(Q) \mid X) \leq -I^{\text{fin}}(Q) + \varepsilon \quad X - a.s. \quad (6.4)$$

On the other hand, for n large enough so that $Q_n \in \mathcal{O}(Q)$, we have from Proposition 4.1 that

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{O}(Q) \mid X) \geq -I^{\text{fin}}(Q_n) \quad X - a.s. \quad (6.5)$$

Combining (6.4–6.5), we get that, for any $\varepsilon > 0$,

$$\liminf_{n \rightarrow \infty} I^{\text{fin}}(Q_n) \geq I^{\text{fin}}(Q) - \varepsilon. \quad (6.6)$$

Now let $\varepsilon \downarrow 0$, to conclude that I^{fin} is lower semicontinuous on $\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ (recall also (5.11)).

3. From (1.16) we have

$$I^{\text{fin}}(Q) \geq H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) \quad \forall Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}}) \quad (6.7)$$

Since $\{Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}): H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) \leq C\}$ is compact for all $C < \infty$ (see, e.g., Dembo and Zeitouni [5], Corollary 6.5.15), it follows that I^{fin} has compact level sets on $\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$.

4. As mentioned at the end of Section 5, I^{que} inherits from I^{fin} that it is lower semicontinuous and has compact level sets. In particular, I^{que} is the lower semicontinuous extension of I^{fin} from $\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$. Moreover, since I^{fin} is affine on $\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ and I^{que} arises as the truncation limit of I^{fin} (recall (5.10)), it follows that I^{que} is affine on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$.

5. It is immediate from (1.15–1.16) that $q_{\rho, \nu}^{\otimes \mathbb{N}}$ is the unique zero of I^{que} . □

7 Proof of Theorem 1.4

Proof. The extension is an easy generalisation of the proof given in Sections 3–4.

(a) Assume that ρ satisfies (1.1) with $\alpha = 1$. Since the LDP upper bound holds by the *annealed* LDP (compare (1.8) and (1.16)), it suffices to prove the LDP lower bound. To achieve this, we first show that for any $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ and $\varepsilon > 0$ there exists an open neighbourhood $\mathcal{O}(Q) \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ of Q such that

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{O}(Q) | X) \geq -I^{\text{ann}}(Q) - \varepsilon \quad X\text{-a.s.} \quad (7.1)$$

After that, the extension from $\mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ follows the argument in Section 5.

In order to verify (7.1), observe that, by our assumption on $\rho(\cdot)$, for any $\alpha' > 1$ there exists a $C_{\alpha'} > 0$ such that

$$\frac{\rho(n)}{n^{\alpha'}} \geq C_{\alpha'} \quad \forall n \in \text{supp}(\rho). \quad (7.2)$$

Picking α' so close to 1 that $(\alpha' - 1)m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) < \varepsilon/2$, we can trace through the proof of Proposition 4.1 in Section 4 to construct an open neighbourhood $\mathcal{O}(Q) \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ of Q satisfying

$$\begin{aligned} \liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(R_N \in \mathcal{O}(Q) | X) \\ \geq -H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) - (\alpha' - 1)m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}) - \varepsilon/2 \geq -I^{\text{ann}}(Q) - \varepsilon \quad X\text{-a.s.}, \end{aligned} \quad (7.3)$$

which is (7.1).

(b) We only give a sketch of the argument. Assume $\alpha = \infty$ in (1.1). For $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$, the lower bound (which is non-zero only when $Q \in \mathcal{R}_{\nu}$) follows from Birkner [2], Proposition 2, or can alternatively be obtained from the argument in Section 4. Now consider a $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ with $m_Q = \infty$, $H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) < \infty$ and $\lim_{\text{tr} \rightarrow \infty} m_{[Q]_{\text{tr}}} H(\Psi_{[Q]_{\text{tr}}} | \nu^{\otimes \mathbb{N}}) = 0$, let $\mathcal{O}(Q) \subset \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ be an open neighbourhood of Q . For simplicity, we assume $\text{supp}(\rho) = \mathbb{N}$. Fix $\varepsilon > 0$. We can find a sequence $\delta_N \downarrow 0$ such that

$$\max \left\{ -\frac{1}{N} \log \rho(n) : n \leq \lceil N\delta_N \rceil \right\} \leq \varepsilon. \quad (7.4)$$

Furthermore,

$$\frac{1}{N} h \left(Q|_{\mathcal{F}_N} | q_{\rho, \nu}^{\otimes N} \right) \geq H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) - \varepsilon \quad (7.5)$$

for $N \geq N_0 = N_0(\varepsilon, Q)$, and we can find $\text{tr}_0 \in \mathbb{N}$ such that

$$\frac{1}{N_0} h \left(([Q]_{\text{tr}})|_{\mathcal{F}_{N_0}} | q_{\rho, \nu}^{\otimes N_0} \right) \geq \frac{1}{N_0} h \left(Q|_{\mathcal{F}_{N_0}} | q_{\rho, \nu}^{\otimes N_0} \right) - \varepsilon \quad (7.6)$$

for $\text{tr} \geq \text{tr}_0$. Hence

$$H([Q]_{\text{tr}} | q_{\rho, \nu}^{\otimes \mathbb{N}}) \geq H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) - 2\varepsilon \quad \text{for } \text{tr} \geq \text{tr}_0. \quad (7.7)$$

We may also assume that $[Q]_{\text{tr}} \in \mathcal{O}(Q)$ for $\text{tr} \geq \text{tr}_0$. For a given $N \geq N_0$, pick $\text{tr}(N) \geq \text{tr}_0$ so large that $m_{[Q]_{\text{tr}(N)}} H(\Psi_{[Q]_{\text{tr}(N)}} | \nu^{\otimes \mathbb{N}}) \leq \delta_N/2$. Using the strategy described at the beginning of Section 4, we can construct a neighbourhood $\mathcal{O}_N \subset \mathcal{O}(Q)$ of $[Q]_{\text{tr}(N)}$ such that the conditional probability $\mathbb{P}(R_N \in \mathcal{O}_N | X)$ is bounded below by

$$\exp[-N(H([Q]_{\text{tr}} | q_{\rho, \nu}^{\otimes \mathbb{N}}) - \varepsilon)] \times \text{the cost of the first jump}, \quad (7.8)$$

where the first jump takes us to a region of size $\approx Nm_{[Q]_{\text{tr}(N)}}$ on which the medium looks “ $\Psi_{[Q]_{\text{tr}(N)}}$ -typical”. Since, in a typical medium, the size of the first jump will be

$$\approx \exp[Nm_{[Q]_{\text{tr}(N)}} H(\Psi_{[Q]_{\text{tr}(N)}} | \nu^{\otimes \mathbb{N}})] \leq \exp[N\delta_N], \quad (7.9)$$

we obtain from (7.4) and (7.7–7.9) that

$$\mathbb{P}(R_N \in \mathcal{O}(Q) | X) \geq \exp[-N(H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) + 4\varepsilon)] \quad (7.10)$$

for N large enough.

For the upper bound we can argue as follows: For $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ put

$$r(Q) := \limsup_{\text{tr} \rightarrow \infty} m_{[Q]_{\text{tr}(N)}} H(\Psi_{[Q]_{\text{tr}(N)}} | \nu^{\otimes \mathbb{N}}). \quad (7.11)$$

Since ρ satisfies the bound (3.3) for any $\alpha > 1$, we obtain from the upper bound in Theorem 1.2 that the rate function at Q is at least

$$\limsup_{\text{tr} \rightarrow \infty} I^{\text{fin}}([Q]_{\text{tr}}) = H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) + (\alpha - 1)r(Q), \quad (7.12)$$

hence equals ∞ if $r(Q) > 0$. On the other hand, if $r(Q) = 0$, then this is simply the annealed bound. \square

8 Proof of Corollary 1.6

Proof. Let E be a Polish space with metric d_E (equipped with its Borel- σ -algebra \mathcal{B}_E). We can choose a sequence of nested finite partitions $\mathcal{A}_c = \{A_{c,1}, \dots, A_{c,n_c}\}$, $c \in \mathbb{N}$, of E with the property that

$$\forall x \in E : \lim_{c \rightarrow \infty} \text{diam}(\langle x \rangle_c) = 0, \quad (8.1)$$

where the *coarse-graining map* $\langle \cdot \rangle_c$ maps an element of E to the element of \mathcal{A}_c it is contained in. Each $\mathcal{A}_c = \langle E \rangle_c$ is a finite set, which we equip with the discrete metric d_c . Extend $\langle \cdot \rangle_c$ to $\langle E \rangle_{c'}$ for each $c' > c$ via $\langle A_{c',i'} \rangle_c = A_{c,i}$ if $A_{c',i'} \subset A_{c,i}$. Then the collection $\mathcal{A}_c, \langle \cdot \rangle_c, c \in \mathbb{N}$, forms a projective family, and the projective limit

$$F = \{(\xi_1, \xi_2, \dots) : \xi_c \in \mathcal{A}_c, \langle \xi_{c'} \rangle_c = \xi_c, 1 \leq c < c'\} \quad (8.2)$$

is again a Polish space with the metric

$$d_F((\xi_1, \xi_2, \dots), (\eta_1, \eta_2, \dots)) := \sum_{c=1}^{\infty} 2^{-c} d_c(\xi_c, \eta_c). \quad (8.3)$$

We equip F with its Borel- σ -algebra \mathcal{B}_F . We can identify E with a subset of F via $\iota : x \mapsto (\langle x \rangle_c)_{c \in \mathbb{N}}$, since ι is injective by (8.1). Note that $\iota(E)$ is a measurable subset of F (in general $\iota(E) \neq F$; it

is easy to see that $\iota(E)$ is a closed subset of F when E is compact; for non-compact E use the one-point compactification of E).

Note that the topology generated by d_F on $\iota(E)$ is finer than the original topology generated by d_E : By (8.1), for each $x \in E$ and $\varepsilon > 0$, there is an $\varepsilon' > 0$ such that the d_F -ball of radius ε' around x is contained in the d -ball of radius ε around x . We will make use of the fact that

$$\text{the trace of } \mathcal{B}_F \text{ on } \iota(E) \text{ agrees with the image of } \mathcal{B}_E \text{ under } \iota. \quad (8.4)$$

To check this, note that for any $x \in E$, the function

$$F \ni \xi \mapsto \begin{cases} d_E(\iota^{-1}(\xi), x), & \xi \in \iota(E), \\ \infty, & \text{otherwise,} \end{cases} \quad (8.5)$$

can be pointwise approximated by functions that are constant on $\iota(A_{c,i})$, $i = 1, \dots, n_c$, and is therefore \mathcal{B}_F -measurable.

We extend $\langle \cdot \rangle_c$ in the obvious way to E^N and \tilde{E}^N , $N \in \mathbb{N} \cup \{\infty\}$ (via coordinate-wise coarse-graining), and then to $\mathcal{P}(E^N)$, $\mathcal{P}(\tilde{E}^N)$, $N \in \mathbb{N}$, and finally to $\mathcal{P}^{\text{inv}}(E^{\mathbb{N}})$ and $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ (by taking image measures). Note that $\langle \cdot \rangle_c$ and $[\cdot]_{\text{tr}}$ commute, and

$$m_Q = m_{\langle Q \rangle_c}, \quad \langle \Psi_Q \rangle_c = \Psi_{\langle Q \rangle_c}, \quad Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}}). \quad (8.6)$$

By Theorem 1.2, for each $c \in \mathbb{N}$ the family

$$\mathbb{P}(\langle R_N \rangle_c \in \cdot \mid X), \quad N \in \mathbb{N}, \quad (8.7)$$

X -a.s. satisfies the LDP with deterministic rate function

$$I_c^{\text{que}}(Q) = \begin{cases} I_c^{\text{fin}}(Q) := H(Q \mid \langle q_{\rho, \nu}^{\otimes \mathbb{N}} \rangle_c) + (\alpha - 1)m_Q H(\Psi_Q \mid \langle \nu^{\otimes \mathbb{N}} \rangle_c), & Q \in \mathcal{P}^{\text{inv, fin}}(\langle \tilde{E} \rangle_c^{\mathbb{N}}), \\ \lim_{\text{tr} \rightarrow \infty} I_c^{\text{fin}}([Q]_{\text{tr}}), & \text{if } m_Q = \infty. \end{cases} \quad (8.8)$$

Hence, by the Dawson-Gärtner *projective limit theorem* (see Dembo and Zeitouni [5], Theorem 4.6.1), the family $\mathbb{P}(R_N \in \cdot \mid X)$, $N \in \mathbb{N}$, X -a.s. satisfies the LDP on $\mathcal{P}^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ with rate function

$$I_F^{\text{que}}(Q) = \sup_{c \in \mathbb{N}} I_c^{\text{que}}(\langle Q \rangle_c), \quad Q \in \mathcal{P}^{\text{inv}}(\tilde{F}^{\mathbb{N}}). \quad (8.9)$$

The following lemma follows from Deuschel and Stroock [6], Lemma 4.4.15.

Lemma 8.1. *Let G be a Polish space, let $\mathcal{A}_c = \{A_{c,1}, \dots, A_{c,n_c}\}$, $c = 1, 2, \dots$ be a sequence of nested finite partitions of G such that $\lim_{c \rightarrow \infty} \text{diam}(\langle x \rangle_c) = 0$ for all $x \in G$ (with a coarse-graining map defined as above). Then we have, for $\mu, \nu \in \mathcal{P}(G)$,*

$$h(\langle \mu \rangle_c \mid \langle \nu \rangle_c) \nearrow h(\mu \mid \nu) \quad \text{as } c \rightarrow \infty. \quad (8.10)$$

Let

$$\mathcal{P}_E^{\text{inv}}(F^{\mathbb{N}}) := \left\{ \Phi \in \mathcal{P}^{\text{inv}}(F^{\mathbb{N}}) : \pi_1 \Phi(\iota(E)) = 1 \right\}, \quad (8.11)$$

$$\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}}) := \left\{ Q \in \mathcal{P}^{\text{inv}}(\tilde{F}^{\mathbb{N}}) : \pi_1 Q(\iota(\tilde{E})) = 1 \right\}. \quad (8.12)$$

Note that (8.4) allows to view each $\Phi \in \mathcal{P}^{\text{inv}}(E^{\mathbb{N}})$ as an element of $\mathcal{P}_E^{\text{inv}}(F^{\mathbb{N}})$ and each $Q \in \mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ as an element of $\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ via the identification of E and $\iota(E) \subset F$. In particular, we can view $\nu^{\otimes \mathbb{N}}$ as an element of $\mathcal{P}_E^{\text{inv}}(F^{\mathbb{N}})$ and $q_{\rho, \nu}^{\otimes \mathbb{N}}$ as an element of $\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$. We will make use of the fact that,

since each real-valued d_E -continuous function on $\iota(E)$ is automatically d_F -continuous, the weak topology on $\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ is finer than the weak topology on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$.

Fix $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{F}^{\mathbb{N}})$. Note that the functions

$$(N, c) \mapsto \frac{1}{N} h(\langle \pi_N Q \rangle_c \mid \langle q_{\rho, \nu}^{\otimes N} \rangle_c) \quad \text{and} \quad (L, c) \mapsto \frac{1}{L} h(\langle \pi_L \Psi_Q \rangle_c \mid \langle \nu^{\otimes L} \rangle_c) \quad (8.13)$$

are non-decreasing in both coordinates. Then deduce from (8.9) and (1.16) that

$$\begin{aligned} I_F^{\text{que}}(Q) &= \sup_{c \in \mathbb{N}} \left\{ H(\langle Q \rangle_c \mid \langle q_{\rho, \nu}^{\otimes N} \rangle_c) + (\alpha - 1) m_{\langle Q \rangle_c} H(\Psi_{\langle Q \rangle_c} \mid \langle \nu^{\otimes N} \rangle_c) \right\} \\ &= \sup_{c \in \mathbb{N}} \left\{ \sup_{N \in \mathbb{N}} \frac{1}{N} h(\langle \pi_N Q \rangle_c \mid \langle q_{\rho, \nu}^{\otimes N} \rangle_c) + (\alpha - 1) m_Q \sup_{L \in \mathbb{N}} \frac{1}{L} h(\langle \pi_L \Psi_Q \rangle_c \mid \langle \nu^{\otimes L} \rangle_c) \right\} \\ &= \sup_{N \in \mathbb{N}} \frac{1}{N} \sup_{c \in \mathbb{N}} h(\langle \pi_N Q \rangle_c \mid \langle q_{\rho, \nu}^{\otimes N} \rangle_c) + (\alpha - 1) m_Q \sup_{L \in \mathbb{N}} \frac{1}{L} \sup_{c \in \mathbb{N}} h(\langle \pi_L \Psi_Q \rangle_c \mid \langle \nu^{\otimes L} \rangle_c) \\ &= \sup_{N \in \mathbb{N}} \frac{1}{N} h(\pi_N Q \mid q_{\rho, \nu}^{\otimes N}) + (\alpha - 1) m_Q \sup_{L \in \mathbb{N}} \frac{1}{L} h(\pi_L \Psi_Q \mid \nu^{\otimes L}) \\ &= H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) + (\alpha - 1) m_Q H(\Psi_Q \mid \nu^{\otimes \mathbb{N}}), \end{aligned} \quad (8.14)$$

where we have used Lemma 8.1 in the fourth line. Note that in the third line interchanging the suprema and splitting out the supremum over the sum is justified because of (8.13).

For $Q \in \mathcal{P}^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ with $m_Q = \infty$ we see from (8.9), (1.15) and (8.14) that

$$\begin{aligned} I_F^{\text{que}}(Q) &= \sup_{c \in \mathbb{N}} I_c^{\text{que}}(\langle Q \rangle_c) = \sup_{c \in \mathbb{N}} \sup_{\text{tr} \in \mathbb{N}} \left\{ H([\langle Q \rangle_c]_{\text{tr}} \mid \langle q_{\rho, \nu}^{\otimes N} \rangle_c) + (\alpha - 1) m_{[Q]_{\text{tr}}} H(\Psi_{[Q]_{\text{tr}}} \mid \langle \nu^{\otimes N} \rangle_c) \right\} \\ &= \sup_{\text{tr} \in \mathbb{N}} \left\{ H([Q]_{\text{tr}} \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) + (\alpha - 1) m_{[Q]_{\text{tr}}} H(\Psi_{[Q]_{\text{tr}}} \mid \nu^{\otimes \mathbb{N}}) \right\}. \end{aligned} \quad (8.15)$$

Note that $\sup_{\text{tr} \in \mathbb{N}}$ can be replaced by $\text{tr} \rightarrow \infty$ by arguments analogous to Step 3 in the proof of Theorem 1.2.

Finally, we transfer the LDP from $\mathcal{P}^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ to $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$. To this end, we first verify that the rate function is concentrated on $\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$. Put

$$F'' := \{y \in \tilde{F} : y \text{ contains at least one letter from } F \setminus \iota(E)\}. \quad (8.16)$$

Then $q_{\rho, \nu}(F'') = 0$. For $Q \in \mathcal{P}^{\text{inv}}(\tilde{F}^{\mathbb{N}}) \setminus \mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ we have $\pi_1 Q(F'') > 0$, and hence

$$I_F^{\text{que}}(Q) \geq H(Q \mid q_{\rho, \nu}^{\otimes \mathbb{N}}) \geq h(\pi_1 Q \mid q_{\rho, \nu}) = \infty. \quad (8.17)$$

Thus, by Dembo and Zeitouni [5], Lemma 4.1.5, the family $\mathbb{P}(R_N \in \cdot \mid X)$ satisfies for $\nu^{\otimes \mathbb{N}}$ -a.s. all X an LDP on $\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ with rate N and with rate function given by (1.15–1.16).

To conclude the proof, observe that we can identify $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$ and $\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$, and that the weak topology on $\mathcal{P}^{\text{inv}}(\tilde{E}^{\mathbb{N}})$, which is ‘built’ on d_E , is not finer than that which $\mathcal{P}_E^{\text{inv}}(\tilde{F}^{\mathbb{N}})$ inherits from $\mathcal{P}^{\text{inv}}(\tilde{F}^{\mathbb{N}})$, which is ‘built’ on d_F (recall the discussion following (8.11–8.12)). Consequently, the LDP carries over. \square

A Appendix: Continuity under truncation limits

The following lemma implies (1.17).

Lemma A.1. For all $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$,

$$\begin{aligned} \lim_{\text{tr} \rightarrow \infty} H([Q]_{\text{tr}} | q_{\rho, \nu}^{\otimes \mathbb{N}}) &= H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}), \\ \lim_{\text{tr} \rightarrow \infty} m_{[Q]_{\text{tr}}} H(\Psi_{[Q]_{\text{tr}}} | \nu^{\otimes \mathbb{N}}) &= m_Q H(\Psi_Q | \nu^{\otimes \mathbb{N}}). \end{aligned} \quad (\text{A.1})$$

Proof. The proof is not quite standard, because Q and $[Q]_{\text{tr}}$, respectively, Ψ_Q and $\Psi_{[Q]_{\text{tr}}}$ are not “ \bar{d} -close” when tr is large, so that we cannot use the fact that entropy is “ \bar{d} -continuous” (see Shields [10]).

Lower semi-continuity yields $\liminf_{\text{tr} \rightarrow \infty} \text{l.h.s.} \geq \text{r.h.s.}$ for both limits, so we need only prove the reverse inequality. Note that, for all $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$,

$$H(Q) \leq h(Q|_{\mathcal{F}_1}) \leq h(\mathcal{L}_Q(\tau_1)) + m_Q \log |E| < \infty, \quad H(\Psi_Q) \leq \log |E| < \infty, \quad H(Q | q_{\rho, \nu}^{\otimes \mathbb{N}}) < \infty. \quad (\text{A.2})$$

For Z a random variable, we write $\mathcal{L}_Q(Z)$ to denote the law of Z under Q .

A.1 Proof of first half of (A.1)

Proof. Since $q_{\rho, \nu}^{\otimes \mathbb{N}}$ is a product measure, we have for, any $\text{tr} \in \mathbb{N}$,

$$\begin{aligned} H([Q]_{\text{tr}} | q_{\rho, \nu}^{\otimes \mathbb{N}}) &= -H([Q]_{\text{tr}}) - \mathbb{E}_{[Q]_{\text{tr}}}[\log \rho(\tau_1)] - \mathbb{E}_{[Q]_{\text{tr}}} \left[\sum_{i=1}^{\tau_1} \log \nu(Y_i^{(1)}) \right] \\ &= -H([Q]_{\text{tr}}) - \mathbb{E}_Q[\log \rho(\tau_1 \wedge \text{tr})] - \mathbb{E}_Q \left[\sum_{i=1}^{\tau_1 \wedge \text{tr}} \log \nu(Y_i^{(1)}) \right]. \end{aligned} \quad (\text{A.3})$$

By dominated convergence, using that $m_Q < \infty$ and $\log \rho(n) \leq C \log(n+1)$ for some $C < \infty$, we see that as $\text{tr} \rightarrow \infty$ the last two terms in the second line converge to

$$- \mathbb{E}_Q[\log \rho(\tau_1)] - \mathbb{E}_Q \left[\sum_{i=1}^{\tau_1} \log \nu(Y_i^{(1)}) \right]. \quad (\text{A.4})$$

Thus, it remains to check that

$$\lim_{\text{tr} \rightarrow \infty} H([Q]_{\text{tr}}) = H(Q). \quad (\text{A.5})$$

Obviously, $H([Q]_{\text{tr}}) \leq H(Q)$ for all $\text{tr} \in \mathbb{N}$ (indeed, $h([Q]_{\text{tr}}|_{\mathcal{F}_N}) \leq h(Q|_{\mathcal{F}_N})$ for all $N, \text{tr} \in \mathbb{N}$, because $[Q]_{\text{tr}}$ is the image measure of Q under the truncation map). For the asymptotic converse, we argue as follows. A decomposition of entropy gives

$$h(Q|_{\mathcal{F}_N}) = h([Q]_{\text{tr}}|_{\mathcal{F}_N}) + \int_{[\tilde{E}]_{\text{tr}}^N} h(\mathcal{L}_Q(\pi_N Y | \pi_N[Y]_{\text{tr}} = z)) (\pi_N[Q]_{\text{tr}})(dz), \quad (\text{A.6})$$

where π_N is the projection onto the first N words, and $\mathcal{L}_Q(\pi_N Y | \pi_N[Y]_{\text{tr}} = z)$ is the conditional distribution of the first N words given their truncations. We have

$$h(\mathcal{L}_Q(\pi_N Y | \pi_N[Y]_{\text{tr}} = z)) \leq \sum_{i=1}^N h(\mathcal{L}_Q(Y_i | \pi_N[Y]_{\text{tr}} = z)) \quad (\text{A.7})$$

and

$$\begin{aligned} &\int_{[\tilde{E}]_{\text{tr}}^N} h(\mathcal{L}_Q(Y_i | \pi_N[Y]_{\text{tr}} = z)) (\pi_N[Q]_{\text{tr}})(dz) \\ &\leq \int_{[\tilde{E}]_{\text{tr}}^N} h(\mathcal{L}_Q(Y_i | [Y]_{\text{tr}} = z_i)) (\pi_N[Q]_{\text{tr}})(dz) \\ &= \int_{[\tilde{E}]_{\text{tr}}} h(\mathcal{L}_Q(Y_1 | [Y]_{\text{tr}} = y)) (\pi_1[Q]_{\text{tr}})(dy), \quad 1 \leq i \leq N, \end{aligned} \quad (\text{A.8})$$

where the inequality in the second line comes from the fact that conditioning on less increases entropy, and the third line uses the shift-invariance. Combining (A.6–A.8) and letting $N \rightarrow \infty$, we obtain

$$H(Q) \leq H([Q]_{\text{tr}}) + \int_{[\tilde{E}]_{\text{tr}}} h\left(\mathcal{L}_Q(Y_1 \mid [Y_1]_{\text{tr}} = y)\right) (\pi_1[Q]_{\text{tr}})(dy), \quad (\text{A.9})$$

and so it remains to check that the second term in the right-hand side vanishes as $\text{tr} \rightarrow \infty$.

Note that this term equals (write ε for the empty word and $w \cdot w'$ for the concatenation of words w and w')

$$\begin{aligned} & - \sum_{\substack{w \in \tilde{E} \\ \tau(w) = \text{tr}}} [Q]_{\text{tr}}(w) \sum_{w' \in \tilde{E} \cup \{\varepsilon\}} \frac{Q(w \cdot w')}{[Q]_{\text{tr}}(w)} \log \left[\frac{Q(w \cdot w')}{[Q]_{\text{tr}}(w)} \right] \\ & = - \sum_{\substack{w'' \in \tilde{E} \\ \tau(w'') \geq \text{tr}}} Q(w'') \log Q(w'') + \sum_{\substack{w'' \in \tilde{E} \\ \tau(w'') \geq \text{tr}}} Q(w'') \log [Q]_{\text{tr}}([w'']_{\text{tr}}). \end{aligned} \quad (\text{A.10})$$

But

$$0 \geq \sum_{\substack{w'' \in \tilde{E} \\ \tau(w'') \geq \text{tr}}} Q(w'') \log [Q]_{\text{tr}}([w'']_{\text{tr}}) \geq \sum_{\substack{w'' \in \tilde{E} \\ \tau(w'') \geq \text{tr}}} Q(w'') \log Q(w''), \quad (\text{A.11})$$

and so the right-hand side of (A.10) vanishes as $\text{tr} \rightarrow \infty$. \square

A.2 Proof of second half of (A.1)

Note that $\lim_{\text{tr} \rightarrow \infty} m_{[Q]_{\text{tr}}} = m_Q$ and $w - \lim_{\text{tr} \rightarrow \infty} \Psi_{[Q]_{\text{tr}}} = \Psi_Q$ by dominated convergence, implying that

$$\liminf_{\text{tr} \rightarrow \infty} H(\Psi_{[Q]_{\text{tr}}} \mid \nu^{\otimes \mathbb{N}}) \geq H(\Psi_Q \mid \nu^{\otimes \mathbb{N}}). \quad (\text{A.12})$$

So it remains to check the reverse inequality. Since $\nu^{\otimes \mathbb{N}}$ is product measure, we have

$$H(\Psi_{[Q]_{\text{tr}}} \mid \nu^{\otimes \mathbb{N}}) = -H(\Psi_{[Q]_{\text{tr}}}) - \frac{1}{m_{[Q]_{\text{tr}}}} \mathbb{E}_Q \left[\sum_{i=1}^{\tau_1 \wedge \text{tr}} \log \nu \left(Y_i^{(1)} \right) \right]. \quad (\text{A.13})$$

By dominated convergence, as $\text{tr} \rightarrow \infty$ the second term converges to

$$\frac{1}{m_Q} \mathbb{E}_Q \left[\sum_{i=1}^{\tau_1} \log \nu \left(Y_i^{(1)} \right) \right] = \int_E \Psi_Q(dx) \log \nu(x). \quad (\text{A.14})$$

Thus, it remains to check that

$$\lim_{\text{tr} \rightarrow \infty} H(\Psi_{[Q]_{\text{tr}}}) = H(\Psi_Q). \quad (\text{A.15})$$

We will first prove (A.15) for ergodic Q , in which case $[Q]_{\text{tr}}$, Ψ_Q , $\Psi_{[Q]_{\text{tr}}}$ are ergodic (Birkner [2], Remark 5).

For $\Psi \in \mathcal{P}^{\text{erg}}(E^{\mathbb{N}})$ and $\varepsilon \in (0, 1)$, let

$$\mathcal{N}_n(\Psi, \varepsilon) = \min \{ \#A : A \subset E^n, \Psi(A \times E^\infty) \geq \varepsilon \} \quad (\text{A.16})$$

be the (n, ε) covering number of Ψ . For any $\varepsilon \in (0, 1)$, we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mathcal{N}_n(\Psi, \varepsilon) = H(\Psi) \quad (\text{A.17})$$

(see Shields [10], Theorem I.7.4). The idea behind (A.15) is that there are $\approx \exp[nH(\Psi_Q)]$ “ Ψ_Q -typical” sequences of length n , and that a “ $\Psi_{[Q]_{\text{tr}}}$ -typical” sequence arises from a “ Ψ_Q -typical”

sequence by eliminating a fraction δ_{tr} of the letters, where $\delta_{\text{tr}} \rightarrow 0$ as $\text{tr} \rightarrow \infty$. Hence $\mathcal{N}_n(\Psi_Q, \varepsilon)$ cannot be much larger than $\mathcal{N}_n(\Psi_{[Q]_{\text{tr}}}, \varepsilon)$ (on an exponential scale), implying that $H(\Psi_Q) - H(\Psi_{[Q]_{\text{tr}}})$ must be small.

To make this argument precise, fix $\varepsilon > 0$ and pick N_0 so large that

$$Q(|\kappa(Y^{(1)}, \dots, Y^{(N)})| \in Nm_Q[1 - \varepsilon, 1 + \varepsilon]) > 1 - \varepsilon \quad \text{for } N \geq N_0. \quad (\text{A.18})$$

Pick $\text{tr}_0 \in \mathbb{N}$ so large that for $\text{tr} \geq \text{tr}_0$ and $N \geq N_0$,

$$Q\left(\sum_{i=1}^N (\tau_i - \text{tr})_+ < N\varepsilon\right) > 1 - \varepsilon/2, \quad Q(\tau_1 \leq \text{tr}) > 1 - \varepsilon/2, \quad m_{[Q]_{\text{tr}}} > (1 - \varepsilon)m_Q. \quad (\text{A.19})$$

For $n \geq \lceil N_0/m_Q \rceil$, we will construct a set $B \subset E^n$ such that

$$\Psi_Q(B \times E^\infty) \geq \frac{1}{2}, \quad |B| \leq \exp[n(H(\Psi_{[Q]_{\text{tr}}}) + \delta)], \quad (\text{A.20})$$

where δ can be made arbitrarily small by choosing ε small in (A.18–A.19). Hence, by the asymptotic cover property (A.17), we have $H(\Psi_Q) \leq (1 + \delta)H(\Psi_{[Q]_{\text{tr}}})$ and

$$\liminf_{\text{tr} \rightarrow \infty} H(\Psi_{[Q]_{\text{tr}}}) \geq H(\Psi_Q), \quad (\text{A.21})$$

completing the proof of (A.15).

We verify (A.20) as follows. Put $N := \lceil nm_Q(1 + 2\varepsilon) \rceil$. By (A.18–A.19) and the asymptotic cover property (A.17) for $\Psi_{[Q]_{\text{tr}}}$, there is a set $A \subset \tilde{E}^N$ such that

$$\mathbb{E}_Q[\tau_1 \mathbb{1}_A(Y^{(1)}, \dots, Y^{(N)})] > (1 - \varepsilon)m_Q \quad (\text{A.22})$$

and

$$\begin{aligned} |\kappa(y^{(1)}, \dots, y^{(N)})| &\geq n(1 + \varepsilon), \quad \tau(y^{(1)}) \leq \text{tr}, \quad \sum_{i=1}^N (\tau(y^{(i)}) - \text{tr})_+ < N\varepsilon, \\ \forall (y^{(1)}, \dots, y^{(N)}) &\in A, \end{aligned} \quad (\text{A.23})$$

while the set

$$B' := \{\kappa([y^{(1)}]_{\text{tr}}, \dots, [y^{(N)}]_{\text{tr}})|_{(0, \lceil (1-\varepsilon)n \rceil]} : (y^{(1)}, \dots, y^{(N)}) \in A\} \subset E^{\lceil (1-\varepsilon)n \rceil} \quad (\text{A.24})$$

satisfies

$$|B'| \leq \exp[n(H(\Psi_{[Q]_{\text{tr}}}) + \varepsilon)]. \quad (\text{A.25})$$

Put

$$B := \{\kappa(y^{(1)}, \dots, y^{(N)})|_{(0, n]} : (y^{(1)}, \dots, y^{(N)}) \in A\} \subset E^n. \quad (\text{A.26})$$

Observe that each $x' \in B'$ corresponds to at most

$$|E|^{\varepsilon n} \binom{n}{\varepsilon n} \leq \exp[-n(\varepsilon \log \varepsilon + (1 - \varepsilon) \log(1 - \varepsilon)) + n\varepsilon \log |E|] \quad (\text{A.27})$$

different $x \in B$, so that

$$|B| \leq |B'| \exp[-n(\varepsilon \log \varepsilon + (1 - \varepsilon) \log(1 - \varepsilon)) + n\varepsilon \log |E|]. \quad (\text{A.28})$$

We have

$$\begin{aligned}
m_Q \Psi_Q(B \times E^\infty) &\geq \mathbb{E}_Q \left[\sum_{k=0}^{\tau_1-1} \mathbb{1}_{B \times E^\infty}(\theta^k \kappa(Y)) \mathbb{1}_A(Y^{(1)}, \dots, Y^{(N)}) \right] \\
&= \mathbb{E}_Q \left[\sum_{k=0}^{\tau_1 \wedge \text{tr}-1} \mathbb{1}_{B' \times E^\infty}(\theta^k \kappa([Y]_{\text{tr}})) \mathbb{1}_A(Y^{(1)}, \dots, Y^{(N)}) \right] \\
&\geq \mathbb{E}_Q \left[\sum_{k=0}^{\tau_1 \wedge \text{tr}-1} \mathbb{1}_{B' \times E^\infty}(\theta^k \kappa([Y]_{\text{tr}})) \right] - \varepsilon m_Q \\
&= m_{[Q]_{\text{tr}}} \Psi_{[Q]_{\text{tr}}}(B' \times E^\infty) - \varepsilon m_Q,
\end{aligned} \tag{A.29}$$

so that, finally,

$$\Psi_Q(B \times E^\infty) \geq \frac{m_{[Q]_{\text{tr}}}}{m_Q} \Psi_{[Q]_{\text{tr}}}(B' \times E^\infty) - \varepsilon \geq \frac{1}{2}. \tag{A.30}$$

Combining (A.25), (A.28) and (A.30), we obtain (A.20) with

$$\delta = -(\varepsilon \log \varepsilon + (1 - \varepsilon) \log(1 - \varepsilon)) + \varepsilon(1 + \log |E|). \tag{A.31}$$

Since $\limsup_{\text{tr} \rightarrow \infty} H(\Psi_{[Q]_{\text{tr}}}) \leq H(\Psi_Q)$ by upper semi-continuity of H (see e.g. Georgii [7], Proposition. 15.14), this concludes the proof of (A.15) for ergodic Q .

For general $Q \in \mathcal{P}^{\text{inv,fin}}(\tilde{E}^{\mathbb{N}})$, we recall the ergodic decomposition formulas stated in (6.1–6.2). These yield

$$\Psi_{[Q]_{\text{tr}}} = \int_{\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})} \frac{m_{[Q']_{\text{tr}}}}{m_{[Q]_{\text{tr}}}} \Psi_{[Q']_{\text{tr}}} W_Q(dQ') \tag{A.32}$$

and

$$H(\Psi_{[Q]_{\text{tr}}}) = \int_{\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})} \frac{m_{[Q']_{\text{tr}}}}{m_{[Q]_{\text{tr}}}} H(\Psi_{[Q']_{\text{tr}}}) W_Q(dQ'), \tag{A.33}$$

because specific entropy is affine. The integrand inside (A.33) is non-negative and, by the above, converges to $\frac{m_{Q'}}{m_Q} H(\Psi_{Q'})$ as $\text{tr} \rightarrow \infty$. Hence, by Fatou's lemma,

$$\liminf_{\text{tr} \rightarrow \infty} H(\Psi_{[Q]_{\text{tr}}}) \geq \int_{\mathcal{P}^{\text{erg,fin}}(\tilde{E}^{\mathbb{N}})} \frac{m_{Q'}}{m_Q} H(\Psi_{Q'}) W_Q(dQ') = H(\Psi_Q), \tag{A.34}$$

which concludes the proof. \square

References

- [1] M. Birkner, *Particle Systems with Locally Dependent Branching: Long-Time Behaviour, Genealogy and Critical Parameters*, Dissertation, Johann Wolfgang Goethe-Universität Frankfurt am Main, 2003.
<http://publikationen.ub.uni-frankfurt.de/volltexte/2003/314/>
- [2] M. Birkner, Conditional large deviations for a sequence of words, *Stoch. Proc. Appl.* 118 (2008) 703–729.
- [3] M. Birkner, A. Greven, F. den Hollander, Large deviations for the collision local time of transient random walks and intermediate phases in interacting stochastic systems, preprint 2008.
- [4] F. Comets, Large deviation estimates for a conditional probability distribution. Applications to random interaction Gibbs measures, *Probab. Theory Relat. Fields* 80 (1989) 407–432.

- [5] A. Dembo and O. Zeitouni, *Large Deviations Techniques and Applications*, 2nd Edition, Springer, 1998.
- [6] J.-D. Deuschel and D.W. Stroock, *Large Deviations*, Academic Press, Boston, 1989.
- [7] H.-O. Georgii, *Gibbs Measures and Phase Transitions*, de Gruyter Studies in Mathematics 9, Walter de Gruyter, Berlin, 1988.
- [8] F. den Hollander, *Large Deviations*, Fields Institute Monographs 14, American Mathematical Society, Providence, RI, 2000.
- [9] R. Holley and T. M. Liggett, Generalized potlatch and smoothing processes, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 55 (1981) 165-195.
- [10] P.C. Shields, *The Ergodic Theory of Discrete Sample Paths*, American Mathematical Society, Providence, RI, 1996.
- [11] V. Strassen, The existence of probability measures with given marginals, *Ann. Math. Statist.* 36 (1965) 423-439.
- [12] F. Toninelli, Disordered pinning models and copolymers: beyond annealed bounds, *Ann. Appl. Probab.* 18 (2008) 1569-1587.
- [13] P. Walters, *An Introduction to Ergodic Theory*, Graduate Texts in Mathematics 79, Springer, New York, 1982.