



Universiteit  
Leiden  
The Netherlands

## Discussion of five-colour observations of stars of high velocity

Oosterhoff, P.T.; Ponsen, J.

### Citation

Oosterhoff, P. T., & Ponsen, J. (1966). Discussion of five-colour observations of stars of high velocity. *Bulletin Of The Astronomical Institutes Of The Netherlands*, 18, 150. Retrieved from <https://hdl.handle.net/1887/6106>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/6106>

**Note:** To cite this publication please use the final published version (if applicable).

## DISCUSSION OF FIVE-COLOUR OBSERVATIONS OF STARS OF HIGH VELOCITY

P. TH. OOSTERHOFF and J. PONSEN†

Received 14 June 1965

A discussion is presented of the five-colour measures of stars of high velocity published by PONSEN and OOSTERHOFF (1965). The accidental errors and night errors are analysed. The final magnitudes and colours with their mean errors are given in table 3.

The individual photo-electric measures in five colours of stars of high velocity and of several pulsating variables made with the light-collector of the Leiden southern station in South Africa are published by PONSEN and OOSTERHOFF (1965). In the present article we shall refer to this publication as Paper I.

The stars of high velocity were selected from the finding list by BUSCOMBE and MORRIS (1958). The number of stars from this list which have been measured is 53. Most of these stars were observed more than once in the same night. From the differences between such observations we have derived the accidental mean errors of a single observation. The stars were first divided into three groups according to brightness, i.e. according to their value of  $V$ . The resulting mean errors are given in table 1. The numbers in brackets indicate the number of differences used.  $V_J$  is the visual magnitude in the  $UBV$  system, computed with the formula of Paper I. Brightness and colours as well as the mean errors are expressed in the logarithm of the intensities. Within each brightness group the colour ( $V-B$ ) has the highest accuracy. The colours ( $B-L$ ) and ( $B-U$ ) are about of equal accuracy and are only slightly less accurate than the colour ( $V-B$ ). However, the mean error of the ( $U-W$ ) colour is about two and a half times as large as that for the colours ( $B-L$ ) and ( $B-U$ ). For the stars in this programme the intensity in the  $W$ -band is very much smaller than in any of the other bands and furthermore the extinction coefficient is considerably larger than for the other

bands. These two effects no doubt explain the larger mean errors. Most of the stars observed show an excess in the  $L$ ,  $U$  and  $W$  intensities. A two-colour diagram is shown in figure 1. A strong correlation is found to exist between the ultraviolet excess and the spacial motion. This correlation is shown in figure 2.

Studying the observations in the main table of Paper I we found that the differences between the observations of the same star made in different nights are larger than would be expected from the accidental errors derived above. Not all the nights were photometrically perfect. In some the extinction coefficients showed considerable variations. The value of these coefficients, which we used, were interpolated in time and assumed to be the same all over the sky. Therefore it is not sur-

TABLE 1  
Accidental errors

Group	Mean error	
$2 < V < 3$ $9.35 < V_J < 11.85$	$V$	$\pm 0.0047$ (53)
	$V-B$	$\pm 0.0029$ (53)
	$B-U$	$\pm 0.0044$ (53)
	$U-W$	$\pm 0.0102$ (52)
	$B-L$	$\pm 0.0043$ (53)
$3 < V < 4$ $6.85 < V_J < 9.35$	$V$	$\pm 0.0029$ (53)
	$V-B$	$\pm 0.0014$ (53)
	$B-U$	$\pm 0.0021$ (53)
	$U-W$	$\pm 0.0056$ (53)
	$B-L$	$\pm 0.0024$ (53)
$4 < V$ $V_J < 6.85$	$V$	$\pm 0.0035$ (35)
	$V-B$	$\pm 0.0012$ (35)
	$B-U$	$\pm 0.0012$ (35)
	$U-W$	$\pm 0.0036$ (35)
	$B-L$	$\pm 0.0014$ (35)

TABLE 2  
Night errors

	Mean error	
$V$	$\pm 0.0056$	(44)
$V-B$	$\pm 0.0021$	(44)
$B-U$	$\pm 0.0031$	(44)
$U-W$	$\pm 0.0059$	(43)
$B-L$	$\pm 0.0011$	(44)

prising that the true errors are larger than the accidental errors of table 1. From the differences between observations made in different nights we have computed night errors taking due account of the accidental errors already derived. The resulting mean night errors are shown in table 2. In this case we made no distinction between bright and faint stars.

The final results for the 53 stars measured are presented in table 3. The stars are indicated by their number from Buscombe's finding list (further identifications at the end of the paper). The final results for  $V$  and for the four colours are weighted mean values. The mean errors have been computed from the accidental and night errors given in the tables 2 and 3. In the last two columns of this table we have added the visual magnitude in the  $UBV$  system as computed with the formula from Paper I, and the spectral type as given in the finding list.

Three of our stars occur also in the catalogue of high-velocity stars by ROMAN (1955). These stars are

Buscombe 128	A5 = Roman 62	sdF4
Buscombe 821	F5 = Roman 373	sdF5
Buscombe 1258	sdF8 = Roman 551	sdF8.

For star 128 our measures confirm the spectral type given by Miss Roman. For the two stars 1247 and 1248, for which Buscombe and Morris give spectral type K0, our measures indicate a spectral type in the F-class.

The numbers 83, 127 and 1258 of our list are identical with the numbers 5, 10 and 49 in the article by PRZYBYLSKI and MORRIS KENNEDY (1965). Furthermore 22 stars of our list also occur in the list of photoelectric magnitudes and colours of southern stars by COUSINS and STOY (1963).

We have compared the values of  $V_j$  in the last column but one of our table 3 with the  $V$  values in the

$UBV$  system as given by the authors mentioned above. The agreement is very close. For the 22 stars of the Cape list the zero-point difference was found to be 0.002 mag and the mean error of one difference  $\pm 0.018$  mag. According to the mean errors given in table 3 we expect a mean error of  $\pm 0.012$  mag for the  $V_j$  values, which would leave an error of  $\pm 0.013$  mag for the Cape magnitudes.

Walraven has made a thorough study of the relations between the colours in his five-colour photometry and the colours given by Cousins and Stoy in the publication mentioned above. His comparisons are based on a much larger number of stars and on a wider spectral range than we have at our disposal. As Walraven intends to publish the results of his investigation in *Bull. Astr. Inst. Netherlands*, it is sufficient to state here that our results in comparing our colours with the Cape colours are in close agreement with the relations derived by Walraven.

Dr. Walraven informed me that among the bright stars which he measured there is one star of high velocity, namely HD 132475 = Buscombe 776. His measures for this star are

$$\begin{aligned} V &= 3.322 & V-B &= +0.2282 \\ V_j &= 8.536 & B-U &= +0.2557 \\ \text{Sp} &= \text{F8} & U-W &= +0.2371 \\ & & B-L &= +0.1738. \end{aligned}$$

In figure 1 we have plotted the observations in the two-colour diagram of  $(V-B)$  against  $(B-U)$ . The curve for the main-sequence stars was taken from Walraven. From type O to A0 the curve is accurately determined. For the subtypes A, F and G its determination is less accurate and for the K- and M-type stars the curve is very uncertain.

Many of the points in figure 1 fall well above the main-sequence curve, indicating the existence of an ultraviolet excess. In this respect our diagram is very similar to the two-colour diagrams in the  $UBV$  system given by BECKER (1963) for the stars of Miss Roman's list and by SANDAGE and EGGEN (1959) for a number of subdwarfs.

In figure 1 we first notice one star with an exceptional ultraviolet excess. It is Buscombe 981, a peculiar B-type star at low galactic latitude in the direction of the galactic centre. Consequently it may be strongly reddened and probably does not intrinsically belong

TABLE 3

Final magnitude and colours, expressed in the logarithm of the intensity, of stars of high velocity. The star numbers are from the catalogue by Buscombe and Morris. The visual magnitude in the  $UBV$  system is given in the column headed  $V_J$

Star	V	V-B	B-U	U-W	B-I	$V_J$	Sp
83	2.6952 +46	0.2141 +20	0.2445 +31	0.1666 +66	0.1743 +23	10.105	F1
88	3.4112 -60	0.2369 -23	0.2669 -34	0.1964 -71	0.1912 -20	8.312	F8
95	4.1926 43	0.2102 16	0.2948 23	0.1736 46	0.2118 11	6.362	F5
127	2.5568 46	0.1820 20	0.2542 31	0.1603 66	0.1555 23	10.455	dF1
128	2.8351 46	0.1801 20	0.2410 31	0.1493 66	0.1503 23	9.760	A5
161	2.2558 46	0.1744 20	0.2288 31	0.1584 66	0.1471 23	11.209	sdA8
451	2.8754 38	0.1956 17	0.2330 25	0.1562 54	0.1624 19	9.657	sdF3
456	3.8422 34	0.2069 13	0.3047 20	0.1800 41	0.2108 12	7.239	F8
460	3.8244 42	0.2636 16	0.3029 24	0.2219 50	0.2541 14	7.276	G0
461	3.5480 42	0.2434 16	0.3275 24	0.2180 50	0.2492 14	7.970	F8
462	2.9766 58	0.0640 23	0.4574 35	0.1337 69	0.2084 19	9.420	A0
465	4.3976 43	0.4636 16	0.5642 23	0.3634 46	0.5038 11	5.818	K0
484	3.9538 42	0.2468 16	0.3102 24	0.1988 50	0.2437 14	6.955	G0
493	2.9638 46	0.3025 20	0.3259 31	0.2138 66	0.3041 23	9.423	G5
502	5.2868 66	0.4382 24	0.5961 33	0.3269 69	0.5325 18	3.598	K0
522	4.4609 43	0.2208 16	0.3015 23	0.1978 45	0.2187 10	5.690	F8
531	3.9308 60	0.1706 23	0.3404 34	0.1732 71	0.2061 20	7.022	F2
532	5.4791 66	0.5580 24	0.7214 33	0.4389 69	0.6285 18	3.102	K0
556	4.3087 61	0.0816 22	0.4287 32	0.1508 64	0.2338 15	6.088	F0
565	4.7120 61	0.1820 22	0.3394 32	0.1870 64	0.2018 15	5.067	F2
821	3.8543 60	0.2073 23	0.2414 34	0.1754 71	0.1431 20	7.208	F5
928	3.2478 42	0.1911 16	0.2507 24	0.1580 50	0.1497 14	8.727	F5
934	2.4894 65	0.2554 29	0.2709 44	0.1954 93	0.1785 32	10.615	sdF8
954	3.6950 42	0.2772 16	0.3168 24	0.2114 50	0.2610 14	7.598	G0
975	5.1715 43	0.6016 16	0.8078 23	0.4648 45	0.6906 10	3.866	K0
981	3.6794 63	0.2199 25	0.1357 37	0.1243 81	0.0833 27	7.644	Bp
1057	4.3884 43	0.2467 16	0.3223 23	0.1895 45	0.2519 10	5.868	F8
1067	3.2052 42	0.2728 16	0.3261 24	0.2203 50	0.2732 14	8.823	G0
1080	4.2261 43	0.2336 16	0.3082 23	0.1890 46	0.2204 11	6.276	F8
1085	3.2750 42	0.2302 16	0.2477 24	0.1792 49	0.1898 13	8.654	sdG0
1090	4.6777 43	0.2279 16	0.3109 23	0.1873 45	0.2070 10	5.147	F8
1131	3.7070 42	0.2334 16	0.2938 24	0.1936 50	0.2179 14	7.573	G0
1150	3.6085 42	0.2436 16	0.3386 24	0.2031 50	0.2605 14	7.818	G0
1159	3.1242 42	0.2798 16	0.3306 24	0.2020 50	0.2832 14	9.025	G5
1163	2.5220 46	0.1686 20	0.2274 31	0.1388 66	0.1359 23	10.544	sdF
1168	3.2837 42	0.3704 16	0.4730 24	0.2407 50	0.4403 14	8.614	G5
1177	2.4870 46	0.6088 20	0.6896 31	0.4595 93	0.5645 23	10.576	K4
1186	3.8879 42	0.2429 16	0.3064 24	0.1981 49	0.2332 13	7.120	F8
1197	4.2484 43	0.2763 16	0.3421 23	0.2135 45	0.2860 10	6.214	G0
1204	4.6281 43	0.2901 16	0.2805 23	0.2336 45	0.1991 10	5.263	G0
1209	3.5935 42	0.2604 16	0.3195 24	0.2136 50	0.2554 14	7.854	G0
1220	3.5846 42	0.2738 16	0.3127 24	0.2056 50	0.2670 14	7.874	G0
1237	3.6154 42	0.2515 16	0.3100 24	0.2106 50	0.2442 14	7.800	G0
1246	4.4086 43	0.1941 16	0.3174 23	0.1760 46	0.2071 11	5.824	F8
1247	2.0567 45	0.2163 20	0.3106 30	0.1811 62	0.2316 21	11.701	K0
1248	2.1047 46	0.2186 20	0.3263 31	0.1838 66	0.2120 23	11.581	K0
1258	3.4752 42	0.1982 16	0.2332 24	0.1612 50	0.1602 14	8.157	sdF8
1266	3.1327 42	0.2420 16	0.3120 24	0.1962 49	0.2437 13	9.008	F8
1276	3.0490 60	0.3500 23	0.4601 34	0.2311 71	0.4130 20	9.204	G5
1279	3.8838 42	0.4314 16	0.5698 24	0.3274 49	0.5274 13	7.107	K0
1283	2.6940 65	0.2824 29	0.3222 44	0.1990 93	0.2763 32	10.100	G0
1284	4.2854 61	0.4249 22	0.5328 32	0.3310 64	0.4685 15	6.103	G5
1311	3.1627 42	0.1297 16	0.3117 24	0.1409 50	0.1658 14	8.947	A2




Figure 1. The high-velocity stars in the two-colour diagram. ( $V-B$ ) and ( $B-U$ ) expressed in logarithm of the intensity.

to that part of the diagram where it is found. Next we notice a group of stars with large ultraviolet excess, the excess in  $U$  being about 0.075. Quite separate from this group we find another group of stars with an excess in  $U$  of about 0.030. And finally there are some stars which fall just above or below the main-sequence curve.

We refer here also to the interesting two-colour diagrams for dwarfs and subdwarfs by DIXON (1965), who used them for a study of the past rates of star formation. As our colours deviate from those in the  $UBV$  system a direct comparison of the diagrams cannot be made.

We have computed approximate values of the excesses in the logarithm of the intensities for the  $L$ -,  $U$ - and  $W$ -bands separately, by comparing each programme star with a star on the main sequence with identical value of ( $V-B$ ). The photometric data for these main-sequence stars were read from Walraven's curves in the two-colour diagrams presented at the symposium on Spectral Classification and Multicolour Photometry, held at Saltsjöbaden in 1964. As the red part of the main sequence is very poorly determined, we have not made this computation for the programme stars with values of ( $V-B$ ) larger than  $+0.4$ . The results are presented in table 4. In this table the stars

have been arranged in three groups, with large ultraviolet excess, with moderate excess and without noticeable excess. Within each group the stars have been placed in the order of increasing ( $V-B$ ) values. In the fourth column we have added the spectral type, taken from the catalogue by Buscombe and Morris. The data within brackets were taken from Miss Roman's catalogue. In the last column we have given the space motions in km/sec relative to the local standard of rest, or the observed radial velocities. The latter are given in brackets. These data also were taken from the catalogue by Buscombe and Morris.

In the first group of stars with large ultraviolet excess we find the three stars common with Miss Roman's list and further all the stars of which the subdwarf character has been established spectroscopically. It is clear that the ultraviolet excess affects all three bands  $L$ ,  $U$  and  $W$ . If we form mean values for the 14 stars in this group, we find for  $L + 0.065$ , for  $U + 0.081$  and for  $W + 0.086$ . Therefore the effect is smallest for  $L$  and largest for  $W$ . However the figures in table 4 for this first group indicate a systematic change with colour, or with the value of ( $V-B$ ), in the ratio between the excess for the three bands. For the blue stars the excess in  $L$  is definitely smaller than the excess in  $U$  or  $W$ , but for the reddest stars the excess in  $L$  is larger than that of the other two bands.

There probably are two causes for this systematic shift in the values of these ratios. In computing the excesses we have assumed that the  $B$ -band is not affected by the difference in strength of the metal lines between subdwarfs and normal dwarfs. It is not unlikely that the errors in the computed excesses, introduced by this assumption, depend on the spectral type. Another cause for the systematic shift mentioned above could be errors in the shape of the curves, representing the main-sequence stars, in the various two-colour diagrams. According to Walraven the shape of the curves for the main sequence in the F and G spectral region is not very well determined.

There is little to say about the stars with moderate ultraviolet excess or about the stars near the main sequence. It is clear, however, that the space motions and radial velocities of the stars in these two groups are systematically smaller than those of the stars with large excess.

To study the correlation between ultraviolet excess

TABLE 4

The excess in  $L$ ,  $U$  en  $W$ , expressed in the logarithm of the intensity. The last column shows the space motion in km/sec relative to the local standard of rest and in brackets the radial velocity relative to the Sun

Star	$V-B$	Excess in			Spectrum	Velocity (km/sec)
		$L$	$U$	$W$		
Stars with large ultraviolet excess						
1311	0.1297	+ 0.026	+ 0.055	+ 0.069	A2	
1163	0.1686	+ 0.068	+ 0.104	+ 0.126	sdF	(+103)
161	0.1744	+ 0.058	+ 0.099	+ 0.102	sdA8	(+154)
128	0.1801	+ 0.057	+ 0.085	+ 0.097	A5 (sdF4)	(+ 63)
127	0.1820	+ 0.054	+ 0.071	+ 0.084	dF1	(+234)
928	0.1911	+ 0.064	+ 0.071	+ 0.077	F5	
451	0.1956	+ 0.055	+ 0.097	+ 0.098	sdF3	178
1258	0.1982	+ 0.059	+ 0.087	+ 0.094	sdF8	252
821	0.2073	+ 0.081	+ 0.079	+ 0.075	(sd)F5	231
83	0.2141	+ 0.056	+ 0.088	+ 0.096	F1	
1085	0.2302	+ 0.056	+ 0.079	+ 0.085	sdG0	218
88	0.2369	+ 0.064	+ 0.063	+ 0.056	F8	
934	0.2554	+ 0.096	+ 0.072	+ 0.071	sdF8	182
1204	0.2901	+ 0.111	+ 0.090	+ 0.073	dG0	213
Stars with moderate ultraviolet excess						
456	0.2069	+ 0.014	+ 0.016	+ 0.007	F8	
95	0.2102	+ 0.016	+ 0.026	+ 0.024	F5	69
522	0.2208	+ 0.018	+ 0.023	+ 0.003	F8	86
1090	0.2279	+ 0.037	+ 0.015	+ 0.011	F8	65
1131	0.2334	+ 0.033	+ 0.035	+ 0.027	G0	
1080	0.2336	+ 0.030	+ 0.020	+ 0.017	F8	
1266	0.2420	+ 0.016	+ 0.021	+ 0.016	F8	
1186	0.2429	+ 0.029	+ 0.028	+ 0.020	F8	
484	0.2468	+ 0.024	+ 0.026	+ 0.019	G0	
1237	0.2515	+ 0.026	+ 0.020	+ 0.012	G0	
1209	0.2604	+ 0.025	+ 0.027	+ 0.011	G0	
460	0.2636	+ 0.029	+ 0.044	+ 0.033	G0	85
1067	0.2728	+ 0.019	+ 0.030	+ 0.014	G0	
1220	0.2738	+ 0.026	+ 0.044	+ 0.044	G0	154
954	0.2772	+ 0.035	+ 0.042	+ 0.038	G0	
1159	0.2798	+ 0.017	+ 0.032	+ 0.039	G5	
1283	0.2824	+ 0.025	+ 0.041	+ 0.051	G0	(- 98)
493	0.3025	+ 0.020	+ 0.056	+ 0.065	G5	
Stars near the main sequence						
462	0.0640	- 0.020	- 0.021	- 0.017	A0 SB	
556	0.0816	- 0.046	- 0.007	- 0.017	F0 SB	
531	0.1706	- 0.003	- 0.010	- 0.022	F2	
565	0.1820	+ 0.006	- 0.014	- 0.039	F2	75
1246	0.1941	+ 0.008	0.000	- 0.011	F8	
1247	0.2163	+ 0.001	+ 0.011	+ 0.007	K0	
1248	0.2186	+ 0.023	- 0.003	- 0.010	K0	
461	0.2434	+ 0.013	+ 0.006	- 0.021	F8	(+ 71)
1150	0.2436	+ 0.002	- 0.005	- 0.017	G0	
1057	0.2467	+ 0.015	+ 0.014	+ 0.016	F8	67
1197	0.2763	+ 0.010	+ 0.016	+ 0.010	G0	97
1276	0.3500	- 0.035	- 0.013	+ 0.009	G5	
1168	0.3704	- 0.035	+ 0.009	+ 0.031	G5	90




Figure 2. The excess in  $U$ , expressed in the logarithm of the intensity, plotted against the spacial motion relative to the local standard of rest. Arrows indicate stars for which only radial velocities are known.

and space velocity we have plotted in figure 2 the excess in the  $U$ -band, from table 4, against the space motions from the same table. For stars with unknown spacial motion, but with given radial velocity we have reduced the latter to the local standard of rest. These radial velocities have been indicated in figure 2 by arrows. The coefficient of correlation between the ex-

cess and the space motion is  $+0.90$ . We derived the following relation

$$\text{space motion} = 59 + 1730 \cdot (\text{excess in } U) \\ \pm 8 \quad 230 \text{ (m.e.)}$$

The dispersion of the space velocities relative to this formula is  $\pm 31$  km/sec. All the stars plotted in figure 2, which have a spacial motion larger than 80 km/sec, show an ultraviolet excess. We conclude that for F-type and early G-type stars high-velocity and ultraviolet excess are strongly correlated.

### References

- W. BECKER, 1963, *Basic Astronomical Data*, ed. K. Aa. Strand (Univ. of Chicago Press, Chicago) 241  
 W. BUSCOMBE and P. M. MORRIS, 1958, *Mem. Mount Stromlo Obs.* No. 14  
 A. W. J. COUSINS and R. H. STOY, 1963, *Roy. Obs. Bull.* No. 64  
 M. E. DIXON, 1965, *Mon. Not. Roy. Astr. Soc.* **129** 51  
 J. PONSEN and P. Th. OOSTERHOFF, 1966, *Bull. Astr. Inst. Netherlands Suppl.* 1 1  
 A. PRZYBYLSKI and P. MORRIS KENNEDY, 1965, *Mon. Not. Roy. Astr. Soc.* **129** 63  
 N. G. ROMAN, 1955, *Ap. J. Suppl.* **2** 195  
 A. R. SANDAGE and O. J. EGGEN, 1959, *Mon. Not. Roy. Astr. Soc.* **119** 283

Identification of Buscombe numbers with Henry Draper or Durchmusterung numbers

Busc.	HD or DM	Busc.	HD or DM	Busc.	HD or DM	Busc.	HD or DM
83	-61°282	502	88 284	1067	190 649	1209	212 231
88	10 607	522	91 889	1080	193 307	1220	214 385
95	11 262	531	93 751	1085	193 901	1237	217 276
127	-17°484	532	93 813	1090	196 378	1246	218 630
128	16 031	556	98 088	1131	201 099	1247	218 693
161	Ross 570	565	98 991	1150	203 448	1248	218 750
451	-3°2525	821	140 283	1159	204 670	1258	219 617
456	77 462	928	160 617	1163	-35°148 49	1266	220 536
460	78 558	934	-8°4501	1168	205 855	1276	222 013
461	78 670	954	165 896	1177	-41°146 56	1279	222 237
462	78 753	975	171 443	1186	208 998	1283	222 766
465	80 050	981	172 488	1197	210 918	1284	222 803
484	83 529	1057	189 340	1204	211 998	1311	224 927
493	-13°2948						