



Universiteit
Leiden
The Netherlands

Aria of the Dutch North Sea

Sertlek, H.O.; Sertlek H.O.

Citation

Sertlek, H. O. (2016, June 9). *Aria of the Dutch North Sea*. Retrieved from <https://hdl.handle.net/1887/40158>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/40158>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/40158> holds various files of this Leiden University dissertation

Author: Sertlek, Hüseyin Özkan

Title: Aria of the Dutch North Sea

Issue Date: 2016-06-09

References

A

- [1] A Marine Strategy Directive to save Europe's seas and oceans, http://ec.europa.eu/environment/water/marine/index_en.htm, last accessed 03/11/2010.
- [2] Aarts G. et al, (2012), Comparative interpretation of count, presence-absence and point methods for species distribution models. *Methods in Ecology and Evolution*, 3, 177–187.
- [3] Aarts G. et al. (submitted on 23-11-2015), Movement behaviour influences population level exposure to underwater sound: Impact of explosions on harbor porpoises, *Journal of Animal Ecology*.
- [4] Abramowitz M. and Stegun I., (1972), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, New York: Dover Publications.
- [5] Ainslie, M. A. , (2005). "Effect of wind-generated bubbles on fixed range acoustic attenuation in shallow water at 1–4kHz". *J. Acoust. Soc. Am.* , 118(6), 3513-3523.
- [6] Ainslie, M. A. et al, (2009). "Assessment of natural and anthropogenic sound sources and acoustic propagation in the North Sea." TNO report TNO-DV, C085.
- [7] Ainslie M.A., (2010a), *Principles of Sonar Performance Modeling*, Praxis Publishing, Chichester, UK.
- [8] Ainslie M.A., (2010b), Editorial: Validation of Sonar Performance Assessment Tools, in *Validation of Sonar Performance Assessment Tools (workshop held in memory of D. E. Weston, 7-9 April 2010)*, *Proceedings of the IOA*, Vol.32, Pt.2, pp.1-8.
- [9] Ainslie et al, (2011), An analytical solution for signal, background and signal to background ratio for a low frequency active sonar in a Pekeris waveguide satisfying Lambert's rule, *Proc. 4th International Conf. and Exhibition on Underwater Acoustic Measurements: Technologies and Results*, pp 491-498.
- [10] Ainslie M. A., (2011), Ad hoc European working group on terminology for underwater sound, "Standard for measurement and monitoring of underwater noise, Part I: Physical Quantities and

References

- Their Units”, TNO Report, TNO-DV 2011 C235,.Ainslie M.A. et al, (2012a), “What is the Source Level of Pile-Driving Noise in Water?”, The Effects of Noise on Aquatic Life, pp.445-448, Springer Media.
- [11] Ainslie, M.A. et al, (2012b), Dredger noise during Maasvlakte 2 construction: Noise maps and risk assessment, TNO report 2012 R10818
- [12] Ainslie M.A. et al, (2013), Echo, reverberation and echo to reverberation ratio for isovelocity waveguides related to Weston Memorial workshop Scenarios A2.I AND A2.IV.
- [13] Ainslie M.A. et al, (2014), Practical spreading laws: the snakes and ladders of shallow water acoustics, Proceedings 2nd International Conference and Exhibition on Underwater Acoustics, UA2014, 22-27 June, Rhodes, Greece, 879-886.
- [14] Ainslie M.A., Ellis D.D. and Harrison C.H., (2015), Low frequency bottom reverberation in a Pekeris waveguide with Lambert’s rule, Journal of Computational Acoustics, Vol. 23, 1550022
- [15] Akamatsu t. et al, (2005). New stereo acoustic data logger for free-ranging dolphins and porpoises. Marine Technology Society Journal, 38, 6-12.
- [16] Anon et al, 2014, “Draft joint workshop report: predicting sound fields; global soundscape modelling to inform management of cetaceans and anthropogenic noise”. Available from: “<https://events.iwc.int/index.php/scientific/SC65B/paper/viewFile/802/870/SC-65b-Rep03rev.pdf>”
- [17] APL-UW, (1994), APL-UW high-frequency ocean environmental acoustic models handbook(APL-UW TR 9407, AEAS 9501 October), Applied physics Laboratory, University of Washington, Seattle, WA
- [18] [AQUO], Information on the “Achieve quieter oceans by shipping noise footprint reduction (AQUO)” European project can be found on: <http://www.aquo.eu/>
- [19] [Aquo-Sonic (Madrid)], Meeting report in preparation.
- [20] Arveson P. T. and Vendittis D. J. , (2000). “Radiated noise characteristics of a modern cargo ship”. J. Acoust. Soc. Am. , 107(1), 118-129

B

- [21] Bailey H. et al, (2010), Assessing underwater noise levels during pile-driving at an offshore

References

- wind farm and its potential effects on marine mammals”, Mar. Pol. Bul. ,60 (6), p.888-897.
- [22] Bailey H. et al, (2014) Modelling harbour seal habitat by combining data from multiple tracking systems. Journal of Experimental Marine Biology and Ecology, 450, 30–39.
- [23] Borja A. et al, (2010) Marine management – Towards an integrated implementation of the European Marine Strategy Framework and the Water Framework Directives. Mar Poll Bull 60:2175–2186.
- [24] Borsani J.F., (2014), Impacts of noise and use of propagation models to predict the recipient side of noise, Cefas project report C6082
- [25] Buckland, S.T. et al, (2004) Advanced Distance Sampling. Oxford University press, New York.
- [26] Bouton N. et al, (2015), Hearing and water column use in the North Sea fishes: a review to serve exploration of variation in exposure to vessel sounds among species and species group, Sonic project report (draft), (<http://www.sonic-project.eu/>)

C

- [27] Camphuysen C. J. and Siemensma M. L., (2011). Conservation plan for the Harbour Porpoise *Phocoena phocoena* in The Netherlands: towards a favourable conservation status. (NIOZ Report 2011-07), Royal Netherlands Institute for Sea Research, Texel.
- [28] Camphuysen C. J. (2011). Recent trends and spatial patterns in near-shore sightings of harbour porpoises (*Phocoena phocoena*) in the Netherlands (Southern Bight, North Sea), 1990-2010. Lutra, 54, 39–47.
- [29] Camphuysen C.J. and Krop A., (2011), Maternal care, calf-training and site fidelity in a wild harbour porpoise in the North Sea. Lutra, 54, 123–126.
- [30] [Cet sound], Cet sound workshop report available from : <http://cetsound.noaa.gov/report.html>
- [31] Chapman N. R. (1985), Measurement of the waveform parameters of shallow explosive charges. J. Acoust. Soc. Am., 78, 672–681.
- [32] Clausen, K. T. et al , (2010), Click Communication In Harbour Porpoises *Phocoena Phocoena*. The International Journal Of Animal Sound And Its Recording, 20, 1–28.

References

- [33] Cole R. H., (1965). Underwater explosions. [Originally published in 1948 by Princeton University Press] (Dover, New York).
- [34] Collins M. D. and Evans R. B., (1992), A two-way parabolic equation for acoustic backscattering in the ocean, *J. Acoust. Soc. Am.* 91 ,1357–1368.
- [35] Collins M. D., (1999), RAM user guide, <http://ram.nrl.navy.mil/>
- [36] Collins M. D., (1993), A Split-Step Pade Solution for the Parabolic Equation Method, *J. Acoust. Soc. Am.* 93, 1736-1742,
- [37] Commission Decision No. 2010/477/EU on criteria and methodological standards on good environmental status of marine waters, (2010), *O. J. L* 232/14.
- [38] Culik B. et al, (2001). Reactions of harbor porpoises *Phocoena phocoena* and herring *Clupea harengus* to acoustic alarms. *Mar. Ecol. Prog. Ser.* 211, 255-260.

D

- [39] Dähne M. et al , (2013), Effects of pile-driving on harbour porpoises (*Phocoena phocoena*) at the first offshore wind farm in Germany, *Environmental Research Letters*, 8, 025002.
- [40] Daly JN and Harrison J., (2012), The Marine Mammal Protection Act: a regulatory approach to identifying and minimizing acoustic-related impacts on marine mammals, *Adv. Exp. Med Biol.* 730:537-9,
- [41] Danil K. and St. Leger J. A. , (2011). Seabird and dolphin mortality associated with underwater detonation exercises. *Marine Technology Society Journal*, 6, 89-95.
- [42] de Jong C.A.F. et al , (2010), “Underwater noise of Trailing Suction Hopper Dredgers at Maasvlakte 2: Analysis of source levels and background Noise”. TNO-DV report 2010 C335.
- [43] Denham R.N., Intensity decay laws for near-surface sound sources in the ocean, *J. Acoust. Soc. Am.* 79, 60 (1986)
- [44] Dekeling et al, (2014a), Monitoring Guidance for Underwater Noise in European Seas, Part I: Executive Summary, J.V. JRC Scientific and Policy Report EUR 26557 EN, Publications Office of the European Union, Luxembourg, (<http://publications.jrc.ec.europa.eu/repository/handle/111111111/30979>)

References

- [45] Dekeling et al, (2014b), Monitoring Guidance for Underwater Noise in European Seas, Part II: Monitoring Guidance Specifications, JRC Scientific and Policy Report EUR 26555 EN, Publications Office of the European Union, Luxembourg, <http://publications.jrc.ec.europa.eu/repository/handle/111111111/30973>)
- [46] Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (marine strategy framework directive), (2015), (<http://eur-lex.europa.eu/legal-content/en/txt/?uri=celex:32008l0056>”, accessed 18 september 2015).
- [47] Duren R E, 1988, A theory for marine source arrays, *Geophysics* 53 650–8
- [48] Dreschler J. et al ,(2009), “Measurements of underwater background noise Maasvlakte 2”, TNO-DV 2009 C212
- [49] Dragoset B., “Introduction to Air Guns and Air-Gun Arrays,” *The Leading Edge*, pp.892-897, (August 2000)

E

- [50] [EC, 2010] Commission Decision No. 2010/477/EU on criteria and methodological standards on good environmental status of marine waters, 2010 O. J. L 232/14.
- [51] Ellison, W. T. et al, (2012). A New Context-Based Approach to Assess Marine Mammal Behavioral Responses to Anthropogenic Sounds. *Conservation Biology*, 1, 21-28.
- [52] Endangered Species Act of 1973, “<http://www.fws.gov/laws/lawsdigest/ESACT.html>”, last accessed 25 April 2012.
- [53] Emodnet, 2014, public Bathymetry database available online at: <http://www.emodnet-hydrography.eu/>)
- [54] Erbe, C. and King, A. (2009), Modeling cumulative sound exposure around marine seismic surveys, *J. Acoust. Soc. Am.* ,125 (4): pp. 2443-2451.
- [55] Erbe C. et al, (2012), Mapping cumulative noise from shipping to inform marine spatial planning. , *J. Acoust. Soc. Am.* , 132 (5): pp. 432-428.
- [56] Erbe C. et al, (2014), Identifying Modeled Ship Noise Hotspots for Marine Mammals of Canada’s Pacific Region. *PLoS ONE*. 9 (3): e89820 .

References

[57] Etopo, 2014, public Bathymetry database available online:<http://www.ngdc.noaa.gov/mgg/global/global.html>

[58] Evans R.B., (1983), A coupled mode solution for acoustic propagation in a waveguide with stepwise depth variations of a penetrable bottom," J. Acoust. Soc. Am. 74, 188-195.

F

[59] Fay, R. R. (2009), Soundscapes and the sense of hearing of fishes. Integr. Zool. 4,26–32.

[60] Finneran J. J. and Schlundt C. E., (2011). Subjective loudness level measurements and equal loudness contours in a bottlenose dolphin (*Tursiops truncatus*). J. Acoust. Soc. Am., 130(5), 3124-3136.

[61] Finneran, J. J. and Jenkins A. K. ,(2012). Criteria and thresholds for U.S. Navy acoustic and 51 explosive effects analysis, San Diego, California: SPAWAR Systems Center Pacific, April 2012, pp. 64.

[62] Finneran J. J. and Schlundt C. E. (2013). Effects of fatiguing tone frequency on temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*). J. Acoust. Soc. Am. , 133(3), 1819-1826.

[63] Fisher, F. H., and Simmons, V. P. (1977), Sound absorption in seawater. J. Acoust. Soc. Am. , 62(3): 558–564.

[64] Fisheries Hydroacoustic Working Group (FHWG),"Agreement in Principle for Interim Criteria for Injury to Fish from Pile Driving Activities",12 June 2008 (Vancouver, Canada, 2008), http://www.dot.ca.gov/hq/env/bio/fisheries_bioacoustics.htm, last accessed 22 April 2012.

[65] Fletcher H. and Munson W. A. ,(1933). Loudness, its definition, measurement and calculation. J. Acoust. Soc. Am. 5, 82-108. 10.1121/1.1915637.

[66] Folegot T, (2012), Ship traffic noise distribution in the Strait of Gibraltar: an exemplary case for monitoring global ocean noise using real-time technology now available for understanding the effects of noise on marine life, Adv Exp Med Biol. 2012;730:601-4.

G

[67] Gauss, R. C. et al, (2002). Broadband models for predicting bistatic bottom, surface, and

References

volume scattering strengths (No. NRL/FR/7100--02-10). NAVAL RESEARCH LAB WASHINGTON DC.

[68] Gebco(+), 2014, public Bathymetry database available online at: <http://www.gebco.net/>

[69] Geelhoed S. et al, (2012). Shortlist Masterplan Wind Aerial surveys of harbour porpoises on the Dutch Continental Shelf (IMARES Report number C103/11). Den Burg, Texel.

[70] Geelhoed, S.C.V. et al, (2013) Abundance of harbour porpoises (*Phocoena phocoena*) on the Dutch Continental Shelf, aerial surveys in July 2010-March 2011. *Lutra*, 56, 45–57.

[71] Geological Survey of the Netherlands , DINO database, (<http://www.geologicalsurvey.nl>)

[72] Gray L. M., and Greeley D. S. , (1980), Source level model for propeller blade rate radiation for the world's merchant fleet, *J. Acoust. Soc. Am.* , 67(2), 516-522.

[73] Groot-Hedlin C., (2004), Criteria for discretization of seafloor bathymetry when using a staircase approximation: application to computation of T-phase seismograms, *J. Acoust. Soc. Am.*, 115 (3), 1103-1113.

H

[74] Haelters J, Dulière V, Vigin L, Degraer S (2015) Towards a numerical model to simulate the observed displacement of harbour porpoises *Phocoena phocoena* due to pile driving in Belgian waters. *Hydrobiologia* 756:105–116

[75] Hall V.H, (2015), An analytical model for the underwater sound pressure waveforms radiated when an offshore pile is driven, *J. Acoust. Soc. Am.* , 138 (2): pp. 795-806.

[76] Hammond P.S. et al, (2013) Cetacean abundance and distribution in European Atlantic shelf waters to inform conservation and management. *Biological Conservation*, 164, 107–122.

[77] Hansen M. et al, (2008). Low-frequency components in harbour porpoise (*Phocoena phocoena*) clicks: communication signal, by-product, or artifacts? *J. Acoust. Soc. Am.* 124, 4059–4068.

[78] Harrison C.H., (2005), Closed form bistatic reverberation and target echoes with variable bathymetry and sound speed, *IEEE J. Oceanic Eng.* 30, 660–675.

[79] Harrison C.H. and Nielsen P.L. , (2007), "Multipath pulse shapes in shallow water: Theory and simulation", *J. Acoust. Soc. Am.*, 121 (3), p.1362-1373

References

- [80] Harrison C. H. and Ainslie M. A. ,(2010), Fixed time versus fixed range reverberation calculation: Analytical solution, *J. Acoust. Soc. Am.* 128,28–38
- [81] Harrison C.H. , (2010), An approximate form of the Rayleigh reflection loss and its phase: application to reverberation calculation. *J Acoust Soc Am.*, 128(1), 50-7
- [82] Harrison, C. H. ,(2013). Ray convergence in a flux-like propagation formulation. *J. Acoust. Soc. Am.* , 133(6), 3777-3789.
- [83] Harwood J. et al, (2013). A protocol for implementing the interim population consequences of disturbance (PCOD) approach: quantifying and assessing the effects of UK offshore renewable energy developments on marine mammal populations (report SMRUL-TCE-2013-014 Scottish Marine and Freshwater Science 5(2)).
- [84] Hatch LT and Fristrup KM, (2009), No barrier at the boundaries: implementing regional frameworks for noise management in protected natural areas, *Marine Ecology Progress Series* 395, 223-244
- [85] Heaney K. D. and Campbell R. L., (2015), Three-dimensional parabolic equation modelling of mesoscale eddy deflection, *J. Acoust. Soc. Am.*, submitted August 11 2015.
- [86] Heinis F. et al, (2013), “Monitoring programme for the Maasvlakte 2, part III—the effects of underwater sound”
- [87] Hiby L. and Lovell P. (1998) Using Aircraft in Tandem Formation to Estimate Abundance of Harbour Porpoise. *Biometrics*, 54, 1280–1289.
- [88] Hildebrand J.A., (2009), Anthropogenic and natural sources of ambient noise in the ocean”, *Mar. Ecol. Prog. Ser.*, 395: p.5–20.
- [89] Holland C. W., (2010), Propagation in a waveguide with range dependent seabed properties, *J. Acoust. Soc. Am.* 128 (5), 2596-2607.
- [90] Horne B van (1983) Density as a Misleading Indicator of Habitat Quality. *J Wildl Manage* 47:893–901
- [91] Huijbers CM et al, (2008) Shallow patch reefs as alternative habitats for early juveniles of some mangrove/seagrass-associated fish species in Bermuda. *Revista de Biologia Tropical* 56 (Suppl. 1): 161-169

I

- [92] IEC 1995 (EN 61260), Electroacoustics - Octave-band and fractional-octave-band filters, International Electrotechnical Commission, Geneva, Switzerland, 1996.
- [93] IEC 60050:1994, International Electrotechnical Vocabulary, part 801: Acoustics and Electroacoustics, (section 801-32 covers terms for underwater acoustics), International Electrotechnical Commission (IEC), Geneva.
- [94] IEC60565: 2006 Underwater acoustics-Hydrophones - Calibration in the frequency range 0.01 Hz to 1 MHz, IEC 60565 - 2006 (EN 60565: 2007, BS60565:2007), International Electrotechnical Commission, Geneva, 2006
- [95] IJsseldijk L.L. et al, (2015) Going with the flow: tidal influence on the occurrence of the harbour porpoise (*Phocoena phocoena*) in the Marsdiep area, The Netherlands. *Journal of Sea Research*, 103,129–137.
- [96] ISO/DIS 18405,Underwater acoustics terminology, International Organization for Standardization, Geneva, Switzerland
- [97] ISO/CD 18405 Underwater acoustics terminology, International Organization for Standardization, Geneva, Switzerland
- [98] [ISO - GUM], JCGM 100 series, Guides to the expression of uncertainty in measurement (GUM series). Available from: <http://www.iso.org/sites/JCGM/GUM-introduction.htm>
- [99] ISO 266:1997, Acoustics - Preferred frequencies (1997)
- [100] ISO 80000-8:2007, Quantities and units – part 8: Acoustics, International Organization for Standardisation (ISO), Geneva.

J

- [101] Jackson, D. R. (1994). APL-UW high-frequency ocean environmental acoustic models handbook, Applied Physics Laboratory, University of Washington, Technical Report, 9407.
- [102] Jensen F.B. and Ferla C.M.,(1990), Numerical solutions of range-dependent benchmark problems in ocean acoustics, *J. Acoust. Soc. Am*, 87 (4) , 1499-1510.
- [103] Jensen F.B. et al. ,(1994), *Computational Ocean Acoustics*, American Institute of Physics, New

References

York

[104] Jackson D.R., (1994). "APL-UW high-frequency ocean environmental acoustic models handbook". Applied Physics Laboratory, University of Washington, Technical Report, 9407.

[105] Jones AR et al, (2014) Fine-scale hydrodynamics influence the spatio-temporal distribution of harbour porpoises at a coastal hotspot. *Prog Oceanogr* 128:30–48

K

[106] Kastelein, R.A. (1997) Food Consumption and Growth of Marine Mammals. Landbouwniversiteit Wageningen, the Netherlands.

[107] Kastelein, R. A. et al, (2002). Audiogram of a harbor porpoise (*Phocoena phocoena*) measured with narrow-band frequency-modulated signals. *J. Acoust. Soc. Am.* , 112, 334-344.

[108] Kastelein, R.A. et al , (2007) The influence of signal parameters on the sound source localization ability of a harbor porpoise (*Phocoena phocoena*). *J. Acoust. Soc. Am.* , 122, 1238–1248.

[109] Kastelein, R. A. et al, (2012a). Temporary threshold shifts and recovery in a harbor porpoise (*Phocoena phocoena*) after octave-band noise at 4 kHz . *J. Acoust. Soc. Am.* 132 , 3525.

[110] Kastelein, R. A. et al, (2012b). The hearing threshold of a harbor porpoise (*Phocoena phocoena*) for impulsive sounds (L). *J. Acoust. Soc. Am.* , 132 (2), 607-610.

[111] Kastelein, R. A. et al, (2013). Hearing frequency thresholds of a harbour porpoise (*Phocoena phocoena*) temporarily affected by a continuous 1.5 kHz tone. *J. Acoust. Soc. Am.* , 134, 2286-2292.

[112] Kastelein, R. A. et al, (2014a). Effect of level, duration, and inter-pulse interval of 1-2 kHz sonar signal exposures on harbour porpoise hearing. *J. Acoust. Soc. Am.* , 136, 412-422.

[113] Kastelein, R. A. et al, (2014b). Frequency of greatest temporary hearing threshold shift in harbor porpoises (*Phocoena phocoena*) depends on the noise level. *J. Acoust. Soc. Am.* , 136, 1410-1418.

[114] Kastelein, R. A. et al, (2015a) Hearing frequencies of a harbor porpoise (*Phocoena phocoena*) temporarily affected by played back offshore pile-driving sounds. *J. Acoust. Soc. Am.* , 137, 556.

[115] Kastelein, R. A. et al, (2015b). Effects of exposure to intermittent and continuous 6–7 kHz sonar sweeps on harbor porpoise (*Phocoena phocoena*) hearing. *J. Acoust. Soc. Am.* , 137 (4),

References

1623–1633.

[116] Ketten D. R. et al, (1993). Blast injury in humpback whale ears: Evidence and implications. *J. Acoust. Soc. Am.* , 94(3/2), 1849-1850.

[117] Ketten D. R. (2004). Experimental Measures of Blast and Acoustic Trauma in Marine Mammals. (ONR Final Report: N000149711030).

[118] Ketten D. R. (2012). Marine Mammal Auditory System Noise Impacts: Evidence and Incidence. In A. N. Popper & A. Hawkins (Eds.) *The Effects of Noise on Aquatic Life*. New York: Springer.

[119] King, S.L. et al (2015) An interim framework for assessing the population consequences of disturbance (ed C Kurle). *Methods in Ecology and Evolution*, 6, n/a–n/a.

[120] Kornhauser E.T. and Raney W.P.,(1955) Attenuation in shallow-water propagation due to an absorbing bottom, *J. Acoust. Soc. Am*, 27(4), 689-692.

[121] Koschinski S. and Koch K. H., (2009), Underwater Unexploded Ordnance – Methods for a Cetacean-friendly Removal of Explosives as Alternatives to Blasting, IWC, Scientific Committee Paper SC/61 E:1-13.

[122] Koschinski S. (2011). Underwater noise pollution from munitions clearance and disposal, possible effects on marine vertebrates, and its mitigation. *Marine Technology Society Journal*, 45, 80-88.

[123] Kryter K. D. et al, (1966). Hazardous exposure to intermittent and steady state noise. *J. Acoust. Soc. Am.* , 39, 451-464.

[124] Kryter K. D., (1994). *The handbook of hearing and the effects of noise*. New York: Academic Press. 673

[125] Kujiwa S. G., and Liberman M. C. (2009). Adding insult to injury: cochlear nerve degeneration after “temporary” noise-induced hearing loss. *J. Neurosci.* 20, 14077-14085.

[126] Kuperman W.A. and Ferla M.C., (1985) ,A shallow water experiment to determine the source spectrum level of wind-generated noise, *J. Acoust. Soc. Am.* ,77: pp. 2067-2073.

[127] KNMI, The Royal Netherlands Meteorological Institute (KNMI) Databases <http://www.knmi.nl>

L

References

- [128] Laws R.M. et al.,1990,Computer modeling of clustered air guns: First Break, 18(9), 331–338
- [129] Lewis, J. A. (1996). Effects of underwater explosions on life in the sea. Melbourne, Victoria: Dept. of Defence, Defence Science and Technology Organisation, 38 pp.
- [130] Li Guo-Fa et al,2010, Modeling air gun signatures in marine seismic exploration considering multiple physical factors, Applied Geophysics, Vol.7, No.2 , P. 158 – 165
- [131] Lippert et al,2015, COMPILE - A Generic Benchmark Case for Predictions of Marine Pile Driving Noise", accepted for publication by IEEE J Oceanic Eng, December 2015.
- [132] Lucke K. et al (2009). Temporary shift in masked hearing thresholds in a harbor porpoise (*Phocoena phocoena*) after exposure to seismic airgun stimuli. J. Acoust. Soc. Am. , 125 (6), 4060-4070.
- [133] Lucke et al, (2014), Report of the Workshop on International Harmonisation of Approaches to Define Underwater Noise Exposure Criteria (Budapest, Hungary 17th August 2013)
- [134] Lurton, X. (2002). An introduction to underwater acoustics: principles and applications. springer.
- [135] Lusseau D. et al (2012). Assessing the risks to marine mammal populations from renewable energy devices – an interim approach. Final Workshop report, 29 pp.

M

- [136] [Maasvlakte 2, 2014], Website on the extension of the Rotterdam port <https://www.maasvlakte2.com/en/index/>
- [137] MacGillivray A.O., (2006), An acoustic study of seismic airgun noise in Queen Charlotte Basin, MSc Thesis.
- [138] MacGillivray A.O. and Chapman N.R., (2012), Modeling Underwater Sound propagation from an airgun array using the parabolic equation method, Canadian Acoustics, Vol.40, No.1, 19-25 .
- [139] Macpherson and Daintith, (1967)Practical model of shallow-water acoustic propagation, J. Acoust. Soc. Am., 41(4), 850-854.
- [140] Madsen, P. A, (2005). Marine mammals and noise: Problems with root mean square sound pressure levels for transients. J. Acoust. Soc. Am. , 117(6), 3952–3957.

References

- [141] Marine Mammal Protection Act (MMPA) of 1972, "<http://www.nmfs.noaa.gov/pr/laws/mmpa/>", last accessed 25 April 2012.
- [142] Malme C.I. et al, (1983), Investigations of the potential effects of underwater noise from petroleum industry activities on migrating gray whale behaviour. Bolt Beranek & Newman Report Number 5366, Cambridge, Massachusetts, Bolt Beranek and Newman Inc.
- [143] McCarthy E. et al, (2011) Changes in spatial and temporal distribution and vocal behavior of Blainville's beaked whales (*Mesoplodon densirostris*) during multiship exercises with mid-frequency sonar. *Marine Mammal Science*, 27, E206–E226.
- [144] Mennitt D. et al, (2014), A geospatial model of ambient sound pressure levels in the contiguous United States, *J. Acoust. Soc. Am.* 135 (5), 2746–2764
- [145] Møhl B and Andersen S. , (1973). Echolocation: high-frequency component in click of harbor porpoise (*Phocoena-Ph. L.*). *J. Acoust. Soc. Am.* 54, 1368–1372.
- [146] Mobley Curtis D., (1994), *Light and Water: Radiative Transfer in Natural Waters*, Academic Press; 0125027508
- [147] Montgomery et al. 2006; Sound as an orientation cue for the pelagic larvae of reef fishes and decapod crustaceans. *Adv. Mar. Biol.* 2006;51:143–96.
- [148] McLennan DA (2003), The importance of olfactory signals in the gasterosteid mating system: sticklebacks go multimodal, *Biol J Linnean Soc*, 80:555–572.
- [149] Myrberg AA Jr, Fuiman LA. 2002. The sensory world of reef fishes. *Coral reef fishes*. New York: Elsevier Science. p 123–160.

N

- [150] Nabe-Nielsen, J. et al, (2013) How a simple adaptive foraging strategy can lead to emergent home ranges and increased food intake. *Oikos*, 122, 1307–1316.
- [151] Nabe-Nielsen, J. et al, (2014). Effects of noise and by-catch on a Danish harbour porpoise population. *Ecological Modelling*, 272, 242–251
- [152] National Research Council, (2003), *Ocean noise and marine mammals*, Ocean Study Board,

References

Washington, DC: The National Academies Press.

[153] New, L. et al, (2014), Using short-term measures of behaviour to estimate long-term fitness of southern elephant seals. *Marine Ecology Progress Series*, 496, 99–108.

[154] Nowacek, D.P. et al, (2007), Responses of cetaceans to anthropogenic noise, *Mammal Review*, 37, 81-115.

[155] [NPL GPG 133 2014], Good Practice Guide for Underwater Noise Measurement, National Measurement Office, Marine Scotland, The Crown Estate, Robinson, S.P., Lepper, P. A. and Hazelwood, R.A., NPL Good Practice Guide No. 133, 2014.

[156] Nützel, B. (2008). Untersuchungen zum Schutz von Schweinswalen vor Schockwellen. (Forschungsanstalt der Bundeswehr für Wasserschall und geophysik (FWG), Technischer Bericht TB 2008-7).

O

[157] Ocean Acoustic Library (OALIB), (2013), <http://oalib.hlsresearch.com>, last accessed 31 March 2013

[158] Olbers H.W.M., 1826, Ueber die Durchsichtigkeit des Weltraums in Bode's Jahrbuch für 1826, pp. 110–121.

[159] OSPAR Commission (2010). Quality status report, Assessment of the impact of dumped conventional and chemical munitions. OSPAR Commission, London, U.K.

[160] Otani, S. et al, (2001) Oxygen consumption and swim speed of the harbor porpoise *Phocoena phocoena*. *Fisheries Science*, 67, 894–898.

P

[161] Parks SE et al, (2005), The gunshot sound produced by male north Atlantic right whales (*Eubalaena glacialis*) and its potential function in reproductive advertisement", *Marine Mammal Science* 21, 458-475

[162] Pfeiffer, F. (2014). Changes in Properties of Explosives Due to Prolonged Seawater Exposure.

References

Marine Technology Society Journal, 102-110.

[163] Poppe P. M. and Wijers C. M. J., ACM, (1990), Transactions on Mathematical Software 16, 38-46,

[164] Popper, A. N., and Hastings, M. C. (2009). The effects of anthropogenic sources of sound on fishes. *J. Fish Biol.* 75, 455–489.

[165] Porter M. B., (1990), The KRAKEN Normal Mode Program, SACLANT Undersea Research Centre Tech. Rep., 1990.

[166] Porter M.B., Jensen F.B. and Ferla C.M., (1991), The problem of energy conservation in one-way models, *J. Acoust. Soc. Am.* 89, 1058-1067.

[167] Porter M. B. and Henderson L. J., (2013), Global Ocean Soundscapes, Proceedings of the International Congress on Acoustics 2013, Vol. 19, 010050 (Proceedings of Meetings on Acoustics) Montreal, Canada.

[168] Porter M.B., (2013), Bellhop Manual and User Guide, <http://oalib.hlsresearch.com/Rays/>, last accessed 31.03.2013

[169] Preston J.R. and Ellis D.D., (2012), Adiabatic normal mode reverberation and clutter modelling. Proc. of 11th European Conference on Underwater Acoustics (ECUA), 239-246, Edinburgh, UK.

[170] Prosperetti A., (1984), Bubble phenomena in sound fields: Part one, *Ultrasonics* (22), 69-77

[171] Prosperetti A. et al, (1988), Nonlinear bubble dynamics, *J. Acoust. Soc. Am.* 83(2), 502-514

R

[172] Radford C.A. et al , (2011), Modelling a reef as an extended sound source increases the predicted range at which reef noise may be heard by fish larvae, *Mar. Ecol. Prog. Ser.* , Vol. 438, 167-174

[173] Radford, A. N., Kerridge, A., and Simpson, S. D. (2014). Acoustic communication in a noisy world: can fish compete with anthropogenic noise? *Behav. Ecol.* 25, 1022–1030.

[174] [RD Coordinates], <https://nl.wikipedia.org/wiki/Rijksdriehoekskoordinaten>

References

- [175] Reeder D.M. and Kramer K.M. (2005) ,Stress in free-ranging mammals: integrating physiology, ecology, and natural history. *Journal of Mammalogy*, 86, 225–235.
- [176] Reinhall P.G. and Dahl P.H., Underwater Mach wave radiation from impact pile driving: Theory and observation, *J. Acoust. Soc. Am.* , 130 (3), p.1209-1216 , (2011).
- [177] Richardson W. J et al, (1995). *Marine Mammals and Noise*. Academic Press, San Diego, CA, 576 pp.
- [178] Robertson J. S., (1999), Sound propagation over a large wedge a comparison between the geometrical theory of diffraction, *J. Acoust. Soc. Am.* 106 (1), 113-119.
- [179] Rossington, K. et al, (2013) Eco-hydro-acoustic modeling and its use as an EIA tool. *Marine pollution bulletin*, 75, 235–43.
- [180] Ryan K.L.,(2009), NATO Undersea Research Centre Marine Mammal Risk Mitigation Rules and Procedures, NURC-SP-2009-02

S

- [181] Scheidat, M. et al , (2008) Harbour porpoise *Phocoena phocoena* abundance in the southwestern Baltic Sea. *Endangered Species Research*, 5, 215–223.
- [182] Scheidat, M. et al ,(2012) Using aerial surveys to estimate density and distribution of harbour porpoises in Dutch waters. *Journal of Sea Research*, 69, 1-7.
- [183] Schecklman, S. et al, 2015, A computational method to predict and study underwater noise due to pile driving , *J. Acoust. Soc. Am.* , 138(1), 258-266.
- [184] Schuknecht, H.F. (1993). *Pathology of the Ear*, 2nd edn., Lea-Felbiger, Philadelphia.
- [185] [Scrimger & Heitmeyer, 1989], Scrimger, P., & Heitmeyer, R. M. (1991). Acoustic source-level measurements for a variety of merchant ships. *J. Acoust. Soc. Am.* , 89(2), 691-699.
- [186] Slabbekoorn H and Bouton N., (2008), Soundscape orientation: a new field in need of sound investigation. *Animal Behaviour* 76: e5-e8.
- [187] Slabbekoorn, H et al, 2010, A noisy spring: the impact of globally rising underwater sound levels on fish. *Trends Ecol. Evol.* 25, 419–427.

References

- [188] Sertlek H.O. and Aksoy S., (2010), An Alternative Approach for Treatment of Lossy Seabed In Fluid-Solid Waveguide Problems, ECUA 2010, Istanbul, Turkey.
- [189] Sertlek H.Ö, Aksoy S., (2012), Benchmarking Of Acoustic Pulse Propagation Problems In An Isovelocity Waveguide By An Analytical Time Domain Normal Mode Method, 11th European Conference on Underwater Acoustics (ECUA), Edinburgh, The Scotland, (2-6 July 2012)
- [190] Sertlek H.O., Ainslie M.A. (2013), "Propagation loss model comparisons on selected scenarios from the Weston memorial workshop." UA2013 1st international Conference and Exhibition on underwater acoustics.
- [191] Sertlek, H. Ö., & Ainslie, M. A. (2014a). A depth-dependent formula for shallow water propagation. J. Acoust. Soc. Am. , 136(2), 573-582.
- [192] Sertlek H.O., Ainslie M.A. (2014b), "A fast algorithm for the computation of incoherent propagation loss for variable water depth: a validation study" UA2014 2nd International Conference and Exhibition on underwater acoustics.
- [193] Sertlek, H. Ö. et al, (under review), Range-dependent propagation equations and results for Pekeris waveguide, under review in JASA.
- [194] Sertlek et al,(2015b), Mapping underwater sound in the Dutch part of the North Sea, Vol.875 , Chap.124, Advances in Experimental Medicine and Biology, pp 1001-100
- [195] Slabbekoorn H. et al, (2010)., "A noisy spring: the impact of globally rising underwater sound levels on fish", Trends Ecol Evol. , 25(7), 419-27
- [196] Soloway A. G. and Dahl P. H., (2014). Peak sound pressure and sound exposure level from underwater explosions in shallow water. J. Acoust. Soc. Am., 136, 3, EL219-EL223.
- [197] [SONIC], Information on the "Suppression Of underwater Noise Induced by Cavitation (SONIC)" European project can be found on: <http://www.sonic-project.eu/>
- [198] Southall B. L. et al, (2007), Marine Mammal Noise Exposure Criteria: Initial Scientific Recommendations, Aquatic Mammals, 33 (4), pp. 411-509.
- [199] Stevens, M. (2013), Sensory Ecology, Behaviour, and Evolution, Oxford University Press.
- [200] Sutton et al, (2011), Mapping the Spatio-temporal Distribution of Underwater Noise in Irish Waters, EPA STRIVE Report, 2011-W-MS-7, (<http://erc.epa.ie/safer/reports>, last accessed on 09/12/2015)

References

[201] Suzuki, Y. and Takeshima, H. (2004). Equal loudness-level contours for pure tones. *J. Acoust. Soc. Am.* 116, 918-933.

T

[202] Teilmann, J. et al, (2013) Geographic, seasonal, and diurnal surface behavior of harbor porpoises. *Marine Mammal Science*, 29, E60–E76.

[203] Thompson, P.M. et al, (2013a) Short-term disturbance by a commercial two-dimensional seismic survey does not lead to long-term displacement of harbour porpoises. *Proceedings. Biological sciences / The Royal Society*, 280, 20132001.

[204] Thompson P.M. et al, (2013b) Framework for assessing impacts of pile-driving noise from offshore wind farm construction on a harbour seal population. *Environmental Impact Assessment Review*, 43, 73–85.

[205] Thorp, W. H. (1967). Analytic Description of the Low-Frequency Attenuation Coefficient. *J. Acoust. Soc. Am.* , 42(1), 270-270.

[206] Thorsos E.I. and Perkins J.S., (2007), Overview of the reverberation modeling workshops, *J. Acoust. Soc. Am.* Volume 122, Issue 5, pp. 3074-3074.

[207] Tindle C.T., (1979), The equivalence of bottom loss and mode attenuation per cycle in underwater acoustic, *J. Acoust. Soc. Am.*, 66(1) , 250-255.

[208] Tindle C.T. and Weston D.E., (1980), Connection of acoustic beam displacement, cycle distances and attenuations for rays and normal modes, *J. Acoust. Soc. Am.*, 67 , 1614-1622.

[209] Tougaard J et al, (2009), “Pile driving zone of responsiveness extends beyond 20 km for harbor porpoises (*Phocoena phocoena* (L.))”, *J. Acoust. Soc. Am.* 126, pp.11-14.

[210] Tougaard J. et al, (2009). Pile-driving zone of responsiveness extends beyond 20 km for harbour porpoises (*Phocoena phocoena*, (L.)). *J. Acoust. Soc. Am.* , 126(1), 11-14.

[211] Tougaard J. et al, (2014) Cetacean noise criteria revisited in the light of proposed exposure limits for harbour porpoises. *Marine Pollution Bulletin*, 90, 196–208.

[212] Tougaard J. et al, (2015). Cetacean noise criteria revisited in the light of proposed exposure limits for harbour porpoises. *Marine Pollution Bulletin*, 90, 1–2, 196-208.

[213] Trees et al. 2005, Monitoring water transparency and diver visibility in ports and harbors

References

using aircraft hyperspectral remote sensing , Proc. SPIE 5780, Photonics for Port and Harbor Security, 91 (May 25, 2005)

[214] Tyack, P. L. (2008). Implications for marine mammals of large-scale changes in the marine acoustic environment. *Journal of Mammalogy*, 89(3), 549–558.

[215] Tyack P. et al, (2015), International Quiet Ocean Experiment plan, (http://www.scor-int.org/IQOE/IQOE_Science_Plan-Final.pdf, Last accessed 09/12/2015)

U

[216] Urlick R.J., “Principles of underwater sound for engineers”, McGraw-Hill, New York, (1975).

V

[217] Van Horne B., (1983) Density as a Misleading Indicator of Habitat Quality. *The Journal of Wildlife Management*, 47, 893–901.

[218] Verboom W. C. and Kastelein R. A. (1997). Structure of harbour porpoise (*Phocoena phocoena*) click train signals. In: *The biology of the Harbour porpoise* (Eds. Read, A.J., Wiepkema, P.R. and Nachtigall, P.E.), De Spil Publishers, Woerden, The Netherlands, 343-362.

[219] Verboom, W. C. and Kastelein, R. A., (2005). Some examples of marine mammal discomfort thresholds in relation to man-made noise. *Proceedings UDT 2005*, Amsterdam, June 2005.

[220] Von Benda-Beckmann A.M. et al, (2014). Assessment of impact of underwater clearance of historical explosives by the Royal Netherlands Navy on harbour porpoises in the North Sea, TNO Report, TNO 2014 R10916.

[221] Von Benda-Beckmann A. M. et al, (2015), Assessing the Impact of Underwater Clearance of Unexploded Ordnance on Harbour Porpoises (*Phocoena phocoena*) in the Southern North Sea, *Aquatic Mammals*, 2015, 41(4), 503-523.

W

[222] Wales S.C. and Heitmeyer R.M.(2002), An ensemble source spectra model for merchant ship-

References

radiated noise , J. Acoust. Soc. Am., Vol 111, Issue 3: 1211-1231

[223] Wang L. et al, (2014), "review of underwater acoustic propagation models", NPL report AIR (RES) 086

[224] Wenz. G.M., (1962), Acoustic ambient noise in the ocean: Spectra and sources, J. Acoust. Soc. Am. vol. 34, p. 1936–1956.

[225] Weston D.E.,(1959), Guided propagation in a slowly varying medium, Proc.Phys.Soc., 73,365–384

[226] Weston, D. E. (1960), Underwater explosions as acoustic sources, Proc.Phys.Soc., 76, 233-249.

[227] Weston D.E., (1960), A Moiré fringe analog of sound propagation in shallow water, J. Acoust. Soc. Am. 32-6, 647-654.

[228] Weston, D. E. (1962), Explosive sources in Underwater Acoustics (Plenum Press, New York), pp 51-66.

[229] Weston D.E., (1971),Intensity-range relations in oceanographic acoustics, J. Sound Vib.,18, pp.271-287

[230] Weston D.E., (1976), Propagation in water with uniform sound velocity but variable-depth lossy bottom, J. Sound Vib., 47, pp.473-483.

[231] Weston D.E. and Tindle C.T., (1979),Reflection loss and mode attenuation in a Pekeris model, J. Acoust. Soc. Am., 66(3) , 872-879

[232] Weston D. E., (1980), Acoustic flux methods for oceanic waveguides, J.Acoust.Soc.Am.,68(1), 287-296.

[233] Weston D.E. ,(1980), Wave theory peaks in range averaged channels of uniform sound velocity, J. Acoust. Soc. Am. ,68 (1), 282 -286

[234] Weston D. E. and Ching P. A., (1989). Wind effects in shallow-water acoustic transmission. J. Acoust. Soc. Am. , 86(4), 1530-1545.

[235] Weston D.E., (1991), Diffusion-like growth of decoupling and enhancement depths in shallow-water acoustic propagation, Proc. I.O.A. 13 (3), 85-92.

[236] Weston D.E., (1994), Wave shifts, beam shifts, and their role in modal and adiabatic propagation, J. Acoust. Soc. Am. 96, 406-416

References

[237] World Ocean Atlas 2013 (WOA13) is a set of objectively analyzed (1° grid) climatological fields of in situ temperature, salinity, dissolved oxygen, Apparent Oxygen Utilization (AOU), percent oxygen saturation, phosphate, silicate, and nitrate at standard depth levels for annual, seasonal, and monthly compositing periods for the World Ocean. Available from: <http://www.nodc.noaa.gov/OC5/woa13/>

[238] Wensveen, P. J. et al, (2014), Equal latency contours and auditory weighting functions for the harbour porpoise (*Phocoena phocoena*). *The Journal of Experimental Biology*, 217, 359-369.

[239] Westgate, A. J. et al, (1995). Diving behaviour of harbour porpoises, *Phocoena phocoena*. *Canadian Journal of Fisheries and Aquatic Science*, 52, 1065-1073.

[240] Wouter H. and Slabbekoorn H, (2015) , Pollution going multimodal: the complex impact of the human-altered sensory environment on animal perception and performance *Biol. Lett.* 2015 11 20141051.

Y

Z

[241] Zampolli M. et al, Scenarios for benchmarking range-dependent active sonar performance models, *Proceedings of the IOA*, Vol. 32, Pt. 2, pp 53-63,2010.

[242] Zampolli et al, (2013) , Validation of finite element computations for the quantitative prediction of underwater noise from impact pile driving, *J Acoust Soc Am.*, 133(1):72-81.

[243] Ziolkowski, A., (1970), A method for calculating the output pressure waveform from an air-gun: *Geophys. J. R. Astr. Soc.*, 21, 137 – 161.

[244] Ziolkowski A. et al, (1982), The signature of an air-gun array: Computation from near-field measurements including interactions: *Geophysics*, 47(10), 1413- 1421.

References