



Universiteit
Leiden
The Netherlands

Homogeneous nucleation for Glauber and Kawasaki dynamics in large volumes at low temperatures

Bovier, A.; Hollander, W.T.F. den; Spitoni, C.

Citation

Bovier, A., Hollander, W. T. F. den, & Spitoni, C. (2010). Homogeneous nucleation for Glauber and Kawasaki dynamics in large volumes at low temperatures. *Annals Of Probability*, 38(2), 661-713. doi:10.1214/09-AOP492

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/60055>

Note: To cite this publication please use the final published version (if applicable).

HOMOGENEOUS NUCLEATION FOR GLAUBER AND KAWASAKI DYNAMICS IN LARGE VOLUMES AT LOW TEMPERATURES

BY ANTON BOVIER¹, FRANK DEN HOLLANDER² AND CRISTIAN SPITONI

Rheinische Friedrich–Wilhelms-Universität Bonn, Leiden University and EURANDOM and Leiden University and EURANDOM

In this paper, we study metastability in large volumes at low temperatures. We consider both Ising spins subject to Glauber spin-flip dynamics and lattice gas particles subject to Kawasaki hopping dynamics. Let β denote the inverse temperature and let $\Lambda_\beta \subset \mathbb{Z}^2$ be a square box with periodic boundary conditions such that $\lim_{\beta \rightarrow \infty} |\Lambda_\beta| = \infty$. We run the dynamics on Λ_β , starting from a random initial configuration where all of the droplets (clusters of plus-spins and clusters of particles, respectively) are small. For large β and for interaction parameters that correspond to the metastable regime, we investigate how the transition from the metastable state (with only small droplets) to the stable state (with one or more large droplets) takes place under the dynamics. This transition is triggered by the appearance of a single *critical droplet* somewhere in Λ_β . Using potential-theoretic methods, we compute the *average nucleation time* (the first time a critical droplet appears and starts growing) up to a multiplicative factor that tends to 1 as $\beta \rightarrow \infty$. It turns out that this time grows as $K e^{\Gamma\beta} / |\Lambda_\beta|$ for Glauber dynamics and as $K\beta e^{\Gamma\beta} / |\Lambda_\beta|$ for Kawasaki dynamics, where Γ is the local canonical (resp. grand-canonical) energy, to create a critical droplet and K is a constant reflecting the geometry of the critical droplet, provided these times tend to infinity (which puts a growth restriction on $|\Lambda_\beta|$). The fact that the average nucleation time is inversely proportional to $|\Lambda_\beta|$ is referred to as *homogeneous nucleation* because it says that the critical droplet for the transition appears essentially independently in small boxes that partition Λ_β .

1. Introduction and main results.

1.1. *Background.* In a recent series of papers, Gaudillière et al. [12–14] study a system of lattice gas particles subject to Kawasaki hopping dynamics in a large box at low temperature and low density. Using the so-called *pathwise approach* to

Received June 2008; revised May 2009.

¹Supported by DFG and NWO through the Dutch–German Bilateral Research Group on “Random Spatial Models from Physics and Biology” (2003–2009).

²Supported by NWO through Grant no. 613.000.556.

AMS 2000 subject classifications. 60K35, 82C26.

Key words and phrases. Glauber dynamics, Kawasaki dynamics, critical droplet, metastable transition time, last-exit biased distribution, Dirichlet principle, Berman–Konsowa principle, capacity, flow, cluster expansion.

metastability (see Olivieri and Vares [23]), they show that the transition time between the metastable state (the gas phase with only small droplets) and the stable state (the liquid phase with one or more large droplets) is inversely proportional to the volume of the large box, provided that the latter does not grow too fast with the inverse temperature. This type of behavior is called *homogeneous nucleation* because it corresponds to the situation where the critical droplet triggering the nucleation appears essentially independently in small boxes that partition the large box. The *nucleation time* (the first time a critical droplet appears and starts growing) is computed up to a multiplicative error that is small on the scale of the exponential of the inverse temperature. The techniques developed in [12–14] center around the idea of approximating the low temperature and low density Kawasaki lattice gas by an ideal gas without interaction and showing that this ideal gas stays close to equilibrium while exchanging particles with droplets that are growing and shrinking. In this way, the large system is shown to behave essentially like the union of many small independent systems, leading to homogeneous nucleation. The proofs are long and complicated, but they provide considerable detail about the typical trajectory of the system prior to and shortly after the onset of nucleation.

In the present paper, we consider the same problem, both for Ising spins subject to Glauber spin-flip dynamics and for lattice gas particles subject to Kawasaki hopping dynamics. Using the potential-theoretic approach to metastability (see Bovier [5]), we improve upon an aspect of the results in [12–14], namely, we compute the *average* nucleation time up to a multiplicative error that tends to 1 as the temperature tends to 0, thereby providing a very sharp estimate of the time at which the gas starts to condensate.

We have no results about the typical time it takes for the system to grow a large droplet after the onset of nucleation. This is a difficult problem that will be addressed in future work. All that we can prove is that the dynamics has a negligible probability to shrink down a supercritical droplet once it has managed to create one. At least this shows that the appearance of a single critical droplet indeed represents the threshold for nucleation, as was shown in [12–14]. A further restriction is that we need to draw the initial configuration according to a *class* of initial distributions on the set of subcritical configurations, called the *last-exit biased distributions* since these are particularly suitable for the use of potential theory. It remains a challenge to investigate to what extent this restriction can be relaxed. This problem is addressed with some success in [12–14] and will also be tackled in future work.

Our results are an extension to large volumes of the results for small volumes obtained in Bovier and Manzo [8] (resp. Bovier, den Hollander and Nardi [7]). In large volumes, even at low temperatures, entropy is competing with energy because the metastable state and the states that evolve from it under the dynamics have a highly nontrivial structure. Our main goal in the present paper is to extend the potential-theoretic approach to metastability in order to be able to deal with large volumes. This is part of a broader program where the objective is to adapt the potential-theoretic approach to situations where entropy cannot be neglected.

In the same direction, Bianchi, Bovier, and Ioffe [3] study the dynamics of the random field Curie–Weiss model on a finite box at a fixed positive temperature.

As we will see, the basic difficulty in estimating the nucleation time is to obtain sharp upper and lower bounds on *capacities*. Upper bounds follow from the Dirichlet variational principle, which represents a capacity as an infimum over a class of test functions. In [3], a new technique is developed, based on a variational principle due to Berman and Konsowa [2], which represent a capacity as a supremum over a class of unit flows. This technique allows for lower bounds to be obtained and it will also be exploited here.

1.2. *Ising spins subject to Glauber dynamics.* We will study models in finite boxes, Λ_β , in the limit as both the inverse temperature, β , and the volume of the box, $|\Lambda_\beta|$, tend to infinity. Specifically, we let $\Lambda_\beta \subset \mathbb{Z}^2$ be a square box with odd side length, centered at the origin with periodic boundary conditions. A spin configuration is denoted by $\sigma = \{\sigma(x) : x \in \Lambda_\beta\}$, with $\sigma(x)$ representing the spin at site x , and is an element of $\mathcal{X}_\beta = \{-1, +1\}^{\Lambda_\beta}$. It will frequently be convenient to identify a configuration σ with its *support*, defined as $\text{supp}[\sigma] = \{x \in \Lambda_\beta : \sigma(x) = +1\}$.

The interaction is defined by the the usual Ising Hamiltonian

$$(1.1) \quad H_\beta(\sigma) = -\frac{J}{2} \sum_{\substack{(x,y) \in \Lambda_\beta \\ x \sim y}} \sigma(x)\sigma(y) - \frac{h}{2} \sum_{x \in \Lambda_\beta} \sigma(x), \quad \sigma \in \mathcal{X}_\beta,$$

where $J > 0$ is the pair potential, $h > 0$ is the magnetic field and $x \sim y$ means that x and y are nearest neighbors. The *Gibbs measure* associated with H_β is

$$(1.2) \quad \mu_\beta(\sigma) = \frac{1}{Z_\beta} e^{-\beta H_\beta(\sigma)}, \quad \sigma \in \mathcal{X}_\beta,$$

where Z_β is the normalizing partition function.

The dynamics of the model will the continuous-time Markov chain, $(\sigma(t))_{t \geq 0}$, with state space \mathcal{X}_β , whose transition rates are given by

$$(1.3) \quad c_\beta(\sigma, \sigma') = \begin{cases} e^{-\beta[H_\beta(\sigma') - H_\beta(\sigma)]_+}, & \text{for } \sigma' = \sigma^x \text{ for some } x \in \Lambda_\beta, \\ 0, & \text{otherwise,} \end{cases}$$

where σ^x is the configuration obtained from σ by flipping the spin at site x and $[\cdot]_+$ denotes the positive part. We refer to this Markov process as *Glauber dynamics*. It is ergodic and reversible with respect to its unique invariant measure, μ_β , that is,

$$(1.4) \quad \mu_\beta(\sigma)c_\beta(\sigma, \sigma') = \mu_\beta(\sigma')c_\beta(\sigma', \sigma) \quad \forall \sigma, \sigma' \in \mathcal{X}_\beta.$$

Glauber dynamics exhibits metastable behavior in the regime

$$(1.5) \quad 0 < h < 2J, \quad \beta \rightarrow \infty.$$

To understand this, let us briefly recall what happens in a finite β -independent box $\Lambda \subset \mathbb{Z}^2$. Let \boxminus_Λ and \boxplus_Λ denote the configurations where all spins in Λ are -1




FIG. 1. A critical droplet for Glauber dynamics on Λ . The shaded area represents the $(+1)$ -spins; the unshaded area represents the (-1) -spins [see (1.6)].

(resp. $+1$). As was shown by Neves and Schonmann [22], for Glauber dynamics restricted to Λ with periodic boundary conditions and subject to (1.5), the *critical droplets* for the crossover from \boxminus_{Λ} to \boxplus_{Λ} are elements of the set of all those configurations where the $(+1)$ -spins form an $\ell_c \times (\ell_c - 1)$ quasi-square (in either of the both orientations) with a protuberance attached to one of its longest sides, where

$$(1.6) \quad \ell_c = \left\lceil \frac{2J}{h} \right\rceil$$

(see Figures 1 and 2; for nondegeneracy reasons, it is assumed that $2J/h \notin \mathbb{N}$). The quasi-squares without the protuberance are called *proto-critical droplets*.

Let us now return to our setting with finite β -dependent volumes $\Lambda_{\beta} \subset \mathbb{Z}^2$. We will start our dynamics on Λ_{β} from initial configurations in which all droplets are “sufficiently small.” To make this notion precise, let $C_B(\sigma)$, $\sigma \in \mathcal{X}_{\beta}$, be the




FIG. 2. A nucleation path from \boxminus_{Λ} to \boxplus_{Λ} for Glauber dynamics. Γ in (1.10) is the minimal energy barrier the path has to overcome under the local variant of the Hamiltonian in (1.1).

configuration that is obtained from σ by a “bootstrap percolation map,” that is, by circumscribing all of the droplets in σ with rectangles and continuing to do so in an iterative manner until a union of disjoint rectangles is obtained (see Kotecký and Olivieri [19]). We call $C_B(\sigma)$ *subcritical* if all of its rectangles fit inside proto-critical droplets and are at distance ≥ 2 from each other (i.e., are noninteracting).

- DEFINITION 1.1. (a) $\mathcal{S} = \{\sigma \in \mathcal{X}_\beta : C_B(\sigma) \text{ is subcritical}\}$;
 (b) $\mathcal{P} = \{\sigma \in \mathcal{S} : c_\beta(\sigma, \sigma') > 0 \text{ for some } \sigma' \in \mathcal{S}^c\}$;
 (c) $\mathcal{C} = \{\sigma' \in \mathcal{S}^c : c_\beta(\sigma, \sigma') > 0 \text{ for some } \sigma \in \mathcal{S}\}$.

We refer to \mathcal{S} , \mathcal{P} and \mathcal{C} as the set of *subcritical*, *proto-critical* and *critical* configurations, respectively. Note that, for every $\sigma \in \mathcal{X}_\beta$, each step in the bootstrap percolation map $\sigma \rightarrow C_B(\sigma)$ decreases the energy and therefore the Glauber dynamics moves from σ to $C_B(\sigma)$ in a time of order 1. This is why $C_B(\sigma)$, rather than σ , appears in the definition of \mathcal{S} .

For $\ell_1, \ell_2 \in \mathbb{N}$, let $R_{\ell_1, \ell_2}(x) \subset \Lambda_\beta$ be the $\ell_1 \times \ell_2$ rectangle whose lower-left corner is x . We always take $\ell_1 \leq \ell_2$ and allow for both orientations of the rectangle. For $L = 1, \dots, 2\ell_c - 3$, let $Q_L(x)$ denote the L th element in the *canonical* sequence of growing squares and quasi-squares

$$(1.7) \quad R_{1,2}(x), R_{2,2}(x), R_{2,3}(x), R_{3,3}(x), \dots, R_{\ell_c-1, \ell_c-1}(x), R_{\ell_c-1, \ell_c}(x).$$

In what follows, we will choose to start the dynamics in a way that is suitable for the use of potential theory, as follows. First, we take the initial law to be concentrated on one of the sets $\mathcal{S}_L \subset \mathcal{S}$ defined by

$$(1.8) \quad \mathcal{S}_L = \{\sigma \in \mathcal{S} : \text{each rectangle in } C_B(\sigma) \text{ fits inside } Q_L(x) \text{ for some } x \in \Lambda_\beta\},$$

where L is any integer satisfying

$$(1.9) \quad L^* \leq L \leq 2\ell_c - 3$$

with $L^* = \min \left\{ 1 \leq L \leq 2\ell_c - 3 : \lim_{\beta \rightarrow \infty} \frac{\mu_\beta(\mathcal{S}_L)}{\mu_\beta(\mathcal{S})} = 1 \right\}$.

In words, \mathcal{S}_L is the subset of those subcritical configurations whose droplets fit inside a square or quasi-square labeled L , with L chosen large enough so that \mathcal{S}_L is typical within \mathcal{S} under the Gibbs measure μ_β as $\beta \rightarrow \infty$ (our results will not depend on the choice of L subject to these restrictions). Second, we take the initial law to be biased according to the last exit of \mathcal{S}_L for the transition from \mathcal{S}_L to a target set in \mathcal{S}^c . (Different choices will be made for the target set and the precise definition of the biased law will be given in Section 2.2.) This is a *highly specific* choice, but clearly one of physical interest.

REMARKS. (1) Note that $\mathcal{S}_{2\ell_c-3} = \mathcal{S}$, which implies that the range of L -values in (1.9) is nonempty. The value of L^* depends on how fast Λ_β grows with β . In Appendix C.1, we will show that, for every $1 \leq L \leq 2\ell_c - 4$, $\lim_{\beta \rightarrow \infty} \mu_\beta(\mathcal{S}_L)/\mu_\beta(\mathcal{S}) = 1$ if and only if $\lim_{\beta \rightarrow \infty} |\Lambda_\beta|e^{-\beta\Gamma_{L+1}} = 0$, with Γ_{L+1} the energy needed to create a droplet $Q_{L+1}(0)$ at the origin. Thus, if $|\Lambda_\beta| = e^{\theta\beta}$, then $L^* = L^*(\theta) = (2\ell_c - 3) \wedge \min\{L \in \mathbb{N} : \Gamma_{L+1} > \theta\}$, which increases stepwise from 1 to $2\ell_c - 3$ as θ increases from 0 to Γ defined in (1.10).

(2) If we draw the initial configuration σ_0 from some subset of \mathcal{S} that has a strong recurrence property under the dynamics, then the choice of initial distribution on this subset should not matter. This issue will be addressed in future work.

To state our main theorem for Glauber dynamics, we need some further notation. The key quantity for the nucleation process is

$$(1.10) \quad \Gamma = J[4\ell_c] - h[\ell_c(\ell_c - 1) + 1],$$

which is the energy needed to create a critical droplet of (+1)-spins at a given location in a sea of (-1)-spins (see Figures 1 and 2). For $\sigma \in \mathcal{X}_\beta$, let \mathbb{P}_σ denote the law of the dynamics starting from σ and, for ν a probability distribution on \mathcal{X} , put

$$(1.11) \quad \mathbb{P}_\nu(\cdot) = \sum_{\sigma \in \mathcal{X}_\beta} \mathbb{P}_\sigma(\cdot)\nu(\sigma).$$

For a nonempty set $\mathcal{A} \subset \mathcal{X}_\beta$, let

$$(1.12) \quad \tau_{\mathcal{A}} = \inf\{t > 0 : \sigma_t \in \mathcal{A}, \sigma_{t-} \notin \mathcal{A}\}$$

denote the first time that the dynamics enters \mathcal{A} . For nonempty and disjoint sets $\mathcal{A}, \mathcal{B} \subset \mathcal{X}_\beta$, let $\nu_{\mathcal{A}}^{\mathcal{B}}$ denote the *last-exit biased distribution on \mathcal{A} for the crossover to \mathcal{B}* defined in (2.9) in Section 2.2. Put

$$(1.13) \quad N_1 = 4\ell_c, \quad N_2 = \frac{4}{3}(2\ell_c - 1).$$

For $M \in \mathbb{N}$ with $M \geq \ell_c$, define

$$(1.14) \quad \mathcal{D}_M = \{\sigma \in \mathcal{X}_\beta : \exists x \in \Lambda_\beta \text{ such that } \text{supp}[C_B(\sigma)] \supset R_{M,M}(x)\},$$

that is, the set of configurations containing a supercritical droplet of size M . For our results below to be valid, we need to assume that

$$(1.15) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| = \infty, \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta|e^{-\beta\Gamma} = 0.$$

THEOREM 1.2. *In the regime (1.5), subject to (1.9) and (1.15), the following hold:*

(a)

$$(1.16) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta|e^{-\beta\Gamma} \mathbb{E}_{\nu_{\mathcal{S}_L}^{\mathcal{S}^c}}(\tau_{\mathcal{S}^c}) = \frac{1}{N_1};$$

(b)

$$(1.17) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| e^{-\beta\Gamma} \mathbb{E}_{\nu_{S_L}^{\mathcal{S}^c \setminus \mathcal{C}}}(\tau_{\mathcal{S}^c \setminus \mathcal{C}}) = \frac{1}{N_2};$$

(c)

$$(1.18) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| e^{-\beta\Gamma} \mathbb{E}_{\nu_{S_L}^{\mathcal{D}_M}}(\tau_{\mathcal{D}_M}) = \frac{1}{N_2} \quad \forall \ell_c \leq M \leq 2\ell_c - 1.$$

The proof of Theorem 1.2 will be given in Section 3. Part (a) says that the average time to create a critical droplet is $[1 + o(1)]e^{\beta\Gamma}/N_1|\Lambda_\beta|$. Parts (b) and (c) say that the average time to go beyond this critical droplet and to grow a droplet that is twice as large is $[1 + o(1)]e^{\beta\Gamma}/N_2|\Lambda_\beta|$. The factor N_1 counts the number of *shapes* of the critical droplet, while $|\Lambda_\beta|$ counts the number of *locations*. The average times to create a critical and a supercritical droplet differ by a factor $N_2/N_1 < 1$. This is because once the dynamics is “on top of the hill” \mathcal{C} , it has a positive probability to “fall back” to \mathcal{S} . On average, the dynamics makes $N_1/N_2 > 1$ attempts to reach the top \mathcal{C} before it finally “falls over” to $\mathcal{S}^c \setminus \mathcal{C}$. After that, it rapidly grows a large droplet (see Figure 2).

REMARKS. (1) The second condition in (1.15) will not actually be used in the proof of Theorem 1.2(a). If this condition fails, then there is a positive probability to see a proto-critical droplet in Λ_β under the starting measure $\nu_{S_L}^{\mathcal{S}^c}$ and nucleation sets in immediately. Theorem 1.2(a) continues to be true, but it no longer describes metastable behavior.

(2) In Appendix D, we will show that the *average* probability under the Gibbs measure μ_β of destroying a supercritical droplet and returning to a configuration in S_L is exponentially small in β . Hence, the crossover from S_L to $\mathcal{S}^c \setminus \mathcal{C}$ represents the true threshold for nucleation and Theorem 1.2(b) represents the true nucleation time.

(3) We expect Theorem 1.2(c) to hold for values of M that grow with β as $M = e^{o(\beta)}$. As we will see in Section 3.3, the necessary capacity estimates carry over, but the necessary equilibrium potential estimates are not yet available. This problem will be addressed in future work.

(4) Theorem 1.2 should be compared with the results in Bovier and Manzo [8] for the case of a finite β -independent box Λ (large enough to accommodate a critical droplet). In that case, if the dynamics starts from \boxtimes_Λ , then the average time it needs to hit \mathcal{C}_Λ (the set of configurations in Λ with a critical droplet), respectively, \boxplus_Λ equals

$$(1.19) \quad K e^{\beta\Gamma} [1 + o(1)] \quad \text{with } K = K(\Lambda, \ell_c) = \frac{1}{N} \frac{1}{|\Lambda|} \text{ for } N = N_1 \text{ (resp. } N_2).$$

(5) Note that in Theorem 1.2, we compute the first time when a critical droplet appears anywhere (!) in the box Λ_β . It is a different issue to compute the first

time when the plus-phase appears near the origin. This time, which depends on how a supercritical droplet grows and eventually invades the origin, was studied by Dehghanpour and Schonmann [10, 11], Shlosman and Schonmann [24] and, more recently, by Cerf and Manzo [9].

1.3. *Lattice gas subject to Kawasaki dynamics.* We next consider the lattice gas subject to Kawasaki dynamics and state a similar result for homogeneous nucleation. Some aspects are similar to what we have seen for Glauber dynamics, but there are notable differences.

A lattice gas configuration is denoted by $\sigma = \{\sigma(x) : x \in \Lambda_\beta\}$, with $\sigma(x)$ representing the number of particles at site x , and is an element of $\mathcal{X}_\beta = \{0, 1\}^{\Lambda_\beta}$. The Hamiltonian is given by

$$(1.20) \quad H_\beta(\sigma) = -U \sum_{\substack{(x,y) \in \Lambda_\beta \\ x \sim y}} \sigma(x)\sigma(y), \quad \sigma \in \mathcal{X}_\beta,$$

where $-U < 0$ is the binding energy and $x \sim y$ means that x and y are neighboring sites. Thus, we are working in the *canonical ensemble*, that is, there is no term analogous to the second term in (1.1). The number of particles in Λ_β is

$$(1.21) \quad n_\beta = \lceil \rho_\beta |\Lambda_\beta| \rceil,$$

where ρ_β is the particle density, which is chosen to be

$$(1.22) \quad \rho_\beta = e^{-\beta\Delta}, \quad \Delta > 0.$$

Put

$$(1.23) \quad \mathcal{X}_\beta^{(n_\beta)} = \{\sigma \in \mathcal{X}_\beta : |\text{supp}[\sigma]| = n_\beta\},$$

where $\text{supp}[\sigma] = \{x \in \Lambda_\beta : \sigma(x) = 1\}$.

REMARK. If we were to work in the *grand-canonical ensemble*, then we would have to consider the Hamiltonian

$$(1.24) \quad H^{gc}(\sigma) = -U \sum_{\substack{(x,y) \in \Lambda_\beta \\ x \sim y}} \sigma(x)\sigma(y) + \Delta \sum_{x \in \Lambda_\beta} \sigma(x), \quad \sigma \in \mathcal{X}_\beta,$$

where $\Delta > 0$ is an activity parameter taking over the role of h in (1.1). The second term would mimic the presence of an infinite gas reservoir with density ρ_β outside Λ_β . Such a Hamiltonian was used in earlier work on Kawasaki dynamics, when a finite β -independent box with open boundaries was considered (see, e.g., den Hollander, Olivieri and Scoppola [18], den Hollander et al. [17] and Bovier, den Hollander and Nardi [7]).

The dynamics of the model will be the continuous-time Markov chain, $(\sigma_t)_{t \geq 0}$, with state space $\mathcal{X}_\beta^{(n_\beta)}$, whose transition rates are

$$(1.25) \quad c_\beta(\sigma, \sigma') = \begin{cases} e^{-\beta[H_\beta(\sigma') - H_\beta(\sigma)]_+}, & \text{for } \sigma' = \sigma^{x,y} \text{ for some } x, y \in \Lambda_\beta \text{ with } x \sim y, \\ 0, & \text{otherwise,} \end{cases}$$

where $\sigma^{x,y}$ is the configuration obtained from σ by interchanging the values at sites x and y . We refer to this Markov process as *Kawasaki dynamics*. It is ergodic and reversible with respect to the canonical Gibbs measure

$$(1.26) \quad \mu_\beta(\sigma) = \frac{1}{Z_\beta^{(n_\beta)}} e^{-\beta H_\beta(\sigma)}, \quad \sigma \in \mathcal{X}_\beta^{(n_\beta)},$$

where $Z_\beta^{(n_\beta)}$ is the normalizing partition function. Note that the dynamics preserves particles, that is, it is *conservative*.

Kawasaki dynamics exhibits metastable behavior in the regime

$$(1.27) \quad U < \Delta < 2U, \quad \beta \rightarrow \infty.$$

This is again inferred from the behavior of the model in a finite β -independent box $\Lambda \subset \mathbb{Z}^2$. Let \square_Λ (resp. \blacksquare_Λ) denote the configurations where all of the sites in Λ are vacant (resp. occupied). For Kawasaki dynamics on Λ with an *open boundary*, where particles are annihilated at rate 1 and created at rate $e^{-\Delta\beta}$, it was shown in den Hollander, Olivieri, and Scoppola [18] and in Bovier, den Hollander, and Nardi [7] that, subject to (1.27) and for the Hamiltonian in (1.24), the critical droplets for the crossover from \square_Λ to \blacksquare_Λ are the set of all those configurations where the particles form either:

- (1) an $(\ell_c - 2) \times (\ell_c - 2)$ square with four bars attached to the four sides with total length $3\ell_c - 3$; or
- (2) an $(\ell_c - 1) \times (\ell_c - 3)$ rectangle with four bars attached to the four sides with total length $3\ell_c - 2$,

plus a *free particle* anywhere in the box, where

$$(1.28) \quad \ell_c = \left\lceil \frac{U}{2U - \Delta} \right\rceil$$

[see Figures 3 and 4; for nondegeneracy reasons, it is assumed that $U/(2U - \Delta) \notin \mathbb{N}$].

Let us now return to our setting with finite β -dependent volumes. We define a reference distance, L_β , as

$$(1.29) \quad L_\beta^2 = e^{(\Delta - \delta_\beta)\beta} = \frac{1}{\rho_\beta} e^{-\delta_\beta\beta}$$




FIG. 3. A critical droplet for Kawasaki dynamics on Λ (a proto-critical droplet plus a free particle). The shaded area represents the particles; the unshaded area represents the vacancies [see (1.28)]. The proto-critical droplet for Kawasaki dynamics drawn in the figure has the same shape as the critical droplet for Glauber dynamics, but there are other shapes as well [see (1) and (2) below (1.27)]. A proto-critical droplet for Kawasaki dynamics becomes critical when a free particle is added.

with

$$(1.30) \quad \lim_{\beta \rightarrow \infty} \delta_\beta = 0, \quad \lim_{\beta \rightarrow \infty} \beta \delta_\beta = \infty,$$




FIG. 4. A nucleation path from \square_Λ to \blacksquare_Λ for Kawasaki dynamics on Λ with open boundary. Γ in (1.33) is the minimal energy barrier the path has to overcome under the local variant of the grand-canonical Hamiltonian in (1.24).

that is, L_β is marginally below the typical inter-particle distance. We assume L_β to be odd and write $B_{L_\beta, L_\beta}(x)$, $x \in \Lambda_\beta$, for the square box with side length L_β whose center is x .

DEFINITION 1.3. (a) $\mathcal{S} = \{\sigma \in \mathcal{X}_\beta^{(n_\beta)} : |\text{supp}[\sigma] \cap B_{L_\beta, L_\beta}(x)| \leq \ell_c(\ell_c - 1) + 1 \forall x \in \Lambda_\beta\}$;

(b) $\mathcal{P} = \{\sigma \in \mathcal{S} : c_\beta(\sigma, \sigma') > 0 \text{ for some } \sigma' \in \mathcal{S}^c\}$;

(c) $\mathcal{C} = \{\sigma' \in \mathcal{S}^c : c_\beta(\sigma, \sigma') > 0 \text{ for some } \sigma \in \mathcal{S}\}$;

(d) $\mathcal{C}^- = \{\sigma \in \mathcal{C} : \exists x \in \Lambda_\beta \text{ such that } B_{L_\beta, L_\beta}(x) \text{ contains a proto-critical droplet plus a free particle at distance } L_\beta\}$;

(e) $\mathcal{C}^+ =$ the set of configurations obtained from \mathcal{C}^- by moving the free particle to a site at distance 2 from the proto-critical droplet, that is, next to its boundary;

(f) $\tilde{\mathcal{C}} =$ the set of configurations “interpolating” between \mathcal{C}^- and \mathcal{C}^+ , that is, the free particle is somewhere between the boundary of the proto-critical droplet and the boundary of the box of size L_β around it.

As before, we refer to \mathcal{S} , \mathcal{P} and \mathcal{C} as the set of *subcritical*, *proto-critical* and *critical* configurations, respectively. Note that, for every $\sigma \in \mathcal{S}$, the number of particles in a box of size L_β does not exceed the number of particles in a proto-critical droplet. These particles do not have to form a cluster or to be near each other because the Kawasaki dynamics brings them together in a time of order $L_\beta^2 = o(1/\rho_\beta)$.

The initial law will again be concentrated on sets $\mathcal{S}_L \subset \mathcal{S}$, this time defined by

$$(1.31) \quad \mathcal{S}_L = \{\sigma \in \mathcal{X}_\beta^{(n_\beta)} : |\text{supp}[\sigma] \cap B_{L_\beta, L_\beta}(x)| \leq L \forall x \in \Lambda_\beta\},$$

where L is any integer satisfying

$$(1.32) \quad L^* \leq L \leq \ell_c(\ell_c - 1) + 1$$

$$\text{with } L^* = \min\left\{1 \leq L \leq \ell_c(\ell_c - 1) + 1 : \lim_{\beta \rightarrow \infty} \frac{\mu_\beta(\mathcal{S}_L)}{\mu_\beta(\mathcal{S})} = 1\right\}.$$

In words, \mathcal{S}_L is the subset of those subcritical configurations for which no box of size L_β carries more than L particles, with L again chosen such that \mathcal{S}_L is typical within \mathcal{S} under the Gibbs measure μ_β as $\beta \rightarrow \infty$.

REMARK. Note that $\mathcal{S}_{\ell_c(\ell_c-1)+1} = \mathcal{S}$. As for Glauber dynamics, the value of L^* depends on how fast Λ_β grows with β . In Appendix C.2, we will show that for every $1 \leq L \leq \ell_c(\ell_c - 1)$, $\lim_{\beta \rightarrow \infty} \mu_\beta(\mathcal{S}_L)/\mu_\beta(\mathcal{S}) = 1$ if and only if $\lim_{\beta \rightarrow \infty} |\Lambda_\beta| e^{-\beta(\Gamma_{L+1} - \Delta)} = 0$, with Γ_{L+1} the energy needed to create a droplet of $L + 1$ particles, closest in shape to a square or quasi-square, in $B_{L_\beta, L_\beta}(0)$ under the grand-canonical Hamiltonian on this box. Thus, if $|\Lambda_\beta| = e^{\theta\beta}$, then $L^* = L^*(\theta) = [\ell_c(\ell_c - 1) + 1] \wedge \min\{L \in \mathbb{N} : \Gamma_{L+1} - \Delta > \theta\}$, which increases stepwise from 1 to $\ell_c(\ell_c - 1) + 1$ as θ increases from Δ to Γ defined in (1.33).

Set

$$(1.33) \quad \Gamma = -U[(\ell_c - 1)^2 + \ell_c(\ell_c - 1) + 1] + \Delta[\ell_c(\ell_c - 1) + 2],$$

which is the energy of a critical droplet at a given location with respect to the *grand-canonical* Hamiltonian given by (1.24) (see Figures 3 and 4). Put $N = \frac{1}{3}\ell_c^2(\ell_c^2 - 1)$. For $M \in \mathbb{N}$ with $M \geq \ell_c$, define

$$(1.34) \quad \mathcal{D}_M = \{\sigma \in \mathcal{X}_\beta : \exists x \in \Lambda_\beta \text{ such that } \text{supp}[(\sigma)] \supset R_{M,M}(x)\},$$

that is, the set of configurations containing a supercritical droplet of size M . For our results below to be valid, we need to assume that

$$(1.35) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| \rho_\beta = \infty, \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| e^{-\beta\Gamma} = 0.$$

This first condition says that the number of particles tends to infinity and ensures that the formation of a critical droplet somewhere does not globally deplete the surrounding gas.

THEOREM 1.4. *In the regime (1.27), subject to (1.32) and (1.35), the following hold:*

(a)

$$(1.36) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| \frac{4\pi}{\beta\Delta} e^{-\beta\Gamma} \mathbb{E}_{\nu_{S_L}^{\mathcal{S}^c \setminus \tilde{\mathcal{C}} \cup \mathcal{C}^+}}(\tau_{(\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+}) = \frac{1}{N};$$

(b)

$$(1.37) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| \frac{4\pi}{\beta\Delta} e^{-\beta\Gamma} \mathbb{E}_{\nu_{S_L}^{\mathcal{D}_M}}(\tau_{\mathcal{D}_M}) = \frac{1}{N} \quad \forall \ell_c \leq M \leq 2\ell_c - 1.$$

The proof of Theorem 1.4, which is an analog of Theorem 1.2, will be given in Section 4. Part (a) says that the average time to create a critical droplet is $[1 + o(1)](\beta\Delta/4\pi)e^{\beta\Gamma} N |\Lambda_\beta|$. The factor $\beta\Delta/4\pi$ comes from the simple random walk that is performed by the free particle “from the gas to the proto-critical droplet” (i.e., the dynamics goes from \mathcal{C}^- to \mathcal{C}^+), which slows down the nucleation. The factor N counts the number of shapes of the proto-critical droplet (see Bovier, den Hollander and Nardi [7]). Part (b) says that once the critical droplet is created, it rapidly grows to a droplet that has twice the size.

REMARKS. (1) As for Theorem 1.2(c), we expect Theorem 1.4(b) to hold for values of M that grow with β as $M = e^{o(\beta)}$. See Section 4.2 for more details.

(2) In Appendix D, we will show that the *average* probability under the Gibbs measure μ_β of destroying a supercritical droplet and returning to a configuration in S_L is exponentially small in β . Hence, the crossover from S_L to $\mathcal{S}^c \setminus \tilde{\mathcal{C}} \cup \mathcal{C}^+$

represents the true threshold for nucleation and Theorem 1.4(a) represents the true nucleation time.

(3) It was shown in Bovier, den Hollander and Nardi [7] that the average crossover time in a finite box Λ equals

$$(1.38) \quad Ke^{\beta\Gamma} [1 + o(1)] \quad \text{with } K = K(\Lambda, \ell_c) \sim \frac{\log |\Lambda|}{4\pi} \frac{1}{N|\Lambda|}, \Lambda \rightarrow \mathbb{Z}^2.$$

This matches the $|\Lambda_\beta|$ -dependence in Theorem 1.4, with the logarithmic factor in (1.38) accounting for the extra factor $\beta\Delta$ in Theorem 1.4 compared to Theorem 1.2. Note that this factor is particularly interesting since it says that the *effective box size* responsible for the formation of a critical droplet is L_β .

1.4. *Outline.* The remainder of this paper is organized as follows. In Section 2, we present a brief sketch of the basic ingredients of the potential-theoretic approach to metastability. In particular, we exhibit a relation between average crossover times and capacities and we state two variational representations for capacities, the first of which is suitable for deriving upper bounds and the second for deriving lower bounds. Section 3 contains the proof of our results for the case of Glauber dynamics. Technically, this will be relatively easy and will give a first flavor of how our method works. In Section 4, we deal with Kawasaki dynamics. Here, we will encounter several rather more difficult issues, all coming from the fact that Kawasaki dynamics is *conservative*. The first issue is to understand why the constant Γ , representing the local energetic cost to create a critical droplet, involves the grand-canonical Hamiltonian, even though we are working in the canonical ensemble. This mystery will, of course, be resolved by the observation that the formation of a critical droplet reduces the entropy of the system: the precise computation of this entropy loss yields Γ via *equivalence of ensembles*. The second problem is to control the probability of a particle moving from the gas to the proto-critical droplet at the last stage of the nucleation. This nonlocality issue will be dealt with via upper and lower estimates. Appendices A–D collect some technical lemmas that are needed in Sections 3 and 4.

The extension of our results to higher dimensions is limited only by the combinatorial problems involved in the computation of the number of critical droplets (which is hard in the case of Kawasaki dynamics) and of the probability for simple random walk to hit a critical droplet of a given shape when coming from far. We will not pursue this generalization here. The relevant results for a β -independent box in \mathbb{Z}^3 can be found in Ben Arous and Cerf [1] (Glauber) and den Hollander, Nardi, Olivieri and Scoppola [17] (Kawasaki). For recent overviews on droplet growth in metastability, we refer the reader to den Hollander [15, 16] and Bovier [4, 5]. A general overview on metastability is given in the monograph by Olivieri and Vares [23].

2. Basic ingredients of the potential-theoretic approach. The proofs of Theorems 1.2 and 1.4 use the *potential-theoretic approach* to metastability developed in Bovier et al. [6]. This approach is based on the following three ob-

servations. First, most quantities of physical interest can be represented in terms of Dirichlet problems associated with the generator of the dynamics. Second, the Green function of the dynamics can be expressed in terms of capacities and equilibrium potentials. Third, capacities satisfy variational principles that allow upper and lower bounds to be obtained in a flexible way. We will see that in the current setting, the implementation of these observations provides very sharp results.

2.1. *Equilibrium potential and capacity.* The fundamental quantity in the theory is the *equilibrium potential*, $h_{\mathcal{A},\mathcal{B}}$, associated with two nonempty disjoint sets of configurations, $\mathcal{A}, \mathcal{B} \subset \mathcal{X}$ (\mathcal{X}_β or $\mathcal{X}_\beta^{(n_\beta)}$), which, probabilistically, is given by

$$(2.1) \quad h_{\mathcal{A},\mathcal{B}}(\sigma) = \begin{cases} \mathbb{P}_\sigma(\tau_{\mathcal{A}} < \tau_{\mathcal{B}}), & \text{for } \sigma \in (\mathcal{A} \cup \mathcal{B})^c, \\ 1, & \text{for } \sigma \in \mathcal{A}, \\ 0, & \text{for } \sigma \in \mathcal{B}, \end{cases}$$

where

$$(2.2) \quad \tau_{\mathcal{A}} = \inf\{t > 0 : \sigma_t \in \mathcal{A}, \sigma_{t-} \notin \mathcal{A}\},$$

$(\sigma_t)_{t \geq 0}$ is the continuous-time Markov chain with state space \mathcal{X} and \mathbb{P}_σ is its law starting from σ . This function is harmonic and is the unique solution of the *Dirichlet problem*

$$(2.3) \quad \begin{aligned} (Lh_{\mathcal{A},\mathcal{B}})(\sigma) &= 0, & \sigma \in (\mathcal{A} \cup \mathcal{B})^c, \\ h_{\mathcal{A},\mathcal{B}}(\sigma) &= 1, & \sigma \in \mathcal{A}, \\ h_{\mathcal{A},\mathcal{B}}(\sigma) &= 0, & \sigma \in \mathcal{B}, \end{aligned}$$

where the generator is the matrix with entries

$$(2.4) \quad L(\sigma, \sigma') = c_\beta(\sigma, \sigma') - \delta_{\sigma, \sigma'} c_\beta(\sigma), \quad \sigma, \sigma' \in \mathcal{X},$$

where $c_\beta(\sigma)$ is the total rate at which the dynamics leaves σ ,

$$(2.5) \quad c_\beta(\sigma) = \sum_{\sigma' \in \mathcal{X} \setminus \{\sigma\}} c_\beta(\sigma, \sigma'), \quad \sigma \in \mathcal{X}.$$

A related quantity is the *equilibrium measure* on \mathcal{A} , which is defined as

$$(2.6) \quad e_{\mathcal{A},\mathcal{B}}(\sigma) = -(Lh_{\mathcal{A},\mathcal{B}})(\sigma), \quad \sigma \in \mathcal{A}.$$

The equilibrium measure also has a probabilistic meaning, namely,

$$(2.7) \quad \mathbb{P}_\sigma(\tau_{\mathcal{B}} < \tau_{\mathcal{A}}) = \frac{e_{\mathcal{A},\mathcal{B}}(\sigma)}{c_\beta(\sigma)}, \quad \sigma \in \mathcal{A}.$$

The key object we will work with is the *capacity*, which is defined as

$$(2.8) \quad \text{CAP}(\mathcal{A}, \mathcal{B}) = \sum_{\sigma \in \mathcal{A}} \mu_\beta(\sigma) e_{\mathcal{A},\mathcal{B}}(\sigma).$$

2.2. *Relation between crossover time and capacity.* The first important ingredient of the potential-theoretic approach to metastability is a formula for the average crossover time from \mathcal{A} to \mathcal{B} . To state this formula, we define the probability measure $\nu_{\mathcal{A}}^{\mathcal{B}}$ on \mathcal{A} which we already referred to in Section 1, namely,

$$(2.9) \quad \nu_{\mathcal{A}}^{\mathcal{B}}(\sigma) = \begin{cases} \frac{\mu_{\beta}(\sigma)e_{\mathcal{A},\mathcal{B}}(\sigma)}{\text{CAP}(\mathcal{A}, \mathcal{B})}, & \text{for } \sigma \in \mathcal{A}, \\ 0, & \text{for } \sigma \in \mathcal{A}^c. \end{cases}$$

The following proposition is proved in, for example, Bovier [5].

PROPOSITION 2.1. *For any two nonempty disjoint sets $\mathcal{A}, \mathcal{B} \subset \mathcal{X}$,*

$$(2.10) \quad \sum_{\sigma \in \mathcal{A}} \nu_{\mathcal{A}}^{\mathcal{B}}(\sigma) \mathbb{E}_{\sigma}(\tau_{\mathcal{B}}) = \frac{1}{\text{CAP}(\mathcal{A}, \mathcal{B})} \sum_{\sigma \in \mathcal{B}^c} \mu_{\beta}(\sigma) h_{\mathcal{A},\mathcal{B}}(\sigma).$$

REMARKS. (1) Due to (2.7) and (2.8), the probability measure $\nu_{\mathcal{A}}^{\mathcal{B}}(\sigma)$ can be written as

$$(2.11) \quad \nu_{\mathcal{A}}^{\mathcal{B}}(\sigma) = \frac{\mu_{\beta}(\sigma)c_{\beta}(\sigma)}{\text{CAP}(\mathcal{A}, \mathcal{B})} \mathbb{P}_{\sigma}(\tau_{\mathcal{B}} < \tau_{\mathcal{A}}), \quad \sigma \in \mathcal{A},$$

and thus has the flavor of a last-exit biased distribution. Proposition 2.1 explains why our main results on average crossover times stated in Theorem 1.2 and 1.4 are formulated for this initial distribution. Note that

$$(2.12) \quad \mu_{\beta}(\mathcal{A}) \leq \sum_{\sigma \in \mathcal{B}^c} \mu_{\beta}(\sigma) h_{\mathcal{A},\mathcal{B}}(\sigma) \leq \mu_{\beta}(\mathcal{B}^c).$$

We will see that in our setting, $\mu_{\beta}(\mathcal{B}^c \setminus \mathcal{A}) = o(\mu_{\beta}(\mathcal{A}))$ as $\beta \rightarrow \infty$ so that the sum in the right-hand side of (2.10) is $\sim \mu_{\beta}(\mathcal{A})$ and the computation of the crossover time reduces to the estimation of $\text{CAP}(\mathcal{A}, \mathcal{B})$.

(2) For a fixed target set \mathcal{B} , the choice of the starting set \mathcal{A} is free. It is tempting to choose $\mathcal{A} = \{\sigma\}$ for some $\sigma \in \mathcal{X}$. This was done for the case of a finite β -independent box Λ . However, in our case (and, more generally, in cases where the state space is large) such a choice would give intractable numerators and denominators in the right-hand side of (2.10). As a rule, to make use of the identity in (2.10), \mathcal{A} must be so large that the harmonic function $h_{\mathcal{A},\mathcal{B}}$ “does not change abruptly near the boundary of \mathcal{A} ” for the target set \mathcal{B} under consideration.

As noted above, average crossover times are essentially governed by capacities. The usefulness of this observation comes from the computability of capacities, as will be explained next.

2.3. *The Dirichlet principle: a variational principle for upper bounds.* The capacity is a boundary quantity because $e_{\mathcal{A},\mathcal{B}} > 0$ only on the boundary of \mathcal{A} .

The analog of Green’s identity relates it to a bulk quantity. Indeed, in terms of the *Dirichlet form* defined by

$$(2.13) \quad \mathcal{E}(h) = \frac{1}{2} \sum_{\sigma, \sigma' \in \mathcal{X}} \mu_\beta(\sigma) c_\beta(\sigma, \sigma') [h(\sigma) - h(\sigma')]^2, \quad h : \mathcal{X} \rightarrow [0, 1],$$

it follows, via (2.1) and (2.7) and (2.8), that

$$(2.14) \quad \text{CAP}(\mathcal{A}, \mathcal{B}) = \mathcal{E}(h_{\mathcal{A}, \mathcal{B}}).$$

Elementary variational calculus shows that the capacity satisfies the *Dirichlet principle*.

PROPOSITION 2.2. *For any two nonempty disjoint sets $\mathcal{A}, \mathcal{B} \subset \mathcal{X}$,*

$$(2.15) \quad \text{CAP}(\mathcal{A}, \mathcal{B}) = \min_{\substack{h : \mathcal{X} \rightarrow [0, 1] \\ h|_{\mathcal{A}} \equiv 1, h|_{\mathcal{B}} \equiv 0}} \mathcal{E}(h).$$

The importance of the Dirichlet principle is that it yields computable upper bounds for capacities by means of suitable choices of the test function h . In metastable systems, with the proper physical insight, it is often possible to guess a reasonable test function. In our setting, this will be seen to be relatively easy.

2.4. *The Berman–Konsowa principle: a variational principle for lower bounds.*

We will describe a little-known variational principle for capacities that is originally due to Berman and Konsowa [2]. Our presentation will follow the argument given in Bianchi, Bovier and Ioffe [3].

In the following, it will be convenient to think of \mathcal{X} as the vertex set of a graph $(\mathcal{X}, \mathcal{E})$ whose edge set \mathcal{E} consists of all pairs (σ, σ') , $\sigma, \sigma' \in \mathcal{X}$, for which $c_\beta(\sigma, \sigma') > 0$.

DEFINITION 2.3. Given two nonempty disjoint sets $\mathcal{A}, \mathcal{B} \subset \mathcal{X}$, a loop-free nonnegative unit flow, f , from \mathcal{A} to \mathcal{B} is a function $f : \mathcal{E} \rightarrow [0, \infty)$ such that:

- (a) $(f(e) > 0 \implies f(-e) = 0) \forall e \in \mathcal{E}$;
- (b) f satisfies Kirchoff’s law, that is,

$$(2.16) \quad \sum_{\sigma' \in \mathcal{X}} f(\sigma, \sigma') = \sum_{\sigma'' \in \mathcal{X}} f(\sigma'', \sigma) \quad \forall \sigma \in \mathcal{X} \setminus (\mathcal{A} \cup \mathcal{B});$$

- (c) f is normalized, that is,

$$(2.17) \quad \sum_{\sigma \in \mathcal{A}} \sum_{\sigma' \in \mathcal{X}} f(\sigma, \sigma') = 1 = \sum_{\sigma'' \in \mathcal{X}} \sum_{\sigma \in \mathcal{B}} f(\sigma'', \sigma);$$

- (d) any path from \mathcal{A} to \mathcal{B} along edges e such that $f(e) > 0$ is self-avoiding.

The space of all loop-free nonnegative unit flows from \mathcal{A} to \mathcal{B} is denoted by $\mathbb{U}_{\mathcal{A}, \mathcal{B}}$.

A natural flow is the *harmonic flow*, which is constructed from the equilibrium potential $h_{\mathcal{A},\mathcal{B}}$ as

$$(2.18) \quad f_{\mathcal{A},\mathcal{B}}(\sigma, \sigma') = \frac{1}{\text{CAP}(\mathcal{A}, \mathcal{B})} \mu_{\mathcal{B}}(\sigma) c_{\mathcal{B}}(\sigma, \sigma') [h_{\mathcal{A},\mathcal{B}}(\sigma) - h_{\mathcal{A},\mathcal{B}}(\sigma')]_+,$$

$\sigma, \sigma' \in \mathcal{X}.$

It is easy to verify that $f_{\mathcal{A},\mathcal{B}}$ satisfies (a)–(d). Indeed, (a) is obvious, (b) uses the harmonicity of $h_{\mathcal{A},\mathcal{B}}$, (c) follows from (2.6) and (2.8), while (d) comes from the fact that the harmonic flow only moves in directions where $h_{\mathcal{A},\mathcal{B}}$ decreases.

A loop-free nonnegative unit flow f is naturally associated with a probability measure \mathbb{P}^f on self-avoiding paths, γ . To see this, define $F(\sigma) = \sum_{\sigma' \in \mathcal{X}} f(\sigma, \sigma')$, $\sigma \in \mathcal{X} \setminus \mathcal{B}$. Then \mathbb{P}^f is the Markov chain $(\sigma_n)_{n \in \mathbb{N}_0}$ with initial distribution $\mathbb{P}^f(\sigma_0) = F(\sigma_0)1_{\mathcal{A}}(\sigma_0)$ and transition probabilities

$$(2.19) \quad q^f(\sigma, \sigma') = \frac{f(\sigma, \sigma')}{F(\sigma)}, \quad \sigma \in \mathcal{X} \setminus \mathcal{B},$$

such that the chain is stopped upon arrival in \mathcal{B} . In terms of this probability measure, we have the following proposition (see [3] for a proof).

PROPOSITION 2.4. *Let $\mathcal{A}, \mathcal{B} \subset \mathcal{X}$ be two nonempty disjoint sets. Then, with the notation introduced above,*

$$(2.20) \quad \text{CAP}(\mathcal{A}, \mathcal{B}) = \sup_{f \in \mathbb{U}_{\mathcal{A},\mathcal{B}}} \mathbb{E}^f \left(\left[\sum_{e \in \gamma} \frac{f(e_l, e_r)}{\mu_{\mathcal{B}}(e_l) c_{\mathcal{B}}(e_l, e_r)} \right]^{-1} \right),$$

where $e = (e_l, e_r)$ and the expectation is with respect to γ . Moreover, the supremum is realized for the harmonic flow $f_{\mathcal{A},\mathcal{B}}$.

The nice feature of this variational principle is that any flow gives a computable lower bound. In this sense, (2.15) and (2.20) complement each other. Moreover, since the harmonic flow is optimal, a good approximation of the harmonic function $h_{\mathcal{A},\mathcal{B}}$ by a test function h leads to a good approximation of the harmonic flow $f_{\mathcal{A},\mathcal{B}}$ by a test flow f after putting h instead of $h_{\mathcal{A},\mathcal{B}}$ into (2.18). Again, in metastable systems, with the proper physical insight, it is often possible to guess a reasonable flow. We will see in Sections 3–4 how this is put to work in our setting.

3. Proof of Theorem 1.2.

3.1. *Proof of Theorem 1.2(a).* To estimate the average crossover time from $S_L \subset \mathcal{S}$ to S^c , we will use Proposition 2.1. With $\mathcal{A} = S_L$ and $\mathcal{B} = S^c$, (2.10) reads

$$(3.1) \quad \sum_{\sigma \in S_L} v_{S_L}^{S^c}(\sigma) \mathbb{E}_{\sigma}(\tau_{S^c}) = \frac{1}{\text{CAP}(S_L, S^c)} \sum_{\sigma \in S} \mu_{\mathcal{B}}(\sigma) h_{S_L, S^c}(\sigma).$$

The left-hand side is the quantity of interest in (1.16). In Sections 3.1.1 and 3.1.2, we estimate $\sum_{\sigma \in \mathcal{S}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c}(\sigma)$ and $\text{CAP}(\mathcal{S}_L, \mathcal{S}^c)$. The estimates will show that

$$(3.2) \quad \text{r.h.s. (3.1)} = \frac{1}{N_1 |\Lambda_\beta|} e^{\beta\Gamma} [1 + o(1)], \quad \beta \rightarrow \infty.$$

3.1.1. Estimate of $\sum_{\sigma \in \mathcal{S}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c}(\sigma)$.

LEMMA 3.1. $\sum_{\sigma \in \mathcal{S}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c}(\sigma) = \mu_\beta(\mathcal{S}) [1 + o(1)]$ as $\beta \rightarrow \infty$.

PROOF. Write, using (2.1),

$$(3.3) \quad \begin{aligned} \sum_{\sigma \in \mathcal{S}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c}(\sigma) &= \sum_{\sigma \in \mathcal{S}_L} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c}(\sigma) + \sum_{\sigma \in \mathcal{S} \setminus \mathcal{S}_L} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c}(\sigma) \\ &= \mu_\beta(\mathcal{S}_L) + \sum_{\sigma \in \mathcal{S} \setminus \mathcal{S}_L} \mu_\beta(\sigma) \mathbb{P}_\sigma(\tau_{\mathcal{S}_L} < \tau_{\mathcal{S}^c}). \end{aligned}$$

The last sum is bounded above by $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L)$, but $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L) = o(\mu_\beta(\mathcal{S}))$ as $\beta \rightarrow \infty$ by our choice of L in (1.9). \square

3.1.2. Estimate of $\text{CAP}(\mathcal{S}_L, \mathcal{S}^c)$.

LEMMA 3.2. $\text{CAP}(\mathcal{S}_L, \mathcal{S}^c) = N_1 |\Lambda_\beta| e^{-\beta\Gamma} \mu_\beta(\mathcal{S}) [1 + o(1)]$ as $\beta \rightarrow \infty$ with $N_1 = 4\ell_c$.

PROOF. The proof proceeds via upper and lower bounds.

Upper bound. We use the Dirichlet principle and a test function that is equal to 1 on \mathcal{S} to get the upper bound

$$(3.4) \quad \begin{aligned} \text{CAP}(\mathcal{S}_L, \mathcal{S}^c) &\leq \text{CAP}(\mathcal{S}, \mathcal{S}^c) = \sum_{\substack{\sigma \in \mathcal{S}, \sigma' \in \mathcal{S}^c \\ c_\beta(\sigma, \sigma') > 0}} \mu_\beta(\sigma) c_\beta(\sigma, \sigma') \\ &= \sum_{\substack{\sigma \in \mathcal{S}, \sigma' \in \mathcal{S}^c \\ c_\beta(\sigma, \sigma') > 0}} [\mu_\beta(\sigma) \wedge \mu_\beta(\sigma')] \leq \mu_\beta(\mathcal{C}), \end{aligned}$$

where the second equality uses (1.4) in combination with the fact that $c_\beta(\sigma, \sigma') \vee c_\beta(\sigma', \sigma) = 1$, by (1.3). Thus, it suffices to show that

$$(3.5) \quad \mu_\beta(\mathcal{C}) \leq N_1 |\Lambda_\beta| e^{-\beta\Gamma} [1 + o(1)] \quad \text{as } \beta \rightarrow \infty.$$

For every $\sigma \in \mathcal{P}$, there are one or more rectangles $R_{\ell_c-1, \ell_c}(x)$, $x = x(\sigma) \in \mathcal{X}_\beta$, that are filled by (+1)-spins in $C_B(\sigma)$. If $\sigma' \in \mathcal{C}$ is such that $\sigma' = \sigma^y$ for some




FIG. 5. $R_{\ell_c-1, \ell_c}(x)$ (shaded box) and $[R_{\ell_c+1, \ell_c+2}(x - (1, 1))]^c$ (complement of dotted box).

$y \in \Lambda_\beta$, then σ' has a (+1)-spin at y situated on the boundary of one of these rectangles. Let

$$(3.6) \quad \begin{aligned} \hat{\mathcal{S}}(x) &= \{\sigma \in \mathcal{S} : \text{supp}[\sigma] \subseteq R_{\ell_c-1, \ell_c}(x)\}, \\ \check{\mathcal{S}}(x) &= \{\sigma \in \mathcal{S} : \text{supp}[\sigma] \subseteq [R_{\ell_c+1, \ell_c+2}(x - (1, 1))]^c\}. \end{aligned}$$

For every $\sigma \in \mathcal{P}$, we have $\sigma = \hat{\sigma} \vee \check{\sigma}$ for some $\hat{\sigma} \in \hat{\mathcal{S}}(x)$ and $\check{\sigma} \in \check{\mathcal{S}}(x)$, uniquely decomposing the configuration into two noninteracting parts inside $R_{\ell_c-1, \ell_c}(x)$ and $[R_{\ell_c+1, \ell_c+2}(x - (1, 1))]^c$ (see Figure 5). We have

$$(3.7) \quad H_\beta(\sigma) - H_\beta(\Xi) = [H_\beta(\hat{\sigma}) - H_\beta(\Xi)] + [H_\beta(\check{\sigma}) - H_\beta(\Xi)].$$

Moreover, for any $y \notin \text{supp}[C_B(\sigma)]$, we have

$$(3.8) \quad H_\beta(\sigma^y) \geq H_\beta(\sigma) + 2J - h.$$

Hence,

$$(3.9) \quad \begin{aligned} \mu_\beta(\mathcal{C}) &= \frac{1}{Z_\beta} \sum_{\sigma \in \mathcal{P}} \sum_{\substack{x \in \Lambda_\beta \\ \sigma^x \in \mathcal{C}}} e^{-\beta H_\beta(\sigma^x)} \\ &\leq \frac{1}{Z_\beta} N_1 e^{-\beta[2J-h-H_\beta(\Xi)]} \sum_{x \in \Lambda_\beta} \sum_{\check{\sigma} \in \check{\mathcal{S}}(x)} e^{-\beta H_\beta(\check{\sigma})} \sum_{\substack{\hat{\sigma} \in \hat{\mathcal{S}}(x) \\ \hat{\sigma} \vee \check{\sigma} \in \mathcal{P}}} e^{-\beta H_\beta(\hat{\sigma})} \\ &\leq [1 + o(1)] \frac{1}{Z_\beta} N_1 |\Lambda_\beta| e^{-\beta\Gamma} \sum_{\check{\sigma} \in \check{\mathcal{S}}(0)} e^{-\beta H_\beta(\check{\sigma})} \\ &= [1 + o(1)] N_1 |\Lambda_\beta| e^{-\beta\Gamma} \mu_\beta(\check{\mathcal{S}}(0)), \end{aligned}$$

where the first inequality uses (3.7)–(3.8), with $N_1 = 2 \times 2\ell_c = 4\ell_c$ counting the number of critical droplets that can arise from a proto-critical droplet via a spin flip (see Figure 1), and the second inequality uses the fact that

$$(3.10) \quad \begin{aligned} \hat{\sigma} \in \hat{\mathcal{S}}(0), \quad \hat{\sigma} \vee \check{\sigma} \in \mathcal{P} \\ \implies H_\beta(\hat{\sigma}) \geq H_\beta(R_{\ell_c-1, \ell_c}(0)) = \Gamma - (2J - h) + H_\beta(\Xi) \end{aligned}$$




FIG. 6. Canonical order to break down a critical droplet.

with equality in the right-hand side if and only if $\text{supp}[\hat{\sigma}] = R_{\ell_c-1, \ell_c}(0)$. Combining (3.4) and (3.9) with the inclusion $\check{\mathcal{S}}(0) \subset \mathcal{S}$, we get the upper bound in (3.5).

Lower bound. We exploit Proposition 2.4 by making a judicious choice for the flow f . In fact, in the Glauber case, this choice will be simple: with each configuration $\sigma \in \mathcal{S}_L$, we associate a configuration in $\mathcal{C} \subset \mathcal{S}^c$ with a unique critical droplet and a flow that, from each such configuration, follows a unique deterministic path along which this droplet is broken down in the *canonical order* (see Figure 6) until the set \mathcal{S}_L is reached, that is, a square or quasi-square droplet with label L is left over [recall (1.7)–(1.8)].

Let $w(\beta)$ be such that

$$(3.11) \quad \lim_{\beta \rightarrow \infty} w(\beta) = \infty, \quad \lim_{\beta \rightarrow \infty} \frac{1}{\beta} \log w(\beta) = 0, \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta|/w(\beta) = \infty$$

and define

$$(3.12) \quad \mathcal{W} = \{\sigma \in \mathcal{S} : |\text{supp}[\sigma]| \leq |\Lambda_\beta|/w(\beta)\}.$$

Let $\mathcal{C}_L \subset \mathcal{C} \subset \mathcal{S}^c$ be the set of configurations obtained by picking any $\sigma \in \mathcal{S}_L \cap \mathcal{W}$ and adding somewhere in Λ_β a critical droplet at distance ≥ 2 from $\text{supp}[\sigma]$. Note that the density restriction imposed on \mathcal{W} guarantees that adding such a droplet is possible almost everywhere in Λ_β for β large enough. Denoting by $P_{(y)}(x)$ the critical droplet obtained by adding a protuberance at y along the longest side of the rectangle $R_{\ell_c-1, \ell_c}(x)$, we may write

$$(3.13) \quad \mathcal{C}_L = \{\sigma \cup P_{(y)}(x) : \sigma \in \mathcal{S} \cap \mathcal{W}, x, y \in \Lambda_\beta, (x, y) \perp \sigma\},$$

where $(x, y) \perp \sigma$ stands for the restriction that the critical droplet $P_{(y)}(x)$ is not interacting with $\text{supp}[\sigma]$, which implies that $H_\beta(\sigma \cup P_{(y)}(x)) = H_\beta(\sigma) + \Gamma$ (see Figures 7 and 8).




FIG. 7. The critical droplet $P_{(y)}(x)$.




FIG. 8. Going from \mathcal{S}_L to \mathcal{C}_L by adding a critical droplet $P_{(y)}(x)$ somewhere in Λ_β .

Now, for each $\sigma \in \mathcal{C}_L$, we let $\gamma_\sigma = (\gamma_\sigma(0), \gamma_\sigma(1), \dots, \gamma_\sigma(K))$ be the canonical path from $\sigma = \gamma_\sigma(0)$ to \mathcal{S}_L along which the critical droplet is broken down, where $K = v(2\ell_c - 3) - v(L)$ with

$$(3.14) \quad v(L) = |Q_L(0)|$$

[recall (1.7)]. We will choose our flow such that

$$(3.15) \quad f(\sigma', \sigma'') = \begin{cases} v_0(\sigma), & \text{if } \sigma' = \sigma, \sigma'' = \gamma_\sigma(1) \text{ for some } \sigma \in \mathcal{C}_L, \\ \sum_{\tilde{\sigma} \in \mathcal{C}_L} f(\gamma_{\tilde{\sigma}}(k-1), \gamma_{\tilde{\sigma}}(k)), & \\ 0, & \text{if } \sigma' = \gamma_\sigma(k), \sigma'' = \gamma_\sigma(k+1) \\ & \text{for some } k \geq 1, \sigma \in \mathcal{C}_L, \\ & \text{otherwise.} \end{cases}$$

Here, v_0 is some initial distribution on \mathcal{C}_L that will turn out to be arbitrary as long as its support is all of \mathcal{C}_L .

From (3.15), we see that the flow increases whenever paths merge. In our case, this happens only after the first step, when the protuberance at y is removed. Therefore, we get the explicit form

$$(3.16) \quad f(\sigma', \sigma'') = \begin{cases} v_0(\sigma), & \text{if } \sigma' = \sigma, \sigma'' = \gamma_\sigma(1) \text{ for some } \sigma \in \mathcal{C}_L, \\ C v_0(\sigma), & \text{if } \sigma' = \gamma_\sigma(k), \sigma'' = \gamma_\sigma(k+1) \\ & \text{for some } k \geq 1, \sigma \in \mathcal{C}_L, \\ 0, & \text{otherwise,} \end{cases}$$

where $C = 2\ell_c$ is the number of possible positions of the protuberance on the

proto-critical droplet (see Figure 6). Using Proposition 2.4, we therefore have

$$\begin{aligned}
 \text{CAP}(\mathcal{S}_L, \mathcal{S}^c) &= \text{CAP}(\mathcal{S}^c, \mathcal{S}_L) \geq \text{CAP}(\mathcal{C}_L, \mathcal{S}_L) \\
 (3.17) \quad &\geq \sum_{\sigma \in \mathcal{C}_L} \nu_0(\sigma) \left[\sum_{k=0}^{K-1} \frac{f(\gamma_\sigma(k), \gamma_\sigma(k+1))}{\mu_\beta(\gamma_\sigma(k))c_\beta(\gamma_\sigma(k), \gamma_\sigma(k+1))} \right]^{-1} \\
 &= \sum_{\sigma \in \mathcal{C}_L} \left[\frac{1}{\mu_\beta(\sigma)c_\beta(\gamma_\sigma(0), \gamma_\sigma(1))} \right. \\
 &\quad \left. + \sum_{k=1}^{K-1} \frac{C}{\mu_\beta(\gamma_\sigma(k))c_\beta(\gamma_\sigma(k), \gamma_\sigma(k+1))} \right]^{-1}.
 \end{aligned}$$

Thus, all we have to do is to control the sum between square brackets.

Because $c_\beta(\gamma_\sigma(0), \gamma_\sigma(1)) = 1$ (removing the protuberance lowers the energy), the term with $k = 0$ equals $1/\mu_\beta(\sigma)$. To show that the terms with $k \geq 1$ are of higher order, we argue as follows. Abbreviate $\Xi = h(\ell_c - 2)$. For every $k \geq 1$ and $\sigma(0) \in \mathcal{C}_L$, we have [see Figure 9 and recall (1.2) and (1.3)]

$$\begin{aligned}
 (3.18) \quad \mu_\beta(\gamma_\sigma(k))c_\beta(\gamma_\sigma(k), \gamma_\sigma(k+1)) &= \frac{1}{Z_\beta} e^{-\beta[H_\beta(\gamma_\sigma(k)) \vee H_\beta(\gamma_\sigma(k+1))]} \\
 &\geq \mu_\beta(\sigma_0) e^{\beta[2J-h-\Xi]} = \mu_\beta(\sigma) e^{\beta\delta},
 \end{aligned}$$

where $\delta = 2J - h - \Xi = 2J - h(\ell_c - 1) > 0$ [recall (1.6)]. Therefore,

$$(3.19) \quad \sum_{k=1}^{K-1} \frac{C}{\mu_\beta(\gamma_\sigma(k))c_\beta(\gamma_\sigma(k), \gamma_\sigma(k+1))} \leq \frac{1}{\mu_\beta(\sigma)} C K e^{-\delta\beta}$$




FIG. 9. Visualization of (3.18).

and so, from (3.17), we get

$$(3.20) \quad \text{CAP}(\mathcal{S}_L, \mathcal{S}^c) \geq \sum_{\sigma \in \mathcal{C}_L} \frac{\mu_\beta(\sigma)}{1 + CK e^{-\beta\delta}} = \frac{\mu_\beta(\mathcal{C}_L)}{1 + CK e^{-\beta\delta}} = [1 + o(1)]\mu_\beta(\mathcal{C}_L).$$

The last step is to estimate, with the help of (3.13),

$$(3.21) \quad \begin{aligned} \mu_\beta(\mathcal{C}_L) &= \frac{1}{Z_\beta} \sum_{\sigma \in \mathcal{C}_L} e^{-\beta H_\beta(\sigma)} = \frac{1}{Z_\beta} \sum_{\sigma \in \mathcal{S}_L \cap \mathcal{W}} \sum_{\substack{x, y \in \Lambda_\beta \\ (x, y) \perp \sigma}} e^{-\beta H_\beta(\sigma \cup P_{(y)}(x))} \\ &= e^{-\beta\Gamma} \frac{1}{Z_\beta} \sum_{\sigma \in \mathcal{S}_L \cap \mathcal{W}} e^{-\beta H_\beta(\sigma)} \sum_{\substack{x, y \in \Lambda_\beta \\ (x, y) \perp \sigma}} 1 \\ &\geq e^{-\beta\Gamma} \mu_\beta(\mathcal{S}_L \cap \mathcal{W}) N_1 |\Lambda_\beta| [1 - (\ell_c + 1)^2/w(\beta)]. \end{aligned}$$

The last inequality uses the fact that $|\Lambda_\beta|(\ell_c + 1)^2/w(\beta)$ is the maximal number of sites in Λ_β where it is not possible to insert a noninteracting critical droplet [recall (3.12) and note that a critical droplet fits inside an $\ell_c \times \ell_c$ square]. According to Lemma A.1 in Appendix A, we have

$$(3.22) \quad \mu_\beta(\mathcal{S}_L \cap \mathcal{W}) = \mu_\beta(\mathcal{S}_L)[1 + o(1)],$$

while conditions (1.8) and (1.9) imply that $\mu_\beta(\mathcal{S}_L) = \mu_\beta(\mathcal{S})[1 + o(1)]$. Combining the latter with (3.20) and (3.21), we obtain the desired lower bound. \square

3.2. *Proof of Theorem 1.2(b).* We use the same technique as in Section 3.1, which is why we only give a sketch of the proof.

To estimate the average crossover time from $\mathcal{S}_L \subset \mathcal{S}$ to $\mathcal{S}^c \setminus \mathcal{C}$, we will use Proposition 2.1. With $\mathcal{A} = \mathcal{S}_L$ and $\mathcal{B} = \mathcal{S}^c \setminus \mathcal{C}$, (2.10) reads

$$(3.23) \quad \sum_{\sigma \in \mathcal{S}_L} \nu_{\mathcal{S}_L}^{\mathcal{S}^c \setminus \mathcal{C}}(\sigma) \mathbb{E}_\sigma(\tau_{\mathcal{S}^c \setminus \mathcal{C}}) = \frac{1}{\text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C})} \sum_{\sigma \in \mathcal{S} \cup \mathcal{C}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}}(\sigma).$$

The left-hand side is the quantity of interest in (1.17).

In Sections 3.2.1 and 3.2.2 we estimate $\sum_{\sigma \in \mathcal{S} \cup \mathcal{C}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}}(\sigma)$ and $\text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C})$. The estimates will show that

$$(3.24) \quad \text{the right-hand side of (3.23)} = \frac{1}{N_2 |\Lambda_\beta|} e^{\beta\Gamma} [1 + o(1)], \quad \beta \rightarrow \infty.$$

3.2.1. *Estimate of $\sum_{\sigma \in \mathcal{S} \cup \mathcal{C}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}}(\sigma)$.*

LEMMA 3.3. $\sum_{\sigma \in \mathcal{S} \cup \mathcal{C}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}}(\sigma) = \mu_\beta(\mathcal{S})[1 + o(1)]$ as $\beta \rightarrow \infty$.

PROOF. Write, using (2.1),

$$\begin{aligned}
 (3.25) \quad & \sum_{\sigma \in \mathcal{S} \cup \mathcal{C}} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}}(\sigma) \\
 &= \mu_\beta(\mathcal{S}_L) + \sum_{\sigma \in (\mathcal{S} \setminus \mathcal{S}_L) \cup \mathcal{C}} \mu_\beta(\sigma) \mathbb{P}_\sigma(\tau_{\mathcal{S}_L} < \tau_{\mathcal{S}^c \setminus \mathcal{C}}).
 \end{aligned}$$

The last sum is bounded above by $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L) + \mu_\beta(\mathcal{C})$. As before, $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L) = o(\mu_\beta(\mathcal{S}))$ as $\beta \rightarrow \infty$. But (1.35) and (3.9) imply that $\mu_\beta(\mathcal{C}) = o(\mu_\beta(\mathcal{S}))$ as $\beta \rightarrow \infty$. \square

3.2.2. Estimate of $\text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C})$.

LEMMA 3.4. $\text{CAP}(\mathcal{S}, \mathcal{S}^c \setminus \mathcal{C}) = N_2 |\Lambda_\beta| e^{-\beta \Gamma} \mu_\beta(\mathcal{S}) [1 + o(1)]$ as $\beta \rightarrow \infty$ with $N_2 = \frac{4}{3}(2\ell_c - 1)$.

PROOF. The proof is similar to that of Lemma 3.2, except that it takes care of the transition probabilities away from the critical droplet.

Upper bound. Recalling (2.13)–(2.15) and noting that Glauber dynamics does not allow transitions within \mathcal{C} , we have, for all $h : \mathcal{C} \rightarrow [0, 1]$,

$$\begin{aligned}
 (3.26) \quad & \text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}) \leq \text{CAP}(\mathcal{S}, \mathcal{S}^c \setminus \mathcal{C}) \\
 & \leq \sum_{\sigma \in \mathcal{C}} \mu_\beta(\sigma) [\hat{c}_\sigma (h(\sigma) - 1)^2 + \check{c}_\sigma (h(\sigma) - 0)^2],
 \end{aligned}$$

where $\hat{c}_\sigma = \sum_{\eta \in \mathcal{S}} c_\beta(\sigma, \eta)$ and $\check{c}_\sigma = \sum_{\eta \in \mathcal{S}^c \setminus \mathcal{C}} c_\beta(\sigma, \eta)$. The quadratic form in the right-hand side of (3.26) achieves its minimum for $h(\sigma) = \hat{c}_\sigma / (\hat{c}_\sigma + \check{c}_\sigma)$, so

$$(3.27) \quad \text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}) \leq \sum_{\sigma \in \mathcal{C}} C_\sigma \mu_\beta(\sigma)$$

with $C_\sigma = \hat{c}_\sigma \check{c}_\sigma / (\hat{c}_\sigma + \check{c}_\sigma)$. We have

$$\begin{aligned}
 (3.28) \quad & \sum_{\sigma \in \mathcal{C}} C_\sigma \mu_\beta(\sigma) = \frac{1}{Z_\beta} \sum_{\sigma \in \mathcal{P}} \sum_{\substack{x \in \Lambda_\beta \\ \sigma^x \in \mathcal{C}}} C_{\sigma^x} e^{-\beta H_\beta(\sigma^x)} \\
 &= e^{-\beta(2J-h)} \frac{1}{Z_\beta} \sum_{\sigma \in \mathcal{P}} e^{-\beta H_\beta(\sigma)} 2 \left(\frac{1}{2} 4 + \frac{2}{3} (2\ell_c - 4) \right) \\
 &= e^{-\beta(2J-h)} \mu_\beta(\mathcal{P}) N_2 = \frac{1}{N_1} \mu_\beta(\mathcal{C}) N_2,
 \end{aligned}$$

where, in the second line, we use the fact that $C_\sigma = \frac{1}{2}$ if σ has a protuberance in a corner (2×4 choices) and $C_\sigma = \frac{2}{3}$ otherwise [$2 \times (2\ell_c - 4)$ choices].




FIG. 10. Canonical order to break down a proto-critical droplet plus a double protuberance. In the first step, the double protuberance has probability $\frac{1}{2}$ to be broken down in either of the two possible ways. The subsequent steps are deterministic, as in Figure 6.

Lower bound. In analogy with (3.13), denoting by $P_{(y)}^2(x)$ the droplet obtained by adding a double protuberance at y along the longest side of the rectangle $R_{\ell_c-1, \ell_c}(x)$, we define the set $\mathcal{D}_L \subset \mathcal{S}^c \setminus \mathcal{C}$ by

$$(3.29) \quad \mathcal{D}_L = \{\sigma \cup P_{(y)}^2(x) : \sigma \in \mathcal{S}_L \cap \mathcal{W}, x, y \in \Lambda_\beta, (x, y) \perp \sigma\}.$$

As in (3.15), we may choose any starting measure on \mathcal{D}_L . We choose the flow as follows. For the first step, we choose

$$(3.30) \quad f(\sigma', \sigma) = \frac{1}{2} \nu_0(\sigma), \quad \sigma' \in \mathcal{D}_L, \sigma \in \mathcal{C}_L,$$

which reduces the double protuberance to a single protuberance [compare (3.13) and (3.29)]. For all subsequent steps, we follow the deterministic paths γ_σ used in Section 3.1.2, which start from $\gamma_\sigma(0) = \sigma$ (see Figure 10). Note, however, that we get different values for the flows $f(\gamma_\sigma(0), \gamma_\sigma(1))$ depending on whether or not the protuberance sits in a corner. In the former case, it has only one possible antecedent, and so

$$(3.31) \quad f(\gamma_\sigma(0), \gamma_\sigma(1)) = \frac{1}{2} \nu_0(\sigma),$$

while in the latter case it has two antecedents, and so

$$(3.32) \quad f(\gamma_\sigma(0), \gamma_\sigma(1)) = \nu_0(\sigma).$$

This time, the terms $k = 0$ and $k = 1$ are of the same order, while, as in (3.19), all of the subsequent steps give a contribution that is a factor $O(e^{-\delta\beta})$ smaller. Indeed, in analogy with (3.17), we obtain, writing $\sigma \sim \sigma'$ when $c_\beta(\sigma', \sigma) > 0$,

$$(3.33) \quad \begin{aligned} & \text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C}) \\ &= \text{CAP}(\mathcal{S}^c \setminus \mathcal{C}, \mathcal{S}_L) \geq \text{CAP}(\mathcal{D}_L, \mathcal{S}_L) \\ &\geq \sum_{\sigma' \in \mathcal{D}_L} \frac{1}{2} \sum_{\substack{\sigma \in \mathcal{C}_L \\ \sigma \sim \sigma'}} \left[\frac{f(\sigma', \sigma)}{\mu_\beta(\sigma)} + \frac{f(\sigma, \gamma_\sigma(1))}{\mu_\beta(\sigma)} \right. \\ &\quad \left. + \sum_{k=1}^{K-1} \frac{f(\gamma_\sigma(k), \gamma_\sigma(k+1))}{\mu_\beta(\gamma_\sigma(k)) c_\beta(\gamma_\sigma(k), \gamma_\sigma(k+1))} \right]^{-1} \end{aligned}$$

$$\begin{aligned}
 &\geq \sum_{\sigma' \in \mathcal{D}_L} \frac{1}{2} \sum_{\substack{\sigma \in \mathcal{C}_L \\ \sigma \sim \sigma'}} \mu_\beta(\sigma) [f(\sigma', \sigma) + f(\sigma, \gamma_\sigma(1)) + CK e^{-\beta\delta}]^{-1} \\
 &= [1 + o(1)] \mu_\beta(\mathcal{C}_L) \left(\frac{2\ell_c - 4}{2\ell_c} \frac{1}{1 + 1/2} + \frac{1}{2} \frac{4}{2\ell_c} \frac{1}{1/2 + 1/2} \right) \\
 &= [1 + o(1)] \mu_\beta(\mathcal{C}_L) \frac{N_2}{N_1}.
 \end{aligned}$$

Using (3.21) and the remarks following it, we get the desired lower bound. \square

3.3. Proof of Theorem 1.2(c). Write

$$\begin{aligned}
 &\sum_{\sigma \in \mathcal{D}_M^c} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{D}_M}(\sigma) \\
 (3.34) \quad &= \sum_{\sigma \in \mathcal{S}_L} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{D}_M}(\sigma) + \sum_{\sigma \in \mathcal{D}_M^c \setminus \mathcal{S}_L} \mu_\beta(\sigma) h_{\mathcal{S}_L, \mathcal{D}_M}(\sigma) \\
 &= \mu_\beta(\mathcal{S}_L) + \sum_{\sigma \in \mathcal{D}_M^c \setminus \mathcal{S}_L} \mu_\beta(\sigma) \mathbb{P}_\sigma(\tau_{\mathcal{S}_L} < \tau_{\mathcal{D}_M}).
 \end{aligned}$$

The last sum is bounded above by $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L) + \mu_\beta(\mathcal{D}_M^c \setminus \mathcal{S})$. But $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L) = o(\mu_\beta(\mathcal{S}))$, as $\beta \rightarrow \infty$ by our choice of L in (1.9), while $\mu_\beta(\mathcal{D}_M^c \setminus \mathcal{S}) = o(\mu_\beta(\mathcal{S}))$ as $\beta \rightarrow \infty$ because of the restriction $\ell_c \leq M2\ell_c - 1$. Indeed, under that restriction, the energy of a square droplet of size M is strictly larger than the energy of a critical droplet.

PROOF OF THEOREM 1.2(c). The proof follows along the same lines as that of Theorems 1.2(a) and (b) in Sections 3.1 and 3.2. The main point is to prove that $\text{CAP}(\mathcal{S}_L, \mathcal{D}_M) = [1 + o(1)] \text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C})$. Since $\text{CAP}(\mathcal{S}_L, \mathcal{D}_M) \leq \text{CAP}(\mathcal{S}_L, \mathcal{S}^c \setminus \mathcal{C})$, which was estimated in Section 3.2, we need only prove a lower bound on $\text{CAP}(\mathcal{S}_L, \mathcal{D}_M)$. This is done by using a flow that breaks down an $M \times M$ droplet to a square or quasi-square droplet Q_L in the canonical way, which takes $M^2 - v(L)$ steps [recall Figure 6 and (3.14)]. The leading terms are still the proto-critical droplet with a single and a double protuberance. With each $M \times M$ droplet is associated a unique critical droplet so that the pre-factor in the lower bound is the same as in the proof of Theorem 1.2(b).

Note that we can even allow M to grow with β as $M = e^{o(\beta)}$. Indeed, (3.11) and (3.12) show that there is enough room to add a droplet of size $e^{o(\beta)}$ almost everywhere in Λ_β and the factor $M^2 e^{-\delta\beta}$ replacing $K e^{-\delta\beta}$ in (3.20) is still $o(1)$. \square

4. Proof of Theorem 1.4.

4.1. *Proof of Theorem 1.4(a).*

4.1.1. *Estimate of $\sum_{\sigma \in \text{SU}(\tilde{\mathcal{C}} \setminus \mathcal{C}^+)} \mu_\beta(\sigma) h_{\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+}(\sigma)$.*

LEMMA 4.1. $\sum_{\sigma \in \text{SU}(\tilde{\mathcal{C}} \setminus \mathcal{C}^+)} \mu_\beta(\sigma) h_{\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+}(\sigma) = \mu_\beta(\mathcal{S})[1 + o(1)]$ as $\beta \rightarrow \infty$.

PROOF. Write, using (2.1),

$$(4.1) \quad \begin{aligned} & \sum_{\sigma \in \text{SU}(\tilde{\mathcal{C}} \setminus \mathcal{C}^+)} \mu_\beta(\sigma) h_{\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+}(\sigma) \\ &= \mu_\beta(\mathcal{S}_L) + \sum_{\sigma \in (\mathcal{S} \setminus \mathcal{S}_L) \cup (\tilde{\mathcal{C}} \setminus \mathcal{C}^+)} \mu_\beta(\sigma) \mathbb{P}_\sigma(\tau_{\mathcal{S}_L} < \tau_{(\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+}). \end{aligned}$$

The last sum is bounded above by $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L) + \mu_\beta(\tilde{\mathcal{C}} \setminus \mathcal{C}^+)$, but $\mu_\beta(\mathcal{S} \setminus \mathcal{S}_L) = o(\mu_\beta(\mathcal{S}))$ as $\beta \rightarrow \infty$ by our choice of L in (1.32). In Lemma B.3 in Appendix B.3, we will show that $\mu_\beta(\tilde{\mathcal{C}} \setminus \mathcal{C}^+) = o(\mu_\beta(\mathcal{S}))$ as $\beta \rightarrow \infty$. \square

4.1.2. *Estimate of $\text{CAP}(\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+)$.*

LEMMA 4.2. $\text{CAP}(\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+) = N |\Lambda_\beta|^{\frac{4\pi}{\beta\Delta}} e^{-\beta\Gamma} \mu_\beta(\mathcal{S})[1 + o(1)]$ as $\beta \rightarrow \infty$ with $N = \frac{1}{3} \ell_c^2 (\ell_c^2 - 1)$.

PROOF. The argument is in the same spirit as that in Section 3.1.2. However, a number of additional obstacles that arise from the conservative nature of Kawasaki dynamics need to be overcome. The proof proceeds via upper and lower bounds and takes up quite a bit of space.

Upper bound. The proof consists of seven steps.

1. *Proto-critical droplet (see Figure 11) and free particle.* We have

$$(4.2) \quad \begin{aligned} & \text{CAP}(\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+) \\ & \leq \text{CAP}(\mathcal{S} \cup \mathcal{C}^-, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+) \\ & = \min_{\substack{h: \mathcal{X}_\beta^{(n_\beta)} \rightarrow [0,1] \\ h|_{\mathcal{S} \cup \mathcal{C}^-} \equiv 1, h|_{(\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+} \equiv 0}} \frac{1}{2} \sum_{\sigma, \sigma' \in \mathcal{X}_\beta^{(n_\beta)}} \mu_\beta(\sigma) c_\beta(\sigma, \sigma') [h(\sigma) - h(\sigma')]^2. \end{aligned}$$

Split the right-hand side into a contribution coming from $\sigma, \sigma' \in \tilde{\mathcal{C}}$ and the rest:

$$(4.3) \quad \text{right-hand side of (4.2)} = I + \gamma_1(\beta),$$




FIG. 11. Schematic picture of the sets S, C^-, C^+ defined in Definition 1.3 and the set \tilde{C} interpolating between C^- and C^+ .

where

$$(4.4) \quad I = \min_{\substack{h: \tilde{C} \rightarrow [0,1] \\ h|_{C^-} \equiv 1, h|_{C^+} \equiv 0}} \frac{1}{2} \sum_{\sigma, \sigma' \in \tilde{C}} \mu_\beta(\sigma) c_\beta(\sigma, \sigma') [h(\sigma) - h(\sigma')]^2$$

and $\gamma_1(\beta)$ is an error term that will be estimated in Step 7. This term will turn out to be small because $\mu_\beta(\sigma) c_\beta(\sigma, \sigma')$ is small when either $\sigma \in \mathcal{X}_\beta^{(n_\beta)} \setminus \tilde{C}$ or $\sigma' \in \mathcal{X}_\beta^{(n_\beta)} \setminus \tilde{C}$. Next, partition \tilde{C}, C^-, C^+ into sets $\tilde{C}(x), C^-(x), C^+(x), x \in \Lambda_\beta$, by requiring that the lower-left corner of the proto-critical droplet is in the center of the box $B_{L_\beta, L_\beta}(x)$. Then, because $c_\beta(\sigma, \sigma') = 0$ when $\sigma \in \tilde{C}(x)$ and $\sigma' \in \tilde{C}(x')$ for some $x \neq x'$, we may write

$$(4.5) \quad I = |\Lambda_\beta| \min_{\substack{h: \tilde{C}(0) \rightarrow [0,1] \\ h|_{C^-(0)} \equiv 1, h|_{C^+(0)} \equiv 0}} \frac{1}{2} \sum_{\sigma, \sigma' \in \tilde{C}(0)} \mu_\beta(\sigma) c_\beta(\sigma, \sigma') [h(\sigma) - h(\sigma')]^2.$$

2. Decomposition of configurations. Define [cf. (3.6)]

$$(4.6) \quad \begin{aligned} \hat{C}(0) &= \{\sigma \mathbb{1}_{B_{L_\beta, L_\beta}(0)} : \sigma \in \tilde{C}(0)\}, \\ \check{C}(0) &= \{\sigma \mathbb{1}_{[B_{L_\beta, L_\beta}(0)]^c} : \sigma \in \tilde{C}(0)\}. \end{aligned}$$

Every $\sigma \in \tilde{C}(0)$ can then be uniquely decomposed as $\sigma = \hat{\sigma} \vee \check{\sigma}$ for some $\hat{\sigma} \in \hat{C}(0)$ and $\check{\sigma} \in \check{C}(0)$. Note that $\hat{C}(0)$ has $K = \ell_c(\ell_c - 1) + 2$ particles and $\check{C}(0)$ has $n_\beta - K$ particles [and recall that, by the first half of (1.35), $n_\beta \rightarrow \infty$ as $\beta \rightarrow \infty$]. Define

$$(4.7) \quad C^{fp}(0) = \{\sigma \in \tilde{C}(0) : H_\beta(\sigma) = H_\beta(\hat{\sigma}) + H_\beta(\check{\sigma})\},$$

that is, the set of configurations consisting of a proto-critical droplet and a free particle inside $B_{L_\beta, L_\beta}(0)$ not interacting with the particles outside $B_{L_\beta, L_\beta}(0)$. Write $C^{fp,-}(0)$ [resp. $C^{fp,+}(0)$] to denote the subsets of $C^{fp}(0)$ where the free particle is

at distance L_β (resp. 2) from the proto-critical droplet. Split the right-hand side of (4.5) into a contribution coming from $\sigma, \sigma' \in \mathcal{C}^{\text{fp}}(0)$ and the rest:

$$(4.8) \quad \text{right-hand side of (4.5)} = |\Lambda_\beta| [II + \gamma_2(\beta)],$$

where

$$(4.9) \quad II = \min_{\substack{h: \mathcal{C}^{\text{fp}}(0) \rightarrow [0,1] \\ h|_{\mathcal{C}^{\text{fp},-}(0)} \equiv 1, h|_{\mathcal{C}^{\text{fp},+}(0)} \equiv 0}} \frac{1}{2} \sum_{\sigma, \sigma' \in \mathcal{C}^{\text{fp}}(0)} \mu_\beta(\sigma) c_\beta(\sigma, \sigma') [h(\sigma) - h(\sigma')]^2$$

and $\gamma_2(\beta)$ is an error term that will be estimated in Step 6. This term will turn out to be small because of loss of entropy when the particle is at the boundary.

3. *Reduction to capacity of simple random walk.* Estimate

$$(4.10) \quad \begin{aligned} II &= \min_{\substack{h: \mathcal{C}^{\text{fp}}(0) \rightarrow [0,1] \\ h|_{\mathcal{C}^{\text{fp},-}(0)} \equiv 1, h|_{\mathcal{C}^{\text{fp},+}(0)} \equiv 0}} \frac{1}{2} \sum_{\check{\sigma}, \check{\sigma}' \in \check{\mathcal{C}}(0)} \sum_{\substack{\hat{\sigma}, \hat{\sigma}' \in \hat{\mathcal{C}}(0): \\ \hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma}' \in \mathcal{C}^{\text{fp}}(0)}} \mu_\beta(\hat{\sigma} \vee \check{\sigma}) \\ &\quad \times c_\beta(\hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma}') \\ &\quad \times [h(\hat{\sigma} \vee \check{\sigma}) - h(\hat{\sigma}' \vee \check{\sigma}')]^2 \\ &\leq \min_{\substack{g: \hat{\mathcal{C}}(0) \rightarrow [0,1] \\ g|_{\hat{\mathcal{C}}^-(0)} \equiv 1, g|_{\hat{\mathcal{C}}^+(0)} \equiv 0}} \frac{1}{2} \sum_{\check{\sigma} \in \check{\mathcal{C}}(0)} \sum_{\substack{\hat{\sigma}, \hat{\sigma}' \in \hat{\mathcal{C}}(0): \\ \hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma}' \in \mathcal{C}^{\text{fp}}(0)}} \mu_\beta(\hat{\sigma} \vee \check{\sigma}) c_\beta(\hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma}') \\ &\quad \times [g(\hat{\sigma}) - g(\hat{\sigma}')]^2, \end{aligned}$$

where $\hat{\mathcal{C}}^-(0)$ [resp. $\hat{\mathcal{C}}^+(0)$] denote the subsets of $\hat{\mathcal{C}}(0)$ where the free particle is at distance L_β (resp. 2) from the proto-critical droplet and the inequality comes from substituting

$$(4.11) \quad h(\hat{\sigma} \vee \check{\sigma}) = g(\hat{\sigma}), \quad \hat{\sigma} \in \hat{\mathcal{C}}(0), \check{\sigma} \in \check{\mathcal{C}}(0),$$

and afterwards replacing the double sum over $\check{\sigma}, \check{\sigma}' \in \check{\mathcal{C}}(0)$ by the single sum over $\check{\sigma} \in \check{\mathcal{C}}(0)$ because $c_\beta(\hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma}') > 0$ only if either $\hat{\sigma} = \hat{\sigma}'$ or $\check{\sigma} = \check{\sigma}'$ (the dynamics updates one site at a time). Next, estimate

$$(4.12) \quad \begin{aligned} &\text{right-hand side of (4.10)} \\ &\leq \sum_{\check{\sigma} \in \check{\mathcal{C}}(0)} \frac{1}{Z_\beta^{(n_\beta)}} e^{-\beta H_\beta(\check{\sigma})} \\ &\quad \times \min_{\substack{g: \hat{\mathcal{C}}(0) \rightarrow [0,1] \\ g|_{\hat{\mathcal{C}}^-(0)} \equiv 1, g|_{\hat{\mathcal{C}}^+(0)} \equiv 0}} \frac{1}{2} \sum_{\substack{\hat{\sigma}, \hat{\sigma}' \in \hat{\mathcal{C}}(0) \\ \hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma}' \in \mathcal{C}^{\text{fp}}(0)}} e^{-\beta H_\beta(\hat{\sigma})} c_\beta(\hat{\sigma}, \hat{\sigma}') \\ &\quad \times [g(\hat{\sigma}) - g(\hat{\sigma}')]^2, \end{aligned}$$

where we have used $H_\beta(\sigma) = H_\beta(\hat{\sigma}) + H_\beta(\check{\sigma})$ from (4.7) and have written $c_\beta(\hat{\sigma}, \hat{\sigma}')$ to denote the transition rate associated with the Kawasaki dynamics restricted to $B_{L_\beta, L_\beta}(0)$, which clearly equals $c_\beta(\hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma})$ for every $\check{\sigma} \in \check{\mathcal{C}}(0)$ such that $\hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma} \in \mathcal{C}^{\text{fp}}(0)$ because there is no interaction between the particles inside and outside $B_{L_\beta, L_\beta}(0)$. The minimum in the right-hand side of (4.12) can be estimated from above by

$$(4.13) \quad \text{minimum in (4.12)} \leq \sum_{\sigma \in \mathcal{P}(0)} \mathcal{V}_\beta(\sigma)$$

with $\mathcal{P}(0)$ the set of proto-critical droplets with lower-left corner at 0 and

$$(4.14) \quad \mathcal{V}_\beta(\sigma) = \min_{\substack{f: \mathbb{Z}^2 \rightarrow [0,1] \\ f|_{P_\sigma(0)} \equiv 1, f|_{[B_{L_\beta, L_\beta}(0)]^c} \equiv 0}} \frac{1}{2} \sum_{\substack{x, x' \in \mathbb{Z}^2 \\ x \sim x'}} [f(x) - f(x')]^2,$$

where $P_\sigma(0)$ is the support of the proto-critical droplet in σ and $x \sim x'$ means that x and x' are neighboring sites. Indeed, (4.13) is obtained from the expression in (4.12) by dropping the restriction $\hat{\sigma} \vee \check{\sigma}, \hat{\sigma}' \vee \check{\sigma} \in \mathcal{C}^{\text{fp}}(0)$, substituting

$$(4.15) \quad g(P_\sigma(0) \cup \{x\}) = f(x), \quad \sigma \in \mathcal{P}(0), x \in B_{L_\beta, L_\beta}(0) \setminus P_\sigma(0),$$

and noting that $c_\beta(P_\sigma(0) \cup \{x\}, P_\sigma(0) \cup \{x'\}) = 1$ when $x \sim x'$ and 0 otherwise. What (4.14) says is that

$$(4.16) \quad \mathcal{V}_\beta(\sigma) = \text{CAP}(P_\sigma(0), [B_{L_\beta, L_\beta}(0)]^c)$$

is the capacity of simple random walk between the proto-critical droplet $P_\sigma(0)$ in σ and the exterior of $B_{L_\beta, L_\beta}(0)$. Now, define

$$(4.17) \quad \check{Z}_\beta^{(n-K)}(0) = \sum_{\check{\sigma} \in \check{\mathcal{C}}(0)} e^{-\beta H_\beta(\check{\sigma})}.$$

We then obtain from (4.12) and (4.13) that

$$(4.18) \quad \text{right-hand side (4.12)} \leq e^{-\beta \Gamma^*} \frac{\check{Z}_\beta^{(n-K)}(0)}{Z_\beta^{(n\beta)}} \sum_{\sigma \in \mathcal{P}(0)} \mathcal{V}_\beta(\sigma),$$

where $\Gamma^* = -U[(\ell_c - 1)^2 + \ell_c(\ell_c - 1) + 1]$ is the binding energy of the proto-critical droplet [cf. (1.33)].

4. *Capacity estimate.* For future reference, we state the following estimate on capacities for simple random walk.

LEMMA 4.3. *Let $U \subset \mathbb{Z}^2$ be any set such that $\{0\} \subset U \subset B_{k,k}(0)$, with $k \in \mathbb{N} \cup \{0\}$ independent of β . Let $V \subset \mathbb{Z}^2$ be any set such that $[B_{KL_\beta, KL_\beta}(0)]^c \subset V \subset [B_{L_\beta, L_\beta}(0)]^c$, with $K \in \mathbb{N}$ independent of β . Then*

$$(4.19) \quad \begin{aligned} \text{CAP}(\{0\}, [B_{KL_\beta, KL_\beta}(0)]^c) &\leq \text{CAP}(U, V) \\ &\leq \text{CAP}(B_{k,k}(0), [B_{L_\beta, L_\beta}(0)]^c). \end{aligned}$$

Moreover, via (1.29) and (1.30),

$$\begin{aligned}
 \text{CAP}(B_{k,k}(0), [B_{KL_\beta, KL_\beta}(0)]^c) &= [1 + o(1)] \frac{2\pi}{\log(KL_\beta) - \log k} \\
 (4.20) \qquad \qquad \qquad &= [1 + o(1)] \frac{4\pi}{\beta\Delta}, \quad \beta \rightarrow \infty.
 \end{aligned}$$

PROOF. The inequalities in (4.19) follow from standard monotonicity properties of capacities. The asymptotic estimates in (4.20) for capacities of concentric boxes are standard (see, e.g., Lawler [20], Section 2.3) and also follow by comparison to Brownian motion. \square

We can apply Lemma 4.3 to estimate $\mathcal{V}_\beta(\sigma)$ in (4.16) since the proto-critical droplet with lower-left corner in 0 fits inside the box $B_{2\ell_c, 2\ell_c}(0)$. This gives

$$(4.21) \qquad \mathcal{V}_\beta(\sigma) = \frac{4\pi}{\beta\Delta} [1 + o(1)] \quad \forall \sigma \in \mathcal{P}(0), \beta \rightarrow \infty.$$

Moreover, from Bovier, den Hollander and Nardi [7], Lemmas 3.4.2 and 3.4.3, we know that $N = |\mathcal{P}(0)|$, the number of shapes of the proto-critical droplet, equals $N = \frac{1}{3}\ell_c^2(\ell_c^2 - 1)$.

5. *Equivalence of ensembles.* According to Lemma B.1 in Appendix B, we have

$$(4.22) \qquad \frac{\check{Z}_\beta^{(n_\beta - K)}(0)}{Z_\beta^{(n_\beta)}} = (\rho_\beta)^K \mu_\beta(\mathcal{S}) [1 + o(1)], \quad \beta \rightarrow \infty.$$

This is an ‘‘equivalence of ensembles’’ property relating the probabilities to find $n_\beta - K$ (resp. n_β) particles inside $[B_{L_\beta, L_\beta}(0)]^c$ [recall (4.6)]. Combining (4.2), (4.3), (4.5), (4.8), (4.10), (4.12), (4.18), (4.21) and (4.22), we get

$$(4.23) \qquad \text{CAP}(\mathcal{S}, \mathcal{C}^+) \leq \gamma_1(\beta) + |\Lambda_\beta| \gamma_2(\beta) + N |\Lambda_\beta| \frac{4\pi}{\beta\Delta} e^{-\beta\Gamma} \mu_\beta(\mathcal{S}) [1 + o(1)],$$

$\beta \rightarrow \infty,$

where we have used the fact that $\Gamma^* + \Delta K = \Gamma$ defined in (1.33). This completes the proof of the upper bound, provided that the error terms $\gamma_1(\beta)$ and $\gamma_2(\beta)$ are negligible.

6. *Second error term.* To estimate the error term $\gamma_2(\beta)$, note that the configurations in $\tilde{\mathcal{C}}(0) \setminus \mathcal{C}^{\text{fp}}(0)$ are those for which, inside $B_{L_\beta, L_\beta}(0)$, there is a proto-critical droplet whose lower-left corner is at 0 and a particle that is at the boundary and attached to some cluster outside $B_{L_\beta, L_\beta}(0)$. Recalling (4.5)–(4.9), we therefore have

$$\begin{aligned}
 \gamma_2(\beta) &\leq \sum_{\sigma \in \tilde{\mathcal{C}}(0) \setminus \mathcal{C}^{\text{fp}}(0)} \sum_{\sigma' \in \tilde{\mathcal{C}}(0)} \mu_\beta(\sigma) c_\beta(\sigma, \sigma') [h(\sigma) - h(\sigma')]^2 \\
 (4.24) \qquad \qquad &\leq 6\mu_\beta(\tilde{\mathcal{C}}(0) \setminus \mathcal{C}^{\text{fp}}(0)),
 \end{aligned}$$

where we have used the facts that $h : \tilde{\mathcal{C}}(0) \rightarrow [0, 1]$, $\mu_\beta(\sigma)c_\beta(\sigma, \sigma') = \mu_\beta(\sigma) \wedge \mu_\beta(\sigma')$ and there are six possible transitions from $\tilde{\mathcal{C}}(0) \setminus \mathcal{C}^{\text{fp}}(0)$ to $\tilde{\mathcal{C}}(0)$: three through a move by the particle at the boundary of $B_{L_\beta, L_\beta}(0)$ and three through a move by a particle in the cluster outside $B_{L_\beta, L_\beta}(0)$. Since

$$(4.25) \quad H_\beta(\sigma) \geq H_\beta(\hat{\sigma}) + H_\beta(\check{\sigma}) - U, \quad \sigma \in \tilde{\mathcal{C}}(0) \setminus \mathcal{C}^{\text{fp}}(0),$$

it follows from the same argument as in Steps 3 and 5 that

$$(4.26) \quad \mu_\beta(\tilde{\mathcal{C}}(0) \setminus \mathcal{C}^{\text{fp}}(0)) \leq N e^{-\beta\Gamma^*} (\rho_\beta)^{K+1} \mu_\beta(\mathcal{S}) e^{\beta U} 4(K-1)[1 + o(1)],$$

where $(\rho_\beta)^{K+1}$ comes from the fact that $n_\beta - (K + 1)$ particles are outside $B_{L_\beta-1, L_\beta-1}(0)$ (again, use Lemma B.1 in Appendix B), $e^{\beta U}$ comes from the gap in (4.25) and $4(K - 1)$ counts the maximal number of places at the boundary of $B_{L_\beta, L_\beta}(0)$ where the particle can interact with particles outside $B_{L_\beta, L_\beta}(0)$ due to the constraint that defines \mathcal{S} [recall Definition 1.3(a)]. Since $\rho_\beta e^{\beta U} = o(1)$ by (1.27), we therefore see that $\gamma_2(\beta)$ is indeed small compared to the main term of (4.23).

7. *First error term.* To estimate the error term $\gamma_1(\beta)$, we define the sets of pairs of configurations

$$(4.27) \quad \begin{aligned} \mathcal{I}_1 &= \{(\sigma, \eta) \in [\mathcal{X}_\beta^{(n_\beta)}]^2 : \sigma \in \mathcal{S}, \eta \in \mathcal{S}^c \setminus \tilde{\mathcal{C}}\}, \\ \mathcal{I}_2 &= \{(\sigma, \eta) \in [\mathcal{X}_\beta^{(n_\beta)}]^2 : \sigma \in \tilde{\mathcal{C}}, \eta \in \mathcal{S}^c \setminus \tilde{\mathcal{C}}\} \end{aligned}$$

and estimate

$$(4.28) \quad \gamma_1(\beta) \leq \frac{1}{2} \sum_{i=1}^2 \sum_{(\sigma, \eta) \in \mathcal{I}_i} \mu_\beta(\sigma)c_\beta(\sigma, \eta) = \frac{1}{2} \Sigma(\mathcal{I}_1) + \frac{1}{2} \Sigma(\mathcal{I}_2).$$

The sum $\Sigma(\mathcal{I}_1)$ can be written as

$$(4.29) \quad \begin{aligned} \Sigma(\mathcal{I}_1) &= |\Lambda_\beta| \sum_{\sigma \in \mathcal{P}} \sum_{\eta \in \mathcal{S}^c \setminus \tilde{\mathcal{C}}} c_\beta(\eta, \sigma) 1\{|\text{supp}[\eta] \cap B_{L_\beta, L_\beta}(0)| = K\} \\ &\quad \times \frac{1}{Z_\beta^{(n_\beta)}} e^{-\beta H_\beta(\eta)}, \end{aligned}$$

where we have used the facts that $\mu_\beta(\sigma)c_\beta(\sigma, \eta) = \mu_\beta(\eta)c_\beta(\eta, \sigma)$, $\sigma, \eta \in \mathcal{X}_\beta^{(n_\beta)}$ and $c_\beta(\eta, \sigma) = 0$, $\eta \in \mathcal{S}^c \setminus \tilde{\mathcal{C}}$, $\sigma \notin \mathcal{P}$ [recall Definition 1.3(b)]. We have

$$(4.30) \quad H_\beta(\eta) \geq H_\beta(\hat{\eta}) + H_\beta(\check{\eta}) - kU, \quad \eta \in \mathcal{S}^c \setminus \tilde{\mathcal{C}},$$

where k counts the number of pairs of particles interacting across the boundary of $B_{L_\beta, L_\beta}(0)$. Moreover, since $\eta \notin \tilde{\mathcal{C}}$, we have

$$(4.31) \quad H_\beta(\hat{\eta}) \geq \Gamma^* + U.$$

Inserting (4.30) and (4.31) into (4.29), we obtain

$$\begin{aligned}
 \Sigma(\mathcal{I}_1) &\leq |\Lambda_\beta| e^{-\beta\Gamma^*} \mu_\beta(\mathcal{S}) [1 + o(1)] \sum_{k=0}^K (\rho_\beta)^{K+k} [4(K-1)]^k e^{\beta(k-1)U} \\
 (4.32) \qquad &= |\Lambda_\beta| e^{-\beta\Gamma} \mu_\beta(\mathcal{S}) [1 + o(1)] e^{-\beta U},
 \end{aligned}$$

where $(\rho_\beta)^{K+k}$ comes from the fact that $n_\beta - (K + k)$ particles are outside $B_{L_\beta-1, L_\beta-1}(0)$ (again, use Lemma B.1 in Appendix B) and the inequality again uses an argument similar to the arguments used in Steps 3 and 5. Therefore, $\Sigma(\mathcal{I}_1)$ is small compared to the main term of (4.23). The sum $\Sigma(\mathcal{I}_2)$ can be estimated as

$$\begin{aligned}
 \Sigma(\mathcal{I}_2) &= \sum_{\sigma \in \tilde{\mathcal{C}}} \sum_{\eta \in \mathcal{S}^c \setminus \tilde{\mathcal{C}}} \mu_\beta(\sigma) c_\beta(\sigma, \eta) \\
 (4.33) \qquad &= |\Lambda_\beta| \sum_{\sigma \in \tilde{\mathcal{C}}(0)} \mu_\beta(\sigma) \sum_{\eta \in \mathcal{S}^c \setminus \tilde{\mathcal{C}}(0)} c_\beta(\sigma, \eta) \\
 &\leq |\Lambda_\beta| \mu_\beta(\tilde{\mathcal{C}}(0)) \{e^{-\beta U} + (4L_\beta)\rho_\beta[1 + o(1)]\},
 \end{aligned}$$

where the first term comes from detaching a particle from the critical droplet and the second term from a extra particle entering $B_{L_\beta, L_\beta}(0)$. The term between braces is $o(1)$. Moreover, $\mu_\beta(\tilde{\mathcal{C}}(0)) = \mu_\beta(\mathcal{C}^{\text{fp}}(0)) + \mu_\beta(\tilde{\mathcal{C}}(0) \setminus \mathcal{C}^{\text{fp}}(0))$. The second term was estimated in (4.26) and the first term can again be estimated as in Steps 3 and 5:

$$\begin{aligned}
 \mu_\beta(\mathcal{C}^{\text{fp}}(0)) &= \sum_{\hat{\sigma} \in \hat{\mathcal{C}}(0)} \sum_{\check{\sigma} \in \check{\mathcal{C}}(0)} \mu_\beta(\hat{\sigma} \vee \check{\sigma}) = N e^{-\beta\Gamma^*} \frac{\check{Z}_\beta^{(n_\beta-K)}(0)}{Z_\beta^{(n_\beta)}} \\
 (4.34) \qquad &= N e^{-\beta\Gamma} \mu_\beta(\mathcal{S}) [1 + o(1)].
 \end{aligned}$$

Therefore, $\Sigma(\mathcal{I}_2)$ is also small compared to the main term of (4.23).

Lower bound. The proof of the lower bound follows the same line of argument as for Glauber dynamics, in that it relies on the construction of a suitable unit flow. This flow will, however, be considerably more difficult. In particular, we will no longer be able to get away with choosing a deterministic flow and the full power of the Berman–Konsowa variational principle has to be harnessed. The proof consists of five steps.

For future reference, we state the following property of the harmonic function for simple random walk on \mathbb{Z}^2 .

LEMMA 4.4. *Let g be the harmonic function of simple random walk on $B_{2L_\beta, 2L_\beta}(0)$ (which is equal to 1 on $\{0\}$ and 0 on $[B_{2L_\beta, 2L_\beta}(0)]^c$). There then exists a constant $C < \infty$ such that*

$$(4.35) \qquad \sum_e [g(z) - g(z + e)]_+ \leq C/L_\beta \qquad \forall z \in [B_{L_\beta, L_\beta}(0)]^c.$$

PROOF. See, for example, Lawler, Schramm and Werner [21], Lemma 5.1. The proof can be given via the estimates in Lawler [20], Section 1.7, or via a coupling argument. \square

1. *Starting configurations.* We start our flow on a subset of the configurations in \mathcal{C}^+ that is sufficiently large and sufficiently convenient. Let $\mathcal{C}_2^+ \subset \mathcal{C}^+$ denote the set of configurations having a proto-critical droplet with lower-left corner at some site $x \in \Lambda_\beta$, a free particle at distance 2 from this proto-critical droplet, no other particles in the box $B_{2L_\beta, 2L_\beta}(x)$ and satisfying the constraints in \mathcal{S}_L , that is, all other boxes of size $2L_\beta$ carry no more particles than there are in a proto-critical droplet. This is the same as \mathcal{C}^+ , except that the box around the proto-critical droplet has size $2L_\beta$ rather than L_β .

Let $K = \ell_c(\ell_c - 1) + 2$ be the volume of the critical droplet and let $\mathcal{S}_2^{(n_\beta - K)}$ be the analog of \mathcal{S} when the total number of particles is $n_\beta - K$ and the boxes in which we count particles have size $2L_\beta$ (cf. Definition 1.3). Similarly as in (3.17), our task is to derive a lower bound for $\text{CAP}(\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+) = \text{CAP}((\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+, \mathcal{S}_L) \geq \text{CAP}(\mathcal{C}_L, \mathcal{S}_L)$, where $\mathcal{C}_L \subset \mathcal{C}_2^+ \subset \mathcal{C}^+$ defined by

$$(4.36) \quad \mathcal{C}_L = \{ \sigma \cup P_{(y)}(x, z) : \sigma \in \mathcal{S}_2^{(n_\beta - K)}, x, y \in \Lambda_\beta, (x, y, z) \perp \sigma \}$$

is the analog of (3.13), namely, the set of configurations obtained from $\mathcal{S}_2^{(n_\beta - K)}$ by adding a critical droplet somewhere in Λ_β (lower-left corner at x , protuberance at y , free particle at z) such that it does not interact with the particles in σ and has an empty box of size $2L_\beta$ around it. Note that the $n_\beta - K$ particles can block at most $n_\beta(2L_\beta)^2 = o(|\Lambda_\beta|)$ sites from being the center of an empty box of size $2L_\beta$, and so the critical particle can be added at $|\Lambda_\beta| - o(|\Lambda_\beta|)$ locations.

We partition \mathcal{C}_L into sets $\mathcal{C}_L(x)$, $x \in \Lambda_\beta$, according to the location of the proto-critical droplet. It suffices to consider the case where the critical droplet is added at $x = 0$ because the union over x trivially produces a factor $|\Lambda_\beta|$.

2. *Overall strategy.* Starting from a configuration in $\mathcal{C}_L(0)$, we will successively pick $K - L$ particles from the critical droplet (starting with the free particle at z at distance 2) and move them out of the box $B_{L_\beta, L_\beta}(0)$, placing them essentially uniformly in the annulus $B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0)$. Once this has been achieved, the configuration is in \mathcal{S}_L . Each such move will produce an entropy of order L_β^2 , which will be enough to compensate for the loss of energy in tearing down the droplet (recall Figure 4). The order in which the particles are removed follows the canonical order employed in the lower bound for Glauber dynamics (recall Figure 6). As with Glauber dynamics, we will use Proposition 2.4 to estimate

$$(4.37) \quad \text{CAP}(\mathcal{C}_L, \mathcal{S}_L) \geq |\Lambda_\beta| \sum_{\sigma \in \mathcal{C}_L(0)} \sum_{\gamma : \gamma_0 = \sigma} \mathbb{P}^f(\gamma) \sum_{k=0}^{\tau(\gamma)} \left[\frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k) c_\beta(\gamma_k, \gamma_{k+1})} \right]^{-1}$$

for a suitably constructed flow f and associated path measure \mathbb{P}^f , starting from some initial distribution on $\mathcal{C}_L(0)$ (which, as with Glauber, will be irrelevant), and $\tau(\gamma)$ being the time at which the last of the $K - L$ particles exits the box $B_{L_\beta, L_\beta}(0)$.

The difference between Glauber and Kawasaki is that, while in Glauber the droplet can be torn down via single spin-flips, in Kawasaki, after we have detached a particle from the droplet, we need to move it out of the box $B_{L_\beta, L_\beta}(0)$, which takes a large number of steps. Thus, $\tau(\gamma)$ is the sum of $K - L$ stopping times, all but the first of which are themselves sums of two stopping times, one to detach the particle and one to move it out of the box $B_{L_\beta, L_\beta}(0)$. With each motion of a single particle, we need to gain an entropy factor of order close to $1/\rho_\beta$. This will be done by constructing a flow that involves only the motion of this single particle, based on the harmonic function of the simple random walk in the box $B_{2L_\beta, 2L_\beta}(0)$ up to the boundary of the box $B_{L_\beta, L_\beta}(0)$. Outside $B_{L_\beta, L_\beta}(0)$, the flow becomes more complex: we modify it in such a way that a small fraction of the flow, of order $L_\beta^{-1+\varepsilon}$ for some $\varepsilon > 0$ small enough, is going in the direction of removing the next particle from the droplet. The reason for this choice is that we want to make sure that the flow becomes sufficiently small, of order $L_\beta^{-2+\varepsilon}$, so that this can compensate for the fact that the Gibbs weight in the denominator of the lower bound in (2.20) is reduced by a factor $e^{-\beta U}$ when the protuberance is detached. The reason for the extra ε is that we want to make sure that, along most of the paths, the protuberance is detached before the first particle leaves the box $B_{2L_\beta, 2L_\beta}(0)$.

Once the protuberance detaches itself from the proto-critical droplet, the first particle stops and the second particle moves in the same way as the first particle did when it moved away from the proto-critical droplet, and so on. This is repeated until no more than L particles remain in $B_{L_\beta, L_\beta}(0)$, by which time we have reached S_L . As we will see, the only significant contribution to the lower bound comes from the motion of the first particle (as with Glauber dynamics), and this coincides with the upper bound established earlier. The details of the construction are, to some extent, arbitrary and there are many other choices imaginable.

3. *First particle.* We first construct the flow that moves the particle at distance 2 from the proto-critical droplet to the boundary of the box $B_{L_\beta, L_\beta}(0)$. This flow will consist of independent flows for each fixed shape and location of the critical droplet. This first part of the flow will be seen to produce the essential contribution to the lower bound.

We label the configurations in $\mathcal{C}_L(0)$ by σ , describing the shape of the critical droplet, as well as the configuration outside the box $B_{2L_\beta, 2L_\beta}(0)$, and we label the position of the free particle in σ by $z_1(\sigma)$.

Let g be the harmonic function for simple random walk with boundary conditions 0 on $[B_{2L_\beta, 2L_\beta}(0)]^c$ and 1 on the critical droplet. We then choose our flow to be

$$(4.38) \quad f(\sigma(z), \sigma(z')) = \begin{cases} C_1[g(z) - g(z + e)]_+, & \text{if } z' = z + e, \|e\| = 1, \\ 0, & \text{otherwise,} \end{cases}$$

where $\sigma(z)$ is the configuration obtained from σ by placing the first particle at site z . The constant C_1 is chosen to ensure that f defines a unit flow in the sense of Definition 2.3, that is,

$$(4.39) \quad \begin{aligned} \sum_{\sigma \in \mathcal{C}_L(0)} C_1 \sum_{z_1(\sigma), e} [g(z_1(\sigma)) - g(z_1(\sigma) + e)] \\ = C_1 \sum_{\sigma \in \mathcal{C}_L(0)} \text{CAP}(P_\sigma(0), [B_{2L_\beta, 2L_\beta}(0)]^c) = 1, \end{aligned}$$

where $P_\sigma(0)$ denotes the support of the proto-critical droplet in σ and the capacity refers to the simple random walk.

Now, let $z^1(k)$ be the location of the first particle at time k , and

$$(4.40) \quad \tau^1 = \inf\{k \in \mathbb{N} : z^1(k) \in [B_{L_\beta, L_\beta}(0)]^c\}$$

be the first time when, under the Markov chain associated with the flow f , it exits $B_{L_\beta, L_\beta}(0)$. Let γ be a path of this Markov chain. Then, by (4.38) and (4.39), we have

$$(4.41) \quad \sum_{k=0}^{\tau^1} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})} = \frac{C_1[g(z^1(0)) - g(z^1(\tau^1))]}{\mu_\beta(\gamma_0)},$$

where the sum over the g 's is telescoping because only paths along which the g -function decreases carry positive probability, and $c_\beta(\gamma_k, \gamma_{k+1}) = 1$ for all $0 \leq k \leq \tau^1$ because the first particle is free. We have $g(z^1(0)) = 1$, while, by Lemma 4.4, there exists a $C < \infty$ such that

$$(4.42) \quad g(x) \leq C/\log L_\beta, \quad x \in [B_{L_\beta, L_\beta}(0)]^c.$$

Therefore,

$$(4.43) \quad \sum_{k=0}^{\tau^1} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})} = \frac{C_1}{\mu_\beta(\gamma_0)}[1 + o(1)].$$

Next, by Lemma 4.3, we have

$$(4.44) \quad \text{CAP}(P_\sigma(0), [B_{2L_\beta, 2L_\beta}(0)]^c) = \frac{4\pi}{\beta\Delta}[1 + o(1)], \quad \sigma \in \mathcal{C}_L(0), \beta \rightarrow \infty$$

[because $\{0\} \subset P_\sigma(0) \subset B_{2\ell_c, 2\ell_c}(0)$ for all $\sigma \in \mathcal{C}_L(0)$]. Since $N = |\mathcal{C}_L(0)|$, it follows from (4.39) that

$$(4.45) \quad \frac{1}{C_1} = N \frac{4\pi}{\beta\Delta}[1 + o(1)],$$

and so (4.43) becomes

$$(4.46) \quad \left[\sum_{k=0}^{\tau^1} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})} \right]^{-1} = \mu_\beta(\gamma_0) N \frac{4\pi}{\beta\Delta}[1 + o(1)], \quad \beta \rightarrow \infty.$$

This is the contribution we want because when we sum (4.46) over $\gamma_0 = \sigma \in \mathcal{C}_L(0)$ [recall (4.37)], we get a factor

$$(4.47) \quad \mu_\beta(\mathcal{C}_L(0)) = e^{-\beta\Gamma} \mu_\beta(\mathcal{S})[1 + o(1)].$$

To see why (4.47) is true, recall from (4.36) that $\mathcal{C}_L(0)$ is obtained from $\mathcal{S}_2^{(n_\beta-K)}$ by adding a critical droplet with lower-left corner at the origin that does not interact with the $n_\beta - K$ particles elsewhere in Λ_β . Hence,

$$(4.48) \quad \mu_\beta(\mathcal{C}_L(0)) = e^{-\beta\Gamma^*} \frac{\tilde{Z}_\beta^{(n_\beta-K)}(0)}{Z_\beta^{(n_\beta)}},$$

where $\tilde{Z}_\beta^{(n_\beta-K)}(0)$ is the analog of $\check{Z}_\beta^{(n_\beta-K)}(0)$ [defined in (4.17)] obtained by requiring that the $n_\beta - K$ particles are in $[R_{\ell_c, \ell_c}(0)]^c$ instead of $[B_{L_\beta, L_\beta}(0)]^c$. However, it will follow from the proofs of Lemmas B.1 and B.2 in Appendix B that, just as in (4.22),

$$(4.49) \quad \frac{\tilde{Z}_\beta^{(n_\beta-K)}(0)}{Z_\beta^{(n_\beta)}} = (\rho_\beta)^K \mu_\beta(\mathcal{S})[1 + o(1)], \quad \beta \rightarrow \infty,$$

which yields (4.47) because $\Gamma = \Gamma^* + K\Delta$. For the remaining part of the construction of the flow, it therefore suffices to ensure that the sum beyond τ^1 gives a smaller contribution.

4. *Second particle.* Once the first particle (i.e., the free particle) has left the box $B_{L_\beta, L_\beta}(0)$, we need to allow the second particle (i.e., the protuberance) to detach itself from the proto-critical droplet and to move out of $B_{L_\beta, L_\beta}(0)$ as well. The problem is that detaching the second particle reduces the Gibbs weight appearing in the denominator by $e^{-U\beta}$, while the increments of the flow are reduced to only about $1/L_\beta$. Thus, we cannot immediately detach the second particle. Instead, we only do this with probability L_β .

The idea is that once the first particle is outside $B_{L_\beta, L_\beta}(0)$, we *leak* some of the flow that drives the motion of the first particle into a flow that detaches the second particle. To do this, we have to first construct a *leaky flow* in $B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0)$ for simple random walk. This is done as follows.

Let $p(z, z + e)$ denote the transition probabilities of simple random walk driven by the harmonic function g on $B_{2L_\beta, 2L_\beta}(0)$. Put

$$(4.50) \quad \tilde{p}(z, z + e) = \begin{cases} p(z, z + e), & \text{if } z \in B_{L_\beta, L_\beta}(0), \\ (1 - L_\beta^{-1+\varepsilon})p(z, z + e), & \text{if } z \in B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0). \end{cases}$$

Use the transition probabilities $\tilde{p}(z, z + e)$ to define a path measure \tilde{P} . This path measure describes simple random walk driven by g , but with a killing probability

$L_\beta^{-1+\varepsilon}$ inside the annulus $B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0)$. Put

$$(4.51) \quad k(z, z + e) = \sum_\gamma \tilde{P}(\gamma) \mathbb{1}_{(z, z+e) \in \gamma}, \quad z \in B_{2L_\beta, 2L_\beta}(0).$$

This edge function satisfies the following equations:

$$(4.52) \quad \begin{aligned} &\bullet k(z, z + e) = [g(z) - g(z + e)]_+ \\ &\quad \text{if } z \in B_{L_\beta, L_\beta}(0); \\ &\bullet k(z, z + e) = 0 \\ &\quad \text{if } z \in B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0) \text{ and } [g(z) - g(z + e)]_+ = 0; \\ &\bullet (1 - L_\beta^{-1+\varepsilon}) \sum_e k(z + e, z) \mathbb{1}_{g(z+e) - g(z) > 0} \\ &= \sum_e k(z, z + e) \mathbb{1}_{g(z) - g(z+e) > 0} \\ &\quad \text{if } z \in B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0). \end{aligned}$$

Note that inside the annulus $B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0)$ at each site, the flow out is less than the flow in by a leaking factor $1 - L_\beta^{-1+\varepsilon}$. We pick $\varepsilon > 0$ sufficiently small that

$$(4.53) \quad e^{\beta U} \text{ is exponentially smaller in } \beta \text{ than } L_\beta^{2-\varepsilon}$$

[which is possible by (1.27) and (1.29) and (1.30)]. The important fact for us is that this leaky flow is dominated by the harmonic flow associated with g , in particular, the flow in satisfies

$$(4.54) \quad \sum_e k(z + e, z) \leq \sum_e [g(z + e) - g(z)]_+ \quad \forall z \in B_{2L_\beta, 2L_\beta}(0)$$

(and the same applies for the flow out). This inequality holds because g satisfies the same equations as in (4.50) and (4.51), but without the leaking factor $1 - L_\beta^{-1+\varepsilon}$.

Using this leaky flow, we can now construct a flow involving the first two particles, as follows:

$$(4.55) \quad \begin{aligned} &\bullet f(\sigma(z_1, a), \sigma(z_1 + e, a)) = C_1 k(z_1, z_1 + e) \\ &\quad \text{if } z_1 \in B_{2L_\beta, 2L_\beta}(0); \\ &\bullet f(\sigma(z_1, a), \sigma(z_1, b)) = C_1 L_\beta^{-1+\varepsilon} \sum_e k(z_1, z_1 + e) \\ &\quad \text{if } z_1 \in B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0); \\ &\bullet f(\sigma(z_1, z_2), \sigma(z_1, z_2 + e)) = \left\{ C_1 L_\beta^{-1+\varepsilon} \sum_e k(z_1, z_1 + e) \right\} \\ &\quad \times [g(z_2) - g(z_2 + e)]_+ \\ &\quad \text{if } z_1 \in B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0), z_2 \in B_{L_\beta, L_\beta}(0) \setminus P_\sigma(0). \end{aligned}$$

Here, we write a and b for the respective locations of the second particle prior to and after it detaches itself from the proto-critical droplet, and $\sigma(z_1, z_2)$ for the configuration obtained from σ by placing the first particle (that was at distance 2 from the proto-critical droplet) at site z_1 and the second particle (that was the protuberance) at site z_2 . The flow for other motions is zero and the constant C_1 is the same as in (4.38)–(4.39).

We next define two further stopping times, namely,

$$(4.56) \quad \zeta^2 = \inf\{k \in \mathbb{N} : z^2(\gamma_k) = b\},$$

that is, the first time the second particle (the protuberance) detaches itself from the proto-critical droplet and

$$(4.57) \quad \tau^2 = \inf\{k \in \mathbb{N} : z^2(\gamma_k) \in [B_{L_\beta, L_\beta}(0)]^c\},$$

that is, the first time the second particle exits the box $B_{L_\beta, L_\beta}(0)$. Note that since we choose the leaking probability to be $L^{-1+\varepsilon}$, the probability that ζ^2 is larger than the first time the first particle exits $B_{2L_\beta, 2L_\beta}(0)$ is of order $\exp[-L^\varepsilon]$ and is hence negligible. We will disregard the contributions of such paths in the lower bound. Paths with this property will be called *good*.

We will next show that (4.41) also holds if we extend the sum along any path of positive probability up to ζ^2 . The reason for this lies in Lemma 4.4. Let γ be a path that has a positive probability under the path measure \mathbb{P}^f associated with f stopped at τ^2 . We will assume that this path is good in the sense described above. To that end, we decompose

$$(4.58) \quad \begin{aligned} & \sum_{k=0}^{\tau^2} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})} \\ &= \sum_{k=0}^{\tau^1} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})} + \sum_{k=\tau^1+1}^{\zeta^2-2} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})} \\ & \quad + \sum_{k=\zeta^2-1}^{\tau^2} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})} \\ &= I + II + III. \end{aligned}$$

The term I was already estimated in (4.41)–(4.47). To estimate II , we use (4.42), (4.54) and (4.55) to bound [cf. (4.41)]

$$(4.59) \quad II \leq C_1 \frac{g(z^1(\zeta^2)) - g(z^1(\tau^1))}{\mu_\beta(\gamma_0)} \leq C_1 \frac{[C/\log L_\beta]}{\mu_\beta(\gamma_0)},$$

which is negligible compared to I , due to the factor $C/\log L_\beta$. It remains to estimate III . Note that

$$(4.60) \quad III = \frac{f(\gamma_{\zeta^2-1}, \gamma_{\zeta^2})}{\mu_\beta(\gamma_{\zeta^2-1})c_\beta(\gamma_{\zeta^2-1}, \gamma_{\zeta^2})} + \sum_{k=\zeta^2}^{\tau^2} \frac{f(\gamma_k, \gamma_{k+1})}{\mu_\beta(\gamma_k)c_\beta(\gamma_k, \gamma_{k+1})}.$$

The first term corresponds to the move when the protuberance detaches itself from the proto-critical droplet. Its numerator is given by $f(\sigma(z_1, a), \sigma(z_1, b))$ (for some $z_1 \in [B_{L_\beta, L_\beta}(0)]^c$), which, by Lemma 4.4, (4.54) and (4.55), is smaller than $C_1 L_\beta^{-1+\varepsilon} C L_\beta^{-1} = C_1 C L_\beta^{-2+\varepsilon}$. On the other hand, its denominator is given by

$$(4.61) \quad \mu(\gamma_{\zeta^2-1})c_\beta(\gamma_{\zeta^2-1}, \gamma_{\zeta^2}) = \mu_\beta(\gamma_0)e^{-U\beta}.$$

The same holds for the denominators in all the other terms in III , while the numerators in these terms satisfy the bound

$$(4.62) \quad f(\gamma_k, \gamma_{k+1}) \leq C_1 C L_\beta^{-2+\varepsilon} [g(z^2(\gamma_k)) - g(z^2(\gamma_{k+1}))].$$

Adding up the various terms, we get that

$$(4.63) \quad \begin{aligned} III &\leq \frac{C_1}{\mu_\beta(\gamma_0)} L_\beta^{-2+\varepsilon} e^{\beta U} (1 + [g(z^2(\zeta^2)) - g(z^2(\tau^2))]) \\ &\leq \frac{2C_1}{\mu_\beta(\gamma_0)} L_\beta^{-2+\varepsilon} e^{\beta U}. \end{aligned}$$

The right-hand side is smaller than I by a factor $L_\beta^{-2+\varepsilon} e^{\beta U}$, which, by (4.53), is exponentially small in β .

5. Remaining particles. The lesson from the previous steps is that we can construct a flow with the property that each time we remove a particle from the droplet, we gain a factor $L_\beta^{-2+\varepsilon}$, that is, almost $e^{-\Delta\beta}$. (This entropy gain corresponds to the gain from the magnetic field in Glauber dynamics or from the activity in Kawasaki dynamics on a finite open box.) We can continue our flow by tearing down the critical droplet in the same order as we did for Glauber dynamics. Each removal corresponds to a flow that is built in the same way as described in Step 4 for the second particle. There will be some minor modifications involving a negligible fraction of paths where a particle hits a particle that was moved out earlier, but this is of no consequence. As a result of the construction, the sums along the remainders of these paths will give only negligible contributions.

Thus, we have shown that the lower bound coincides, up to a factor $1 + o(1)$, with the upper bound, and so the lemma is proved. \square

4.2. *Proof of Theorem 1.4(b).* The same observation holds as in (3.34).

PROOF OF THEOREM 1.4(b). The proof follows along the same lines as that of Theorem 1.4(a). The main point is to prove that $\text{CAP}(\mathcal{D}_M, \mathcal{S}_L) = [1 +$

$o(1)]\text{CAP}(\mathcal{C}^+, \mathcal{S}_L)$. Since $\text{CAP}(\mathcal{S}_L, \mathcal{D}_M) \leq \text{CAP}(\mathcal{S}_L, \mathcal{C}^+)$, we need only prove a lower bound on $\text{CAP}(\mathcal{D}_M, \mathcal{S}_L)$. This is done in almost exactly the same way as for Glauber dynamics, by using the construction given there and substituting each Glauber move by a flow involving the motion of just two particles.

Note that as long as $M = e^{o(\beta)}$, an $M \times M$ droplet can be added at $|\Lambda_\beta| - o(|\Lambda_\beta|)$ locations to a configuration $\sigma \in \mathcal{S}$ [cf. (4.36)]. The only novelty is that we have to eventually remove the cloud of particles that is produced in the annulus $B_{2L_\beta, 2L_\beta}(0) \setminus B_{L_\beta, L_\beta}(0)$. This is done in much the same way as before. As long as only $e^{o(\beta)}$ particles have to be removed, potential collisions between particles can be ignored, as they are sufficiently unlikely. \square

APPENDIX A: SPARSENESS OF SUBCRITICAL DROPLETS

Recall Definition 1.1(a), (3.11) and (3.12). In this section, we prove (3.22).

LEMMA A.1. $\lim_{\beta \rightarrow \infty} \frac{1}{\beta} \log \frac{\mu_\beta(\mathcal{S} \setminus \mathcal{W})}{\mu_\beta(\mathcal{S})} = -\infty$.

PROOF. We will prove that $\lim_{\beta \rightarrow \infty} \frac{1}{\beta} \log \mu_\beta(\mathcal{S} \setminus \mathcal{W}) / \mu_\beta(\Xi) = -\infty$. Since $\Xi \in \mathcal{S}$, this will prove the claim.

Let $w(\beta)$ be the function satisfying (3.11). We begin by noting that

$$(A.1) \quad \mu_\beta(\mathcal{S} \setminus \mathcal{W}) \leq \mu_\beta(\mathcal{I}) \quad \text{with } \mathcal{I} = \{\sigma \in \mathcal{S} : |\text{supp}[C_B(\sigma)]| > |\Lambda_\beta|/w(\beta)\}$$

because the bootstrap percolation map increases the number of (+1)-spins. Let $\mathcal{D}(k)$ denote the set of configurations whose support consists on k noninteracting subcritical rectangles. Put $C_1 = (\ell_c + 2)(\ell_c + 1)$. Since the union of a subcritical rectangle and its exterior boundary has at most C_1 sites, it follows that in \mathcal{I} , there are at least $|\Lambda_\beta|/C_1 w(\beta)$ noninteracting rectangles. Thus, we have

$$(A.2) \quad \mu_\beta(\mathcal{I}) \leq \sum_{k=|\Lambda_\beta|/(C_1 w(\beta))}^{K_{\max}} F(k) \quad \text{with } F(k) = \frac{1}{Z_\beta} \sum_{\substack{\sigma \in \mathcal{X}_\beta : \\ C(\sigma) \in \mathcal{D}(k)}} e^{-\beta H_\beta(\sigma)},$$

where $K_{\max} \leq |\Lambda_\beta|$.

Next, note that

$$(A.3) \quad F(k) \leq (2^{C_1})^k \frac{1}{Z_\beta} \sum_{\sigma \in \mathcal{D}(k)} e^{-\beta H_\beta(\sigma)}.$$

Since the bootstrap percolation map is downhill, the energy of a subcritical rectangle is bounded below by $C_2 = 2J - h$ (recall Figure 9) and the number of ways to

place k rectangles in Λ_β is at most $\binom{|\Lambda_\beta|}{k}$, it follows that

$$\begin{aligned}
 (A.4) \quad F(k) &\leq 2^{C_1 k} \binom{|\Lambda_\beta|}{k} \mu_\beta(\boxminus) e^{-C_2 \beta k} \leq 2^{C_1 k} (C_1 e w(\beta))^k \mu_\beta(\boxminus) e^{-C_2 \beta k} \\
 &\leq \mu_\beta(\boxminus) \exp\left[-\frac{1}{2} C_2 \beta k\right],
 \end{aligned}$$

where the second inequality uses the fact that $k! \geq k^k e^{-k}$, $k \in \mathbb{N}$, and the third inequality uses the fact that $w(\beta) = e^{o(\beta)}$. We thus have

$$(A.5) \quad \sum_{k=|\Lambda_\beta|/(C_1 w(\beta))}^{K_{\max}} F(k) \leq 2 \mu_\beta(\boxminus) w(\beta) \frac{|\Lambda_\beta|}{w(\beta)} \exp\left[-\frac{1}{2} \frac{C_2}{C_1} \beta \frac{|\Lambda_\beta|}{w(\beta)}\right],$$

which is the desired estimate because $|\Lambda_\beta|/w(\beta)$ tends to infinity. \square

APPENDIX B: EQUIVALENCE OF ENSEMBLES AND TYPICALITY OF HOLES

For $m \in \mathbb{N}$, let

$$(B.1) \quad \mathcal{S}^{(m)} = \{\sigma \in \mathcal{X}_\beta^{(m)} : |\text{supp}[\sigma] \cap B_{L_\beta, L_\beta}(x)| \leq \ell_c(\ell_c - 1) + 1 \ \forall x \in \Lambda_\beta\}$$

and

$$\begin{aligned}
 (B.2) \quad \check{\mathcal{C}}^{(m)}(0) &= \{\sigma \mathbb{1}_{\in B_{L_\beta, L_\beta}(0)} : \sigma \in \mathcal{S}^{(m)}\}, \\
 \check{Z}_\beta^{(m)}(0) &= \sum_{\sigma \in \check{\mathcal{C}}^{(m)}(0)} e^{-\beta H(\sigma)}.
 \end{aligned}$$

The latter is the partition sum restricted to $B_{L_\beta, L_\beta}(0)$ when it carries m particles. In Appendix B.1, we prove a lemma about ratios of partition sums that was used in (4.22), (4.26), (4.32) and (4.49). In Appendix B.2, we prove that $\lim_{\beta \rightarrow \infty} \mu_\beta(\check{\mathcal{S}}(0))/\mu_\beta(\mathcal{S}) = 1$, which is needed in the proof of this lemma.

B.1. Equivalence of ensembles. Recall (1.22), (4.6) and (4.17).

LEMMA B.1. $\check{Z}_\beta^{(n_\beta - s)}(0)/Z_\beta^{(n_\beta)} = (\rho_\beta)^s \mu_\beta(\mathcal{S})[1 + o(1)]$ as $\beta \rightarrow \infty$ for all $s \in \mathbb{N}$.

PROOF. The proof proceeds via upper and lower bounds.

Upper bound. Let

$$(B.3) \quad \check{\mathcal{S}}(0) = \{\sigma \in \mathcal{S} : \text{supp}[\sigma] \cap B_{L_\beta, L_\beta}(0) = \emptyset\}.$$

Write

$$\begin{aligned}
 \mu_\beta(\check{\mathcal{S}}(0)) &= \frac{1}{Z_\beta^{(n_\beta)}} \\
 (B.4) \quad &\times \sum_{\check{\sigma} \in \check{\mathcal{C}}(0)} \sum_{\substack{\zeta \subset [B_{L_\beta, L_\beta}(0)]^c \setminus \text{supp}[\check{\sigma}] \\ |\zeta|=s}} \binom{n_\beta}{s}^{-1} \mathbb{1}_{\{\check{\sigma} \vee \zeta \in \check{\mathcal{S}}(0)\}} e^{-\beta H_\beta(\check{\sigma} \vee \zeta)}.
 \end{aligned}$$

This relation simply says that n_β particles can be placed outside $B_{L_\beta, L_\beta}(0)$ by first placing $n_\beta - s$ particles and then placing another s particles. Because the interaction is attractive, we have

$$(B.5) \quad H_\beta(\check{\sigma} \vee \zeta) \leq H_\beta(\check{\sigma}) + H_\beta(\zeta) \quad \text{and} \quad H_\beta(\zeta) \leq 0 \quad \forall \check{\sigma}, \zeta.$$

Consequently,

$$\begin{aligned}
 \mu_\beta(\check{\mathcal{S}}(0)) &\geq \binom{n_\beta}{s}^{-1} \frac{1}{Z_\beta^{(n_\beta)}} \\
 (B.6) \quad &\times \sum_{\check{\sigma} \in \check{\mathcal{C}}(0)} e^{-\beta H_\beta(\check{\sigma})} \sum_{\substack{\zeta \subset [B_{L_\beta, L_\beta}(0)]^c \setminus \text{supp}[\check{\sigma}] \\ |\zeta|=s}} \mathbb{1}_{\{\check{\sigma} \vee \zeta \in \check{\mathcal{S}}(0)\}}.
 \end{aligned}$$

We next estimate the second sum, uniformly in $\check{\sigma}$. When we add the s particles, we must make sure not to violate the requirement that all boxes $B_{L_\beta, L_\beta}(x)$, $x \in \Lambda_\beta$, carry not more than K particles [note that $\check{\mathcal{S}}(0) \subset \mathcal{S}$ and recall Definition 1.3(a)]. Partition $\Lambda_\beta \setminus B_{L_\beta, L_\beta}(0)$ into boxes of size L_β . The total number of boxes containing K particles is at most n_β/K . Hence, the total number of sites where we cannot place a particle is at most $(n_\beta/K)(3L_\beta)^2$. Therefore,

$$\begin{aligned}
 &\sum_{\substack{\zeta \subset [B_{L_\beta, L_\beta}(0)]^c \setminus \{\check{\sigma}\} \\ |\zeta|=s}} \mathbb{1}_{\{\check{\sigma} \vee \zeta \in \check{\mathcal{S}}(0)\}} \\
 (B.7) \quad &\geq \binom{|\Lambda_\beta \setminus B_{L_\beta, L_\beta}(0)| - n_\beta - (n_\beta/K)(3L_\beta)^2}{s} \quad \forall \check{\sigma}.
 \end{aligned}$$

But $n_\beta L_\beta^2 = o(n_\beta/\rho_\beta) = o(|\Lambda_\beta|)$ and $L_\beta^2 = o(1/\rho_\beta) = o(|\Lambda_\beta|)$ by (1.22), (1.29) and (1.30), and so the right-hand side of (B.7) equals $[1 + o(1)]|\Lambda_\beta|^s/s!$ as $\beta \rightarrow \infty$. Since the binomial in (B.6) equals $[1 + o(1)](n_\beta)^s/s!$ with $n_\beta = \lceil \rho_\beta |\Lambda_\beta| \rceil$, we end up with [recall (4.17)]

$$(B.8) \quad \mu_\beta(\check{\mathcal{S}}(0)) \geq \frac{\check{Z}_\beta^{(n_\beta-s)}(0)}{Z_\beta^{(n_\beta)}} (\rho_\beta)^{-s} [1 + o(1)]$$

or

$$(B.9) \quad \frac{\check{Z}_\beta^{(n_\beta-s)}(0)}{Z_\beta^{(n_\beta)}} \leq (\rho_\beta)^s \mu_\beta(\check{S}(0)) [1 + o(1)].$$

Since $\check{S}(0) \subset \mathcal{S}$, this gives the desired upper bound.

Lower bound. We return to (B.4). Among the s particles that are added to $[B_{L_\beta, L_\beta}(0)]^c$, let s_1 count the number that interact with the $n_\beta - s$ particles already present and s_2 the number that interact among themselves, where $s_1 + s_2 \leq s$. We can then estimate

$$(B.10) \quad \begin{aligned} \mu_\beta(\check{S}(0)) &\leq \frac{1}{Z_\beta^{(n_\beta)}} \sum_{\check{\sigma} \in \check{C}(0)} \binom{n_\beta}{s}^{-1} e^{-\beta H_\beta(\check{\sigma})} \sum_{\substack{s_1, s_2 \\ 0 \leq s_1 + s_2 \leq s}} \left(\frac{s!}{s_1! s_2!} \right)^{-1} \\ &\quad \times \sum_{\substack{\zeta \subset [B_{L_\beta, L_\beta}(0)]^c \setminus \text{supp}[\check{\sigma}] \\ |\zeta| = s}} e^{-\beta H(\zeta)} \mathbb{1}_{\{|\zeta \cap \partial \check{\sigma}| = s_1\}} \\ &\quad \times \mathbb{1}_{\{s_2 \text{ interacting particles in } \zeta\}} \mathbb{1}_{\{\check{\sigma} \vee \zeta \in \check{S}(0)\}} \\ &\leq [1 + o(1)] \frac{\check{Z}_\beta^{(n_\beta-s)}(0)}{Z_\beta^{(n_\beta)}} (\rho_\beta)^{-s} \\ &\quad + \frac{1}{Z_\beta^{(n_\beta)}} \sum_{\check{\sigma} \in \check{C}(0)} \binom{n_\beta}{s}^{-1} e^{-\beta H_\beta(\check{\sigma})} \\ &\quad \times \sum_{\substack{s_1, s_2 \\ 1 \leq s_1 + s_2 \leq s}} \sum_{\substack{\zeta \subset [B_{L_\beta, L_\beta}(0)]^c \setminus \text{supp}[\check{\sigma}] \\ |\zeta| = s}} e^{-\beta H(\zeta)} \mathbb{1}_{\{|\zeta \cap \partial \check{\sigma}| = s_1\}} \\ &\quad \times \mathbb{1}_{\{s_2 \text{ interacting particles in } \zeta\}} \\ &\quad \times \mathbb{1}_{\{\check{\sigma} \vee \zeta \in \check{S}(0)\}}, \end{aligned}$$

where, in the second inequality, we estimate the term with $s_1 = s_2 = 0$ by using the result for the upper bound. We will show that the other terms are exponentially small.

For fixed $\check{\sigma}$, we may estimate the last sum in (B.10) as

$$(B.11) \quad \begin{aligned} &\sum_{\substack{\zeta \subset [B_{L_\beta, L_\beta}(0)]^c \setminus \text{supp}[\check{\sigma}] \\ |\zeta| = s}} e^{-\beta H(\zeta)} \mathbb{1}_{\{|\zeta \cap \partial \check{\sigma}| = s_1\}} \\ &\quad \times \mathbb{1}_{\{s_2 \text{ interacting particles in } \zeta\}} \mathbb{1}_{\{\check{\sigma} \vee \zeta \in \check{S}(0)\}} \\ &\leq |\Lambda_\beta|^{s-s_1-s_2} (4n_\beta)^{s_1} \sum_{\sigma \in \mathcal{S}^{(s_2)}} e^{-\beta H(\sigma)} \mathbb{1}_{\{s_2 \text{ interacting particles in } \sigma\}} \cdot \end{aligned}$$

Indeed, $|\Lambda_\beta|^{s-s_1-s_2}$ bounds the number of ways to place $s - s_1 - s_2$ noninteracting particles, and $(4n_\beta)^{s_1}$ the number of ways to place s_1 particles that are interacting with the $n_\beta - s$ particles already present. Next, we write

$$\begin{aligned}
 & \sum_{\sigma \in \mathcal{S}(s_2)} e^{-\beta H(\sigma)} \mathbb{1}_{\{s_2 \text{ interacting particles in } \sigma\}} \\
 \text{(B.12)} \quad &= \sum_{m=1}^{s_2} \sum_{j=1}^m \sum_{2 \leq k_1, \dots, k_j \leq K} \sum_{\substack{C = \cup_{i=1}^j C_i \\ \sum_{i=1}^j k_i = m \quad |C_i| = k_i \forall i}} e^{-\beta \sum_{i=1}^j H(C_i)},
 \end{aligned}$$

which is a cluster expansion of the partition function [with noninteracting clusters C_i , all of which have size $\leq K = \ell_c(\ell_c + 1) + 1$]. By a standard isoperimetric inequality, we have $H(C_i) \geq H_{k_i}$, with the latter denoting the energy of a droplet of $k_i = |C_i|$ particles that is closest to a square or quasi-square. Hence,

$$\begin{aligned}
 & |\Lambda_\beta|^{-s_2} \sum_{\sigma \in \mathcal{S}(s_2)} e^{-\beta H(\sigma)} \mathbb{1}_{\{s_2 \text{ interacting particles in } \sigma\}} \\
 & \leq |\Lambda_\beta|^{-s_2} \sum_{m=1}^{s_2} \sum_{j=1}^m \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = s_2}} e^{-\beta \sum_{i=1}^j H_{k_i}} \left(\sum_{\substack{C = \cup_{i=1}^j C_i \\ |C_i| = k_i \forall i}} 1 \right) \\
 \text{(B.13)} \quad & \leq C |\Lambda_\beta|^{-s_2} \sum_{m=1}^{s_2} \sum_{j=1}^m \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = s_2}} e^{-\beta \sum_{i=1}^j H_{k_i}} |\Lambda_\beta|^j \\
 & \leq C \sum_{m=1}^{s_2} \sum_{j=1}^m \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = s_2}} e^{-\beta \sum_{i=1}^j [H_{k_i} + (k_i - 1)\beta^{-1} \log |\Lambda_\beta|]} \\
 & \leq \sum_{m=1}^{s_2} \sum_{j=1}^m \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = s_2}} e^{-\beta \sum_{i=1}^j [H_{k_i} + (k_i - 1)\Delta]},
 \end{aligned}$$

where, in the last inequality, we insert the bound $\beta^{-1} \log |\Lambda_\beta| \geq \Delta$, which is immediate from (1.22) and (1.35).

Now, $H_{k_i} + k_i \Delta$ is the *grand-canonical* energy of a square or quasi-square with k_i particles. It was shown in the proof of Proposition 2.4.2 in Bovier, den Hollander and Nardi [7] that this energy is greater than or equal to $U\sqrt{k_i}$ for $1 \leq k_i \leq 4K$, that is, for a droplet twice the size of the proto-critical droplet. Since $2U > \Delta$, we therefore have that $H_{k_i} + (k_i - 1)\Delta > 0$ when $k_i \geq 4$. Since $\Delta > U$, $H_2 = -U$

and $H_3 = -2U$, we also have that the terms with $k_i = 2, 3$ are greater than 0. Consequently, there exist $\varepsilon > 0$ and a constant C that is independent of β such that

$$(B.14) \quad |\Lambda_\beta|^{-s_2} \sum_{\sigma \in \mathcal{S}^{(s_2)}} e^{-\beta H(\sigma)} \mathbb{1}_{\{s_2 \text{ interacting particles in } \sigma\}} e^{-\beta H(\sigma)} \leq C e^{-\beta \varepsilon}.$$

Combining (B.10), (B.11) and (B.14), we see that the correction term in (B.10) is

$$(B.15) \quad \begin{aligned} & \mu_\beta(\check{\mathcal{S}}(0)) - [1 + o(1)] \frac{\check{Z}_\beta^{(n_\beta - s)}(0)}{Z_\beta^{(n_\beta)}} (\rho_\beta)^{-s} \\ & \leq C [1 + o(1)] \frac{\check{Z}_\beta^{(n_\beta - s)}(0)}{Z_\beta^{(n_\beta)}} (\rho_\beta)^{-s} \sum_{\substack{s_1, s_2 \\ 1 \leq s_1 + s_2 \leq s}} (e^{U\beta} \rho_\beta)^{s_1} e^{-\beta \varepsilon}. \end{aligned}$$

Since $\Delta > U$, we have $e^{U\beta} \rho_\beta \leq 1$ and so the sum is $o(1)$. Hence,

$$(B.16) \quad \frac{\check{Z}_\beta^{(n_\beta - s)}(0)}{Z_\beta^{(n_\beta)}} \geq (\rho_\beta)^s \mu_\beta(\check{\mathcal{S}}(0)) [1 + o(1)].$$

The claim now follows by using Lemma B.2 below. \square

B.2. Typicality of holes.

LEMMA B.2. $\lim_{\beta \rightarrow \infty} \mu_\beta(\check{\mathcal{S}}(0)) / \mu_\beta(\mathcal{S}) = 1.$

PROOF. Since $\check{\mathcal{S}}(0) \subset \mathcal{S}$, we have $\mu_\beta(\check{\mathcal{S}}(0)) \leq \mu_\beta(\mathcal{S})$. It therefore remains to prove the lower bound. Write

$$(B.17) \quad \begin{aligned} \mu_\beta(\mathcal{S}) &= \mu_\beta(\check{\mathcal{S}}(0)) \\ &+ \sum_{m=1}^K \sum_{\eta \in \mathcal{X}_\beta^{(m)}} \sum_{\substack{\zeta \in \mathcal{X}_\beta^{(n_\beta - m)} \\ \eta \vee \zeta \in \mathcal{S}}} \frac{e^{-\beta H(\eta \vee \zeta)}}{Z_\beta^{(n_\beta)}} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \\ &\quad \times \mathbb{1}_{\{\text{supp}[\zeta] \subset [B_{L_\beta, L_\beta}(0)]^c\}} \\ &\leq \mu_\beta(\check{\mathcal{S}}(0)) + \gamma_1(\beta) + \gamma_2(\beta), \end{aligned}$$

where

$$(B.18) \quad \begin{aligned} \gamma_1(\beta) &= \sum_{m=1}^K \sum_{\eta \in \mathcal{X}_\beta^{(m)}} \sum_{\substack{\zeta \in \mathcal{X}_\beta^{(n_\beta - m)} \\ \eta \vee \zeta \in \mathcal{S}}} \frac{e^{-\beta [H(\eta) + H(\zeta)]}}{Z_\beta^{(n_\beta)}} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \\ &\quad \times \mathbb{1}_{\{\text{supp}[\zeta] \subset [B_{L_\beta, L_\beta}(0)]^c\}} \end{aligned}$$

and $\gamma_2(\beta)$ is a term that arises from particles interacting across the boundary of $B_{L_\beta, L_\beta}(0)$. We will show that both $\gamma_1(\beta)$ and $\gamma_2(\beta)$ are negligible.

Estimate, with the help of (B.9) [and recalling (B.1) and (B.2)],

$$\begin{aligned}
 \gamma_1(\beta) &\leq \sum_{m=1}^K \frac{\check{Z}_\beta^{(n_\beta-m)}}{Z_\beta^{(n_\beta)}} \sum_{\eta \in \mathcal{S}^{(m)}} e^{-\beta H(\eta)} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \\
 &= [1 + o(1)] \mu_\beta(\check{\mathcal{S}}(0)) \sum_{m=1}^K (\rho_\beta)^m \sum_{\eta \in \mathcal{S}^{(m)}} e^{-\beta H(\eta)} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \\
 \text{(B.19)} \quad &= [1 + o(1)] \mu_\beta(\check{\mathcal{S}}(0)) \\
 &\quad \times \sum_{m=1}^K (\rho_\beta)^m \sum_{j=1}^m \sum_{2 \leq k_1, \dots, k_j \leq K} \sum_{\substack{C = \bigcup_{i=1}^j C_i \\ \sum_{i=1}^j k_i = m \quad |C_i| = k_i \forall i}} e^{-\beta \sum_{i=1}^j H(C_i)},
 \end{aligned}$$

where the last equality is a cluster expansion, as in (B.12). Again using the isoperimetric inequality, we get [recalling (1.29)]

$$\begin{aligned}
 \gamma_1(\beta) &\leq [1 + o(1)] \mu_\beta(\check{\mathcal{S}}(0)) \\
 &\quad \times \sum_{m=1}^K (\rho_\beta)^m \sum_{j=1}^m \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = m}} e^{-\beta \sum_{i=1}^j H(k_i)} \left(\sum_{\substack{C = \bigcup_{i=1}^j C_i \\ |C_i| = k_i \forall i}} 1 \right) \\
 \text{(B.20)} \quad &\leq C \mu_\beta(\check{\mathcal{S}}(0)) \sum_{m=1}^K (\rho_\beta)^m \sum_{j=1}^m (L_\beta^2)^j \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = m}} e^{-\beta \sum_{i=1}^j H k_i} \\
 &= C \mu_\beta(\check{\mathcal{S}}(0)) \sum_{m=1}^K \sum_{j=1}^m \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = m}} e^{-\beta \sum_{i=1}^j [H k_i + k_i \Delta - (\Delta - \delta_\beta)]} \\
 &\leq C' \mu_\beta(\check{\mathcal{S}}(0)) e^{-\beta \varepsilon}
 \end{aligned}$$

for some $\varepsilon > 0$ and constants $C, C' < \infty$ that are independent of β .

Estimate, with the help of (B.9),

$$\begin{aligned}
 \gamma_2(\beta) &\leq \sum_{m=1}^K \sum_{\eta \in \mathcal{S}^{(m)}} e^{-\beta H(\eta)} \sum_{k=1}^m e^{\beta k U} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \frac{\check{Z}_\beta^{(n_\beta-(m+k))}}{Z_\beta^{(n_\beta)}} \\
 \text{(B.21)} \quad &\leq \sum_{m=1}^K \sum_{\eta \in \mathcal{S}^{(m)}} e^{-\beta H(\eta)} \sum_{k=1}^m e^{\beta k U} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}}
 \end{aligned}$$

$$\begin{aligned} & \times (\rho_\beta)^{m+k} \mu_\beta(\check{\mathcal{S}}(0)) [1 + o(1)] \\ \leq & [1 + o(1)] \mu_\beta(\check{\mathcal{S}}(0)) \\ & \times \sum_{m=1}^K (\rho_\beta)^m \sum_{\eta \in \mathcal{S}^{(m)}} e^{-\beta H(\eta)} \sum_{k=1}^m e^{-\beta k(\Delta-U)} \\ & \times \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \end{aligned}$$

and we can proceed as in (B.19) and (B.20) to show that this term is negligible. \square

B.3. Atypicality of critical droplets. The following lemma was used in Section 4.1.1.

LEMMA B.3. $\lim_{\beta \rightarrow \infty} \mu_\beta(\tilde{\mathcal{C}} \setminus \mathcal{C}^+) / \mu_\beta(\mathcal{S}) = 0$.

PROOF. Similarly as in (B.17), we first write

$$\begin{aligned} \mu_\beta(\tilde{\mathcal{C}} \setminus \mathcal{C}^+) & \leq \mu_\beta(\tilde{\mathcal{C}}) \\ & = |\Lambda_\beta| \gamma(\beta) + |\Lambda_\beta| \sum_{\eta \in \mathcal{X}_\beta^{(K)}} \sum_{\substack{\zeta \in \mathcal{X}_\beta^{(n_\beta - K)} \\ \eta \vee \zeta \in \tilde{\mathcal{C}}}} \frac{e^{-\beta[H(\eta) + H(\zeta)]}}{Z_\beta^{(n_\beta)}} \\ \text{(B.22)} \quad & \times \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \\ & \times \mathbb{1}_{\{\text{supp}[\zeta] \subset [B_{L_\beta, L_\beta}(0)]^c\}}, \end{aligned}$$

with $\gamma(\beta)$ a negligible error term that arises from particles interacting across the boundary of $B_{L_\beta, L_\beta}(0)$. We then proceed as in (B.18)–(B.20), obtaining ($\Gamma = \Gamma^* + K \Delta$)

$$\begin{aligned} & \text{right-hand side of (B.22)} \\ \text{(B.23)} \quad & \leq N |\Lambda_\beta| e^{-\beta \Gamma^*} (\rho_\beta)^K \mu_\beta(\check{\mathcal{S}}(0)) [1 + o(1)] \\ & = N |\Lambda_\beta| e^{-\beta \Gamma} \mu_\beta(\mathcal{S}) [1 + o(1)], \quad \beta \rightarrow \infty, \end{aligned}$$

which is $o(\mu_\beta(\mathcal{S}))$, by (1.35). \square

APPENDIX C: TYPICALITY OF STARTING CONFIGURATIONS

In Sections C.1 and C.2, we prove the claims made in the remarks below (1.9) and (1.32), respectively.

C.1. Glauber.

PROOF. Split

$$(C.1) \quad \mathcal{S} = \mathcal{S}_L \cup (\mathcal{S} \setminus \mathcal{S}_L) = \mathcal{S}_L \cup \mathcal{U}_{>L},$$

where $\mathcal{U}_{>L} \subset \mathcal{S}$ are those configurations σ for which $C_B(\sigma)$ has at least one rectangle that is larger than $Q_L(0)$. We have

$$(C.2) \quad C_B(\sigma) = \bigcup_{x \in X(\sigma)} R_{\ell_1(x), \ell_2(x)}(x),$$

where $X(\sigma)$ is the set of lower-left corners of the rectangles in $C_B(\sigma)$, which, in turn, can be split as

$$(C.3) \quad X(\sigma) = X^{>L}(\sigma) \cup X^{\leq L}(\sigma),$$

where $X^{>L}(\sigma)$ labels the rectangles that are larger than $Q_L(0)$ and $X^{\leq L}(\sigma)$ labels the rest.

Let $\sigma|_A$ denote the restriction of σ to the set $A \subset \mathbb{Z}^2$. Then, for any $x \in X(\sigma)$, we have

$$(C.4) \quad H(\sigma) = H(\sigma|_{R_{\ell_1(x), \ell_2(x)}(x)}) + H(\sigma|_{R_{\ell_2(x), \ell_2(x)}^c(x)})$$

because the rectangles in $C_B(\sigma)$ are noninteracting. Since, for $\sigma \in \mathcal{U}_{>L}$, there is at least one rectangle with lower-left corner in $X^{>L}(\sigma)$, we have

$$(C.5) \quad \begin{aligned} \mu_\beta(\mathcal{U}_{>L}) &\leq \sum_{x \in \Lambda_\beta} \sum_{\sigma \in \mathcal{S}} \mathbb{1}_{\{x \in X^{>L}(\sigma)\}} \mu_\beta(\sigma) \\ &= \sum_{x \in \Lambda_\beta} \sum_{\sigma \in \mathcal{S}} \mathbb{1}_{\{x \in X^{>L}(\sigma)\}} \frac{1}{Z_\beta} \\ &\quad \times \exp\{-\beta[H(\sigma|_{R_{\ell_1(x), \ell_2(x)}(x)}) + H(\sigma|_{R_{\ell_1(x), \ell_2(x)}^c(x)})]\} \\ &\leq e^{-\beta\Gamma_{L+1}} \sum_{x \in \Lambda_\beta} \sum_{\sigma \in \mathcal{S}} \mathbb{1}_{\{x \in X^{>L}(\sigma)\}} \frac{1}{Z_\beta} e^{-\beta H(\sigma|_{R_{\ell_1(x), \ell_2(x)}^c(x)})}, \end{aligned}$$

where Γ_{L+1} is the energy of $Q_{L+1}(0)$. In the last step, we use the fact that the bootstrap map is downhill and that the energy of $Q_L(0)$ is increasing with L . Since the energy of a subcritical rectangle is nonnegative, we get

$$(C.6) \quad \mu_\beta(\mathcal{U}_{>L}) \leq N_{L+1} e^{-\beta\Gamma_{L+1}} |\Lambda_\beta| \mu_\beta(\mathcal{S})$$

with N_{L+1} counting the number of configurations with support in $Q_{L+1}(0)$.

On the other hand, by considering only those configurations in $\mathcal{U}_{>L}$ that have a $Q_{L+1}(0)$ droplet, we get

$$(C.7) \quad \mu_\beta(\mathcal{U}_{>L}) \geq N_{L+1} e^{-\beta\Gamma_{L+1}} |\Lambda_\beta| \mu_\beta^{[Q_{L+1}(0)]^c}(\mathcal{S}),$$

where the last factor is the Gibbs weight of the configurations in \mathcal{S} with support outside $[Q_{L+1}(0)]^c$. It is easy to show that $\mu_\beta^{[Q_{L+1}(0)]^c}(\mathcal{S}) = \mu_\beta(\mathcal{S})[1 + o(1)]$ as $\beta \rightarrow \infty$ and so

$$(C.8) \quad \mu_\beta(\mathcal{U}_{>L}) \geq N_{L+1} e^{-\beta\Gamma_{L+1}} |\Lambda_\beta| \mu_\beta(\mathcal{S}) [1 + o(1)], \quad \beta \rightarrow \infty.$$

Combining (C.6) and (C.7), we conclude that $\lim_{\beta \rightarrow \infty} \mu_\beta(\mathcal{U}_{>L}) / \mu_\beta(\mathcal{S}) = 0$ if and only if

$$(C.9) \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| e^{-\Gamma_{L+1}} = 0. \quad \square$$

C.2. Kawasaki.

PROOF. Split

$$(C.10) \quad \mathcal{S} = \mathcal{S}_L \cup (\mathcal{S} \setminus \mathcal{S}_L) = \mathcal{S}_L \cup \mathcal{U}_{>L},$$

where $\mathcal{U}_{>L} \subset \mathcal{S}$ are those configurations σ for which there exists an x such that $|\text{supp}[\sigma] \cap B_{L_\beta, L_\beta}(x)| > L$. Then

$$(C.11) \quad \begin{aligned} \mu_\beta(\mathcal{U}_{>L}) &\leq \sum_{x \in \Lambda_\beta} \sum_{\sigma \in \mathcal{S}} \sum_{m=L+1}^K \mu_\beta(\sigma) \mathbb{1}_{\{|\text{supp}[\sigma] \cap B_{L_\beta, L_\beta}(x)|=m\}} \\ &= |\Lambda_\beta| [\varphi(\beta) + \gamma(\beta)], \end{aligned}$$

where

$$(C.12) \quad \begin{aligned} \varphi(\beta) &= \sum_{m=L+1}^K \sum_{\eta \in \mathcal{X}_\beta^{(m)}} \sum_{\substack{\zeta \in \mathcal{X}_\beta^{(n_\beta-m)} \\ \eta \vee \zeta \in \mathcal{S}}} \frac{e^{-\beta[H(\eta)+H(\zeta)]}}{Z_\beta^{(n_\beta)}} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \\ &\quad \times \mathbb{1}_{\{\text{supp}[\zeta] \subset [B_{L_\beta, L_\beta}(0)]^c\}} \end{aligned}$$

and $\gamma(\beta)$ is an error term arising from particles interacting across the boundary of $B_{L_\beta, L_\beta}(0)$. By the same argument as in (B.21), this term is negligible. Moreover,

$$(C.13) \quad \begin{aligned} \varphi(\beta) &\leq \sum_{m=L+1}^K \frac{\check{Z}_\beta^{(n_\beta-m)}}{Z_\beta^{(n_\beta)}} \left(\sum_{\eta \in \mathcal{S}^{(m)}} e^{-\beta H(\eta)} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \right) \\ &\leq [1 + o(1)] \mu_\beta(\mathcal{S}) \sum_{m=L+1}^K (\rho_\beta)^m \left(\sum_{\eta \in \mathcal{S}^{(m)}} e^{-\beta H(\eta)} \mathbb{1}_{\{\text{supp}[\eta] \subset B_{L_\beta, L_\beta}(0)\}} \right), \end{aligned}$$

where, in the last inequality, we use Lemmas B.1 and B.2. Now proceed as in (B.19) and (B.20), via the cluster expansion, to get

$$\begin{aligned}
 \varphi(\beta) &\leq [1 + o(1)]C\mu(\mathcal{S}) \sum_{m=L+1}^K \sum_{j=1}^m \sum_{\substack{2 \leq k_1, \dots, k_j \leq K \\ \sum_{i=1}^j k_i = m}} e^{-\beta[H_{k_i} + k_i\Delta - (\Delta - \delta_\beta)]} \\
 \text{(C.14)} \quad &\leq [1 + o(1)]C\mu(\mathcal{S})e^{-\beta[\Gamma_{L+1} - (\Delta - \delta_\beta)]},
 \end{aligned}$$

where H_k is the energy of a droplet with k particles that is closest to a square or quasi-square, $\Gamma_{L+1} = H_{L+1} + (L + 1)\Delta$ and the second inequality uses the isoperimetric inequality, together with the fact that $H_k + k\Delta$ is increasing in k for subcritical droplets.

On the other hand, by considering only those configurations in $\mathcal{U}_{>L}$ that have a droplet with $L + 1$ particles, we get

$$\text{(C.15)} \quad \varphi(\beta) \geq [1 + o(1)]C\mu(\mathcal{S})e^{-\beta[\Gamma_{L+1} - (\Delta - \delta_\beta)]}.$$

Combining (C.11), (C.14) and (C.15), we conclude that $\lim_{\beta \rightarrow \infty} \mu_\beta(\mathcal{U}_{>L}) / \mu_\beta(\mathcal{S}) = 0$ if and only if

$$\text{(C.16)} \quad \lim_{\beta \rightarrow \infty} |\Lambda_\beta| e^{-\beta(\Gamma_{L+1} - (\Delta - \delta_\beta))} = 0. \quad \square$$

APPENDIX D: THE CRITICAL DROPLET IS THE THRESHOLD

In this appendix, we show that our estimates on capacities imply that the *average* probability under the Gibbs measure μ_β of destroying a supercritical droplet and returning to a configuration in \mathcal{S}_L is exponentially small in β . We will give the proof for Kawasaki dynamics, the proof for Glauber dynamics being simpler.

Pick $M \geq \ell_c$. Recall from (2.7) that $e_{\mathcal{D}_M, \mathcal{S}_L}(\sigma) = c_\beta(\sigma)\mathbb{P}_\sigma(\tau_{\mathcal{S}_L} < \tau_{\mathcal{D}_M})$ for $\sigma \in \mathcal{D}_M$. Hence, summing over $\sigma \in \partial\mathcal{D}_M$, the internal boundary of \mathcal{D}_M , we get, using (2.8), that

$$\text{(D.1)} \quad \frac{\sum_{\sigma \in \partial\mathcal{D}_M} \mu_\beta(\sigma)c_\beta(\sigma)\mathbb{P}_\sigma(\tau_{\mathcal{S}_L} < \tau_{\mathcal{D}_M})}{\sum_{\sigma \in \partial\mathcal{D}_M} \mu_\beta(\sigma)c_\beta(\sigma)} = \frac{\text{CAP}(\mathcal{S}_L, \mathcal{D}_M)}{\sum_{\sigma \in \partial\mathcal{D}_M} \mu_\beta(\sigma)c_\beta(\sigma)}.$$

Clearly, the left-hand side of (D.1) is the escape probability to \mathcal{S}_L from $\partial\mathcal{D}_M$ *averaged* with respect to the canonical Gibbs measure μ_β conditioned on $\partial\mathcal{D}_M$, weighted by the outgoing rate c_β . To show that this quantity is small, it suffices to show that the denominator is large compared to the numerator.

By Lemma 4.2,

$$\begin{aligned}
 \text{(D.2)} \quad \text{CAP}(\mathcal{S}_L, \mathcal{D}_M) &\leq \text{CAP}(\mathcal{S}_L, (\mathcal{S}^c \setminus \tilde{\mathcal{C}}) \cup \mathcal{C}^+) \\
 &= N|\Lambda_\beta| \frac{4\pi}{\Delta\beta} e^{-\beta\Gamma} \mu_\beta(\mathcal{S})[1 + o(1)].
 \end{aligned}$$

On the other hand, note that $\partial\mathcal{D}_M$ contains all configurations σ for which there is an $M \times M$ droplet somewhere in Λ_β , all L_β -boxes not containing this droplet carry at most K particles and there is a free particle somewhere in Λ_β . The last condition ensures that $c_\beta(\sigma) \geq 1$. Therefore, we can use Lemma B.1 to estimate

$$\begin{aligned} \sum_{\sigma \in \mathcal{D}_M} \mu_\beta(\sigma) c_\beta(\sigma) &\geq |\Lambda_\beta| e^{-\beta H_{M^2}} \frac{\check{Z}_\beta^{(n_\beta - M^2)}}{Z_\beta^{(n_\beta)}} \\ (D.3) \qquad \qquad \qquad &= |\Lambda_\beta| e^{-\beta H_{M^2}} (\rho_\beta)^{M^2} \mu_\beta(\mathcal{S}) [1 + o(1)], \end{aligned}$$

where H_{M^2} is the energy of an $M \times M$ droplet. Combining (D.2) and (D.3), we find that the left-hand side of (D.1) is bounded from above by

$$(D.4) \qquad \left(N \frac{4\pi}{\Delta\beta} \right) \frac{\exp[-\beta\Gamma]}{\exp[-\beta(H_{M^2} + \Delta M^2)]} [1 + o(1)],$$

which is exponentially small in β because $\Gamma > H_{M^2} + \Delta M^2$ for all $M \geq \ell_c$.

Acknowledgments. The authors would like to thank Alessandra Bianchi, Alex Gaudillière, Dima Ioffe, Francesca Nardi, Enzo Olivieri and Elisabetta Scoppola for ongoing discussions on metastability and for sharing their work in progress. C.S. thanks Martin Slowik for stimulating exchanges.

REFERENCES

- [1] BEN AROUS, G. and CERF, R. (1996). Metastability of the three-dimensional Ising model on a torus at very low temperatures. *Electron. J. Probab.* **1** 10. [MR1423463](#)
- [2] BERMAN, K. A. and KONSOWA, M. H. (1990). Random paths and cuts, electrical networks, and reversible Markov chains. *SIAM J. Discrete Math.* **3** 311–319. [MR1061971](#)
- [3] BIANCHI, A., BOVIER, A. and IOFFE, D. (2009). Sharp asymptotics for metastability in the random field Curie–Weiss model. *Electron. J. Probab.* **14** 1541–1603.
- [4] BOVIER, A. (2006). Metastability: A potential theoretic approach. In *International Congress of Mathematicians. Vol. III* 499–518. Eur. Math. Soc., Zürich. [MR2275693](#)
- [5] BOVIER, A. (2009). Metastability. In *Methods of Contemporary Mathematical Statistical Physics* (R. Kotecký, ed.). *Lecture Notes in Math.* **1970**. Springer, Berlin.
- [6] BOVIER, A., ECKHOFF, M., GAYRARD, V. and KLEIN, M. (2002). Metastability and low lying spectra in reversible Markov chains. *Comm. Math. Phys.* **228** 219–255. [MR1911735](#)
- [7] BOVIER, A., DEN HOLLANDER, F. and NARDI, F. R. (2006). Sharp asymptotics for Kawasaki dynamics on a finite box with open boundary. *Probab. Theory Related Fields* **135** 265–310. [MR2218873](#)
- [8] BOVIER, A. and MANZO, F. (2002). Metastability in Glauber dynamics in the low-temperature limit: Beyond exponential asymptotics. *J. Statist. Phys.* **107** 757–779. [MR1898856](#)
- [9] CERF, R. and MANZO, F. (2008). Private communication.
- [10] DEHGHANPOUR, P. and SCHONMANN, R. H. (1997). Metropolis dynamics relaxation via nucleation and growth. *Comm. Math. Phys.* **188** 89–119. [MR1471333](#)
- [11] DEHGHANPOUR, P. and SCHONMANN, R. H. (1997). A nucleation-and-growth model. *Probab. Theory Related Fields* **107** 123–135. [MR1427719](#)

- [12] GAUDILLIÈRE, A., DEN HOLLANDER, F., NARDI, F. R., OLIVIERI, E. and SCOPPOLA, E. (2009). Ideal gas approximation for a two-dimensional rarefied gas under Kawasaki dynamics. *Stochastic Process. Appl.* **119** 737–774. [MR2499857](#)
- [13] GAUDILLIÈRE, A., DEN HOLLANDER, F., NARDI, F. R., OLIVIERI, E. and SCOPPOLA, E. (2009). Droplet dynamics in a two-dimensional rarified gas under Kawasaki dynamics. Unpublished manuscript.
- [14] GAUDILLIÈRE, A., DEN HOLLANDER, F., NARDI, F. R., OLIVIERI, E. and SCOPPOLA, E. (2009). Homogeneous nucleation for two-dimensional Kawasaki dynamics. To appear.
- [15] DEN HOLLANDER, F. (2004). Metastability under stochastic dynamics. *Stochastic Process. Appl.* **114** 1–26. [MR2094145](#)
- [16] DEN HOLLANDER, F. (2009). Three lectures on metastability under stochastic dynamics. In *Methods of Contemporary Mathematical Statistical Physics* (R. Kotecký, ed.). *Lecture Notes in Math.* **1970**. Springer, Berlin.
- [17] DEN HOLLANDER, F., NARDI, F. R., OLIVIERI, E. and SCOPPOLA, E. (2003). Droplet growth for three-dimensional Kawasaki dynamics. *Probab. Theory Related Fields* **125** 153–194. [MR1961341](#)
- [18] DEN HOLLANDER, F., OLIVIERI, E. and SCOPPOLA, E. (2000). Metastability and nucleation for conservative dynamics. *J. Math. Phys.* **41** 1424–1498. [MR1757966](#)
- [19] KOTECKÝ, R. and OLIVIERI, E. (1993). Droplet dynamics for asymmetric Ising model. *J. Statist. Phys.* **70** 1121–1148. [MR1208633](#)
- [20] LAWLER, G. F. (1991). *Intersections of Random Walks*. Birkhäuser, Boston. [MR1117680](#)
- [21] LAWLER, G. F., SCHRAMM, O. and WERNER, W. (2004). Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Ann. Probab.* **32** 939–995. [MR2044671](#)
- [22] NEVES, E. J. and SCHONMANN, R. H. (1991). Critical droplets and metastability for a Glauber dynamics at very low temperatures. *Comm. Math. Phys.* **137** 209–230. [MR1101685](#)
- [23] OLIVIERI, E. and VARES, M. E. (2005). *Large Deviations and Metastability. Encyclopedia of Mathematics and Its Applications* **100**. Cambridge Univ. Press, Cambridge. [MR2123364](#)
- [24] SCHONMANN, R. H. and SHLOSMAN, S. B. (1998). Wulff droplets and the metastable relaxation of kinetic Ising models. *Comm. Math. Phys.* **194** 389–462. [MR1627669](#)

A. BOVIER
 INSTITUT FUER ANGEWANDTE MATHEMATIK
 RHEINISCHE FRIEDRICH–WILHELMS-
 UNIVERSITAET BONN
 ENDENICHER ALLEE 60
 53115 BONN
 GERMANY
 E-MAIL: bovier@uni-bonn.de

F. DEN HOLLANDER
 C. SPITONI
 MATHEMATICAL INSTITUTE
 LEIDEN UNIVERSITY
 P.O. BOX 9512
 2300 RA LEIDEN
 THE NETHERLANDS
 AND
 EURANDOM
 P.O. BOX 513
 5600 MB EINDHOVEN
 THE NETHERLANDS
 E-MAIL: denholla@math.leidenuniv.nl
spitoni@eurandom.tue.nl