

Acromegaly : treatment and follow-up : the Leiden studies Biermasz, N.R.

Citation

Biermasz, N. R. (2005, November 2). *Acromegaly : treatment and follow-up : the Leiden studies*. Retrieved from https://hdl.handle.net/1887/4334

Version:	Corrected Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/4334

Note: To cite this publication please use the final published version (if applicable).

CONTENTS

Int	troduction		
1.	General introduction	9	
	Outline of the thesis	21	
Tre	Treatment outcome in acromegaly		
2.	Ten-year follow-up results of transsphenoidal microsurgery in acromegaly.	27	
	J Clin Endocrinol Metab 2000; 85:4596-4602		
3.	Long -term follow-up results of postoperative radiotherapy in 36 patients with	43	
	acromegaly. J Clin Endocrinol Metab 2000; 85:2476-2482		
4.	Radiotherapy is effective in reducing serum growth hormone concentration to	59	
	safe levels. Clin Endocrinol (Oxf) 2000; 53:321-327		
5.	Direct postoperative and follow-up results of transsphenoidal surgery in 19	71	
	acromegalic patients pretreated with octreotide compared to those in untreated		
	controls. J Clin Endocrinol Metab1999; 84:3551-3555		
6.	Sandostatin LAR in acromegaly: a 6-week injection interval suppresses	83	
	growth hormone secretion as effectively as a 4-week interval.		
	Clin Endocrinol (Oxf) 2003; 58:288-295		
Clinical outcome in acromegaly			
7.	Determinants of survival in treated acromegaly in a single center:	97	
	predictive value of serial insulin-like growth factor I measurements.		
	J Clin Endocrinol Metab. 2004; 89:2789-2796		
8.	Decreased quality of life in patients with acromegaly despite long-term cure of	113	
	growth hormone excess. J Clin Endocrinol Metab.2004; 89:5369-5376		
9.	Morbidity after long-term remission for acromegaly: persisting joint-related	131	
	complaints cause reduced quality of life. J Clin Endocrinol Metab 2005; 90:2731-2739		
10	. Long-term maintenance of the anabolic effects of growth hormone (GH) on the	153	
	skeleton in cured patients with acromegaly. Eur J Endocrinol 2005; 152: 53-60		
Dv	namic tests in acromegaly		
-	. Intravenous octreotide test predicts the long-term outcome of	167	
11	treatment with octreotide-long-acting repeatable in active acromegaly.	107	
	Growth Horm IGF Res. 2005;15:200-206		
17	. Postoperative persistent thyrotropin releasing hormone (TRH)-induced	181	
١Z	growth hormone release predicts recurrences in patients with acromegaly.	101	
	Clin Endocrinol (Oxf) 2002; 56:313-319		

GH regulation during treatment and in rare cases of acromegaly 13. Octreotide represses secretory-burst mass and non-pulsatile secretion, but 193 does not restore event frequency or orderly GH secretion in acromegaly. Am J Physiol Endocrinol Metab 2004; 286:E25-30 14. Acromegaly caused by growth hormone releasing hormone (GHRH)-producing 207 tumors: Long-term observational studies in three patients. Submitted **Current pharmacotherapy** 15. Current Pharmacotherapy for acromegaly: a review. 229 Exp Opin Pharmacother, 2005, in press **Summary and Discussion** 16. General Discussion and Conclusions 253 17. Summary, Nederlandse samenvatting 279 List of abbreviations 287 Nawoord 289 **Curriculum Vitae** 291 **Publications** 293

۲

۲

۲