



Universiteit  
Leiden  
The Netherlands

## Antiquities of the rainforest: evolution of mycoheterotrophic angiosperms growing on Glomeromycota

Mennes, C.B.

### Citation

Mennes, C. B. (2015, October 6). *Antiquities of the rainforest: evolution of mycoheterotrophic angiosperms growing on Glomeromycota*. Retrieved from <https://hdl.handle.net/1887/35731>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/35731>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/35731> holds various files of this Leiden University dissertation

**Author:** Mennes, Constantijn

**Title:** Antiquities of the rainforest : evolution of mycoheterotrophic angiosperms growing on Glomeromycota

**Issue Date:** 2015-10-06

## **References**

- Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In: Second International Symposium on Information Theory. Akademiai Kiado, Budapest. Pp. 267-281.
- Akaike, H. (1974). A new look at the statistical model identification. IEEE Transactions on Automatic Control 19: 716-723.
- Albert, V.A., Struwe, L. (1997). Phylogeny and classification of *Voyria* (saprophytic Gentianaceae). Brittonia 49: 466-479.
- Alley, N.F., Krieg, G.W., Callen, R.A. (1996). Early Tertiary Eyre Formation, lower Nelly Creek, southern Lake Eyre Basin, Australia: palynological dating of macrofloras and silcrete, and palaeoclimatic interpretations. Australian Journal of Earth Sciences 43: 71-84.
- Bailey, C.D., Koch, M.A., Mayer, M., Mummenhoff, K., O'Kane, S.L., Warwick, S.I., Windham, M.D., Al-Shehbaz, I.A. (2006). Toward a global phylogeny of the Brassicaceae. Molecular Biology and Evolution 23: 2142-2160.
- Barker, N.P., Weston, P.H., Rutschmann, F., Sauquet, H. (2007). Molecular dating of the 'Gondwanan' plant family Proteaceae is only partially congruent with the timing of the break-up of Gondwana. Journal of Biogeography 34: 2012-2027.
- Barrett, C.F., Davis, J.I. (2012). The plastid genome of the mycoheterotrophic *Corallorrhiza striata* (Orchidaceae) is in the relatively early stages of degradation. American Journal of Botany 99: 1513-1523.
- Barrett, C.F., Davis, J.I., Leebens-Mack, J., Conran, J.G., Stevenson, D.W. (2013). Plastid genomes and deep relationships among the commelinid monocot angiosperms. Cladistics 29: 65-87.
- Barrett, C.F., Freudenstein, J.V. (2008). Molecular evolution of *rbcL* in the mycoheterotrophic coralroot orchids (*Corallorrhiza* Gagnepin, Orchidaceae). Molecular Phylogenetics and Evolution 47: 665-679.
- Bello, M.A., Bruneau, A., Forest, F., Hawkins, J.A. (2009). Elusive relationships within order Fabales: Phylogenetic analyses using *matK* and *rbcL* sequence data. Systematic Botany 34: 102-114.
- Bello, M.A., Hawkins J.A., Rudall, P.J. (2010). Floral ontogeny in Polygalaceae and its bearing on the homologies of keeled flowers in Fabales. International Journal of Plant Science 171: 482-498.
- Bello, M.A., Rudall, P.J., Hawkins J.A. (2012). Combined phylogenetic analyses reveal interfamilial relationships and patterns of floral evolution in the eudicot order Fabales. Cladistics 28: 393-421.
- Bentham, G. (1855). On the South American Triurideae and leafless Burmanniaceae from the collections of Mr. Spruce. In: Bentham, G., Hooker, J. (eds.). Hooker's Journal of Botany and Kew Garden Miscellany 7: 8-17.
- Bentham, G., Hooker, J.D. (1883) Ordo CLXVIII Burmanniaceae. In: Bentham, G., Hooker J.D. *Genera plantarum*, Vol. 3. Reeve & Co., London, UK. Pp. 455-460.
- Berry, E.W. (1914) *The Upper Cretaceous and Eocene floras of South Carolina and Georgia*. United States Geological Survey, Washington, DC, USA.
- Bidartondo, M.I. (2005). The evolutionary ecology of myco-heterotrophy. New Phytologist 167: 335-352.
- Bidartondo, M.I., Bruns, T.D. (2002). Fine-level mycorrhizal specificity in the Monotropideae (Ericaceae): specificity for fungal species groups. Molecular Ecology 11: 557-569.
- Bidartondo, M.I., Read, D.J., Trappe, J.M., Merckx, V.S.F.T., Ligrone, R., Duckett, J.G. (2011). The dawn of symbiosis between plants and fungi. Biology Letters 7: 574-577.

- Bidartondo, M.I., Redecker, D., Hijri, I., Wiemken, A., Bruns, T.D., Domínguez, L., Sérsic, A., Leake, J.R., Read, D.J. (2002). Epiparasitic plants specialized on arbuscular mycorrhizal fungi. *Nature* 419: 389–392.
- Björkman, E. (1960). *Monotropa hypopitys* L.- an epiparasite on tree roots. *Physiologica Plantarum* 13: 308-327.
- Blume, C.L. (1825). *Sciaphila* Bl. *Bijdragen tot de Flora van Nederlands Indië*. 10de Stuk. Batavia (Lands Drukkerij). Pp. 514-515.
- Böhme, M. (2003). The Miocene Climatic Optimum: evidence from ectothermic vertebrates of Central Europe. *Palaeogeography, Palaeoclimatology, Paleaeoecology* 195: 389-401.
- Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T.G., Wu, C-H., Xie, D., Suchard, M.A., Rambaut, A., Drummond, A.J. (2014). BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Computational Biology* 10: e1003537.
- Bremer, K. (2000). Early Cretaceous lineages of monocot flowering plants. *Proceedings of the National Academy of Sciences USA* 97: 4707–4711.
- Bremer, K., Friis, E., Bremer, B. (2004). Molecular phylogenetic dating of asterid flowering plants shows Early Cretaceous diversification. *Systematic Biology* 53: 496-505 .
- Bush, C.M., Wagstaff, S.J., Fritsch, P.W. Kron, K.A. (2009). The phylogeny, biogeography and morphological evolution of *Gaultheria* (Ericaceae) from Australia and New Zealand. *Australian Systematic Botany* 22: 229–242.
- Caddick, L.R., Rudall, P.J., Wilkin, P., Hedderson, T.A.J., Chase, M.W. (2002). Phylogenetics of Dioscoreales based on combined analyses of morphological and molecular data. *Botanical Journal of the Linnean Society* 138: 123-144.
- Cameron, K.M., Chase, M.W., Rudall, P.J. (2003). Recircumscription of the Monocotyledonous Family Petrosaviaceae to include *Japonolirion*. *Brittonia* 55: 214-225.
- Carpenter, R.J., Jordan, G.J., Hill, R.S. (2007) A toothed Lauraceae leaf from the Early Eocene of Tasmania, Australia. *International Journal of Plant Sciences* 168: 1191–1198.
- Chacón, J., Camargo de Assis, M., Meerow, A.W., Renner, S.S. (2012). From East Gondwana to Central America: historical biogeography of the Alstroemeriaceae. *Journal of Biogeography* 39: 1806–1818.
- Champion, J. (1847). On two new Ceylon plants related to *Sciaphila* of Blume. *Calcutta Journal of Natural History* 7: 463-469.
- Chase, M.W., Fay, M.F., Devey, D.S., Maurin, O., Ronsted, N., Davies, T.J., Pillon, Y., Petersen, G., Seberg, O., Tamura, M.N.,asmussen-Lange, C.B., Kilu, K., Borsch, T., Davis, J., Stevenson, D.W., Pires, J.C., Givnish, T., Sytsma, K.J., McPherson, M.A., Graham, S.W., Rai, H.S. (2006). Multigene analyses of monocot relationships: A summary. *Aliso* 22: 63-75.
- Chase, M.W., Soltis, D.E., Olmstead, R.G., Morgan, D., Les, D.H., Mishler, B.D., Duvall, M.R., Price, R.A., et al. (1993). Phylogenetics of seed plants: an analysis of nucleotide sequences from the plastid gene *rbcL*. *Annals of the Missouri Botanical Garden* 80: 528-548, 550-580.
- Chase, M.W., Soltis, D.E., Soltis, P.S., Rudall, P.J., Fay, M.F., Hahn, W.H., Sullivan, S., Joseph, J., Molvray, M., Kores, P.J., Givnish, T.J., Sytsma, K.J., Pires, J.C. (2000). Higher-level systematics of the monocotyledons: An assessment of current knowledge and a new classification. In: Wilson, K.L. and Morisson, D.A. (eds.). *Monocots: Systematics and Evolution*. CSIRO Publishing, Melbourne, Victoria, Australia. Pp. 3-16.
- Cheek, M. (2003). Kupeaeae, a new tribe of Triuridaceae from Africa. *Kew Bulletin* 58: 939-949.

- Chen, X., Tamura, M.N. (2000). *Petrosavia*. In: Wu, Z.Y., Raven, P.H. (eds.). Flora of China, Vol. 24 (Flagellariaceae through Maranthaceae). Science Press, Beijing, China, and Missouri Botanical Garden Press, St. Louis, USA. P. 77.
- Chodat, R. (1896). Polygalaceae. In: Engler, A. & Prantl, K., Natürliche Pflanzenfamilien, 3, Leipzig, Germany. Pp. 323-345.
- Christopher, R.A. (1979). Normapolles and triporate pollen assemblages from the Raritan and Magothy Formations (Upper Cretaceous) of New Jersey. *Palynology* 3: 73-121.
- Conran, J.G. (1997). *Paracordyline kerguelensis*, an Oligocene monocotyledon macrofossil from the Kerguélen Islands. *Alcheringa* 21: 129-140.
- Conran, J.G., Bannister, J.M., Lee, D.E. (2009b). Earliest orchid macrofossils: Early Miocene *Dendrobium* and *Earina* (Orchidaceae: Epidendroideae) from New Zealand. *American Journal of Botany* 96: 466-474.
- Conran, J.G., Bannister, J.M., Mildenhall, D.C., Lee, D.E., Chacón, J. Renner, S.S. (2014). Leaf fossils of *Luzuriaga* and a monocot flower with in situ pollen of *Liliacidites contortus* Mildenh. & Bannister sp. nov. (Alstroemeriaeae) from the Early Miocene. *American Journal of Botany* 101: 141-155.
- Conran, J.G., Carpenter, R.J., Jordan, G.J. (2009a). Early Eocene *Ripogonum* (Liliales: Ripogonaceae) leaf macrofossils from southern Australia. *Australian Systematic Botany* 22: 219-228.
- Conran, J.G., Christophel, D.C., Cunningham, L. (2003). An Eocene monocotyledon from Nelly Creek, Central Australia, with affinities to Hemerocallidaceae (Lilianeae: Asparagales). *Alcheringa* 27: 107-115.
- Cook, L.G., Crisp, M.D. (2005). Not so ancient: the extant crown group of *Nothofagus* represents a post-Gondwanan radiation. *Proceedings of the Royal Society B: Biological Sciences* 272: 2535-2544.
- Courtillot, V.E., Renne, P.R. (2003). On the ages of flood basalt events. *Comptes Rendus Geoscience* 335: 113-140.
- Couvreur, T.L.P., Forest, F., Baker, W.J. (2011). Origin and global diversification patterns of tropical rain forests: inferences from a complete genus-level phylogeny of palms. *BMC Biology* 9: 44.
- Crayn, D.M., Rossetto, M., Maynard, D.J. (2006). Molecular phylogeny and dating reveals an Oligo-Miocene radiation of dry-adapted shrubs (former Tremandraceae) from rainforest tree progenitors (Elaeocarpaceae) in Australia. *American Journal of Botany* 93: 1328-1342.
- Cribb, P.J., Wilken, P., Clements, M. (1995). Corsiaceae: a new family for the Falkland Islands. *Kew Bulletin* 50: 171-172.
- Cronn, R., Liston, A., Parks, M., Gernandt, D.S., Shen, R., Mockler, T. (2008). Multiplex sequencing of plant chloroplast genomes using Solexa sequencing-by-synthesis technology. *Nucleic Acids Research* 36: e122.
- Cronquist, A. (1968). The evolution and classification of flowering plants. Houghton & Mifflin, Boston, Massachusetts, USA
- Cronquist, A. (1981). An integrated system of classification of flowering plants. Columbia University Press, New York, New York, USA
- Cronquist, A. (1988). The evolution and classification of flowering plants. Second edition. New York Botanical Garden Press, New York, New York, USA.
- Cullings, K.W., Szaro, T.M., Bruns, T.D. (1996). Evolution of extreme specialization within a lineage of ectomycorrhizal epiparasites. *Nature* 379: 63-66.

Dahlgren, R.M.T., Clifford, H.T., Yeo, P.F. (1985). The families of the monocotyledons. Structure, Evolution, and Taxonomy. Springer, Berlin, Germany.

Darriba, D., G.L. Taboada, R. Doallo, D. Posada. (2012). JModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 772.

Davis, C.C., Webb, C.O., Wurdack, K.J., Jaramillo, C.A., Donoghue, M.J. (2005). Explosive Radiation of Malpighiales Supports a Mid-Cretaceous Origin of Modern Tropical Rain Forests. *The American Naturalist* 165: E36-E65.

Davis, J.I., Stevenson, D.W., Peterson, G., Seberg, O., Campbell, L.M., Freudenstein, J.V., Goldman, D.H., Hardy, C.R., Michelangeli, F.A., Simmons, M.P., Specht, C.D., Vergara-Silva, F., Gandolfo, M. (2004). A phylogeny of the monocots, as inferred from *rbcL* and *atpA* sequence variation, and a comparison of methods for calculating Jackknife and Bootstrap values. *Systematic Botany* 29: 467-510.

De Bruyn, M., B. Stelbrink, R.J. Morley, R. Hall, G.R. Carvalho, C.H. Cannon, G. van den Berghe, *et al.* (2014). Borneo and Indochina are Major Evolutionary Hotspots for Southeast Asian Biodiversity. *Systematic Biology* 63: 879-901.

Demessure, B., Sodzi, N., Petit, J. (1995). A set of universal primers for amplification of polymorphic non-coding regions of mitochondrial and chloroplast DNA in plants. *Molecular Ecology* 4: 129-131.

Drummond, A.J., Ashton, B., Buxton, S., Cheung, M., Cooper, A., Duran, C., Field, M., *et al.* (2013). Geneious Pro v6.1.4. Available from <http://www.geneious.com/>, [accessed February 21, 2013].

Drummond, A.J., B. Ashton, S. Buxton, M. Cheung, A. Cooper, C. Duran, M. Field, *et al.* (2013). Geneious Pro v6.1.7. Available from <http://www.geneious.com/> [accessed 19 February 2014].

Drummond, A.J., Suchard, M.A., Xie, D. Rambaut, A. (2012). Bayesian phylogenetics with BEAUTi and the BEAST 1.7. *Molecular Biology and Evolution* 29: 1969–1973.

Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792-1797.

Edgar, R.C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26: 2460-2461.

Edgar, R.C. (2013). UPARSE: Highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* 10: 996-998.

Ellison, A.M., Gotelli, N.J. (2009). Energetics and the evolution of carnivorous plants – Darwin’s ‘most wonderful plants in the world’. *Journal of Experimental Botany* 60: 19-42.

Endlicher, S. (1837). Genera Plantarum Secundum Ordines Naturalis Disposita. Vindobonae. Pp. 282.

Engler, A. (1889a) Burmanniaceae. In: Engler, A. & Prantl, K. (eds.) Die Natürlichen Pflanzenfamilien II. Teil. Wilhelm Engelmann, Leipzig, Germany. Pp. 44–51.

Engler, A. (1889b). Triuridaceae. In: Engler, A. & Prantl, K. Die Natürlichen Pflanzenfamilien, ed. 1, II (1). Verlag von Wilhelm Engelmann, Leipzig, Germany. Pp. 235-238.

Engler, A. (1897). Übersicht über die Unterabteilungen, Klassen, Reihen, Unterreihen und Familien der Embryophyta siphonogama. In: Engler, A. & Prantl, K. Die Natürlichen Pflanzenfamilien, Nachträge zum II.-IV. Teil. Verlag von Wilhelm Engelmann, Leipzig, Germany. Pp. 342.

Engler, A. (1915). Triuridaceae. In: Engler, A. & Prantl, K. Die Natürlichen Pflanzenfamilien, Nachträge IV zu II.1. Verlag von Wilhelm Engelmann, Leipzig, Germany. P. 8.

Erdtman, G. (1952). Pollen morphology and plant taxonomy – angiosperms: an introduction to

- palynology. Almqvist & Wiksell, Stockholm, Sweden.
- Eriksen, B. (1993a). Floral anatomy and morphology in the Polygalaceae. *Plant Systematic and Evolution* 186: 17-32.
- Eriksen, B. (1993b). Phylogeny of the Polygalaceae and its taxonomic implications. *Plant Systematic and Evolution* 186: 33-55.
- Eyre-Walker, A., Gaut, B.G. (1997). Correlated rates of synonymous site evolution across plant genomes. *Molecular Biology and Evolution* 14: 455-460.
- Faith, D.P. (1992). Conservation evaluation and phylogenetic diversity. *Biological Conservation* 61: 1-10.
- Favre, A., Yuan, Y.-M., Küpfer, P., Alvarez, N. (2010). Phylogeny of subtribe Gentianinae (Gentianaceae): Biogeographic inferences despite limitations in temporal calibration points. *Taxon* 59: 1701-1711.
- Fay, M.F., Chase, M.W., Rønsted, N., Devey, D.S., Pillon, Y., Pires, J.C., Petersen, G., Seberg, O., Davis, J.I. (2006). Phylogenetics of Liliales: summarized evidence from combined analyses of five plastid and one mitochondrial loci. *Aldo* 22: 559-565.
- Feild, T.S., Brodribb, T.J. (2005). A unique mode of parasitism in the conifer coral tree *Parasitaxus ustus* (Podocarpaceae). *Plant, Cell & Environment* 28: 1316-1325.
- Felsenstein, J. (1978). Cases in which parsimony or compatibility methods will be positively misleading. *Systematic Zoology* 27: 401-410.
- Felsenstein, J. (1985). Confidence limits on phylogenetics: an approach using the Bootstrap. *Evolution* 39: 783-791.
- Forest, F., Chase, M.W., Persson, C., Crane, P.R., Hawkins, J.A. (2007). The role of biotic and abiotic factors in evolution of ant dispersal in the milkwort family (Polygalaceae). *Evolution* 61: 1675-1694.
- Freudenstein, J.V., Barrett, C.F. (2010). Mycoheterotrophy and diversity in Orchidaceae. In: Seberg, O., Petersen, G., Barfod, A.S., Davis, J.I. (eds.). *Diversity, phylogeny and evolution in the monocotyledons*. Aarhus University Press, Aarhus, Denmark. Pp. 25-37.
- Freudenstein, J.V., Chase, M.W. (2001). Analysis of mitochondrial *nad1b-c* intron sequences in Orchidaceae: utility and coding of length-change characters. *Systematic Botany* 26: 643-657.
- Friis, E.M. (1988). *Spirematospermum chandlerae* sp. nov., an extinct species of Zingiberaceae from the North American Cretaceous. *Tertiary Research* 9: 149-157.
- Friis, E.M., Crane, P.R., Pedersen, K.R. (2011). Early flowers and angiosperm evolution. Cambridge: Cambridge University Press.
- Furness, C.A., Rudall, P.J., Eastman, A. (2002). Contribution of pollen and tapetal characters to the systematics of Triuridaceae. *Plant Systematics and Evolution* 235: 209-218.
- Gallaud, I. 1905. Études sur les mycorhizes endotrophes. *Revue Générale de Botanique* 17: 7-48, 66-85, 123-136, 223-239, 313-325, 423-433, 479-496.
- Gandolfo, M.A., Nixon, K.C., Crepet, W.L. (2002). Triuridaceae fossil flowers from the Upper Cretaceous of New Jersey. *American Journal of Botany* 89: 1940-1957.
- Gandolfo, M.A., Nixon, K.C., Crepet, W.L., Stevenson, D.W., Friis, E.M. (1998). Oldest known fossils of monocotyledons. *Nature* 394: 532-533.
- Gardner, G. (1843). Description of *Peltophyllum*, a new genus of plants allied to Triuris of Miers, with

- remarks on their affinities. *Transactions of the Linnean Society of London* 19: 155–160. (1845).
- Gernhard, T. (2008). The conditioned reconstructed process. *Journal of Theoretical Biology* 253: 769–778.
- Geyer, C.J. (1991). Markov Chain Monte Carlo maximum likelihood. In: Keramidas, E.M. (ed.) Computing science and statistics: Proceedings of the 23<sup>rd</sup> symposium on the interface. Fairfax: Interface Foundation of North America inc. Fairfax Station, Virginia, USA. Pp. 156–163.
- Giesen, H. (1938). Triuridaceae. In: Engler, A. *Das Pflanzenreich. Regni vegetabilis conspectus IV. 18.* Verlag von Wilhelm Engelmann, Leipzig. Pp. 1–84.
- Giret, A., Lameyre, J., Beaux, J.F., Gautier, I., Verdier, O., Chotin, P., Cantagrel, J.M. (1989). Géologie, deux siècles de recherche dans les îles Kerguelen. In: Laubier, L. (ed.) *Actes du Colloque sur la Recherche Française dans les Terres Australes (Kerguelen, Crozet, Saint-Paul et Amsterdam) et à Partir des Navires qui les Desservent*, Strasbourg 1987. Paris: Comité National Français des Recherches Antarctiques. Pp. 345–355.
- Givnish, T.J., Ames, M., McNeal, J.R., McKain, M.R., Steele, P.R., dePamphilis, C.W., Graham, S.W., Pires, J.C., Stevenson, D.W., Zomlefer, W.B., Briggs, B.G., Duvall, M.R., Moore, M.J., Heaney, J.M., Soltis, D.E., Soltis, P.S., Thiele, K., Leebens-Mack, J.H. (2010). Assembling the tree of the monocotyledons: plastome sequence phylogeny and evolution of Poales. *Annals of the Missouri Botanical Garden* 97: 584–616.
- Gohn, G.S., Dowsett, H.J., Sohl, N.F. (1992). Biostratigraphy of the Middendorf Formation (Upper Cretaceous) in a corehole at Myrtle Beach, South Carolina. *U.S. Geological Survey Bulletin* 2030: 1–12.
- Goldblatt, P., Manning, J.C. (2010). *Geosiris albiflora* (Geosiridoideae), a new species from the Comoro Archipelago. *Bothalia* 40: 169–171.
- Goldblatt, P., Rodriguez, A., Powell, M.P., Davies, T.J., Manning, J.C., Van der Bank, M., Savolainen, V. (2008). Iridaceae ‘Out of Australasia’? Phylogeny, Biogeography, and Divergence Time Based on Plastid DNA Sequences. *Systematic Botany* 33: 495–508.
- Graham, A. (1984). *Lisianthius* pollen from the Eocene of Panama. *Annals of the Missouri Botanical Garden* 71: 987–993.
- Graham, S.W., Reeves, P.A., Burns, A.C.E., Olmstead, R.C. (2000). Microstructural changes in noncoding chloroplast DNA: interpretation, evolution, and utility of indels and inversions in basal angiosperm phylogenetic inference. *International Journal of Plant Sciences* 161: S83–S96.
- Graves, G. (1819). In: Curtis W. (1821). *Flora Londinensis: Containing a History of the Plants Indigenous to Great Britain, Illustrated by Figures of the Natural Size, Enlarged by George Graves and William Jackson Hooker.* Vol. IV, plate 48. London, UK: Richard & Arthur Taylor.
- Greuter W., and others (1988). International Code of Botanical Nomenclature. Koelz Scientific, Königstein. *Regnum vegetabile.* P. 118.
- Guindon, S., Gascuel, O. (2003). A simple, fast, and accurate method to estimate large phylogenies by maximum likelihood. *Systematic Biology* 52: 696–704.
- Gustafsson, A.L., Verola, C.F., Antonelli, A. (2010). Reassessing the temporal evolution of orchids with new fossils and a Bayesian relaxed clock, with implications for the diversification of the rare South American genus *Hoffmannseggella* (Orchidaceae: Epidendroideae). *BMC Evolutionary Biology* 10: 177.
- Haggard, K.K., Tiffney, B.H. (1997). The flora of the Early Miocene Brandon Lignite, Vermont, USA. VIII. *Caldesia* (Alismataceae). *American Journal of Botany* 84: 239–252.

## *Antiquities of the rainforest*

- Hall, R. (2009). Southeast Asia's changing palaeogeography. *Blumea* 54: 148-161.
- Harley, M.M. (1996). Palm pollen and the fossil record. Dissertation. Royal Botanic Gardens, Kew, UK.
- Harling, G., Wilder, G.J., Eriksson, R. (1998). Cyathaceae. In: Kubitzki, K. (ed.). The Families and Genera of Vascular Plants. III Flowering Plants: Monocotyledons. Springer, Berlin, Germany. Pp. 202-215.
- Heads, M. (2006). Panbiogeography of *Nothofagus* (Nothofagaceae): analysis of the main species massings. *Journal of Biogeography* 33: 1066-1075.
- Heintze, A. (1927). Cormofyternas Fylogeni (Phylogenie der Cormophyten). Lund, Hakan Ohlssons Boktryckeri. P. 77.
- Helgason, T., Daniell, T.J., Husband, R., Fitter A.H., Young, J.P.W. (1998). Ploughing up the wood-wide web? *Nature* 394: 431.
- Hemsley, W.B. (1907). Two new Triuridaceae, with some remarks on the genus *Sciaphila*, Blume. *Annals of Botany - London* 21: 71-77.
- Hu, J.-M., Lavin, M., Wojciechowski, M.F., Sanderson, M.J. (2000). Phylogenetic systematics of the tribe Millettiaeae (Leguminosae) based on chloroplast *trnK/matK* sequences and its implications for evolutionary patterns in Papilionoideae. *American Journal of Botany* 87: 418-430.
- Hurvich, C.M., Tsai, C-L. (1989). Regression and time series model selection in small samples. *Biometrika* 76: 297-307.
- Hutchinson, J. (1934). The families of flowering plants. II. Monocotyledons. Macmillan and co. London, UK. Pp. 37-38
- Hutchinson, J. (1967). The genera of flowering plants (Angiospermae): dicotyledons, vol. II. Oxford University Press, London, UK. Pp. 659.
- Hynson, N.A., Bruns, T.D. (2009). Evidence of a myco-heterotroph in the plant family Ericaceae that lacks mycorrhizal specificity. *Proceedings of the Royal Society B- Biological Sciences* 276: 4053-4059.
- Hynson, N.A., Bruns, T.D. (2010). Fungal hosts for mycoheterotrophic plants: a nonexclusive, but highly selective club. *New Phytologist* 185: 598-601.
- Hynson, N.A., Madsen, T.P., Selosse, M.-A., Adam, I.K.U., Ogura-Tsujita, Y., Roy, M., Gebauer, G. (2013). The Physiological Ecology of Mycoheterotrophy. In: V.S.F.T. Merckx [ed.], Mycoheterotrophy. The biology of plants living on fungi. Springer. New York. USA. Pp. 297-342.
- Ibisch, P.L., Neinhuis, C. & Rojas N., P. (1996) On the biology, biogeography, and taxonomy of *Arachnitis* Phil. nom. cons. (*Corsiaceae*) in respect to a new record from Bolivia. *Willdenowia* 26: 321-332.
- Iles, W.J.D., Smith, S.Y., Gandolfo, M.A., Graham, S.W. (2015). A review of monocot fossils suitable for molecular dating analyses. *Botanical Journal of the Linnean Society* 178: 346-374.
- Imhof, S. (2007). Specialized mycorrhizal colonization pattern in achlorophyllous *Epirixanthes* spp. (Polygalaceae). *Plant Biology* 9: 786-792.
- Janssen, T., Bremer, K. (2004). The major monocot groups inferred from 800+ *rbcL* sequences. *Botanical Journal of the Linnean Society* 146: 385-398.
- Janssens, S.B., Knox, E.B., Huysmans, S., Smets, E.F., Merckx, V.S.F.T. (2009). Rapid radiation of *Impatiens* (Balsaminaceae) during Pliocene and Pleistocene: Result of a global climate change.

Molecular Phylogenetics and Evolution 52: 806-824 .

Jaramillo, C., Rueda, M.J., Mora, G. (2006). Cenozoic Plant Diversity in the Neotropics. Science 311: 1893-1896.

Jones, D.L., Gray, B. (2008). *Corsia dispar* D.L.Jones & B.Gray (Corsiaceae), a new species from Australia, and a new combination in *Corsia* for a New Guinea taxon. Austrobaileya 7: 717–722.

Jonker, F.P. (1938) *A monograph of the Burmanniaceae*. Mededeelingen van het Botanisch Museum en Herbarium van de Rijks Universiteit te Utrecht 51. University of Utrecht, Utrecht, the Netherlands.

Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D., Blomberg, S.P., Webb C.O. (2010). Picante: R tools for integrating phylogenies and ecology. Bioinformatics 26: 1463-1464.

Kim, J.S., Hong, J.-K., Chase, M.W., Fay, M.F., Kim, J.-H. (2013). Familial relationships of the monocot order Liliales based on a molecular phylogenetic analysis using four plastid loci: *matK*, *rbcL*, *atpB* and *atpF-H*. Botanical Journal of the Linnean Society 172: 5–21.

Kissling, J. (2012). Taxonomy of *Exochaenium* and *Lagenias*: Two resurrected genera of the tribe Exaceae (Gentianaceae). Systematic Botany 37: 238-253.

Kissling, J., Barrett, S.C.H. (2013). Variation and evolution of herkogamy in *Exochaenium* (Gentianaceae): implications for the evolution of distyly. Annals of Botany 112: 95-102.

Kissling, J., Yuan, Y., Küpfer, P., Mansion, G. (2009). The polyphyletic genus *Sebaea* (Gentianaceae): A step forward in understanding the morphological and karyological evolution of the Exaceae. Molecular Phylogenetics and Evolution 53: 734-748.

Koressaar, T., Remm, M. (2007). Enhancements and modifications of primer design program Primer3. Bioinformatics 23: 1289–1291.

Kubitzki, K. (1998a). Stemonaceae. In: Kubitzki, K. (ed.). The Families and Genera of Vascular Plants. III Flowering Plants: Monocotyledons. Springer, Berlin, Germany. Pp. 422-425.

Kubitzki, K. (1998b). Velloziaceae. In: Kubitzki, K. (ed.). The Families and Genera of Vascular Plants. III Flowering Plants: Monocotyledons. Springer, Berlin, Germany. Pp. 459-467.

Lanfear, R., Calcott, B., Ho, S.Y.W., Guindon, S. (2012). PartitionFinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. Molecular Biology and Evolution 29: 1695–1701.

Le Maout, E., Decaisne, J. (1868). Traité général de botanique descriptive et analytique. Librairie de fermin didot frères. Paris, France. Pp. 553.

Leake, J.R. (1994). The biology of myco-heterotrophic ('saprophytic') plants. New Phytologist 127: 171–216.

Lemaire, B., Huysmans, S., Smets, E.F., Merckx, V.S.F.T. (2011). Rate accelerations in nuclear 18S rDNA of mycoheterotrophic and parasitic angiosperms. Journal of Plant Research 124: 561-576.

Les, D.H., Cleland, M.A., Waycott, M. (1997). Phylogenetic Studies in Alismatidae, II: Evolution of Marine Angiosperms (Seagrasses) and Hydrophily. Systematic Botany 22: 443-463.

Linder, H.P. (1987). The evolutionary history of the Poales/ Restionales- a hypothesis. Kew Bulletin 42: 297-318.

Linder, H.P. (2001). Plant diversity and endemism in sub-Saharan tropical Africa. Journal of Biogeography 28: 169-182.

- Lindley, J. (1846). *The Vegetable Kingdom* 213. Bradbury & Evans, London, UK.
- Lindqvist, J.K., Lee, D.E. (2009). High-frequency paleoclimate signals from Foulden Maar, Waipiatia Volcanic Field, southern New Zealand: an Early Miocene varved lacustrine diatomite deposit. *Sedimentary Geology* 222: 98–110.
- Linnaeus, C. (1753). *Species Plantarum, Exhibentes Plantas Rite Cognitas, Ad Genera Relatas, Cum Differentiis Specificis, Nominibus Trivialibus, Synonymis Selectis, Locis Natalibus, Secundum Systema Sexuale Digestas.* 2 volumes. L. Salvius, Stockholm, Sweden.
- Logacheva, M.D., Schelkunov, M.I., Nuraliev, M.S., Samigullin, T.H., Penin, A.A. (2014). The plastid genome of mycoheterotrophic monocot *Petrosavia stellaris* exhibits both gene losses and multiple rearrangements. *Genome Biology and Evolution* 6: 238-246.
- Maas, P.J.M., Maas-van de Kamer, H., Van Benthem, J., Snelders, H.C.M., Rübsamen, T. (1986). Burmanniaceae. In: Organization for Flora Neotropica (eds.) *Flora Neotropica*, Vol. 42. New York Botanical Garden, New York, USA.
- Maas, P.J.M., Rübsamen, T. (1986). Triuridaceae. In: Organization for Flora Neotropica (eds.) *Flora Neotropica*, Vol. 40. New York Botanical Garden, New York, USA.
- Maas, P.J.M., Ruyters, P. (1986). *Voyria* and *Voyriella* (saprophytic Gentianaceae). In: Organization for Flora Neotropica (eds.) *Flora Neotropica*, Vol. 41. New York Botanical Garden, New York, USA.
- Maas-van de Kamer, H. (1995). Triuridiflorae – Gardner's delight? In: Rudall, P.J., Cribb, P.J., Cutler, D.F., Humphries, C.J. (eds.) *Monocotyledons: systematics and evolution*. Royal Botanic Gardens, Kew. Pp. 287-301.
- Maas-van de Kamer, H. (1998). Burmanniaceae. In: Kubitzki, K. (ed.) *The families and genera of flowering plants, volume 3, Lilianae (except Orchidaceae)*. Springer, Berlin Germany. Pp. 154-164.
- Maas-van de Kamer, H., Weustenfeld, T. (1998). Triuridaceae. In: Kubitzki, K. (ed.). *The Families and Genera of Vascular Plants. III Flowering Plants: Monocotyledons*. Springer, Berlin. Pp. 452-458
- Mabberley, D.J. (2008). *Mabberley's Plant Book*. Third Edition. Cambridge University Press. P. 876.
- Maciunas, E., Conran, J.G., Bannister, J.M., Paull, R., Lee, D.E. (2011). Miocene *Astelia* (Asparagales: Asteliaceae) macrofossil from southern New Zealand. *Australian Systematic Botany* 24: 19-31.
- Maddison, W.P., Maddison, D.R. (2007). *Mesquite: a modular system for evolutionary analysis*. Version 2. Available from: <http://mesquiteproject.org/> [accessed 27 April 2015].
- Marcuse, M. (1902). Anatomisch-biologischer Beitrag zur Mycorrhizenfrage. Ph.D. Dissertation, Jena University, Jena, Germany.
- Martínez, E., Ramos, C.H. (1989). Lacandoniaceae (Triuridales): Una nueva familia de Mexico. *Annals of the Missouri Botanical Garden* 76: 128-135.
- Matzke, N. J. (2013). BioGeoBEARS: BioGeography with Bayesian (and Likelihood) Evolutionary Analysis in R Scripts. R package, version 0.2.1, published July 27, 2013. Available from: <http://CRAN.R-project.org/package=BioGeoBEARS> [accessed 27 April 2015].
- Matzke, N.J. (2014). Model selection in historical biogeography reveals that founder-event speciation is a crucial process in island clades. *Systematic Biology*. Available online August 14, 2014. Available from: <http://dx.doi.org/10.1093/sysbio/syu056> and <http://sysbio.oxfordjournals.org/content/early/2014/09/21/sysbio.syu056.full> — Supplementary data on Dryad: <http://datadryad.org/resource/doi:10.5061/dryad.2mc1t> [accessed 27 April 2015].
- Meng, S.W., Chen, Z.D., Li, D.Z., Liang, H.X. (2002). Phylogeny of Saururaceae based on mitochondrial

- matR* gene sequence data. Journal of Plant Research 115: 71-76.
- Mennes, C.B., Lam, V.K.Y., Rudall P.J., Lyon, S.P., Graham, S.W., Smets, E.F., Merckx, V.S.F.T. (2015a). Ancient Gondwana break-up explains the distribution of the mycoheterotrophic family Corsiaceae (Liliales). Journal of Biogeography 42: 1123-1136.
- Mennes, C.B., Moerland, M.S., Rath, M., Smets, E.F., Merckx, V.S.F.T. (2015b). Evolution of mycoheterotrophy in Polygalaceae: The case of *Epirixanthes*. American Journal of Botany 102: 598-608.
- Mennes, C.B., Smets, E.F., Moses, S.N., Merckx, V.S.F.T. (2013). New insights in the long-debated evolutionary history of Triuridaceae (Pandanales). Molecular Phylogenetics and Evolution 69: 994-1004.
- Merckx, V.S.F.T. (2008). Myco-heterotrophy in Dioscoreales: Systematics and Evolution. PhD thesis, Katholieke Universiteit Leuven, Belgium.
- Merckx, V.S.F.T., Bakker, F.T., Huysmans, S., Smets, E.F. (2009a). Bias and conflict in phylogenetic inference of myco-heterotrophic plants: a case study in Thismiaceae. Cladistics 25: 64-77.
- Merckx, V.S.F.T., Bidartondo, M.I. (2008). Breakdown and delayed cospeciation in the arbuscular mycorrhizal mutualism. Proceedings of the Royal Society B- Biological Sciences 275: 1029-1035.
- Merckx, V.S.F.T., Bidartondo, M.I., Hynson, N.A. (2009b). Myco-heterotrophy: when fungi host plants. Annals of Botany 104: 1255-1261.
- Merckx, V.S.F.T., Chatrou, L.W., Lemaire, B., Sainge, M.N., Huysmans, S. & Smets, E.F. (2008). Diversification of myco-heterotrophic angiosperms: evidence from Burmanniaceae. BMC Evolutionary Biology 8: 178.
- Merckx, V.S.F.T., Freudenstein, J.V. (2010). Evolution of mycoheterotrophy in plants: a phylogenetic perspective. New Phytologist 185: 605-609.
- Merckx, V.S.F.T., Freudenstein, J.V., Kissling J., Christenhusz, M.J.M., Stotler, R.E., Crandall-Stotler, B., Wickett, N., Rudall, P.J., Maas-van de Kamer, H., Maas, P.J.M. (2013b). Chapter 2: Taxonomy and classification. In: V.S.F.T. Merckx [ed.], Mycoheterotrophy. The biology of plants living on fungi, Pp. 19-102, Springer. New York. USA.
- Merckx, V.S.F.T., Huysmans, S., Smets, E.F. (2010a). Cretaceous origins of myco-heterotrophic lineages in Dioscoreales. In: Seberg, O., Petersen, G., Barfod, A. S. & Davis, J. (eds.). Diversity, phylogeny, and evolution in the monocotyledons. Aarhus University Press, Denmark. Pp. 39-53.
- Merckx, V.S.F.T., Janssens, S.B., Hynson, N.A., Specht, C.D., Bruns, T.D., Smets E.F. (2012). Mycoheterotrophic interactions are not limited to a narrow phylogenetic range of arbuscular mycorrhizal fungi. Molecular Ecology 21: 1524-1532.
- Merckx, V.S.F.T., Kissling, J., Henrich, H., Janssens, S.B., Mennes, C.B., Specht, C.D., Smets, E.F. (2013c). Phylogenetic relationships of the mycoheterotrophic genus *Voyria* and the implications for the biogeographic history of Gentianaceae. American Journal of Botany 100: 712-721.
- Merckx, V.S.F.T., Mennes, C.B., Peay, K.G., Geml, J. (2013a). Chapter 5: Evolution and Diversification. In: V.S.F.T. Merckx [ed.], Mycoheterotrophy. The biology of plants living on fungi, Pp. 215-244, Springer. New York. USA.
- Merckx V.S.F.T., Schols, P., Maas-van de Kamer, H., Maas, P., Huysmans, S., Smets, E.F. (2006). Phylogeny and evolution of Burmanniaceae (Dioscoreales) based on nuclear and mitochondrial data. American Journal of Botany 93: 1684-1698.
- Merckx, V.S.F.T., Smets, E.F. (2014). *Thismia americana*, the 101st anniversary of a botanical mystery.

International Journal of Plant Sciences 175: 165-175.

Merckx, V.S.F.T., Smets, E.F., Specht, C.D. (2013d). Chapter 3: Biogeography and Conservation. In: V.S.F.T. Merckx [ed.], Mycoheterotrophy. The biology of plants living on fungi, Pp. 103-156, Springer. New York. USA.

Merckx, V.S.F.T., Stöckel, M., Fleischmann, A., Bruns, T.D., Gebauer, G. (2010b).  $^{15}\text{N}$  and  $^{13}\text{C}$  natural abundance of two mycoheterotrophic and a putative partially mycoheterotrophic species associated with arbuscular mycorrhizal fungi. New Phytologist 188: 590-596.

Miers, J. (1841). Description of a new genus of plants from Brazil. Transactions of the Linnean Society of London 19 tab. VII: 77-80(1845).

Miers, J. (1850). On the family of Triuriaceae. Proceedings of the Linnean Society of London 2: 72-77.

Mitchell, A., Li, R., Brown, J.W., Schönberger, I., Wen, J. (2012). Ancient divergence and biogeography of *Raukaua* (Araliaceae) and close relatives in the southern hemisphere. Australian Systematic Botany 25: 432–446.

Morley, R.J. (2000). Origin and Evolution of Tropical Rain Forests. Wiley, New York, USA.

Mower, J.P., Stefanović, S., Hao, W., Gummow, J.S., Jain, K., Ahmed, D., Palmer, J.D. (2010). Horizontal acquisition of multiple mitochondrial genes from a parasitic plant followed by gene conversion with host mitochondrial genes. BMC Biology 8: 150.

Mueller, C. (1858). Triuriaceae. In: Walpers, G.G. (ed.) Annales Botanices Systematicae, tomus V. Lipsiae Sumptibus Ambrosii Abel. Pp. 915-919.

Neinhuis, C., Ibsch, P.L. (1998). Corsiaceae. In: Kubitzki, K. (ed.) The families and genera of vascular plants, Vol. 3, Liliaceae (except Orchidaceae). Springer, Berlin, Germany. Pp. 198–201.

Neyland, R., Hennigan, M. (2003). A phylogenetic analysis of large-subunit (26S) ribosome DNA sequences suggests that the Corsiaceae are polyphyletic. New Zealand Journal of Botany 41: 1–11.

Nikolov, L.A., Tomlinson, P.B., Manickam, S., Endress, P.K., Kramer, E.M., Davis, C.C. (2014). Holoparasitic Rafflesiaceae possess the most reduced endophytes and yet give rise to the world's largest flowers. Annals of Botany 114: 233-242.

Oehler, E. (1927). Entwicklungsgeschichtlich-zytologische Untersuchungen an einige saprophytischen Gentianaceen. Planta 3: 641-733.

Ohashi, H. (2000). *Petrosavia* (Petrosaviaceae) in Taiwan and Hainan. Taiwania 45: 263-269.

Olmstead, R.G., Palmer, J.D. (1994). Chloroplast DNA systematics: a review of methods and data analysis. American Journal of Botany 81: 1205–1224.

Öpik, M., Vanatoa, A., Vanatoa, E., Moora, M., Davison, J., Kalwij, J.M., Reier, Ü, Zobel, M. (2010). The online database *MaarjAM* reveals global and ecosystemic distribution patterns in arbuscular mycorrhizal fungi (Glomeromycota). New Phytologist 188: 223-241.

Pendry, C.A. (2001). Polygalaceae. In: Santisuk, T., Larsen, K. (eds.) Flora of Thailand 7. Royal Forest Department, Bangkok, Thailand. Pp. 498-538.

Pendry, C.A. (2010). *Epirixanthes compressa* Pendry, a new mycoheterotrophic species of Polygalaceae from Thailand. Thai Forest Bulletin, 38: 184-186.

Perkins, M.E., Brown, F.H., Nash, W.P., Williams, S.K., McIntosh, W. (1998). Sequence, age, and source of silicic fallout tuffs in middle to late Miocene basins of the northern Basin and Range province. Geological Society of America Bulletin 110: 344-360.

- Persson, C. (2001). Phylogenetic relationships in Polygalaceae based on plastid DNA sequences from the *trnL-F* region. *Taxon* 50: 763–779.
- Petersen, G., Seberg, O., Davis, J.I. (2013). Phylogeny of the Liliales (Monocotyledons) with special emphasis on data partition congruence and RNA editing. *Cladistics* 29: 274–295.
- Petersen, G., Seberg, O., Davis, J.I., Goldman, D.H., Stevenson, D.W., Campbell, L.M., Michelangeli, F.A., Specht, C.D., Chase, M.W., Fay, M.F., Pires, J.C., Freudenstein, J.V., Hardy, C.R., Simmons, M.P. (2006). Mitochondrial data in monocot phylogenetics. *Aldo* 22: 52–62.
- Plana, V. (2004). Mechanisms and tempo of evolution in the African Guineo-Congolian rainforest. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 359: 1585–1594.
- Poulsen, V.A. (1890). *Triuris major* sp. nov. Et Bidrag til Triuridaceernes Naturhistorie. *Botanisk Tidskrift* 17: 293–306.
- Prasad, V., Strömborg, C.A.E., Leaché, A.D., Samant, B., Patnaik, R., Tang, L., Mohabey, D.M., Ge, S., Sahni, A. (2011). Late Cretaceous origin of the rice tribe provides evidence for early diversification in Poaceae. *Nature Communications* 2: 480.
- Prasad, V., Strömborg, C.A.E., Leaché, A.D., Samant, B., Patnaik, R., Tang, L., Mohabey, D.M., Ge, S., Sahni, A. (2011). Late Cretaceous origin of the rice tribe provides evidence for early diversification in Poaceae. *Nature Communications* 2: 480.
- Qiagen. (2006). DNEasy plant handbook, “Plant Tissue (Mini Protocol)”. Pp. 24–27.
- Qiu, Y.-L., Li, L., Hendry, T.A., Li, R., Taylor, D.W., Issa, M.J., Ronen, A.J., Vekaria, M.L., White, A.M. (2006). Reconstructing the basal angiosperm phylogeny: evaluating information content of mitochondrial genes. *Taxon* 55: 837–856.
- R Core Team. (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available from <http://www.R-project.org/> [accessed 05 June 2014].
- Rabinowitz, P.D., Coffin, M.F., Falvey, D. (1983). The separation of Madagascar and Africa. *Science* 220: 67–69.
- Rai, H.S., O'Brien, H.E., Reeves P.A., Olmstead, R.G., Graham, S.W. (2003). Inference of higher-order relationships in the cycads from a large chloroplast data set. *Molecular Phylogenetics and Evolution* 29: 350–359.
- Rambaut, A. (2009). FigTree v1.3.1. Available from <http://tree.bio.ed.ac.uk/software/figtree/>, [accessed 21 February 2013].
- Rambaut, A. (2012) FigTree v1.4. University of Edinburgh, Edinburgh, UK. Available from <http://tree.bio.ed.ac.uk/software/figtree/> [accessed 28 April 2014].
- Rambaut, A., Drummond, A.J. (2007). Tracer v1.5. Available from <http://beast.bio.ed.ac.uk/Tracer> [accessed 21 February 2013].
- Rambaut, A., Suchard, M.A., Xie, W., Drummond, A.J. (2013). Tracer v1.6. University of Edinburgh, Edinburgh, UK. Available from <http://tree.bio.ed.ac.uk/software/tracer/> [accessed 28 April 2014].
- Ramírez, S.R., Gravendeel, B., Singer, R.B., Marshall, C.R. & Pierce, N.E. (2007). Dating the origin of the Orchidaceae from a fossil orchid with its pollinator. *Nature* 448: 1042–1045.
- Rath, M., Weber, H.C., Imhof, S. (2013). Morpho-anatomical and molecular characterization of the mycorrhizas of European Polygala species. *Plant Biology* 15: 548–557.

- Raubeson, L.A., Jansen, R.K. (2005) Chloroplast genomes of plants. In: Henry, R.J. (ed.) Plant diversity and evolution: genotypic and phenotypic variation in higher plants. CABI, Wallingford, UK. Pp. 45–68.
- Raven, P.H., Axelrod, D.I. (1972). Plate tectonics and Australasian paleobiogeography. *Science* 176: 1379–1386.
- Raven, P.H., Axelrod, D.I. (1974). Angiosperm biogeography and past continental movements. *Annals of the Missouri Botanical Garden* 61: 539–673.
- Raynal-Roques, A. (1967). Sur un *Sebaea* Africain saprophyte (Gentianaceae). *Adansonia* 7: 207–219.
- Ree, R.H., Smith, S.A. (2008). Maximum Likelihood Inference of Geographic Range Evolution by Dispersal, Local Extinction, and Cladogenesis. *Systematic Biology* 57: 4–14.
- Remizowa, M.V., Sokoloff, D.D., Rudall, P.J. (2010). Evolutionary history of the monocot flower. *Annals of the Missouri Botanical Garden* 97: 617–645.
- Reveal, J.L. (1992). Validation of subclass and superordinal names in Magnoliophyta. *Novon* 2: 235–237.
- Rodríguez-de la Rosa, R.A., Cevallos-Ferriz, S.R.S. (1994). Upper Cretaceous Zingiberalean fruits with in situ seeds from southeastern Coahuila, Mexico. *International Journal of Plant Sciences* 155: 786–805.
- Ronquist, F., Teslenko, M., Van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A., Huelsenbeck, J.P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542.
- Roy, M., Gonneau, C., Rocheteau, A., Berveiller, D., Thomas, J.-C., Damesin, C., Selosse, M.-A. (2013). Why do mixotrophic plants stay green? A comparison between green and achlorophyllous orchid individuals in situ. *Ecological Monographs* 83: 95–117.
- Roy, M., Watthana, S., Stier, A., Richard, F., Vessabutr, S., Selosse, M.-A. (2009). Two mycoheterotrophic orchids from Thailand tropical dipterocarpaceous forests associate with a broad diversity of ectomycorrhizal fungi. *BMC Biology* 7: 51.
- Rübsamen, T. (1986). Morphologische, embryologische und systematische Untersuchungen an Burmanniaceae und Corsiaceae (mit Ausblick auf die Orchidaceae-Apostasioideae). *Dissertationes Botanicae* 92. J. Cramer, Berlin, Germany.
- Rudall, P.J. (2002). Homologies of inferior ovaries and septal nectaries in monocotyledons. *International Journal of Plant Sciences* 163: 261–276.
- Rudall, P.J., Bateman, R.M. (2002). Roles of synorganisation, zygomorphy and heterotopy in floral evolution: the gynostemium and labellum of orchids and other lilioid monocots. *Biological Reviews* 77: 403–441.
- Rudall, P.J., Bateman, R.M. (2006). Morphological phylogenetic analysis of Pandanales: Testing contrasting hypotheses of floral evolution. *Systematic Botany* 31: 223–238.
- Rudall, P.J., Eastman, A. (2002). The questionable affinities of *Corsia* (Corsiaceae): evidence from floral anatomy and pollen morphology. *Botanical Journal of the Linnean Society* 138: 315–324.
- Rudall P.J., Stobart, K.L., Hong, W.P., Conran, J.G., Furness, C.A., Kite, G., Chase, M.W. (2000). Consider the lilies: systematics of Liliales. In: Wilson, K.L., Morrison, D.A. (eds.) *Proceedings of the Second International Conference on the Comparative Biology of the Monocots*, Sydney, Vol. 1, Monocots: systematics and evolution. CSIRO, Melbourne, Australia. Pp. 347–359.
- Sage, R.F. (2003). The evolution of C<sub>4</sub> photosynthesis. *New Phytologist* 161: 341–370.

- Singe, M.N., Kenfack, D., Chuyong, G.B. (2013). Two new species of *Afrothismia* (Thismiaceae) from southern Cameroon. Kew Bulletin 68: 591-597.
- Sanmartín, I., Ronquist, F. (2004). Southern Hemisphere biogeography inferred by event-based models: plant versus animal patterns. Systematic Biology 53: 216–243.
- Schlechter, R. (1912). Neue Triuridaceae Papuasiens. Botanische Jahrbücher 49, 70-89.
- Schumann, K., Lauterbach, C. (1905). Nachträge zur Flora der deutschen Schutzgebiete in der Südsee: 54-56.
- Schuster, T.M., Setaro, S.D., Kron, K.A. (2013). Age estimates for the buckwheat family Polygonaceae based on sequence data calibrated by fossils and with a focus on the Amphi-Pacific *Muehlenbeckia*. PLoS ONE 8: e61261.
- Schwarzott, D., Schüßler, A. (2001). A simple and reliable method for SSU rRNA gene DNA extraction, amplification, and cloning from single AM fungal spores. Mycorrhiza 10: 203-207.
- Scotese, C.R., Gahagan, L.M., Larson, R.L. (1988). Plate tectonic reconstructions of the Cretaceous and Cenozoic ocean basins. Tectonophysics 155: 27–48.
- Silvestro, D., Michalak, I. (2012). raxmlGUI: a graphical front-end for RAxML. Organisms Diversity and Evolution 12: 335–337.
- Simon, L. (1996). Phylogeny of the Glomales: deciphering the past to understand the future. New Phytologist 133: 95-101.
- Sirichamorn, Y., Adema, F.A.C.B., Gravendeel, B., Van Welzen, P.C. (2012). Phylogeny of palaeotropic Derris-like taxa (Fabaceae) based on chloroplast and nuclear DNA sequences shows reorganization of (infra)generic classifications is needed. American Journal of Botany 99: 1793-1808.
- Smets E.F., Ronse De Craene, L.P., Caris, P., Rudall, P.J. (2000). Floral nectaries in monocotyledons: distribution and evolution. In: Wilson, K.L., Morrison, D.A. (eds.) Proceedings of the Second International Conference on the Comparative Biology of the Monocots, Sydney, Vol. 1, Monocots: systematics and evolution. CSIRO, Melbourne, Australia. Pp. 230–240.
- Smit, E., Leeflang, P., Glandorf, B., Van Elsas, J.D., Wernars, K. (1999). Analysis of fungal diversity in the wheat rhizosphere by sequencing of cloned PCR-amplified genes encoding 18S rDNA and temperature gradient gel electrophoresis. Applied Environmental Microbiology 65: 2614-2621.
- Smith, S.E., Read, D.J. (2008). Mycorrhizal Symbiosis. Third Edition. Academic Press, London, UK.
- Sohl, N.F., Owens, J.P. (1991). Cretaceous stratigraphy of the Carolina coastal plain. In: Horton, J.W., Zullo, V.A. (eds.) The geology of the Carolinas: Carolina Geological Society fiftieth anniversary volume. University of Tennessee Press, Knoxville, Tennessee, USA. Pp. 191–220.
- Soltis, D.E., Smith, S.A., Cellinese, N., Wurdack, K.J., Tank, D.C., Brockington, S.F., Refulio-Rodriguez, N.F., Walker, J.B., Moore, M.J., Carlsward, B.S., Bell, C.D., Latvis, M., Crawley, S., Black, C., Diouf, D., Xi, Z., Rushworth, C.A., Gitzendanner, M.A., Sytsma, K.J., Qui, Y.L., Hilu, K.W., Davis, C.C., Sanderson, M.J., Beaman, R.S., Olmstead, R.G., Judd, W.S., Donoghue, M.J., Soltis, P.S. (2011). Angiosperm phylogeny: 17 genes, 640 taxa. American Journal of Botany 98: 704-730.
- Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. Bioinformatics 22: 2688–2690.
- Stevens, P.F. (2001 onwards). Angiosperm Phylogeny Website. Version 12, July 2012 [and more or less continuously updated since], Website <http://www.mobot.org/MOBOT/research/APweb/> [accessed 29 October 2014].

- Stevenson, D.W., Loconte, H. (1995). Cladistic analysis of monocot families. In: Rudall, P.J., Cribb, P.J., Cutler D.F., Humphries C.J. (eds.) *Monocotyledons: systematics and evolution*. Royal Botanic Gardens, Kew, UK. Pp. 543–578.
- Stone, B.C., Huynh, K.-L., Poppendieck, H.-H. (1998). Pandanaceae. In: Kubitzki, K. (ed.). *The Families and Genera of Vascular Plants. III Flowering Plants: Monocotyledons*. Springer, Berlin, Germany. Pp. 397–404.
- Struwe, L., Kadereit, J.W., Klackenberg, J., Nilsson, S., Thiv, M., Von Hagen, K.B., Albert, V.A. (2002). Systematics, character evolution, and biogeography of Gentianaceae, including a new tribal and subtribal classification. In: Struwe, L., Albert, V.A. (eds.). *Gentianaceae, systematics and natural history*. Cambridge University Press, Cambridge, UK. Pp. 21–301.
- Swenson, U., Nylander, S., Munzinger, J. (2014). Sapotaceae biogeography supports New Caledonia being an old Darwinian island. *Journal of Biogeography* 41: 797–809.
- Takahashi, H., Nishio, E., Hayashi, H. (1993). Pollination biology of the saprophytic species *Petrosavia sakuraii* (Makino) van Steenis in Central Japan. *Journal of Plant Research* 106: 213–217.
- Thorne, R. F. (1968). Synopsis of a putatively phylogenetic classification of the flowering plants. *Aliso* 6: 57–66.
- Thorne, R.F. (1983). Proposed new realignments in the angiosperms. *Nordic Journal of Botany* 3: 85–117.
- Thwaites, G.H.K. (1861). *Enumerario Plantarum Zeylaniae*. Dulau & Co. London, UK. Pp. 294.
- Tidwell, W.D., Parker, L.R. (1990). *Protowyucca shadishii* gen. et sp. nov., an arborescent monocotyledon with secondary growth from the middle Miocene of northwestern Nevada, U.S.A. *Review of Palaeobotany and Palynology* 62: 79–95.
- Tiffney, B.H. (1994). Re-evaluation of the age of the Brandon Lignite (Vermont, USA) based on plant megafossils. *Review of Palaeobotany and Palynology* 82: 299–315.
- Traverse, A. (1994). Palynofloral geochronology of the Brandon Lignite of Vermont, USA. *Review of Palaeobotany and Palynology* 82: 265–297.
- Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M., Rozen, S.G. (2012). Primer3 – new capabilities and interfaces. *Nucleic Acids Research* 40: e115.
- Van de Meerendonk, J.P.M. (1984). Triuridaceae. In: Van Steenis C.G.G.J. (ed.). *Flora Malesiana*, Ser. I., 10, 1984, Martinus Nijhoff, The Hague, the Netherlands. Pp. 109–121.
- Van der Meijden, R. (1988). Polygalaceae. In: C.G.G.J. van Steenis and W.J.J.O. de Wilde (eds.), *Flora Malesiana*, 10, Kluwer, Dordrecht, the Netherlands. Pp. 455–539.
- Van Royen, P. (1972). Sertulum Papuanum 17: Corsiaceae of New Guinea and surrounding areas. *Webbia* 27: 223–255.
- Veevers, J.J., Powell, C.M., Roots, S.R. (1991). Review of sea floor spreading around Australia. I. Synthesis of the patterns of spreading. *Australian Journal of Earth Sciences* 38: 373–389.
- Vergara-Silva, F., Espinosa-Matías, S., Ambrose, B.A., Vázquez-Santana, S., Martínez-Mena, A., Márquez-Guzmán, J., Martínez, E., Meyerowitz, E.M., Alvarez-Buylla, E.R. (2003). Inside-out flowers characteristic of *Lacandonia schismatica* evolved at least before its divergence from a closely related taxon, *Triuris brevistylis*. *International Journal of Plant Sciences* 164: 345–357.
- Vinnersten, A., Bremer, K. (2001). Age and biogeography of major clades in Liliaceae. *American Journal of Botany* 88: 1695–1703.

Von Hagen, K.B., Kadereit, J.W. (2002). Phylogeny and flower evolution of the Swertiaeae (Gentianaceae-Gentianeae): Homoplasy and the principle of variable proportions. Systematic Botany 27: 548-572.

Waterman, R.J., Klooster, M.R., Henrich, H., Bidartondo, M.I. (2013). Chapter 7: Species Interactions of Mycoheterotrophic Plants: Specialization and its Potential Consequences. In: V.S.F.T. Merckx (ed.) Mycoheterotrophy. The biology of plants living on fungi, Springer. New York. USA. Pp. 267-296.

White, T.J., Bruns, T.D., Lee, S., Taylor, J.W. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis, M.A., Gelfand, D.H., Sninsky, J.J., White T.P. (eds.) PCR protocols: A guide to methods and applications. Academic Press, Orlando, Florida, USA. Pp. 315-322.

Wicke, S., Schneeweiss, G.M., dePamphilis, C.W., Müller, K.F., Quandt, D. (2011). The evolution of the plastid chromosome in land plants: gene content, gene order, gene function. Plant Molecular Biology 76: 273-297.

Wirz, H. (1910). Beiträge Entwicklungsgeschichte von *Sciaphila* spec. und von *Epirrhizanthes elongata* Bl. Flora 101: 395-446.

Wojciechowski, M.F., Sanderson, M.J., Baldwin, B.G., Donoghue, M.J. (1993). Monophyly of aneuploid *Astragalus* (Fabaceae). Evidence from nuclear ribosomal RNA genes for phylogenetics. American Journal of Botany 80: 711-722.

Woodburne, M.O., Case, J.A. (1996). Dispersal, vicariance, and the late Cretaceous to early Tertiary land mammal biogeography from South America to Australia. Journal of Mammalian Evolution 3: 121-161.

Wyman, S.K., Jansen, R.K., Boore, J.L. (2004). Automatic annotation of organellar genomes with DOGMA. Bioinformatics 20: 3252-3255.

Yamato, M., Ogura-Tsujita, Y., Takahashi, H., Yukawa, T. (2014). Significant difference in mycorrhizal specificity between an autotrophic and its sister mycoheterotrophic plant species of Petrosaviaceae. Journal of Plant Research 127: 685-693.

Yamato, M., Yagame, T., Iwase, K. (2011b). Arbuscular mycorrhizal fungi in roots of non-photosynthetic plants, *Sciaphila japonica* and *Sciaphila tosaensis* (Triuridaceae). Mycoscience 52: 217-233.

Yamato, M., Yagame, T., Shimomura, N., Iwase, K., Takahashi, H., Ogura-Tsujita, Y., Yukawa, T. (2011a). Specific arbuscular mycorrhizal fungi associated with non-photosynthetic *Petrosavia sakuraii* (Petrosaviaceae). Mycorrhiza 21: 631-639.

Yuan, Y.M., Wohlhauser, S., Möller, M., Chassot, P., Mansion, G., Grant, J., Küpfer, P., Klackenberg, J. (2003). Monophyly and relationships of the tribe Exaceae (Gentianaceae) inferred from nuclear ribosomal and chloroplast DNA sequences. Molecular Phylogenetics and Evolution 28: 500-517.

Yuan, Y.M., Wohlhauser, S., Möller, M., Klackenberg, J., Callmander, M.W., Küpfer, P. (2005). Phylogeny and biogeography of *Exacum* (Gentianaceae): A disjunctive distribution in the Indian Ocean basin resulted from long distance dispersal and extensive radiation. Systematic Biology 54: 21-34.

Zhang, D., Saunders, R.M.K., Hu, C-M. (1999). *Corsiopsis chinensis* gen. et sp. nov. (Corsiaceae): first record of the family in Asia. Systematic Botany 24: 311-314.

Zhang, D.X., Saunders, R.M.K. (1999). *Burmannia larseniana* (Burmanniaceae): a new species from Thailand. Nordic Journal of Botany 19: 241-244.

Zhang, D.X., Saunders, R.M.K. (2000). Reproductive biology of a mycoheterotrophic species, *Burmannia wallichii* (Burmanniaceae). Botanical Journal of the Linnean Society 132: 359-367.

*Antiquities of the rainforest*

Zhu, X.Y., Chase, M.W., Qiu, Y.L., Kong, H.Z., Dilcher, D.L., Li, J.H., Chen, Z.D. (2007). Mitochondrial *matR* sequences help to resolve deep phylogenetic relationships in rosids. BMC Evolutionary Biology 7: 217.

Zwickl, D.J. (2006). Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion. PhD Thesis, University of Texas, Austin, Texas, USA.