



Universiteit  
Leiden  
The Netherlands

## Gauge theory and nematic order : the rich landscape of orientational phase transition

Liu, K.

### Citation

Liu, K. (2016, September 6). *Gauge theory and nematic order : the rich landscape of orientational phase transition*. Casimir PhD Series. Retrieved from <https://hdl.handle.net/1887/42793>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/42793>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/42793> holds various files of this Leiden University dissertation.

**Author:** Liu, K.

**Title:** Gauge theory and nematic order : the rich landscape of orientational phase transition

**Issue Date:** 2016-09-06

# Bibliography

- [1] W. Heisenberg, “Mehrkörperproblem und resonanz in der quantenmechanik,” Zeitschrift für Physik **38**, 411–426 (1926).
- [2] L. D. Landau, “On the theory of phase transitions. i.” Zh. Eksp. Teor. Fiz. **11**, 19 (1937).
- [3] L. Landau, “Theory of the superfluidity of helium ii,” Phys. Rev. **60**, 356–358 (1941).
- [4] A. J. Leggett, “A theoretical description of the new phases of liquid  $^3\text{He}$ ,” Rev. Mod. Phys. **47**, 331–414 (1975).
- [5] L. D. Landau and V. Ginzburg, “On the theory of superconductivity,” Zh. Eksp. Teor. Fiz. **20**, 1064 (1950).
- [6] L. P. Gor’kov, “Microscopic derivation of the ginzburg-landau equations in the theory of superconductivity,” Sov. Phys. JETP **9**, 1364–1367 (1959).
- [7] P. W. Anderson, “Plasmons, gauge invariance, and mass,” Phys. Rev. **130**, 439–442 (1963).
- [8] F. Englert and R. Brout, “Broken symmetry and the mass of gauge vector mesons,” Phys. Rev. Lett. **13**, 321–323 (1964).
- [9] P. W. Higgs, “Broken symmetries and the masses of gauge bosons,” Phys. Rev. Lett. **13**, 508–509 (1964).
- [10] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, “Global conservation laws and massless particles,” Phys. Rev. Lett. **13**, 585–587 (1964).
- [11] P. de Gennes and J. Prost, *The Physics of Liquid Crystals*, International Series of Monographs on Physics (Clarendon Press, 1995).
- [12] S. A. Kivelson, E. Fradkin, and V. J. Emery, “Electronic liquid-crystal phases of a doped mott insulator,” Nature **393**, 550–553 (1998).

- [13] E. Fradkin, S. A. Kivelson, M. J. Lawler, J. P. Eisenstein, and A. P. Mackenzie, “Nematic fermi fluids in condensed matter physics,” Annual Review of Condensed Matter Physics **1**, 153–178 (2010).
- [14] J. Zaanen, Z. Nussinov, and S. Mukhin, “Duality in 2+1D quantum elasticity: superconductivity and quantum nematic order,” Ann. Phys. **310**, 181–260 (2004), arXiv:cond-mat/0309397 .
- [15] A. Andreev and I. Grishchuk, “Spin nematics,” Sov. Phys. JETP **60**, 267 (1984).
- [16] D. Podolsky and E. Demler, “Properties and detection of spin nematic order in strongly correlated electron systems,” New Journal of Physics **7**, 59 (2005).
- [17] P. Anderson, *Basic Notions of Condensed Matter Physics*, Basic Notions of Condensed Matter Physics Series (Benjamin/Cummings, 1984).
- [18] D. R. Nelson and J. Toner, “Bond-orientational order, dislocation loops, and melting of solids and smectic-a liquid crystals,” Physical Review B **24**, 363 (1981).
- [19] N. D. Mermin, “The topological theory of defects in ordered media,” Rev. Mod. Phys. **51**, 591–648 (1979).
- [20] L. Michel, “Symmetry defects and broken symmetry. configurations hidden symmetry,” Rev. Mod. Phys. **52**, 617–651 (1980).
- [21] O. Lehmann, “Über fliessende krystalle,” Z. phys. Chem **4**, 462–472 (1889).
- [22] G. Friedel, “The mesomorphic states of matter,” Annales de Physique **18**, 273 (1922).
- [23] K. Liu, J. Nissinen, R.-J. Slager, K. Wu, and J. Zaanen, “Generalized liquid crystals: giant fluctuations and the vestigial chiral order of I, O and T matter,” arXiv preprint arXiv:1512.07822 (2015).
- [24] C. N. Yang and R. L. Mills, “Conservation of isotopic spin and isotopic gauge invariance,” Phys. Rev. **96**, 191–195 (1954).

- [25] P. E. Lammert, D. S. Rokhsar, and J. Toner, “Topology and nematic ordering,” Phys. Rev. Lett. **70**, 1650–1653 (1993).
- [26] P. E. Lammert, D. S. Rokhsar, and J. Toner, “Topology and nematic ordering. i. a gauge theory,” Phys. Rev. E **52**, 1778–1800 (1995).
- [27] E. Fradkin and S. H. Shenker, “Phase diagrams of lattice gauge theories with higgs fields,” Phys. Rev. D **19**, 3682–3697 (1979).
- [28] S. Elitzur, “Impossibility of spontaneously breaking local symmetries,” Phys. Rev. D **12**, 3978–3982 (1975).
- [29] K. Liu, J. Nissinen, R.-J. Slager, K. Wu, and J. Zaanen, “Classification of point-group-symmetric orientational ordering tensors,” arXiv preprint arXiv:1603.04794 (2016).
- [30] K. Liu, J. Nissinen, Z. Nussinov, R.-J. Slager, K. Wu, and J. Zaanen, “Classification of nematic order in  $2+1$  dimensions: Dislocation melting and  $o(2)/Z_N$  lattice gauge theory,” Phys. Rev. B **91**, 075103 (2015).
- [31] A. J. Beekman, J. Nissinen, K. Wu, K. Liu, R.-J. Slager, Z. Nussinov, V. Cvetkovic, and J. Zaanen, “Dual gauge field theory of quantum liquid crystals in two dimensions,” arXiv preprint arXiv:1603.04254 (2016).
- [32] K. Liu, J. Nissinen, J. de Boer, R.-J. Slager, and J. Zaanen, “Hierarchy of orientational phases and axial anisotropies in the gauge theoretical description of generalized nematics,” arXiv preprint arXiv:1606.04507 (2016).
- [33] H. Kleinert, *Gauge Fields in Condensed Matter, Vol.II Stress and Defects* (World Scientific, Singapore, 1989).
- [34] M. Kleman and J. Friedel, “Disclinations, dislocations, and continuous defects: A reappraisal,” Rev. Mod. Phys. **80**, 61–115 (2008).
- [35] J. M. Burgers, *Some considerations on the fields of stress connected with dislocations in a regular crystal lattice. I* (Royal Dutch Academy of Sciences, 1939).
- [36] F. C. Frank, “I. liquid crystals. on the theory of liquid crystals,” Discuss. Faraday Soc. **25**, 19–28 (1958).

- [37] B. Halperin and D. Nelson, “Theory of two-dimensional melting,” Phys. Rev. Lett. **41**, 121–124 (1978).
- [38] H. Kleinert and J. Zaanen, “Nematic world crystal model of gravity explaining absence of torsion in spacetime,” Phys. Lett. A **324**, 361–365 (2004), arXiv:cond-mat/0309379 .
- [39] V. Cvetkovic, Z. Nussinov, and J. Zaanen, “Topological kinematic constraints: dislocations and the glide principle,” Philosophical Magazine **86**, 2995–3020 (2006).
- [40] P. Ehrenfest, “Bemerkung über die angenäherte Gültigkeit der klassischen Mechanik innerhalb der Quantenmechanik,” Zeit. Phys. **45**, 455–457 (1927).
- [41] M. Fisher, P. Weichman, G. Grinstein, and D. Fisher, “Boson localization and the superfluid–insulator transition,” Phys. Rev. B **40**, 546–570 (1989).
- [42] M. Fisher and D. Lee, “Correspondence between two-dimensional bosons and a bulk superconductor in a magnetic field,” Phys. Rev. B **39**, 2756–2759 (1989).
- [43] V. Cvetkovic, *Quantum Liquid Crystals*, Ph.D. thesis, Leiden University (2006), <http://hdl.handle.net/1887/4456> .
- [44] J. Goldstone, A. Salam, and S. Weinberg, “Broken symmetries,” Physical Review **127**, 965–970 (1962).
- [45] A. J. Beekman, K. Wu, V. Cvetkovic, and J. Zaanen, “Deconfining the rotational Goldstone mode: The superconducting quantum liquid crystal in (2+1) dimensions,” Phys. Rev. B **88**, 024121 (2013).
- [46] J.-M. Park and T. C. Lubensky, “Topological defects on fluctuating surfaces: General properties and the kosterlitz-thouless transition,” Phys. Rev. E **53**, 2648–2664 (1996).
- [47] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, Princeton, NJ, 1992).
- [48] X. Wen, *Quantum Field Theory of Many-Body Systems: From the Origin of Sound to an Origin of Light and Electrons*, Oxford Graduate Texts (OUP Oxford, 2004).

- [49] F. J. Wegner, “Duality in generalized ising models and phase transitions without local order parameters,” Journal of Mathematical Physics **12**, 2259–2272 (1971).
- [50] J. B. Kogut, “An introduction to lattice gauge theory and spin systems,” Rev. Mod. Phys. **51**, 659–713 (1979).
- [51] D. Horn, M. Weinstein, and S. Yankielowicz, “Hamiltonian approach to  $Z_N$  lattice gauge theories,” Phys. Rev. D **19**, 3715–3731 (1979).
- [52] G. Bhanot and M. Creutz, “Phase diagram of  $z(n)$  and  $u(1)$  gauge theories in three dimensions,” Phys. Rev. D **21**, 2892–2902 (1980).
- [53] O. Borisenko, V. Chelnokov, G. Cortese, M. Gravina, A. Papa, and I. Surzhikov, “Critical behavior of 3d lattice gauge theories at zero temperature,” Nuclear Physics B **879**, 80 – 97 (2014).
- [54] K. Gregor, D. A. Huse, R. Moessner, and S. L. Sondhi, “Diagnosing deconfinement and topological order,” New Journal of Physics **13**, 025009 (2011).
- [55] W. Janke and H. Kleinert, “How good is the villain approximation?” Nuclear Physics B **270**, 135 – 153 (1986).
- [56] J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, “Renormalization, vortices, and symmetry-breaking perturbations in the two-dimensional planar model,” Phys. Rev. B **16**, 1217–1241 (1977).
- [57] Villain, J., “Theory of one- and two-dimensional magnets with an easy magnetization plane. ii. the planar, classical, two-dimensional magnet,” J. Phys. France **36**, 581–590 (1975).
- [58] Z. Nussinov and J. Zaanen, “Stripe fractionalization i: The generation of ising local symmetry,” J. Phys. IV France **12**, 245–250 (2002).
- [59] J. Zaanen and Z. Nussinov, “Stripe fractionalization: the quantum spin nematic and the abrikosov lattice,” physica status solidi (b) **236**, 332–339 (2003).

- [60] T. Senthil and M. P. A. Fisher, “ $Z_2$  gauge theory of electron fractionalization in strongly correlated systems,” Phys. Rev. B **62**, 7850–7881 (2000).
- [61] T. Senthil and M. P. A. Fisher, “Fractionalization, topological order, and cuprate superconductivity,” Phys. Rev. B **63**, 134521 (2001).
- [62] T. Grover and T. Senthil, “Quantum phase transition from an antiferromagnet to a spin liquid in a metal,” Phys. Rev. B **81**, 205102 (2010).
- [63] S. V. Isakov, R. G. Melko, and M. B. Hastings, “Universal signatures of fractionalized quantum critical points,” Science **335**, 193–195 (2012).
- [64] L. Onsager, “The effects of shape on the interaction of colloidal particles,” Annals of the New York Academy of Sciences **51**, 627–659 (1949).
- [65] W. Maier and A. Saupe, “Eine einfache molekular-statistische theorie der nematischen kristallinflüssigen phase. teil 11.” Zeitschrift für Naturforschung A **14**, 882–889 (1959).
- [66] P. A. Lebwohl and G. Lasher, “Nematic-liquid-crystal order?a monte carlo calculation,” Physical Review A **6**, 426 (1972).
- [67] L. Yu and A. Saupe, “Observation of a biaxial nematic phase in potassium laurate-1-decanol-water mixtures,” Physical Review Letters **45**, 1000 (1980).
- [68] K. Severing and K. Saalwächter, “Biaxial nematic phase in a thermotropic liquid-crystalline side-chain polymer,” Physical review letters **92**, 125501 (2004).
- [69] B. R. Acharya, A. Primak, and S. Kumar, “Biaxial nematic phase in bent-core thermotropic mesogens,” Phys. Rev. Lett. **92**, 145506 (2004).
- [70] L. A. Madsen, T. J. Dingemans, M. Nakata, and E. T. Samulski, “Thermotropic biaxial nematic liquid crystals,” Physical review letters **92**, 145505 (2004).

- [71] G. R. Luckhurst and T. J. Sluckin, eds., *Biaxial Nematic Liquid Crystals: Theory, Simulation and Experiment* (John Wiley & Sons, 2015).
- [72] M. J. Freiser, “Ordered states of a nematic liquid,” *Phys. Rev. Lett.* **24**, 1041–1043 (1970).
- [73] R. Alben, “Phase transitions in a fluid of biaxial particles,” *Physical Review Letters* **30**, 778 (1973).
- [74] J. P. Straley, “Ordered phases of a liquid of biaxial particles,” *Physical Review A* **10**, 1881 (1974).
- [75] F. Biscarini, C. Chiccoli, P. Pasini, F. Semeria, and C. Zannoni, “Phase diagram and orientational order in a biaxial lattice model: A monte carlo study,” *Phys. Rev. Lett.* **75**, 1803–1806 (1995).
- [76] E. Matijevic, “Monodispersed metal (hydrous) oxides - a fascinating field of colloid science,” *Accounts of Chemical Research* **14**, 22–29 (1981).
- [77] F. Li, D. P. Josephson, and A. Stein, “Colloidal assembly: The road from particles to colloidal molecules and crystals,” *Angewandte Chemie International Edition* **50**, 360–388 (2011).
- [78] A. G. Mark, J. G. Gibbs, T.-C. Lee, and P. Fischer, “Hybrid nanocolloids with programmed three-dimensional shape and material composition,” *Nat Mater* **12**, 802–807 (2013).
- [79] M. Huang, C.-H. Hsu, J. Wang, S. Mei, X. Dong, Y. Li, M. Li, H. Liu, W. Zhang, T. Aida, W.-B. Zhang, K. Yue, and S. Z. D. Cheng, “Selective assemblies of giant tetrahedra via precisely controlled positional interactions,” *Science* **348**, 424–428 (2015).
- [80] S. C. Glotzer and M. J. Solomon, “Anisotropy of building blocks and their assembly into complex structures,” *Nature materials* **6**, 557–562 (2007).
- [81] P. F. Damasceno, M. Engel, and S. C. Glotzer, “Predictive self-assembly of polyhedra into complex structures,” *Science* **337**, 453–457 (2012).

- [82] G. van Anders, D. Klotsa, N. K. Ahmed, M. Engel, and S. C. Glotzer, “Understanding shape entropy through local dense packing,” *Proceedings of the National Academy of Sciences* **111**, E4812–E4821 (2014).
- [83] V. N. Manoharan, “Colloidal matter: Packing, geometry, and entropy,” *Science* **349** (2015), 10.1126/science.1253751.
- [84] L. E. Hough, M. Spannuth, M. Nakata, D. A. Coleman, C. D. Jones, G. Dantlgraber, C. Tschierske, J. Watanabe, E. Kiščíkova, D. M. Walba, J. E. MacLennan, M. A. Glaser, and N. A. Clark, “Chiral isotropic liquids from achiral molecules,” *Science* **325**, 452–456 (2009).
- [85] C. Dressel, T. Reppe, M. Prehm, M. Brautzsch, and C. Tschierske, “Chiral self-sorting and amplification in isotropic liquids of achiral molecules,” *Nature chemistry* **6**, 971–977 (2014).
- [86] D. W. Allender, M. A. Lee, and N. Hafiz, “Landau theory of biaxial and uniaxial nematic liquid crystals,” *Molecular Crystals and Liquid Crystals* **124**, 45–52 (1985).
- [87] E. F. Gramsbergen, L. Longa, and W. H. de Jeu, “Landau theory of the nematic-isotropic phase transition,” *Physics Reports* **135**, 195–257 (1986).
- [88] M. V. Jarić, “Landau theory of long-range orientational order,” *Nuclear Physics B* **265**, 647–670 (1986).
- [89] L. G. Fel, “Tetrahedral symmetry in nematic liquid crystals,” *Phys. Rev. E* **52**, 702–717 (1995).
- [90] L. Radzhovsky and T. C. Lubensky, “Fluctuation-driven 1st-order isotropic-to-tetrahedratic phase transition,” *EPL (Europhysics Letters)* **54**, 206 (2001).
- [91] T. C. Lubensky and L. Radzhovsky, “Theory of bent-core liquid-crystal phases and phase transitions,” *Phys. Rev. E* **66**, 031704 (2002).
- [92] B. Mettout, “Macroscopic and molecular symmetries of unconventional nematic phases,” *Phys. Rev. E* **74**, 041701 (2006).

- [93] D. Allender and L. Longa, “Landau–de gennes theory of biaxial nematics reexamined,” *Physical Review E* **78**, 011704 (2008).
- [94] M. Jarić, L. Michel, and R. Sharp, “Zeros of covariant vector fields for the point groups: invariant formulation,” *Journal de Physique* **45**, 1–27 (1984).
- [95] L. Michel and B. Zhilinskii, “Symmetry, invariants, topology. basic tools,” *Physics Reports* **341**, 11–84 (2001).
- [96] S. Romano, “Computer simulation study of a simple cubatic mesogenic lattice model,” *Physical Review E* **74**, 011704 (2006).
- [97] S. Romano, “Computer simulation study of a simple tetrahedratic mesogenic lattice model,” *Physical Review E* **77**, 021704 (2008).
- [98] K. Merkel, A. Kocot, J. Vij, R. Korlacki, G. Mehl, and T. Meyer, “Thermotropic biaxial nematic phase in liquid crystalline organo-siloxane tetrapodes,” *Physical review letters* **93**, 237801 (2004).
- [99] C. Tschierske and D. J. Photinos, “Biaxial nematic phases,” *J. Mater. Chem.* **20**, 4263–4294 (2010).
- [100] J. Toner, “Ordered phases that scatter light strongly,” *Phys. Rev. A.* **27**, 1157 (1983).
- [101] J. R. Podgornik, H. H. Strey, K. Gawrisch, D. C. Rau, A. Rupprecht, and V. A. Parsegian, “Bond orientational order, molecular motion, and free energy of high-density dna mesophasesy,” *PNAS* **93**, 4261 (1996).
- [102] H. Takezoe and Y. Takanishi, “Bent-core liquid crystals: Their mysterious and attractive world,” *Japanese Journal of Applied Physics* **45**, 597 (2006).
- [103] N. W. Ashcroft and N. D. Mermin, *Solid State physics* (Saunders, Philadelphia, 1976).
- [104] J. Friedel, *Dislocations* (Pergamon, 1964).
- [105] L. Longa, G. Pajak, and T. Wydro, “Chiral symmetry breaking in bent-core liquid crystals,” *Physical Review E* **79**, 040701 (2009).

- [106] P. R. Gerber and M. E. Fisher, “Critical temperatures of classical n-vector models on hypercubic lattices,” *Physical Review B* **10**, 4697 (1974).
- [107] R. M. Fernandes, A. V. Chubukov, and J. Schmalian, “What drives nematic order in iron-based superconductors?” *Nat. Phys.* **10**, 97–104 (2014).
- [108] Y. Kamiya, N. Kawashima, and C. Batista, “Dimensional crossover in the quasi-two-dimensional ising-o (3) model,” *Physical Review B* **84**, 214429 (2011).
- [109] D. W. Johnson and K. N. Raymond, “The self-assembly of a  $[\text{Ga}_4\text{L}_6]^{12-}$  tetrahedral cluster thermodynamically driven by host-guest interactions.” *Inorg. Chem.* **40**, 5157–5161 (2001).
- [110] E. K. Honarmand, E. Bill, P. L. Hagedoorn, and W. R. Hagen, “The catalytic center of ferritin regulates iron storage via fe(ii)-fe(iii) displacement.” *Nat. Chem. Biol.* **8**, 941–948 (2012).
- [111] R. Zandi, D. Reguera, R. F. Bruinsma, W. M. Gelbart, and J. Rudnick, “Origin of icosahedral symmetry in viruses.” *PNAS* **101**, 15556–15560 (2004).
- [112] M. Kumar and D. Blaas, “Human rhinovirus subviral a particle binds to lipid membranes over a twofold axis of icosahedral symmetry,” *J. Virol* **87**, 11309–11312 (2013).
- [113] A. Kitaev, “Anyons in an exactly solved model and beyond,” *Annals of Physics* **321**, 2 – 111 (2006), january Special Issue.
- [114] X. G. Wen, “Mean-field theory of spin-liquid states with finite energy gap and topological orders,” *Phys. Rev. B* **44**, 2664–2672 (1991).
- [115] R. S. Andrist, H. G. Katzgraber, H. Bombin, and M. A. Martin-Delgado, “Tricolored lattice gauge theory with randomness: fault tolerance in topological color codes,” *New Journal of Physics* **13**, 083006 (2011).
- [116] C. Xu and A. W. W. Ludwig, “Topological quantum liquids with quaternion non-abelian statistics,” *Phys. Rev. Lett.* **108**, 047202 (2012).

- [117] T. Frankel, *The geometry of physics: an introduction* (Cambridge University Press, 2011).
- [118] P. Chaikin and T. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, 2000).
- [119] E. Fradkin, S. A. Kivelson, M. J. Lawler, J. P. Eisenstein, and A. P. Mackenzie, “Nematic Fermi fluids in condensed matter physics,” *Ann. Rev. Cond. Mat. Phys.* **1**, 153–178 (2010).
- [120] R. Rosso, “Orientational order parameters in biaxial nematics: Polymeric notation,” *Liquid Crystals* **34**, 737–748 (2007).
- [121] P. J. Steinhardt, D. R. Nelson, and M. Ronchetti, “Icosahedral bond orientational order in supercooled liquids,” *Phys. Rev. Lett.* **47**, 1297–1300 (1981).
- [122] P. J. Steinhardt, D. R. Nelson, and M. Ronchetti, “Bond-orientational order in liquids and glasses,” *Phys. Rev. B* **28**, 784–805 (1983).
- [123] M. V. Jarić, “Long-range icosahedral orientational order and quasicrystals,” *Phys. Rev. Lett.* **55**, 607–610 (1985).
- [124] A. Haji-Akbari and S. C. Glotzer, “Strong orientational coordinates and orientational order parameters for symmetric objects,” *Journal of Physics A: Mathematical and Theoretical* **48**, 485201 (2015).
- [125] Y. Sun and Y. Xia, “Shape-controlled synthesis of gold and silver nanoparticles,” *Science* **298**, 2176–2179 (2002).
- [126] M. H. Huang and P.-H. Lin, “Shape-controlled synthesis of polyhedral nanocrystals and their facet-dependent properties,” *Advanced Functional Materials* **22**, 14–24 (2012).
- [127] L. Messio, C. Lhuillier, and G. Misguich, “Lattice symmetries and regular magnetic orders in classical frustrated antiferromagnets,” *Phys. Rev. B* **83**, 184401 (2011).
- [128] T. Grover and T. Senthil, “Non-abelian spin liquid in a spin-one quantum magnet,” *Phys. Rev. Lett.* **107**, 077203 (2011).

- [129] R. Barnett, A. Turner, and E. Demler, “Classifying novel phases of spinor atoms,” Phys. Rev. Lett. **97**, 180412 (2006).
- [130] T. Zibold, V. Corre, C. Frapolli, A. Invernizzi, J. Dalibard, and F. Gerbier, “Spin-nematic order in antiferromagnetic spinor condensates,” Phys. Rev. A **93**, 023614 (2016).
- [131] S. Sternberg, *Group theory and physics* (Cambridge University Press, 1995).
- [132] D. Litvin, “The icosahedral point groups,” Acta Crystallographica Section A: Foundations of Crystallography **47**, 70–73 (1991).
- [133] X. Zheng and P. Palffy-Muhoray, “Eigenvalue decomposition for tensors of arbitrary rank,” electronic-Liquid Crystal Communications (2007).
- [134] S. S. Turzi and F. Bisi, “Determination of the symmetry classes of orientational ordering tensors,” arXiv preprint arXiv:1602.06413 (2016).
- [135] D. R. J. Chillingworth, R. Lauterbach, and S. S. Turzi, “Molien series and low-degree invariants for a natural action of  $\text{SO}(3) \wr \mathbb{Z}_2$ ,” Journal of Physics A: Mathematical and Theoretical **48**, 015203 (2015).
- [136] S. S. Turzi, “On the cartesian definition of orientational order parameters,” Journal of Mathematical Physics **52**, 053517 (2011).
- [137] J. Nye, *Physical Properties of Crystals: Their Representation by Tensors and Matrices*, Oxford science publications (Clarendon Press, 1985).
- [138] S. Romano and G. De Matteis, “Orientationally ordered phase produced by fully antinematic interactions: A simulation study,” Physical Review E **84**, 011703 (2011).
- [139] F. Bisi, G. De Matteis, and S. Romano, “Calamitic and antineamic orientational order produced by the generalized straley lattice model,” Physical Review E **88**, 032502 (2013).

- [140] A. M. Sonnet, E. G. Virga, and G. E. Durand, “Dielectric shape dispersion and biaxial transitions in nematic liquid crystals,” *Phys. Rev. E* **67**, 061701 (2003).
- [141] G. De Matteis and E. G. Virga, “Tricritical points in biaxial liquid crystal phases,” *Physical Review E* **71**, 061703 (2005).
- [142] G. De Matteis, S. Romano, and E. G. Virga, “Bifurcation analysis and computer simulation of biaxial liquid crystals,” *Physical Review E* **72**, 041706 (2005).
- [143] G. Volovik and V. Mineev, “Investigation of singularities in superfluid he3 in liquid crystals by the homotopic topology methods,” *Zh. Eksp. Teor. Fiz.* **72**, 22562274 (1977).
- [144] D. F. Mross and T. Senthil, “Stripe melting and quantum criticality in correlated metals,” *Phys. Rev. B* **86**, 115138 (2012).
- [145] J. Toner, P. E. Lammert, and D. S. Rokhsar, “Topology and nematic ordering. ii. observable critical behavior,” *Phys. Rev. E* **52**, 1801–1810 (1995).

