



Universiteit
Leiden
The Netherlands

The dependence of the A_V prior for SN Ia on host mass and disc inclination

Holwerda, B.W.; Keel, W.C.; Kenworthy, M.A.; Mack, K.J.

Citation

Holwerda, B. W., Keel, W. C., Kenworthy, M. A., & Mack, K. J. (2015). The dependence of the A_V prior for SN Ia on host mass and disc inclination. *Monthly Notices Of The Royal Astronomical Society*, 451(3), 2390-2398. doi:10.1093/mnras/stv1125

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/48987>

Note: To cite this publication please use the final published version (if applicable).

The dependence of the A_V prior for SN Ia on host mass and disc inclination

B. W. Holwerda,¹★ W. C. Keel,² M. A. Kenworthy¹ and K. J. Mack^{3,4,5}

¹University of Leiden, Sterrenwacht Leiden, Niels Bohrweg 2, NL-2333 CA Leiden, the Netherlands

²Department of Physics and Astronomy, University of Alabama, Box 870324, Tuscaloosa, AL 35487, USA

³School of Physics (David Caro Building), University of Melbourne, VIC 3010, Australia

⁴ARC Centre of Excellence for All-sky Astrophysics (CAASTRO), The University of Sydney, NSW 2006, Australia

⁵ARC Centre of Excellence for Particle Physics at Terascale (CoEPP), School of Physics, The University of Melbourne, VIC 3010, Australia

Accepted 2015 May 12. Received 2015 April 15; in original form 2014 September 23

ABSTRACT

Type Ia supernovae (SNe Ia) are used as ‘standard candles’ for cosmological distance scales. To fit their light-curve shape–absolute luminosity relation, one needs to assume an intrinsic colour and a likelihood of host galaxy extinction or a convolution of these, a colour distribution prior. The host galaxy extinction prior is typically assumed to be an exponential drop-off for the current supernova programmes ($P(A_V) \propto e^{-A_V/\tau_0}$). We explore the validity of this prior using the distribution of extinction values inferred when two galaxies accidentally overlap (an occulting galaxy pair). We correct the supernova luminosity distances from the SDSS-III supernova projects (SDSS-SN) by matching the host galaxies to one of three templates from occulting galaxy pairs based on the host galaxy mass and the A_V -bias–prior-scale (τ_0) relation from Jha et al. We find that introducing an A_V prior that depends on host mass results in lowered luminosity distances for the SDSS-SN on average but it does not reduce the scatter in individual measurements. This points, in our view, to the need for many more occulting galaxy templates to match to SN Ia host galaxies to rule out this possible source of scatter in the SN Ia distance measurements. We match occulting galaxy templates based on both mass and projected radius and we find that one should match by stellar mass first with radius as a secondary consideration. We discuss the caveats of the current approach: the lack of enough radial coverage, the small sample of priors (occulting pairs with *HST* data), the effect of gravitationally interacting as well as occulting pairs, and whether an exponential distribution is appropriate. Our aim is to convince the reader that a library of occulting galaxy pairs observed with *HST* will provide sufficient priors to improve (optical) SN Ia measurements to the next required accuracy in cosmology.

Key words: dust, extinction – galaxies: general – galaxies: ISM – galaxies: structure – distance scale.

1 INTRODUCTION

Type Ia supernovae (SNe Ia) are a prime cosmological tool (Riess et al. 1998; Perlmutter et al. 1999). Their relation between light curve and intrinsic luminosity makes them excellent standard candles. Part of the exquisite precision required is a good understanding of the expected dust extinction and known reddening-dimming relation, the extinction law (Cardelli, Clayton & Mathis 1989; Calzetti, Kinney & Storchi-Bergmann 1994). With the dark energy detection result in hand, the supernova community now looks for the next step in accuracy (e.g. the 3 per cent solution; Riess et al. 2011). A much better model for the dust extinction in the supernova’s host galaxy

will be critical. Host galaxy dust is already identified by the Dark Energy Task Force as a principal unknown (Albrecht et al. 2006) and this remains the case, together with the photometric uncertainty across surveys (e.g. Conley et al. 2011; Betoule et al. 2014; Scolnic et al. 2014).

These issues are being addressed by the current generation of supernovae searches, e.g. the Supernova Legacy Survey (Astier et al. 2006), the Equation of State: SuperNovae trace Cosmic Expansion (Wood-Vasey et al. 2007), and the Sloan Digital Sky Survey SuperNova survey (SDSS-SN; Kessler et al. 2009; Sako et al. 2014). The supernovae community has been systematically exploring what could affect the observed relation between peak luminosity and light-curve width (Phillips 1993) for SNe Ia.

A prime candidate is the correlation with properties of the galaxy hosting the SN Ia (Gallagher et al. 2005; Mannucci, Della Valle &

★ E-mail: benne.holwerda@gmail.com

Panagia 2006; Sullivan et al. 2006; Kistler et al. 2013; Pan et al. 2014). Two populations of SNe Ia appear associated with two populations of stars: blue, star-forming galaxies host higher rates of fast declining supernova and red, passive galaxies host predominantly more slowly-declining SNe Ia (Hamuy et al. 1996; Howell 2001; van den Bergh, Li & Filippenko 2005; Mannucci, Della Valle & Panagia 2006; Schawinski 2009; Lampeitl et al. 2010; Wang et al. 2013). The difference in light-curve characteristics may be completely attributable to the different progenitor stellar populations.

However, simultaneously, the issue of host galaxy extinction has come to the fore, either the applicability of the appropriate extinction law (Riess, Press & Kirshner 1996; Phillips et al. 1999; Altavilla et al. 2004; Reindl et al. 2005; Conley et al. 2007; Jha, Riess & Kirshner 2007), the validity of the prior of extinction values (Jha et al. 2007; Wood-Vasey et al. 2007; Gupta et al. 2011), or the possibility that the geometry of the dusty interstellar medium (ISM) in (and hence the extinction distribution) may change with galaxy evolution (Holwerda 2008; Holwerda et al. 2015a). Host galaxies appear to be indistinguishable from normal field galaxies (Childress et al. 2013). After photometric calibration, this host galaxy extinction issue is likely to return as a dominant issue standing in the way of the community’s final goal of 1 per cent precision for SN I (e.g. Riess et al. 2011; Kelly et al. 2015).

There is now an opportunity to probe the dependences of the extinction prior through two projects. First, the SDSS-SN project has generated a uniform sample of SNe Ia and their host galaxy properties of exquisite quality. Secondly, occulting galaxy pairs promise to map extinction in spiral discs using the *Hubble Space Telescope* (*HST*) imaging, enabling a way to generate an extinction prior for SN Ia fit. The first of these extinction maps are now available.

In this paper, we explore the effects of applying the extinction priors from occulting galaxy pairs to the SDSS-SN sample and gauge the effects. We opt for the SDSS-SN sample among all the supernova samples available now as it has uniform properties of not only the supernovae, as measured with both light-curve packages, but also on the host galaxies. The latter is important if one wants to correct for the effects of host galaxy properties. The SDSS-SN project uses two light-curve fitting packages, *MLCS2k2* and *SALT2*. We use the former’s output as it uses an explicit prior for host galaxy extinction, making a direct comparison with the occulting galaxy technique practical, even though *SALT2* and its derivative *BASALT* are now often the more preferred package (e.g. Guy et al. 2010; Betoule et al. 2014; Rest et al. 2014). We should note here that the A_V value used in *MLCS2k2* is not the exact same quantity as measured from galaxy transparency measurements, i.e. occulting galaxy measurements. *MLCS2k2*’s A_V parameter is the difference in supernova colour from an assumed intrinsic one, translated to an extinction via an assumed reddening law (in the case of the SDSS-SN; Kessler et al. 2009, a greyer one than Milky Way). The reddening law in this instant is an implicit input setting, one which has been tweaked to minimize cosmological fit residuals (e.g. Hicken et al. 2009).

We assume in this paper that the shape of the A_V fit parameter’s prior and the shape of the distribution of A_V values observed in occulting galaxies is to first order the same.

We explore the extinction effects by matching SN Ia host galaxies to those occulting galaxies with *HST* data through their stellar mass. In a previous paper on SN Ia (Holwerda et al. 2015a), we explored the relation between the extinction distribution and inclination and found that an inclination correction using a simple $\cos(i)$ is sufficient to make the A_V distribution found by the SDSS-SN identical in high and low-inclination subsamples. We therefore adopt this inclination




Figure 1. The 2MASX occulting galaxy pair at $z=0.06$. The extinction in the overlap region (black aperture) can be estimated from the complementary apertures: the foreground spiral (F, the green aperture) and the background galaxy (B, red aperture).

correction in this paper as well. We will discuss the caveats and limits of the current approach and sketch out how to proceed from here in the generation of improved extinction priors, tailored to the host galaxy of the observed supernova.

2 OCCULTING GALAXIES

One can accurately measure the extinction of light by interstellar dust using a *known* light source. The most accurate extragalactic method is to use a partially occulting galaxy pair (Fig. 1). Assuming both galaxies are symmetric, one can estimate flux contributions to the overlap region from the complementary regions at the same radius in each galaxy. The reliance on differential photometry enables an extinction measurement (A_V) in a single-colour image without relying on any assumed extinction law. With multiple filters, the effective extinction law itself can be measured (Keel & White 2001a; Holwerda et al. 2009; Keel et al. 2014).

2.1 The occulting galaxies method

Fig. 1 shows an example pair, serendipitously imaged with *HST* for the ANGST survey (Dalcanton et al. 2009), near NGC 253 (2MASX). To determine the extinction in the overlap region (black aperture), one estimates the relative flux contributions by both galaxies from the complementary apertures in the foreground (green aperture) and background galaxy (red aperture in Fig. 1), respectively, assuming rotational symmetry (i.e. the image can be rotated and be self-similar). Thus, we estimate the flux contributions to the overlap region from the same image. In the case of ground-based images, each of these apertures would yield single A_V value, averaged over a large portion of the foreground disc. In the case of an *HST* image, it results in a highly accurate map of the dust extinction in the overlap region with hundreds of independent lines of sight. The map is constructed pixel by pixel; for each pixel in the overlap aperture, the corresponding pixel in the background galaxy aperture (red) is found, i.e. the pixel at the same distance from the background galaxy centre but in the opposite direction, as well as the corresponding pixel in the foreground galaxy (same distance with respect to the foreground galaxy centre). We estimate the optical depth ($\bar{\tau}$) or extinction (A_V) from the flux in the overlap ($F + Be^{-\tau}$), a mix of flux from background and foreground galaxies, the flux in the corresponding background pixel (\bar{B}) and foreground pixel (\bar{F} , same colours as Fig. 1):

$$A_V = 1.086 \times \bar{\tau} = -1.086 \times \ln \left(\frac{(F + Be^{-\tau}) - \bar{F}}{\bar{B}} \right). \quad (1)$$




Figure 2. The shape of the A_V distribution of the three occulting galaxies with enough overlap to generate a distribution. These distributions have not yet been corrected for their foreground disc’s inclination. Dashed lines are exponential fits through the distributions. The exponential drop-off can then be used as an SN Ia prior. The derived drop-offs are listed in Fig. 5, with their τ_0 values listed in Table 1.

Thus, a map of dust extinction and a distribution function ($P(A_V)$) such as Fig. 2 can be constructed from an *HST* image. For each type of foreground galaxy, inclination, and radial distance, an appropriate $P(A_V)$ can thus be observed.

2.2 Occulting pairs

The accuracy in individual pairs is limited by the symmetry of both galaxies and image quality. The uncertainty from the assumption of galaxy symmetry can be mitigated by using elliptical galaxies as the background galaxy and by combining the results for several pairs. In the case of *HST* images (e.g. Fig. 1), a single pair contains enough independent lines of sight through the foreground galaxy to construct a $P(A_V)$ such as Fig. 2.

The occulting galaxies technique however has increased steadily in accuracy and usefulness, owing in a large part to the increasing sample sizes. Estimating dust extinction and mass from differential photometry in occulting pairs of galaxies was first proposed by White & Keel (1992). Their technique was then applied to all known pairs using ground-based optical images (Andredakis & van der Kruit 1992; Berlind et al. 1997; Domingue et al. 1999; White, Keel & Conselice 2000) and spectroscopy (Domingue, Keel & White 2000), and later space-based *HST* images (Elmegreen et al. 2001; Keel & White 2001a,b; Holwerda et al. 2009, 2013a). These initial results were limited by sample sizes. More recently, more pairs were found in the SDSS spectroscopic catalogue (86 pairs in Holwerda, Keel & Bolton 2007), and in an ongoing effort in the GalaxyZOO project (Lintott et al. 2008). The current count stands at 1993 pairs (Keel et al. 2013) providing opportunities for follow-up with IFU observations (Holwerda & Keel 2013; Holwerda et al. 2013b) and *GALEX* (Keel et al. 2014).

Highly complete spectroscopic surveys, such the Galaxy and Mass Assembly (GAMA; Driver et al. 2009, 2011; Baldry et al. 2010) are adding many more overlapping pairs through the identification of blended spectra or close on-sky pairs (Holwerda et al. 2015b). Observational results include mean extinction profiles (Domingue et al. 2000; White et al. 2000; Holwerda et al. 2007), an indication that the dust may be fractal (Keel & White 2001a)

Table 1. The properties of the foreground occulting galaxy in three pairs with sufficient *HST* information to form an extinction map. Stellar mass in solar masses and radii in effective (half-light) radii.

Pair	Type	Stellar mass M^*	Radial coverage (R_e)	(τ_0)
UGC 3995	Sa	1.7×10^{10}	0–2	0.22
2MASX	Sd	3.1×10^9	2–5	0.85
AM 0500	Sbc	2.6×10^{10}	~ 3.7	0.67

and that the colour–extinction relation is grey in the ground-based observations (Holwerda et al. 2007). The Galactic extinction law returns as soon as the physical sampling of the overlap region resolves the molecular clouds in the foreground disc (<100 pc; Elmegreen et al. 2001; Keel & White 2001a,b; Holwerda et al. 2009).

The extinction maps of foreground discs made with *HST* are especially useful as a probability map of extinction towards a single object of interest. To date, we have obtained *HST* imaging of three overlapping pairs with a large enough overlap region to generate a reasonable distribution. The normalized distributions of these three are shown in Fig. 2. Radial coverage is very similar to the radii at which SNe Ia are found (Table 1): between 0 and 6 effective radii (converted using the prescriptions in Graham & Driver 2005).

Our previous results on occulting galaxy pairs are consistent with an exponential decline with radius of the average disc attenuation, not dissimilar to the exponential profile of the stellar light (e.g. Domingue et al. 1999; Holwerda 2005; Holwerda et al. 2005a,b). We stress here that we deal here now in the exponential distribution of values in a certain range of radii but this is not the same as the radial exponential decline. For example, the lower values ($A_V \sim 0.1$) are typically, but not exclusively, found at higher radius while the higher values are more typical for spiral arms.

2.3 UGC 3995

This pair of galaxies has been long known to be overlapping and interacting (e.g. Marziani et al. 1999) and we have analysed the extinction properties of both CALIFA DR1 IFU data and archival WPC2 *F606W* imaging in Holwerda & Keel (2013). Radial coverage is from 5–15 arcsec, covering the full 15.60 arcsec Petrosian radius (SDSS-*r*) of UGC3995B, the foreground galaxy. We obtained the foreground galaxy’s mass from the SDSS fluxes of the unocculting side and the prescription from Zibetti, Charlot & Rix (2009).

2.4 2MASX

This pair’s serendipitous discovery was originally reported in Holwerda et al. (2009). The distance and stellar mass was estimated in Holwerda et al. (2013b) and the radial coverage of the overlap region is between 2 and 5 effective (or half-light) radii (R_e).

2.5 AM 500–620

This was a pair identified in White et al. (2000) as an overlapping galaxy pair and the *HST* data were originally reported in Keel & White (2001a). The radial coverage is mostly in the outer disc ($\sim R_{25}$, the 25 mag arcsec $^{-2}$ as defined by de Vaucouleurs et al. 1991). Stellar mass and Hubble type are from Keel & White (2001a).

3 TYPE IA SUPERNOVAE SAMPLE

SNe Ia are the touchstone of cosmological distance measurements. The photometric properties of their light curves are very stable but appear to depend slightly on host galaxy properties (e.g. star formation rate or stellar mass). With this mind, the third incarnation of the SDSS (SDSS-III) included a supernova search with spectroscopic follow-up. This SDSS-Supernova search (SDSS-SN)¹ has yielded a wealth of galaxy and supernova properties.

SDSS-SN is described in Frieman et al. (2008) and in more detail in Sako et al. (2008) and Sako et al. (2014). Observing for three months of the year from 2005 to 2007, SDSS-SN identified hundreds spectroscopically confirmed SNe Ia in the redshift range $0.05 < z < 0.35$.

Sako et al. (2014) present the final data-product of this tremendous observational effort. They include an assessment of SN type based on the light curve (PSNID output) and light-curve fits using the two most commonly used packages, *SALT2* (Guy et al. 2007) and *MLCS2k2* (Jha et al. 2007). Of these two packages, the *SALT2* light-curve fitter is now the most popular. Yet we focus on the *MLCS2k2* package because it explicitly starts with a host galaxy extinction prior, making it easier to directly compare its assumptions to our observations in occulting pairs.

However, we should note that both *SALT2* and *MLCS2k2* essentially measure the reddening of the supernova, not actual attenuation but the convolution of intrinsic supernova colour distribution, scatter by the surrounding interstellar matter and dust attenuation in the host galaxy. The shape of the *MLCS2k2* A_V prior used for the light-curve fits was $P(A_V) \sim e^{-A_V/\tau_0}$, with $\tau_0 = 0.4$. Because it has host galaxy extinction as an explicit prior and a model relation between A_V distribution width (τ_0) and A_V bias, we use the *MLCS2k2* values for our further analysis here. The *MLCS2k2* version used for the SDSS-SN analysis assumed an $R_V = 2.8$ reddening relation to convert reddening to extinction A_V (Kessler et al. 2009), indicating it is not completely host attenuation alone. However, it is not as unphysical as the $R_V = 1.7$ needed elsewhere with *MLCS2k2* (Hicken et al. 2009). For our purposes here, we will explicitly assume that the shape of the prior for *MLCS2k2*'s fit parameter A_V can be taken to be similar to the distribution of actual attenuation values as found in occulting galaxies.

Galaxy properties are those available from the SDSS-DR9 data base and stellar mass and star formation are modelled with two different packages, *FSPS* (Conroy & Gunn 2010) and *PEGASE* (Bruzual A. 2009). For the purpose of this paper, the stellar mass is of interest and we choose the *FSPS* value for the further analysis. Fig. 3 shows the distribution of host mass galaxies with SN Ia. The mean mass is $\sim 10^{10} M_\odot$, similar to AM500. We focus only on those objects that have reasonable chance of being bona fide SNe Ia ($S/N > 5$ and $P(\text{SN Ia}) > 50$ per cent) to ensure the conclusions for the improved prior are based on these only (1698 supernovae in the total sample).

In addition to the values thoughtfully provided by the SDSS-SN project in Sako et al. (2014), we retrieve the axis ratio, position angle, and Petrosian radius from the SDSS server. Using these values, we compute the galactocentric radial distance from the centre of the host galaxy in Petrosian radii. This is a different value than that of the separation presented in Sako et al. (2014), the ‘directional light radius’ (d_{DLR}), which is not deprojected into the plane of the host disc. If we assume discs ($n=1$ Sérsic index), the Petrosian radius corresponds to approximately two effective radii ($R_p = 2.15 \times R_e$;




Figure 3. The distribution of SN Ia host galaxy mass from Sako et al. (2014). They present both *FSPS* (Conroy & Gunn 2010) and *PEGASE* (Bruzual A. 2009) stellar masses. The masses of 2MASX (thick dashed), UGC 3995 (dashed line), and AM500 (dotted line) are marked as well.




Figure 4. The distribution of the projected separation of SNe Ia from their host galaxy in effective radii (R_e) based on the values reported in Sako et al. (2014). For comparison, Sako et al. (2014) plot the distance-to-host or ‘directional light radius’ (d_{DLR}), which is not deprojected into the plane of the host disc.

see Graham & Driver 2005). Fig. 4 shows the radial coverage for the SNe Ia for which SDSS Petrosian radii were available. SNe Ia are typically found by SDSS-SN between 0 and 5 effective radii (< 2.5 Petrosian radii), which correspond approximately to the radii covered by the occulting galaxy pairs (Table 1).

4 ANALYSIS

4.1 A_V distribution and bias

The relation between the exponential drop-off and bias introduced in the A_V value was explored by Jha et al. (2007) using the Riello & Patat (2005) simulations. Fig. 5 shows their relation (fig. 22 in the appendix of Jha et al. 2007). We mark the τ_0 values inferred from

¹ <http://www.sdss.org/supernova/aboutsupernova.html>




Figure 5. The relation between prior exponential scale and the resulting bias in the A_V from Jha et al. (2007). The thick grey line is the MLCS2K2 prior. The values of the three occulting pairs are also marked (not corrected for their inclination).

the occulting galaxy pairs, corrected for inclination ($\tau_0 = \tau / \cos(i)$). To parametrize the A_V bias, we fit the relation in Fig. 5 with

$$\Delta A = -0.5 \times e^{-\tau_0/0.15} + 0.02 \times \tau_0, \quad (2)$$

where τ_0 is the exponential drop-off of the host galaxy prior under consideration.

We note first that none of our three occulting galaxy templates match the drop-off typically used in MLCS2K2 ($\tau_0 = 0.4$) which was used for the SDSS-SN analysis (Sako et al. 2008, 2014). Therefore, assuming the SDSS-SN galaxies adhere to one of these three templates based on their host mass will inevitably introduce a bias in the distribution. Our goal is to explore in which direction and how severe it would be.

4.2 New luminosity distances for the SDSS-SN

Fig. 6 shows the relation between the luminosity distance and redshift for the SDSS-SN SNe Ia. They closely follow the expected




Figure 6. The relation between redshift and MLCS2K2 luminosity distance (top panel) and the difference between the ΛCDM and the SN Ia luminosity distances (bottom panel). The relation for different equations of state ($w = 1/-1$) are shown for illustration only.




Figure 7. The distribution of host galaxy τ_{host} values as calculated from equation (3). There is still a peak at $\tau_0 = 0.35$ as this is the value assigned when no good host estimate was available. The concentration at $\tau_0 = 0$ is due to poorly constrained inclination.

Λ cold dark matter (ΛCDM) relation with scatter around the mean in the residual (bottom panel in Fig. 6). To extract the exact nature of dark energy, one approach is to explore this residual. However, inconsistent or heterogeneous photometry or host galaxy extinction could introduce biases in this residual.

We assign each host galaxy a new exponential template (τ_{OCG}) from one of the three occulting galaxy (OCG) distributions (fit as an exponential drop-off), based on the nearest in stellar mass, and correct for disc inclination:

$$\tau_{\text{host}} = \tau_{\text{OCG}}(M) \times \cos(i_{\text{host}}), \quad (3)$$

where i_{host} is the inclination of the host galaxy as inferred from the minor and major axes ratio:

$$\cos(i_{\text{host}}) = \sqrt{\frac{(b/a)^2 - (b/a)_{\min}^2}{1 - (b/a)_{\min}^2}}, \quad (4)$$

originally from Hubble (1926), where a is the major axis, b the minor axis and we assume $b/a_{\min} = 0.1$. We found in Holwerda et al. (2015b) that the distribution of SDSS-SN A_V values becomes the same for high- and low-inclination galaxies, once the values are corrected for disc inclination using this simple $\cos(i)$ correction (i.e. the screen approximation).

Fig. 7 shows the distribution of resulting τ_{host} for the SDSS-SN SNe Ia sample. The peak around $\tau_{\text{host}} = 0.4$ corresponds to all the host galaxies for which no mass estimate was available in the SDSS-SN catalogue.

Starting from the new τ_{host} exponential values for each host galaxy (equation 3), we can now calculate the A_V bias for each host, based on its stellar mass and inclination (equation 2) and apply this to the luminosity distance. Fig. 8 shows the relative difference with ΛCDM distances as a function of redshift and the mean values (and dispersion) before our A_V bias correction and after (Table 2).

There are two things to note in the corrected values. First is that the mean luminosity distance is slightly lower, i.e. there is a little more extinction in the SN Ia host galaxies if we use individual priors, but there is no less scatter in the values.

The change in the mean is interesting for cosmological study but at present only points to the direction one can expect the luminosity distances to change. The priors may be slightly more tailored but




Figure 8. The relative difference between Λ CDM predicted distance and the SN Ia luminosity distances from SDSS-SN. Individual SN Ia (grey points) and the mean and rms (red points) and once corrected for A_V bias (green points) using the occulting galaxy templates (Fig. 2), matched through host mass. Alternatively, one can match OCG templates by radius (blue points). Two cosmologies are shown as well to illustrate: $w=-0.9$ and -1 , neither a fit to the data. The different cosmologies are shown to illustrate how the choice of dust prior can have an almost similar effect on the inferred cosmology.

Table 2. The average and scatter for Δ_L/D_L in four bins for the original MLCS2k2 analysis, one corrected using the occulting galaxy templates matched with host mass or radius.

z	Δ_L/D_L	Mass-matched	Radius-matched
0.10	0.32 (0.60)	0.29 (0.61)	-0.05 (0.79)
0.20	0.13 (0.24)	0.11 (0.24)	0.06 (0.34)
0.30	0.10 (0.25)	0.08 (0.25)	0.04 (0.34)
0.40	0.17 (0.32)	0.15 (0.32)	0.13 (0.37)

they are hardly appropriate for each host galaxy (see below for the occulting galaxy pair prior caveats). This lack of an appropriate prior is reflected – in our opinion – in the lack of change in dispersion. If an appropriate correction to the prior shape and hence A_V bias had been applied, luminosity distances would adhere closer to the Hubble flow and any remaining scatter can be attributed to individual motion of the galaxies or photometric errors.

Alternatively, one could match the few occulting galaxy templates by their projected radius, not the foreground galaxy’s mass. These are the blue averages points in Fig. 8. The scatter around the new mean residuals is worse than in the case of the mass-matched templates. We can speculate at this point that to match templates, one should select by host mass first and galactocentric radius only as a secondary effect. Once again, we are restricted by the numbers of occulting pair templates we have.

5 CAVEATS

While the radial coverage and host mass range appears reasonably approximated by our three template host galaxies, there are a number of issues outstanding with just using three galaxies as templates. We will focus on the shortcomings of the occulting galaxy pairs presented as priors.

5.1 MLCS2k2 A_V parameter is all dust attenuation

The central assumption in this work has been that the attenuation seen in occulting galaxies is the same as the A_V parameter in MLCS2k2. However, because the A_V is derived assuming an attenuation law, it cannot be ruled out that some measure of the reddening is not from dust attenuation but scatter in, or supernova intrinsic colour scatter.

5.2 Radial coverage

While Table 1 and Fig. 4 seem to suggest that at least the SN Ia occur mostly at the same radii for which we have some extinction distribution from occulting pairs, the coverage is far from complete. For example, for low-mass galaxies, there is only information between 1–2 effective radii.

5.3 Small sample

It is obvious to point out that it is impossible to infer a relation between stellar mass and an extinction distribution with only three galaxy templates. We plan to obtain more templates from future *HST* observations (Holwerda 2014) but for now we can only assign the exponential prior to the closest stellar mass to the host mass. Alternatively, we fitted a linear relation between τ_0 and mass for the occulting galaxy pairs but this resulted in no improvement of the variance of the SN Ia distances.

5.4 Interactions

Occulting galaxies are often interacting and the redshift difference for two of our templates (UGC 3995 and 2MASX) are such that an ongoing interaction cannot be ruled out. It is very likely in the case of UGC 3995. The exact effect of gravitational interaction on the distribution of ISM is unknown but likely to be severe. Is there more dust now at higher radii? Do the shocks and tides move more dust into denser clouds in advance of the burst of star formation associated with gas-rich interactions? The way around this issue again is to observe more and bona fide occulting pairs, well separated in redshift. A selection from a spectroscopic survey such as the GAMA survey (Driver et al. 2009, 2011; Baldry et al. 2010) may well prove to be ideal.

5.5 Is the A_V prior really an exponential?

One strong assumption so far has been that the distribution of extinction values is an exponential drop-off. However, Fig. 2 shows that none of the three distributions is fully described as an exponential. For example, in the more massive galaxies, the peak of the distribution is not close to $A_V = 0$ at all. We noted this in Holwerda & Keel (2013) for both the *HST* observations and the IFU ones. In the case of UGC 3995, one could contribute this to the ongoing interaction. Again the only way forward it to obtain more templates for the same host mass and infer the mean and variance of the distribution function.

Our assumption – and the MLCS2k2 authors’ – is that the shape of the distribution is an exponential. We now strongly suspect the true shape of the extinction distributions may be a lognormal one, similar to what Dalcanton et al. (in preparation) find for the extinction values in front of stars in M31 based on the PHAT project (Dalcanton et al. 2012) or an exponentially modified Gaussian. To illustrate, Fig. 9 shows fits to the A_V distribution using the exponentially modified




Figure 9. The shape of the A_V distribution of the three occulting galaxies with enough overlap to generate a distribution. An exponentially modified Gaussian distribution was used with maximum likelihood to each distribution. The relevant parameters are listed in Table 3.

Table 3. The exponentially modified Gaussian fits to the A_V distributions observed in occulting pairs.

Pair	m	σ	λ
2MASX	0.013	0.015	6.65
UGC3559	0.092	0.092	2.10
AM500	0.013	0.011	2.29
AM1316	1.41	0.50	70.60

Gaussian fit to each distribution using maximum likelihood (fit values in Table 3).

This strongly suggests that MLCS2k2 will now have to be redone using these templates instead of an exponential one.

6 APPLICATION TO SALT2

SALT2 handles the SN Ia colour and dust extinction in a combined parameter, and it may not be obvious how our approach has any easy application to this package. To illustrate, Fig. 10 shows the relation between the observed A_V and the SALT2 colour parameter, c (similar to fig. 9 in Sako et al. 2008). There is a linear relation between the two that follows:

$$c = 0.046 \times A_V + 0.012. \quad (5)$$

Thus, a change in the prior for A_V in the case of MLCS2k2 would translate a change in the prior for SALT2's assumption but in colour-space. To illustrate in Fig. 11, we plot the distribution of c values found in SDSS-III for SN Ia, as well as the values of A_V from MLCS2k2 translated using the above relation. The two distributions are – unsurprisingly – very similar. However, if we then translate the A_V value corrected for the inclination of the disc (inferred from the axis ratio), the distribution of c values changes significantly. In our view, this change in distribution after inclination correction is the original distribution one could adopt as the prior distribution of the parameter c for SALT2 applications, similar to the one to be adopted for MLCS2k2 (since both of them are essentially colour priors).




Figure 10. The relation between extinction as measured by the MLCS2k2 package and the colour parameter from the SALT2 package for the SDSS-SN supernovae. There is a linear relation between the two (fit with maximum likelihood, including the SALT2 colour errors).




Figure 11. The distribution of SALT2 colour parameter c for all SNe Ia (black line) and after converting the A_V values from MLCS2k2 using equation (5) (green histogram). Since the fit is reasonable, there is no big difference between the two. After correction for inclination alone, the A_V values would translate into the blue histogram.

7 DISCUSSION

The SN Ia community is well aware of the effects of host galaxy extinction may have on the results and have actively been searching for both ‘dust-free’ supernovae (i.e. hosted by early-type galaxies, see e.g. Suzuki et al. 2012) and pushed light-curve observations into the near-infrared to mitigate extinction issues.

This paper’s aim is an attempt to map out how the available observational information, and specifically the host galaxy information, can be further used to optimize the SN Ia distance measurement. As a first exercise, we applied the three available templates from occulting galaxies, naively matched to the SN Ia host through the stellar mass or galactocentric radius. Introducing new host extinction templates certainly has an effect but since the scatter in the distances is not markedly reduced, using just these templates would just skew the cosmological result, not improve it. Despite the fact that the A_V values were attained using a more Milky Way-type relation (Kessler et al. 2009), we cannot rule out that none of the current

scatter comes from host attenuation and all of it comes from intrinsic colour scatter in the SN Ia. But assuming much of the scatter is due to host attenuation, when matching host templates through projected radius rather than host mass, the scatter increases more than through mass-matching. This points, in our opinion, in which order one should match the occulting galaxy templates: first through mass and then through galactocentric radius.

Another key issue for SN Ia cosmological measurements will be if the extinction prior will change with redshift. Given that between $z=0$ and 1, specific star formation rates increased by a factor 3–10 (e.g. Noeske et al. 2007), one can expect a much more fractured dusty ISM (thanks to increased turbulence, as we speculated in Holwerda 2008) or a much denser one (to sustain the star formation). Some initial results on dust lanes in L^* edge-ons suggest that the global morphology in disc galaxies remains the same: 80 per cent of these have a dust lane mid-plane (Holwerda et al. 2012b).

Alternatively, if the A_V distribution is significantly different for massive versus lower-mass disc galaxies, any evolution in the mix of SN Ia host galaxies will result in an effective change in the appropriate A_V prior. Dust is strongly suspected to be distributed differently with respect to the stars in lower-mass galaxies. Initial *Spitzer* observations showed very extended dust morphology in a few individual cases of lower-mass galaxies (Hinz et al. 2007, 2009), similar to the occulting pair in Fig. 1. Recently, it has become clear that in the stacked spectral energy distribution of lower-mass galaxies there is virtually no attenuation in the optical and ultraviolet yet Herschel emission clearly show large dust masses at very low temperatures (Popescu et al. 2002; Grossi et al. 2010; Bourne et al. 2012). The combined evidence points to a dependence of star/dust geometry on galaxy mass. Whether this means dust is relatively more extended in radius, height above the plane (e.g. Dalcanton, Yoachim & Bernstein 2004; Holwerda et al. 2012a) or clumpiness, remains to be seen.

Fortunately, global extinction measures can be checked using high-redshift occulting galaxy pairs and both effects can then be accounted for, be it evolution in the mix of host galaxies or in the ISM of the host galaxies themselves. There is, at present, no observed evolution in the colour distribution of observed SNe Ia (Conley et al. 2011; Betoule et al. 2014). However, if there is significant evolution in the $P(A_V)$ distribution with redshifts – either due to ISM evolution in the host galaxies or a significant change in the host galaxy population mix – it may change the relative luminosity distances as a function of redshift, the driver or dark energy cosmology.

We hope to have convinced the reader that the use of occulting galaxy pairs to obtain extinction distribution templates is a good way forward to help reduce the remaining uncertainties in SN Ia distance measurements. The eventual goal is to cover enough range in stellar mass, radial coverage, and inclinations to map the mean and variance of the typical extinction distribution in spiral galaxy discs. This will have many astrophysical applications but the use as a prior for cosmological SN Ia measurements was the principal driver for us to obtain *HST* observations of occulting pairs: GO-13695, *STarlight Absorption Reduction through a Survey of Multiple Occulting Galaxies (STARSMOG)* P.I. B.W. Holwerda, a 150 orbit SNAPshot programme (Holwerda 2014), which is projected to complete a sample of 98 occulting pairs by 2017.

ACKNOWLEDGEMENTS

The authors are very grateful for the insightful comments of the anonymous referee and to the SDSS-SN team who provided the data. KJM is supported by an Australian Research Council Discov-

ery Early Career Researcher Award. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research has made use of NASA's Astrophysics Data System. This research made use of *ASTROPY*, a community-developed core *PYTHON* package for astronomy (Astropy Collaboration et al. 2013). This research made use of *matplotlib*, a *PYTHON* library for publication quality graphics (Hunter 2007). *PYRAF* is a product of the Space Telescope Science Institute, which is operated by AURA for NASA. This research made use of *SCIPY* (Jones et al. 2001).

REFERENCES

- Albrecht A. et al., 2006, preprint ([arXiv:astro-ph/0609591](https://arxiv.org/abs/astro-ph/0609591))
- Altavilla G. et al., 2004, MNRAS, 349, 1344
- Andredakis Y. C., van der Kruit P. C., 1992, A&A, 265, 396
- Astier P. et al., 2006, A&A, 447, 31
- Astropy Collaboration et al., 2013, A&A, 558, A33
- Baldry I. K. et al., 2010, MNRAS, 404, 86
- Berlind A. A., Quillen A. C., Pogge R. W., Sellgren K., 1997, AJ, 114, 107
- Betoule M. et al., 2014, A&A, 568, A22
- Bourne N. et al., 2012, MNRAS, 421, 3027
- Bruzual A. G., 2009, A&A, 500, 521
- Calzetti D., Kinney A. L., Storchi-Bergmann T., 1994, ApJ, 429, 582
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, ApJ, 345, 245
- Childress M. et al., 2013, ApJ, 770, 107
- Conley A., Carlberg R. G., Guy J., Howell D. A., Jha S., Riess A. G., Sullivan M., 2007, ApJ, 664, L13
- Conley A. et al., 2011, ApJS, 192, 1
- Conroy C., Gunn J. E., 2010, ApJ, 712, 833
- Dalcanton J. J., Yoachim P., Bernstein R. A., 2004, ApJ, 608, 189
- Dalcanton J. J. et al., 2009, ApJS, 183, 67
- Dalcanton J. J. et al., 2012, ApJS, 200, 18
- de Vaucouleurs G., de Vaucouleurs A., Corwin H. G., Buta R. J., Paturel G., Fouque P., 1991, Third Reference Catalogue of Bright Galaxies. Volume I–III. Springer-Verlag, Berlin
- Domingue D. L., Keel W. C., Ryder S. D., White R. E., III, 1999, AJ, 118, 1542
- Domingue D. L., Keel W. C. R. E., III, 2000, ApJ, 545, 171
- Driver S. P. et al., 2009, Astron. Geophys., 50, 050000
- Driver S. P. et al., 2011, MNRAS, 413, 971
- Elmegreen D. M., Kaufman M., Elmegreen B. G., Brinks E., Struck C., Klarić M., Thomasson M., 2001, AJ, 121, 182
- Frieman J. A. et al., 2008, AJ, 135, 338
- Gallagher J. S., Garnavich P. M., Berlind P., Challis P., Jha S., Kirshner R. P., 2005, ApJ, 634, 210
- Graham A. W., Driver S. P., 2005, PASA, 22, 118
- Grossi M. et al., 2010, A&A, 518, L52
- Gupta R. R. et al., 2011, ApJ, 740, 92
- Guy J. et al., 2007, A&A, 466, 11
- Guy J. et al., 2010, A&A, 523, A7
- Hamuy M., Phillips M. M., Suntzeff N. B., Schommer R. A., Maza J., Smith R. C., Lira P., Aviles R., 1996, AJ, 112, 2438
- Hicken M., Wood-Vasey W. M., Blondin S., Challis P., Jha S., Kelly P. L., Rest A., Kirshner R. P., 2009, ApJ, 700, 1097
- Hinz J. L., Rieke M. J., Rieke G. H., Willmer C. N. A., Misselt K., Engelbracht C. W., Blaylock M., Pickering T. E., 2007, ApJ, 663, 895
- Hinz J. L., Engelbracht C. W., Willmer C. N. A., Rieke G. H., Rieke M. J., Smith P. S., Blaylock M., Gordon K. D., 2009, in Sheth K., Noriega-Crespo A., Ingalls J., Paladini R., eds, The Fourth Spitzer Science Center Conf., The Evolving ISM in the Milky Way and Nearby Galaxies. Pasadena, CA, p. E7
- Holwerda B. W., 2005, PhD thesis, Rijksuniversiteit Groningen
- Holwerda B. W., 2008, MNRAS, 386, 475

- Holwerda B., 2014, HST Proposal, SNAP-13695, STarlight Absorption Reduction through a Survey of Multiple Occulting Galaxies (STARSMOG)
- Holwerda B. W., Keel W. C., 2013, *A&A*, 556, A42
- Holwerda B. W., González R. A., Allen R. J., van der Kruit P. C., 2005a, *AJ*, 129, 1381
- Holwerda B. W., González R. A., Allen R. J., van der Kruit P. C., 2005b, *AJ*, 129, 1396
- Holwerda B. W., Keel W. C., Bolton A., 2007, *AJ*, 134, 2385
- Holwerda B. W., Keel W. C., Williams B., Dalcanton J. J., de Jong R. S., 2009, *AJ*, 137, 3000
- Holwerda B. W. et al., 2012a, *A&A*, 541, L5
- Holwerda B. W., Dalcanton J. J., Radburn-Smith D., de Jong R. S., Guhathakurta P., Koekemoer A., Allen R. J., Böker T., 2012b, *ApJ*, 753, 25
- Holwerda B. W., Böker T., Dalcanton J. J., Keel W. C., de Jong R. S., 2013b, *MNRAS*, 433, 47
- Holwerda B. W., Dalcanton J. J., Keel W. C., Boeker T., de Jong R. S., 2013a, *MNRAS*, 433, 47
- Holwerda B. W., Reynolds A., Smith M., Kraan-Korteweg R. C., 2015a, *MNRAS*, 446, 3768
- Holwerda B. W. et al., 2015b, *MNRAS*, 449, 4277
- Howell D. A., 2001, *ApJ*, 554, L193
- Hubble E. P., 1926, *ApJ*, 64, 321
- Hunter J. D., 2007, *Comput. Sci. Eng.*, 9, 90
- Jha S., Riess A. G., Kirshner R. P., 2007, *ApJ*, 659, 122
- Jones E. et al., 2001, SciPy: Open Source Scientific Tools for Python available at: <http://www.scipy.org/>.
- Keel W. C., White R. E., III, 2001a, *AJ*, 121, 1442
- Keel W. C., White R. E., III, 2001b, *AJ*, 122, 1369
- Keel W. C., Manning A. M., Holwerda B. W., Mezzoprete M., Lintott C. J., Schawinski K., Gay P., Masters K. L., 2013, *PASP*, 125, 2
- Keel W. C., Manning A. M., Holwerda B. W., Lintott C. J., Schawinski K., 2014, *AJ*, 147, 44
- Kelly P. L., Filippenko A. V., Burke D. L., Hicken M., Ganeshalingam M., Zheng W., 2015, *Science*, 347, 1459
- Kessler R. et al., 2009, *ApJS*, 185, 32
- Kistler M. D., Stanek K. Z., Kochanek C. S., Prieto J. L., Thompson T. A., 2013, *ApJ*, 770, 88
- Lampeitl H. et al., 2010, *ApJ*, 722, 566
- Lintott C. J. et al., 2008, *MNRAS*, 389, 1179
- Mannucci F., Della Valle M., Panagia N., 2006, *MNRAS*, 370, 773
- Marziani P., D’Onofrio M., Dultzin-Hacyan D., Sulentic J. W., 1999, *AJ*, 117, 2736
- Noeske K. G. et al., 2007, *ApJ*, 660, L43
- Pan Y.-C. et al., 2014, *MNRAS*, 438, 1391
- Perlmutter S. et al., 1999, *ApJ*, 517, 565
- Phillips M. M., 1993, *ApJ*, 413, L105
- Phillips M. M., Lira P., Suntzeff N. B., Schommer R. A., Hamuy M., Maza J., 1999, *AJ*, 118, 1766
- Popescu C. C., Tuffs R. J., Völk H. J., Pierini D., Madore B. F., 2002, *ApJ*, 567, 221
- Reindl B., Tammann G. A., Sandage A., Saha A., 2005, *ApJ*, 624, 532
- Rest A. et al., 2014, *ApJ*, 795, 44
- Riello M., Patat F., 2005, *MNRAS*, 362, 671
- Riess A. G., Press W. H., Kirshner R. P., 1996, *ApJ*, 473, 588
- Riess A. G. et al., 1998, *AJ*, 116, 1009
- Riess A. G. et al., 2011, *ApJ*, 730, 119
- Sako M. et al., 2008, *AJ*, 135, 348
- Sako M. et al., 2014, *ApJS*, submitted
- Schawinski K., 2009, *MNRAS*, 397, 717
- Scolnic D. et al., 2014, *ApJ*, 795, 45
- Sullivan M. et al., 2006, *ApJ*, 648, 868
- Suzuki N. et al., 2012, *ApJ*, 746, 85
- van den Bergh S., Li W., Filippenko A. V., 2005, *PASP*, 117, 773
- Wang X., Wang L., Filippenko A. V., Zhang T., Zhao X., 2013, *Science*, 340, 170
- White R. E., Keel W. C., 1992, *Nature*, 359, 129
- White R. E., III, Keel W. C., Conselice C. J., 2000, *ApJ*, 542, 761
- Wood-Vasey W. M. et al., 2007, *ApJ*, 666, 694
- Zibetti S., Charlot S., Rix H., 2009, *MNRAS*, 400, 1181

This paper has been typeset from a \LaTeX file prepared by the author.