

Neutral hydrogen gas, past and future star formation in galaxies in and around the ‘Sausage’ merging galaxy cluster

Andra Stroe,¹★ Tom Oosterloo,^{2,3} Huub J. A. Röttgering,¹ David Sobral,^{1,4,5†} Reinout van Weeren^{6‡} and William Dawson⁷

¹Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, the Netherlands

²ASTRON, Postbus 2, NL-7990 AA Dwingeloo, the Netherlands

³Kapteyn Astronomical Institute, Postbus 800, NL-9700 AV Groningen, the Netherlands

⁴Instituto de Astrofísica e Ciências do Espaço, Universidade de Lisboa, Observatório Astronómico de Lisboa, Tapada da Ajuda, P-1359-018 Lisbon, Portugal

⁵Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Edifício C8, Campo Grande, P-1748-016 Lisbon, Portugal

⁶Harvard Smithsonian Center for Astrophysics (CfA - SAO), 60 Garden Street, Cambridge, MA 02138, USA

⁷Lawrence Livermore National Laboratory, PO Box 808 L-210, Livermore, CA 94551, USA

Accepted 2015 June 30. Received 2015 June 15; in original form 2015 April 28

ABSTRACT

CIZA J2242.8+5301 ($z = 0.188$, nicknamed ‘Sausage’) is an extremely massive ($M_{200} \sim 2.0 \times 10^{15} M_{\odot}$), merging cluster with shock waves towards its outskirts, which was found to host numerous emission line galaxies. We performed extremely deep Westerbork Synthesis Radio Telescope H α observations of the ‘Sausage’ cluster to investigate the effect of the merger and the shocks on the gas reservoirs fuelling present and future star formation (SF) in cluster members. By using spectral stacking, we find that the emission line galaxies in the ‘Sausage’ cluster have, on average, as much H α gas as field galaxies (when accounting for the fact cluster galaxies are more massive than the field galaxies), contrary to previous studies. Since the cluster galaxies are more massive than the field spirals, they may have been able to retain their gas during the cluster merger. The large H α reservoirs are expected to be consumed within ~ 0.75 – 1.0 Gyr by the vigorous SF and active galactic nuclei activity and/or driven out by the outflows we observe. We find that the star formation rate (SFR) in a large fraction of H α emission line cluster galaxies correlates well with the radio broad-band emission, tracing supernova remnant emission. This suggests that the cluster galaxies, all located in post-shock regions, may have been undergoing sustained SFR for at least 100 Myr. This fully supports the interpretation proposed by Stroe et al. and Sobral et al. that gas-rich cluster galaxies have been triggered to form stars by the passage of the shock.

Key words: shock waves – galaxies: active – galaxies: clusters: individual: CIZA J2242.8+5301 – radio continuum: galaxies – radio lines: galaxies.

1 INTRODUCTION

Galaxy cluster environments have a profound impact on the evolution of cluster galaxies. At low redshifts ($z < 0.5$) and focusing on relaxed clusters, the fraction of galaxies which are star forming drops steeply from field environments, to cluster outskirts and cores (Dressler 1980; Balogh et al. 1998; Goto et al. 2003). The morphological transformation of field spirals into cluster ellipticals or S0s has been attributed to a number of processes. The dense intracluster

medium (ICM) could lead to the ram pressure stripping of the gas content of field spirals as they accrete on to the cluster (e.g. Gunn & Gott 1972; Fumagalli et al. 2014). Tidal forces produced by gradients in the cluster gravitational potential or by encounters with other galaxies, can distort infalling galaxies, truncate their halo and disc (harassment; Moore et al. 1996) or remove gas contained in the galaxy and deposit it into the ICM (strangulation; Larson, Tinsley & Caldwell 1980). All these processes ultimately lead to the removal of gas and a truncation of star formation (SF).

The effect of relaxed cluster environments on galaxies is evident using a wide range of diagnostics, which trace different phases and time-scales of SF. Using ultraviolet (UV) data produced by young OB stars, Owers et al. (2012) found galaxies with star-forming trails, which they attribute to gas compression by the high-pressure merger

* E-mail: astroe@strw.leidenuniv.nl

† VENI/IF Fellow.

‡ Einstein Fellow.

environment. The UV radiation coming from massive, short-lived stars excites emission lines. Lines such as H α or [O II] 3727 Å probe SF on time-scales of <10 Myr. Emission line studies confirm that the fraction of star-forming galaxies increase from cluster cores towards field environments (e.g. Balogh et al. 1998; Gavazzi et al. 1998; Finn et al. 2005; Sobral et al. 2011; Darvish et al. 2014). Using far-infrared (FIR) data (tracing dust obscured SF), Rawle et al. (2012) find that the fraction of dusty star-forming galaxies, compared to the total star-forming galaxies, increases from low to high densities, an effect they attribute to dust stripping and heating processes caused by the cluster environment. Similar results are found by Koyama et al. (2013).

Synchrotron emission from supernovae (SNe) traces SF on longer time-scales of about 100 Myr (Condon 1992). Deep radio surveys at GHz frequencies indicate that below 100–200 μ Jy, the number of star-forming galaxies dominates over radio-loud active galactic nuclei (AGN; e.g. Padovani et al. 2011). The number of radio-faint star-forming galaxies (Morrison & Owen 2003) in clusters was found to increase with redshift (Morrison 1999). Radio broad-band (BB) emission from cluster spirals was also found to be enhanced with respect to field counterparts, an effect which can be caused by compression of the magnetic fields (Gavazzi & Jaffe 1985; Andersen & Owen 1995).

In addition to probes of past or current SF, CO rotational transitions can be used as an excellent tracer of molecular gas, which is the raw fuel for future SF episodes (e.g. Leroy et al. 2008). Other gas phases cannot form stars directly, but they have to cool sufficiently to form cold, dense molecular clouds (see review by Carilli & Walter 2013). However, a conversion factor between the CO mass and the total molecular gas is needed, which is highly uncertain (see review by Bolatto, Wolfire & Leroy 2013). Instead of using CO or other direct tracers of molecular gas, many studies use neutral hydrogen H I as a proxy for the molecular gas. Relaxed cluster spirals become increasingly more H I deficient towards cluster cores, an effect which does not depend on cluster global properties such as X-ray luminosity, temperature, velocity dispersion, richness or spiral fraction (e.g. Magri et al. 1988; Cayatte et al. 1990; Solanes et al. 2001; Chung et al. 2009). Oosterloo & van Gorkom (2005) and Scott et al. (2012) have found galaxies with H I tails, knots and filaments, which are possibly caused by ram pressure stripping. Until very recently, H I measurements have been limited to the local Universe ($z \sim 0$). At low redshifts ($z \sim 0.06$), Chengalur, Braun & Wieringa (2001) studied the A3128 cluster and found that the average H I mass for emission line and late-type cluster members is about $(8.6\text{--}8.7) \times 10^8 M_\odot$. They did not find a statistically significant difference between the H I content of emission line galaxies inside and outside the cluster, but the field spirals contain about two times more H I than their cluster counterparts. Pioneering work by Verheijen et al. (2007) and Lah et al. (2007, 2009) used direct detections and stacking to measure the H I content of cluster galaxies up to $z \sim 0.4$. Verheijen et al. (2007) surveyed two $z \sim 0.2$ clusters with very different morphologies: the relaxed, massive galaxy cluster A963 and the low-mass, diffuse cluster A2192. They detect only one H I galaxy within 1 Mpc of the centre of each cluster. By stacking galaxies with known redshifts, they make a clear detection of H I for blue galaxies outside the clusters, but no such detection was made for cluster galaxies. In a detailed study of the cluster A370 at $z \sim 0.37$, Lah et al. (2007) use spectral stacking to measure H I in gas-rich galaxies lying outside or at the outskirts of the cluster.

As discussed previously, relaxed cluster environments are believed to suppress SF by removing cold gas from their host galaxies. At $z < 0.3$ between 10 and 20 per cent of clusters are undergoing

mergers (Katayama et al. 2003; Sanderson, Edge & Smith 2009; Hudson et al. 2010) and this fraction is expected to increase steeply beyond $z = 0.4$ (Mann & Ebeling 2012). The effect of cluster mergers on the SF activity and gas content of galaxies is disputed. Most studies find that cluster mergers trigger SF (Miller & Owen 2003; Umeda et al. 2004; Ferrari et al. 2005; Owen et al. 2005; Johnston-Hollitt et al. 2008; Cedrés et al. 2009; Hwang & Lee 2009; Stroe et al. 2014a, 2015; Sobral et al. 2015; Wegner, Chu & Hwang 2015), but a few studies find they quench it (e.g. Poggianti et al. 2004) or that they have no direct effect (e.g. Chung et al. 2010).

An interesting subset of clusters are those hosting radio relics, extended patches of diffuse radio emission tracing merger-induced shocks (e.g. Ensslin et al. 1998). The H α properties of radio-relic clusters Abell 521, CIZA J2242.8+5301 and 1RXS J0603.3+4214 (Umeda et al. 2004; Stroe et al. 2014a, 2015) indicate that the merger and the passage of the shocks lead to a steep SF increase for <0.5 Gyr. The fast consumption of the gas ultimately quenches the galaxies within a few hundred Myr time-scales (Roediger et al. 2014).

The H α studies of Umeda et al. (2004) and Stroe et al. (2014a, 2015) are tracing instantaneous (averaged over 10 Myr) SF and little is known about SF on longer time-scales and the reservoir of gas that would enable future SF. An excellent test case for studying the gas content of galaxies within merging clusters with shocks is CIZA J2242.8+5301 (Kocevski et al. 2007). For this particular cluster unfortunately, its location in the Galactic plane, prohibits studies of the rest-frame UV or FIR tracing SF on longer time-scales, as the emission is dominated by Milky Way dust. However, the rich multiwavelength data available for the cluster give us an unprecedented detailed view on the interaction of their shock systems with the member galaxies. CIZA J2242.8+5301 is an extremely massive ($M_{200} \sim 2 \times 10^{15} M_\odot$; Dawson et al. 2015; Jee et al. 2015) and X-ray disturbed cluster (Akamatsu & Kawahara 2013; Ogren et al. 2013, 2014) which most likely resulted from a head-on collision of two, equal-mass systems (van Weeren et al. 2011; Dawson et al. 2015). The cluster merger induced relatively strong shocks, which travelled through the ICM, accelerated particles to produce relics towards the north and south of the cluster (van Weeren et al. 2010; Stroe et al. 2013). There is evidence for a few additional smaller shock fronts throughout the cluster volume (Stroe et al. 2013; Ogren et al. 2014). Of particular interest is the northern relic, which earned the cluster the nickname ‘Sausage’. The relic, tracing a shock of Mach number $M \sim 3$ (Stroe et al. 2014c), is detected over a spatial extent of ~ 1.5 Mpc in length and up to ~ 150 kpc in width and over a wide radio frequency range (150 MHz–16 GHz; Stroe et al. 2013, 2014b). There is evidence that the merger and the shocks shape the evolution of cluster galaxies. The radio jets are bent into a head–tail morphology aligned with the merger axis of the cluster. This is probably ram pressure caused by the relative motion of galaxies with respect to the ICM (Stroe et al. 2013). The cluster was also found to host a high fraction of H α emitting galaxies (Stroe et al. 2014a, 2015). The cluster galaxies not only exhibit increased SF and AGN activity compared to their field counterparts, but are also more massive, more metal rich and show evidence for outflows likely driven by SNe (Sobral et al. 2015). Stroe et al. (2015) and Sobral et al. (2015) suggest that these relative massive galaxies (stellar masses of up to $\sim 10^{10.0\text{--}10.7} M_\odot$) retained the metal-rich gas, which was triggered to collapse into dense star-forming clouds by the passage of the shocks, travelling at speeds up to ~ 2500 km s $^{-1}$ (Stroe et al. 2014c), in line with simulations by Roediger et al. (2014).

In this paper we focus on the effect of the massive cluster merger and travelling shocks in the ‘Sausage’ cluster on the H I content

of the galaxies, tracing the gas that may fuel future SF. We place this in context of other phases of SF, averaged over short time-scales (~ 10 Myr, $H\alpha$ data) and averaged over longer time-scales (~ 100 Myr, radio BB data) SF episodes.

The structure of the paper is as follows: in Section 2 we discuss the observations and the reduction of the $H\alpha$, optical and BB data; in Section 3 we discuss the $H\alpha$ detections, stacking, masses and how these correlate with $H\alpha$ and radio luminosities; in Section 4 we discuss the implication of our results for future SF episodes and compare them with other $H\alpha$ studies of clusters. Finally, Section 5 presents a summary of the results, placing them in context of the SF history of the cluster. At the redshift of the ‘Sausage’ cluster, $z \sim 0.188$, 1 arcsec covers a physical scale of 3.18 kpc and the luminosity distance is $d_L \approx 940$ Mpc. All coordinates are in the J2000 coordinate system. We use a Chabrier (2003) initial mass function (IMF) throughout the paper. We correct measurements from other papers accordingly.

2 OBSERVATIONS AND DATA REDUCTION

In our analysis, we combine radio spectral line data tracing $H\alpha$, BB radio data, optical imaging and spectroscopy of passive and star-forming galaxies in and around the cluster.

2.1 $H\alpha$ data

The ‘Sausage’ cluster was observed with the Westerbork Synthesis Radio Telescope (WSRT)¹ in the maxi-short configuration during the second half of 2012, for a total of twenty-six 12-h tracks. The observations were taken in four, slightly overlapping bands of 10 MHz each and central frequencies 1180, 1188, 1195 and 1203 MHz, respectively, therefore fully covering the 1175–1208 MHz range. Recirculation was used to have 512 channels per band (with XX and YY polarizations only). Sources CTD 93 and 3C 147 were used as flux calibrators.

The velocity resolution (after Hanning smoothing), at the redshift of the cluster, is about 20 km s^{-1} , sampled a channel width of $\sim 9.9 \text{ km s}^{-1}$. The velocity range covered is about 4700 km s^{-1} , corresponding to an $H\alpha$ redshift range of 0.184–0.199. The redshift range covers the cluster volume within -1σ to $+3\sigma$ of the cluster redshift $z = 0.188$, where $\sigma = 0.04$ is the cluster velocity dispersion (Dawson et al. 2015). The $H\alpha$ observations cover well the distribution of the $H\alpha$ emitting galaxies ($z = 0.190 \pm 0.010$; Sobral et al. 2015).

Significant radio frequency interference (RFI) is known to be present at the WSRT at frequencies covering the $H\alpha$ redshift range of 0.1–0.25, caused by geopositional systems such as GPS and GLONASS. However, at the time of our observations, the frequencies corresponding to the redshift of the cluster (1185–1200 MHz) were still fairly free of such RFI. With the foreseen deployment of the Galileo geopositional system, the situation in this frequency range will worsen dramatically in the near future.

To remove any residual RFI, we performed our RFI flagging on the Stokes Q (i.e. XX–YY) component on the data. Given the polarized nature of RFI, this removed most of the astronomical signal, but left the RFI mostly intact. The flags found for Stokes Q were then applied to original XX and YY visibilities. Moreover, because the RFI in our data is BB, before flagging we performed a

smoothing in frequency to enhance the ‘sensitivity’ for RFI. These procedures worked very well and the final data cubes do not show any effects of residual RFI, while the noise level ($75 \mu\text{Jy beam}^{-1}$ over 20 km s^{-1}) is very close to what is expected for the integration time.

Once the data were flagged, we applied standard calibration procedures to the data using the miriad software (Sault, Teuben & Wright 1995). The continuum emission was removed from the data by fitting, to all channels, a third-order polynomial to each visibility spectrum (‘UVLIN’). The choice of the order depends on out to which radius there are significant continuum sources and on bandwidth. The higher the order, the better the sources are removed, however the noise, after subtraction, increases. A third-order polynomial fit represents a compromise between sufficient removal of the continuum and little increase in the noise level of the line cube, as shown by Sault (1994). Because this does not take into account the presence of any possible $H\alpha$ emission, the spectra of individual detections, and of the stacked spectra we discuss below, are corrected for this oversubtraction (see Section 3.3).

The synthesized beam of the WSRT observations is $24.9 \times 18.5 \text{ arcsec}^2$ at a position angle of $165^\circ.8$, or $79.3 \times 58.9 \text{ kpc}^2$.

2.2 Optical imaging and spectroscopy

In order to locate the spatial and velocity position of the $H\alpha$ signal, we use the Keck and William Herschel Telescope spectroscopy. Data of both passive and star-forming galaxies in the field of the cluster were presented in Sobral et al. (2015), Dawson et al. (2015) and Stroe et al. (2015).

Galaxies are categorized as passive or emission line based on spectral features. Emission line galaxies were selected through the presence of the $H\alpha$ emission line (with an $H\alpha + [N\text{II}]$ rest-frame equivalent width larger than 13 \AA), tracing hot ionized gas, indicating the presence of SF and/or radio-quiet AGN (broad and narrow line AGN). The spectra of passive galaxies display Balmer absorption features and/or no Balmer series emission lines. The passive galaxies are undetected in $H\alpha$ at the 13 \AA level equivalent width level. The sample was divided into three categories:

- (i) passive galaxies inside the cluster;
- (ii) emission line galaxies within the cluster;
- (iii) emission line galaxies in the field around the cluster.

The number of sources in each subsample can be found in Table 1. The emission line galaxy sample is dominated by star-forming galaxies with a ~ 20 –30 per cent contribution from broad and narrow line optical AGN (see Table 1; Sobral et al. 2015). The cluster members were chosen to be located at a projected distance of less than 1.85 Mpc away from the cluster ‘centre’, in line with the definition of cluster membership from Stroe et al. (2015). Line-emission galaxies outside the cluster were defined to lie outside of the 1.85 Mpc radius. The redshift distribution of the galaxies is plotted in Fig. 1. We note that, as shown by Sobral et al. (2015), the samples of cluster and field line emitters are selected uniformly, down to a similar star formation rate (SFR).

The spectroscopy is supplemented with Subaru Telescope, Canada–France–Hawaii Telescope, William Herschel Telescope and Isaac Newton Telescope BB and narrow-band (NB) imaging tracing the $H\alpha$ emission line in galaxies at the cluster redshift (Stroe et al. 2015). $H\alpha$ luminosities for each source were calculated using the excess of NB emission compared to the i BB filter (for method and details, see Stroe et al. 2015). In the analysis, we

¹ <http://www.astron.nl/radio-observatory/astronomers/observing-wsrt/observing-wsrt>

Table 1. Number of sources with spectroscopic data, separated by galaxy type. Not all spectroscopic sources are covered by the redshift range of the H α observations.

Sample	Total number	Sources within H α z range	Reference
Emission line, field	39	22	Sobral et al. (2015), Stroe et al. (2015)
Emission line, cluster	54	45	Sobral et al. (2015), Stroe et al. (2015)
Cluster star forming	48	39	Sobral et al. (2015)
Cluster AGN	6	6	Sobral et al. (2015)
All emission line	93	67	Sobral et al. (2015), Stroe et al. (2015)
Passive, cluster	184	154	Dawson et al. (2015)
All	277	221	




Figure 1. Redshift distribution of sources in each galaxy sample (emission line cluster members, emission line field galaxies and cluster passive galaxies). The H α redshift coverage of the WSRT data is marked by the vertical dashed lines.

also employ stellar masses derived using the method described in Sobral et al. (2015).

2.3 Broad-band radio data

We identified radio counterparts to the optical sources by cross-matching with a deep, high-resolution (~ 1.5 arcsec) image of the cluster, centred at 1.5 GHz, produced using the upgraded Jansky Very Large Array (JVLA).

Deep JVLA observations of the cluster were taken in the 1–2 GHz L band in A, B, C and D-array configurations. An overview of the observations is given in the Table 2. In total, 16 spectral windows with 64 channels, each covering 64 MHz of bandwidth, were recorded. The data reduction for each observing run was carried out separately, using CASA² version 4.2.

As a first step, the data were Hanning-smoothed and corrections for the antenna positions and elevation-dependent gains were applied. We then obtained an approximate bandpass solution using observations on the primary calibrators (3C 147, 3C 138). We applied the bandpass solutions to the data and flagged RFI in an automated way using the AOFLAGGER (Offringa et al. 2010). This initial bandpass correction was performed to avoid flagging of data due to the bandpass variations across the spectral windows. After flagging, we obtained gain corrections on the primary calibrators using five channels centred at channel 30. These gain solutions were obtained to remove the time-varying gains. We pre-applied these solutions to find delay terms and bandpass solutions. We then redetermined the gain solutions but now using the full channel range pre-applying the bandpass and delay corrections. We then pre-applied these solutions to obtain gain solutions on our secondary calibrator J0542+4951. The cross-hand delays were solved for using the calibrator 3C 138. The channel-dependent polarization leakage terms and polarization

angles were set using 3C 147 and 3C 138, respectively. For observing runs longer than 4 h the leakage terms were determined from scans on the secondary calibrator J0542+4951.³

We bootstrapped the flux-scale from our primary calibrator observations to find the flux density of J0542+4951. The flux-scale was set using the default settings of the task SETJY. As a final step, the calibration tables were applied to the target data. We averaged the target field data by a factor of 3 in time and frequency, to reduce the data volume for imaging.

In the next steps, the calibration solutions were refined using several cycles of phase-only and amplitude and phase self-calibration. For the imaging we employed w-projection (Cornwell, Golap & Bhatnagar 2008, 2005) and MS-MFS clean (Rau & Cornwell 2011) with NTERMS=3. Clean boxes were set by running the Python Blob Detection and Source Measurement (PYBDSM).⁴ A few additional clean regions were added manually for diffuse sources.

After self-calibrating the individual data sets, we combined all the data sets to make one deep image. Two more rounds of amplitude and phase self-calibration (on a 1 h time-scale) were carried out on the combined data set. During the self-calibration the amplitude scale was allowed to drift freely. This was needed to fully align the different data sets and spectral windows and avoid strong artefacts around a few bright sources located in the field of view (FOV).⁵ We made a final image using Briggs (1995; ROBUST=0) weighting and corrected for the primary beam attenuation. We checked the flux-scale of the image against our previous 1.38 GHz WSRT observations of the cluster (van Weeren et al. 2010; Stroe et al. 2013). This was done by checking the integrated fluxes of 10 sources and scaling the fluxes from 1.4 to 1.5 GHz assuming a spectral index of -0.7 , the canonical spectral index for bright radio sources (e.g. Condon 1992). The JVLA fluxes were divided by a correction factor of 1.4 ± 0.1 to realign the flux-scale to the WSRT scales. The resolution of the VLA image is 1.5×1.4 arcsec 2 , with a position angle of 86.5° .

3 RESULTS

3.1 Direct detections

Six galaxies are tentatively detected in H α . The detection criterion is signal above 5σ over at least two consecutive channels. Table 3 lists the redshifts and H α masses of the direct detections. The narrow H α

³ The polarization results will be discussed in a forthcoming paper (van Weeren et al., in preparation).

⁴ <http://dl.dropboxusercontent.com/u/1948170/html/index.html>

⁵ In principle, the global amplitude scale could have been preserved, but such an option is not offered in CASA at the moment.

² <http://casa.nrao.edu>

Table 2. Details of the JVLA observations of the ‘Sausage’ cluster taken in L band (1.5 GHz), combining all four configurations of the telescope for a total of >26 h of observing time.

	<i>L</i> -band A-array	<i>L</i> -band B-array	<i>L</i> -band C-array	<i>L</i> -band D-array
Observation dates	2014 May 11	2013 Oct 31	2013 Sep 2; 2013 Jul 3	2013 Feb 3; 2013 Jan 31; 2013 Jan 27
Total used on source time (h)	~ 6.5	~ 6.5	~ 10	~ 3.5
Integration time (s)	1	3	5	5

Table 3. Properties of the direct detections. The positional uncertainties are 15 arcsec, the error in the redshifts is 0.0005 and the error in the $\text{H}\alpha$ mass about $0.4 \times 10^9 M_\odot$. The stellar masses reported are for the closest face-on spiral found within 7 arcsec.

RA ($^h^m^s$)	Dec. ($^{\circ}^{\prime\prime\prime}$)	z	$M_{\text{H}\alpha}$ ($10^9 M_\odot$)	M_* ($10^9 M_\odot$)	$M_{\text{H}\alpha}/M_*$
22 41 51.1	52 52 55	0.18893	2.1	2.0	1.0
22 42 56.5	52 57 21	0.18916	2.6	^a	—
22 43 14.3	53 04 57	0.18536	1.2	9.7	0.1
22 43 23.2	53 04 41	0.18536	2.2	^b	—
22 41 30.4	53 05 58	0.18486	1.7	^b	—
22 43 43.7	53 09 43	0.18994	1.6	7.9	0.2

^aA bright star overlaps the position of the galaxy, so no reliable counterpart can be found.

^bNo face-on spiral was found nearby the $\text{H}\alpha$ detection indicating the detection is spurious.

profiles (~ 40 km s^{-1}) of the directly detected sources indicate, if real, they are most probably oriented in the plane of the sky. For a given $\text{H}\alpha$ mass, the peak flux for a face on galaxy is higher than for an edge-on galaxy because the same amount of flux divided up in fewer channels. Therefore, we are biased towards detecting face-on galaxies. However, given the very narrow profiles of these tentative detections, they could also be high noise peaks.

We note that none of the six $\text{H}\alpha$ direct detections has counterparts (matches within 10 arcsec) in our spectroscopic data or in the $\text{H}\alpha$ catalogue. This indicates that the sources have faint $\text{H}\alpha$ emission, below the equivalent width detection threshold (13 Å). Their SFR are therefore below $\sim 0.35 M_\odot \text{ yr}^{-1}$.

Given the positional accuracy coupled with the large beam of WSRT finding the right optical counterpart is very challenging. A few (1–6) potential optical hosts are found for the tentative $\text{H}\alpha$ direct detections, but most sources are unresolved and faint (i -band magnitude on average fainter than 20). Therefore reliable photometric redshifts could not be derived and we cannot confirm them as being located at $z \sim 0.2$. However, even if these sources were $z \sim 0.2$ galaxies, they would have small stellar masses. For example, using the closest optical match as galaxy host, we find that their stellar masses are very small ($< 1 \times 10^9 M_\odot$). This would imply unrealistically high gas to stellar mass ratios. Additionally, the morphology of these close optical sources does not match face-on spiral galaxies, as we expect. These arguments indicate that the closest sources are not the correct matches. For three out of the six $\text{H}\alpha$ detections there are large, face-on spiral galaxies in their vicinity (within 7 arcsec), which could be the correct optical counterpart. These three galaxies have stellar masses of $\sim (2\text{--}10) \times 10^9 M_\odot$, which indicates 0.1–1 atomic gas to stellar fractions. For two detections only small sources are located in the vicinity and no obvious face-on galaxies are found nearby the $\text{H}\alpha$, indicating these are spurious detections (noise peaks in adjacent channels mimicking a signal). In one case, a bright star located at the location of the $\text{H}\alpha$ detection prevent correct optical identification.




Figure 2. Top: $\text{H}\alpha$ peak emission flux in the cluster line emitter stack as function of the aperture size used for extracting the individual spectra. The horizontal scale indicates the fraction by which the aperture width and height are scaled as a function of the FWHM. The trend in the peak emission does not change before and after correcting for the continuum oversubtraction (see Section 3.3). Bottom: average $\text{H}\alpha$ mass of cluster line emitters (see Section 3.4), as function of the aperture size used for extracting the individual spectra. The horizontal scale indicates the fraction by which the aperture width and height are scaled as a function of the FWHM.

3.2 $\text{H}\alpha$ stacking

Since only six galaxies were directly detected, we use spectral stacking to measure the average $\text{H}\alpha$ content of the galaxy samples.

We use the optical positions to extract radio spectra for each galaxy, summing the flux within an elliptical aperture equal to the full width at half-maximum (FWHM) of the synthesized beam (24.9×18.5 arcsec 2). This corresponds to a spatial scale of 79.3×58.8 kpc 2 , matched to the physical size of the galaxies, which at $z \sim 0.19$ are unresolved in the $\text{H}\alpha$ observations (Verheijen et al. 2007) (also in line with the galaxy sizes presented in Stroe et al. 2014a, 2015). We further tested the effect of the aperture size on the final stack using apertures of sizes ranging from 0.1 of the FWHM up to two times the FWHM.

To test the effect of aperture size on the final $\text{H}\alpha$ stack, we use elliptical apertures in size equal to a fraction of the FWHM on both the width and height of the synthesized beam. We use apertures from 0.1 to 2 times the FWHM of the radio beam. We follow the procedure described in Section 3.2 to extract spectra at the positions of the cluster emission line members. We find that the peak of the $\text{H}\alpha$ detection remains relatively stable if the aperture is at least 0.9 of the FWHM size (see top panel, Fig. 2).

For each aperture size (from 0.1 to 2.0 times the FWHM of the radio beam, see Fig. 2), we also measure the $\text{H}\alpha$ mass in the way described in Section 3.4. As shown in the bottom panel of Fig. 2, we

Table 4. Peak number of sources in every stack created for the different galaxy samples, the rms noise value obtained for each stack and the parameters of the Gaussian fit to the H₁ signal. H₁ emission is securely detected, at its peak, in the emission line galaxy and all galaxy stacks, but not in the cluster passive galaxies. When integrating the H₁ signal to estimate an H₁ mass, we obtain clear detections for the line emitters.

Sample	Number	σ_{rms} (μJy)	H ₁ peak (μJy)	Peak significance σ_{rms}	H ₁ velocity (km s ⁻¹)	H ₁ width (km s ⁻¹)	M_{H_1} ($10^9 M_{\odot}$)	M_{\star} ($10^9 M_{\odot}$)	M_{H_1}/M_{\star}
Emission line, field	22	21.3	60.4	2.8	1.0	69.3	1.86 ± 1.20	4.8 ± 0.8	0.39 ± 0.26
Emission line, cluster	45	5.9	42.7	7.2	4.6	137.0	2.50 ± 0.62	7.4 ± 0.5	0.34 ± 0.09
All emission line	67	8.2	45.1	5.5	-0.7	104.7	2.00 ± 0.67		
Passive, cluster	154	2.1	5.3	2.5	-97.4	90.1	0.21 ± 0.15	25.6 ± 0.4	$<0.008 \pm 0.006$
All	221	3.1	14.3	4.6	-29.5	107.0	0.60 ± 0.30		

find that the H₁ mass is relatively stable as function of aperture, but it peaks when the size of the aperture is equal to the FWHM of the synthesized beam. The FWHM size of WSRT is also well matched to the expected H₁ disc size of galaxies at $z \sim 0.19$ (Verheijen et al. 2007). We therefore choose to measure H₁ quantities within apertures equal to the FWHM of the synthesized beam.

We extract spectra for all galaxies whose redshift falls within the H₁ redshift range probed by our WSRT observations ($0.184 < z < 0.199$). To study the noise properties, we extract spectra in sky positions shifted by -60 arcsec in the RA direction, but using the same redshifts as the sample of galaxies. This method captures the effect of increasing noise towards the edges of the bandpass. Note however that the noise could be overestimated. Given the large source density of the cluster field, a shift in sky position does not guarantee we will be measuring pure noise, but some apertures could partially fall on undetected source.

Before stacking, we correct the galaxy and noise spectra for the WSRT primary beam, which is a function of distance from the pointing centre and observing frequency:

$$A(r, v) = \cos^6(cvr), \quad (1)$$

where $c = 68$ is a constant, r is the distance from the pointing centre in degrees and v is the observing frequency in GHz. By correcting for the primary beam, we account for the effect of noise increasing towards the FOV edges in both the galaxy and their associated noise spectra.

The spectra are then shifted to the rest-frame velocity using their spectroscopic redshift. We use the radio definition of velocity: $V = c(v_{\text{H}_1} - v_{\text{observed}})/v_{\text{H}_1}$, where c is the speed of light, v_{H_1} is the rest-frame frequency of H₁ and v_{observed} is the observing frequency. The galaxy spectra and the noise spectra are co-added using a weight based on the primary beam ($\propto A^2(r, v)$), which accounts for the fact that the noise for galaxies away from the field centre is larger in proportion to the primary beam. To obtain the correct flux density scale, we normalize the stacked spectrum by the integral of the synthesized beam, integrated over the same spatial region used for extracting the galaxy spectra.

After stacking and normalizing the spectra, we filter the data using a second-order Savitzky–Golay (SG) filter (Savitzky & Golay 1964), which convolves the data with a polynomial filter. Given that our line profiles are resolved (FWHM of the stacks with detection is > 150 km s⁻¹, compared to a channel width of > 20 km s⁻¹, see Table 4), the method reduces the noise, while preserving line profiles (see e.g. Morabito et al. 2014). We tested the method using different window sizes. The effect of the SG filtering with increasing window lengths is shown in Fig. 3. Out of the window size tested, we finally choose a filter window of ~ 300 km s⁻¹, which provides minimal noise, while preserving the width and height of the signal, thus maximizing the signal-to-noise ratio of the possible H₁ detection




Figure 3. Noise reducing properties of the SG filter with increasing filter window width demonstrated on the cluster member galaxy stack. Colours of the lines go from lighter to darker shades in increasing window length. The black line at the foreground of the figure is smoother spectrum after applying the final window length choice, ~ 300 km s⁻¹, which is also the one we finally applied to the data. Similar results are obtained for the galaxy stacks, as well as their associated noise stacks.




Figure 4. Number of sources stacked in each velocity channel, for the separate stacks. The passive cluster galaxy stack (black dashed line) benefits from a factor of > 5 more sources than the stacks for the H_α line emitters (mostly star-forming and some AGN) galaxies inside (dark grey solid line) and outside the cluster (black, solid line). Note how the number of sources dwindles towards the edges of the WSRT H₁ bandpass. The asymmetry in the number of sources is caused by the redshift distribution of the stacked sources (see Fig. 1), preferentially located at redshifts lower than the central WSRT H₁ coverage.

peaks. We also tested other smoothing kernels (e.g. moving boxcar), which resulted in similar results, but with a widening of the profile and reduction of the peak strength.

Separate stacks are produced for the sample of passive cluster galaxies, line emission cluster members and line emission galaxies located within the field environment around the cluster. Line emitters include both star-forming galaxies and optical AGN (see also Section 2.2). Finally, we produce a master stack of all the galaxies available. The number of galaxies for each velocity channel for each stack is shown in Fig. 4 and Table 4. The number of sources in the stack naturally peaks at the 0-velocity position, but dwindles towards higher relative velocities. This effect is governed by where the redshift of each source falls within the WSRT H₁ bandpass.

Because of extensive spectroscopy from Keck aimed at obtaining a dynamical analysis of the cluster that specifically targeted the red sequence, the number of passive cluster galaxies far outnumbers the number of emission line galaxies.

The asymmetry (about the central position) in the number of sources for which data exist, especially visible in the passive cluster galaxy sample, is caused by the discrepancy in the nominal redshift of the cluster ($z = 0.188$, recently derived from more than 200 spectra by Dawson et al. 2015) and the outdated $z = 0.192$ (Kocevski et al. 2007) which was used for creating the WSRT H α set-up. Our spectroscopic measurements are therefore biased towards lower redshifts, compared to the H α coverage of the WSRT data (see Section 2.2 and Fig. 1). If a redshift of a source falls at the middle of the WSRT H α band coverage, the frequency coverage is symmetric about the observed frequency of the H α . However, a lower redshift (than the central redshift) is equivalent to a source having a wider frequency coverage at frequencies lower than the H α and a narrower coverage at higher frequencies. When translating to a rest-frame velocity, there is a preferential data coverage of the positive rest-frame velocities. The missing data at larger absolute rest-frame velocities drive the noise to higher values in those regions.

3.3 Correcting for the oversubtracted continuum emission

As mentioned previously, the bulk of our galaxies do not have a direct detection of the H α line. Hence, the continuum emission subtraction (see Section 2.1) could not take into account the presence of H α emission and leads to an oversubtraction of the continuum where the putative H α line is located. The oversubtraction is not visible (and relevant) in the individual spectra, but it shows up in the stacked spectrum. As expected, the overcorrection of the continuum is evident in the case of the emission line galaxy samples, where the H α signal is located on top of a broad, negative dip. No evident negative trough is present in the passive member sample, where less H α is expected.

We apply a two-step process to correct for the oversubtraction of the continuum. First, to measure the possible extent of the H α in the H α galaxies, we fit Gaussian profiles to their stacked profiles. We select data at least $4\sigma_{\text{Gauss}}$ away from the peak of the Gaussian, where σ_{Gauss} is the dispersion of Gaussian profile. Additionally, we discard the data at relative rest-frame velocities higher than 1000 km s^{-1} . These cuts are employed to exclude any H α signal from the estimation of the continuum, but not include very noisy edge channels, where only a few galaxies are stacked (see Fig. 4). In the case of the passive members, a similar procedure is applied, but since we do not have a clear detection of emission, the velocity range between 250 and 1000 km s^{-1} is used.

We fit the channels free of H α with a second-order polynomial and subtract the fit from the data to correct for the oversubtracted continuum (see example in Fig. 5). The correction is substantial for the cluster H α line galaxies, but the results indicate that the continuum has not been significantly oversubtracted for passive cluster members and those line emitters located in the field environment.

3.4 Measuring the H α signal and its significance

After the stacked spectra are filtered and corrected for the continuum oversubtraction, we fit Gaussian profiles around the 0-velocity position. Results are shown in Fig. 6. The parameters of the Gaussian fits can be found in Table 4. We calculate the rms from the noise stack. Note that even though the number of galaxies in the




Figure 5. Correction for the continuum oversubtraction in the case of the cluster line emitter H α stack.

cluster and field stacks is similar, the noise levels achieved are a factor of ~ 4 different. This is because field galaxies are preferentially located at large distances away from the FOV centre, meaning that the noise levels at their location are higher.

In the case of the H α emission line galaxies, a clear peak is found around the 0-velocity position and a Gaussian profile could clearly be fit. In the cluster line emitter stack, we reach a noise level σ_{rms} of $5.9 \mu\text{Jy}$ and detect H α at a peak significance of 7.2σ . The results are virtually unchanged if the six AGN are removed from the stack. No clear detection of H α is made in the case of the passive galaxies, despite the much larger number (2.5 times) of galaxies stacked, as compared to the line emitter sample (H α measured with a peak of $5.3 \mu\text{Jy}$, with a $\sigma_{\text{rms}} = 2.1 \mu\text{Jy}$). The velocity position of the putative peak ($\sim 100 \text{ km s}^{-1}$), highly offset from 0, is likely a spurious peak and also indicative of a non-detection of H α . The high offset is highly unlikely to be caused by stripping, given we are averaging across 150 passive galaxies within the cluster, which have random motions in the cluster potential. In some of the stacks (e.g. emission cluster galaxy stack and the emission line field galaxy stack), there are additional peaks off-centred from the 0-velocity. In theory, additional peaks in cluster stack, for example, could be caused by tidally stripped tails, pointing in the same redshift direction, such that they add coherently when stacked. However, this is highly unlikely as the cluster galaxies are moving on a range of orbits within the cluster potential. Even if H α tails existed, they would have a variety of orientations. When stacking the galaxies, the tails would therefore not add coherently. Therefore, we believe these to peaks to be caused by noise variations and low-level systematics. Note that at higher rest-frame velocities, the number of sources for which data exist dwindles (as shown in Fig. 4). For example, at $\pm 600 \text{ km s}^{-1}$, the number of sources with data in that velocity channel is already half that at velocity 0. That means that the noise at larger rest-frame velocities will be higher than the noise around the 0-velocity. Therefore the significance of off-centre peaks is actually very low (at least a factor of $\sqrt{2}$ lower than if located at 0-velocity).

The stack using all the galaxies reaches a very low noise level of $3.1 \mu\text{Jy}$. The H α signal peaks at a significance of $4.6\sigma_{\text{rms}}$, value mainly due to the contribution of SF and AGN galaxies.

3.5 H α masses

We use the following relation to convert from radio flux S into H α mass ($M_{\text{H}\alpha}$; Wentzel & van Woerden 1959; Roberts 1962):

$$\frac{M_{\text{H}\alpha}}{M_{\odot}} = \frac{236}{1+z} \left(\frac{D_L}{\text{Mpc}} \right)^2 \left(\frac{\int S_V \text{d}V}{\text{mJy km s}^{-1}} \right), \quad (2)$$

where M_{\odot} is the mass of the Sun, $z = 0.192$ is the mean redshift of the sample of galaxies, $D_L = 940.7 \text{ Mpc}$ is the luminosity distance




Figure 6. Final H I stacks after filtering and correcting for continuum oversubtraction. The solid black lines show the galaxy stacks and the grey lines show the corresponding noise properties. The thin, solid red lines show the Gaussian fits to the profiles located around a 0 rest-frame velocity. The vertical, dashed red line shows the position of the peak, which is labelled with its significance based on the underlying noise properties. The rms noise is calculated using only the data between the vertical, dashed grey lines, where at least 80 per cent of the peak number of sources are stacked. Note clear detections of H I are made for the line emitters, while no detections are made in the case of the cluster passive galaxies.

at that redshift and $\int S_V dV$ is the average of the H I emission over a rest-frame velocity range. As mentioned in Section 3.2, the velocity is defined as $V = (v_{\text{HI}} - v_{\text{observed}})/v_{\text{HI}}$.

For the stacks with H I detections (Fig. 6), the H I mass is averaged over $2\sigma_{\text{Gauss}}$ (two times the Gaussian dispersion) on either side of the peak position. For the passive population, we use the range within 200 km s^{-1} from 0-velocity position. The error in the H I mass is calculated by propagating the rms error through equation (2), using the same velocity range used for the integration of the signal.

Note that even though we use the filtered stacks to calculate the H I masses, similar results would be obtained even if the original data are used. Although the quality of the spectra improve, the errors on the final H I mass do not heavily depend on the filtering. As shown in equation (2), the mass is effectively an integral over the profile. Hence, the effects of SG filtering are reduced because

averaging over a velocity range equivalent to smoothing down to a resolution equal to that velocity range.

We find that the average H I mass for the emission line cluster galaxies ($M_{\text{HI}} = (2.50 \pm 0.62) \times 10^9 M_{\odot}$) is a factor of 1.3 higher than the mass of cold neutral gas in their field counterparts ($M_{\text{HI}} = (1.86 \pm 1.20) \times 10^9 M_{\odot}$). However the difference is not significant. Additionally, the cluster galaxies are on average ~ 1.5 times more massive than their field counterparts (see also Table 5). The average stellar masses (M_{\star}) are calculated over the same galaxies stacked for the H I analysis and the error reported is the standard deviation of the sample. Therefore, when accounting for the differences in stellar masses between the cluster and the field galaxies, the cluster line emitters are consistent with being as gas rich as the field counterparts. The fraction of neutral atomic gas to stellar mass M_{HI}/M_{\star} is 0.39 ± 0.26 and 0.34 ± 0.09 for field and cluster galaxies, respectively. This is a surprising result, as cluster

Table 5. Optical properties of each galaxy stack. The average $H\alpha$ luminosities for each stack are calculated from data from Stroe et al. (2015). The SFRs are calculated using the Kennicutt (1998) conversion. Stellar masses are obtained using the method from Sobral et al. (2015).

Sample	Number	$\log_{10}(L_{H\alpha})$ (erg s $^{-1}$)	SFR (M $_{\odot}$ yr $^{-1}$)	M_{\star} (10 9 M $_{\odot}$)
Emission line, field	22	41.53	1.49	4.8 ± 0.8
Emission line, cluster	45	41.45	1.23	7.4 ± 0.5
Passive, cluster	154	<40.9	<0.35	25.6 ± 0.5

galaxies have been found to contain less H i than galaxies in the field (e.g. Solanes et al. 2001).

The H i masses of the directly detected sources (Table 3) are in the same range as the values for the average emission line stacks (Table 4). However, we do not directly detect any of the emission line systems. This indicates that we are biased towards detecting face-on sources, while the galaxies used for stacking have random orientation (with the bulk being oblique to edge on).

Within our noise limits, we do not detect any significant H i within the passive cluster galaxy population ($M_{\text{Hi}} = (0.21 \pm 0.15) \times 10^9 \text{ M}_{\odot}$). The passive population has at least nine times less H i gas than field line emitters (although not statistically significant) and about 12 times less than the cluster star-forming galaxies and AGNs (2.8 σ significance, where σ is calculated as the errors on the passive galaxy mass and the cluster line emitter mass, added in quadrature). Note, however, that the velocity range used for the integration of the H i signal is larger for the passive cluster members than what was used for the line emitters. The cluster emission line systems have on average much lower masses compared to the passive galaxies. Therefore, the ratio of H i to stellar mass for the passive cluster galaxies is less than 0.008, a factor of ∼30 times lower than the cluster line emitters. Taking into account the errors on the gas fractions for the two populations summed in quadrature, the difference between the gas fraction in passive and cluster active galaxies is significant at the 4 σ level.

3.6 H α –H i correlation

We compare the amount of cold gas and ionized content in each galaxy stack by investigating their H α ($L_{H\alpha}$) and H i (L_{Hi}) luminosities.

The H i luminosity is calculated from the peak flux. The H α luminosity is calculated from the H α flux estimated from the NB observations (see Section 2.2), after correcting for extinction by Galactic dust, as well as for 1 mag for intrinsic dust attenuation (Sobral et al. 2012) within each galaxy. We also remove the contribution of the adjacent [N II] line from the line flux (for details see Stroe et al. 2014a, 2015). We average the corrected H α luminosities

for galaxies within each stack to obtain a mean value for the ionized gas content as function of galaxy type.

The luminosities are calculated in the following way:

$$L_{\text{Hi}, H\alpha} = 4\pi d_L F_{\text{Hi}, H\alpha}, \quad (3)$$

where F is the H α total flux and H i peak flux, respectively, and $d_L = 940$ Mpc is the luminosity distance at the redshift of the cluster.

The H α can be converted into a SFR value using the relationship from Kennicutt (1998), which we correct for the Chabrier (2003) IMF, according to Salim et al. (2007):

$$\frac{\text{SFR}}{\text{M}_{\odot} \text{yr}^{-1}} = \frac{4.4 \times 10^{-42} L_{H\alpha}}{\text{erg s}^{-1}}. \quad (4)$$

Note that not all H i stacked galaxies have an H α flux measurement (see Table 6 for numbers). This may be because the sources have too faint H α line fluxes, below the limits of our NB H α survey. To test how the full H i sample differs from the H i sample with H α measurements, we followed the procedure outlined in Sections 3.2–3.4 and stacked only the H i sources with H α measurements. We find that the peak H i fluxes and the average H i masses for subsamples with H α counterparts matches their parent sample within the error. Therefore, the subsamples with H α measurements are representative of the parent sample. Given the more robust measurement of the average H i properties for the full H i samples (driven by the higher number statistics), in comparing the H α and H i properties, we use average H i properties derived for the full samples.

As Fig. 7 shows, the H α line emitters tend to be more luminous in H i . This is equivalent to galaxies which are more star forming possessing larger reservoirs of atomic gas (Fig. 8).

3.7 H α –radio correlation

We extract sources from the Very Large Array (VLA) 1.5 GHz image using PYBDSM at the positions of the passive and H α line emitter galaxies with spectra (for number of sources see Table 6). The software detects single sources as islands and fits the flux distribution with Gaussians and also calculates the background noise levels using emission-free regions of the sky nearby each source. We assign a source a radio flux density by summing up the flux from all Gaussians belonging to its island. We cross-match radio sources with optical counterparts in our optical spectroscopic catalogue, using a maximum search radius of 5 arcsec, to account for the positional accuracy of the optical and radio images as well as any extent the radio sources may have.

In case a source is not detected in the radio map, we assign it an upper limit flux equivalent to 3 σ_{rms} , where the σ_{rms} is calculated from the noise level at the position of the source. Note that the FOV

Table 6. As Table 5 but only for sources with H α measurements. Optical and BB radio properties of the sources in the line emitter and passive stacks. The average H α luminosities for each stack are calculated from data from Stroe et al. (2015). The SFRs are calculated using the Kennicutt (1998) conversion, with a Chabrier (2003) IMF.

Sample	Number H i sources with H α counterparts	$\log_{10}(L_{H\alpha})^a$ (erg s $^{-1}$)	SFR a (M $_{\odot}$ yr $^{-1}$)	Number H α sources	Number H α sources with radio counterparts	$\log_{10}(L_{H\alpha})^b$ (erg s $^{-1}$)
Emission line, field	20 (~91 per cent)	41.53	1.49	39	3 (~8 per cent)	41.30
Emission line, cluster	29 (~65 per cent)	41.45	1.23	54	24 (~44 per cent)	41.20
Passive, cluster	79 (~51 per cent)	<40.9	<0.35	90	11 (~12 per cent)	40.15

a Average over the H i stacked sources with H α counterparts.

b Average over all spectroscopic sources with H α measurements.




Figure 7. Relationship between H α luminosity and peak H I luminosity. As expected the cluster passive galaxies, selected to be non-line emitters contain less H I than emission line galaxies. Line emitters are selected to be H α luminous, indicating the presence of vigorous SF and/or AGN activity.




Figure 8. Relationship between SFR and integrated H I mass. All SFRs are calculated with a Chabrier IMF. For the passive population, the SFR is an upper limit. Overplotted are the data points for cluster Abell 370 at $z = 0.37$ (Lah et al. 2009) and a blank field at $z = 0.24$ (Lah et al. 2007), together with the SFR- M_{HI} relationship for the local Universe (Doyle & Drinkwater 2006). Note that it is difficult to compare our data set with other work because of the different ways of measuring the H I mass and the different SF tracers used (H α , [O II] or IR).

of the VLA image is large enough (FWHM of ~ 30 arcmin) that it covers all the optical source positions.

We calculate observed 1.4 GHz measurements from the 1.5 GHz values assuming a -0.7 radio spectral index value, and then convert the values to rest frame 1.4 GHz measurements.

A plot showing the relationship between the H α luminosities and 1.5 GHz luminosities (calculated using equation 3) can be found in Fig. 9. The fluxes of the radio counterparts and their morphologies can be found in Appendix A in Table A1. The emission line galaxies have, on average, 1.5 GHz luminosities 1–2 orders of magnitude lower than the passive galaxies. Interestingly, even though the emission line cluster members have similar H α luminosities, and hence




Figure 9. Relationship between H α luminosity and rest frame 1.4 GHz radio luminosity. The vertical black dashed line marks the rms noise at the centre of the radio image, while the grey line represents the detection limit towards the edges of the FOV. The horizontal, solid, black line indicates the detection limit of the H α NB survey. Passive galaxies, undetected in H α , are only plotted if they have a detection in the radio. Error bars are plotted but in most cases they are smaller than the symbol size. On average, the galaxies undergoing SF and optical AGN episodes have lower radio luminosities, probably driven by SNRs, while the passive cluster galaxies host powerful jetted and tailed radio sources. It is noteworthy that, in the case of line emission sources with radio detections the H α luminosity correlates well with the radio luminosity. We overplot for comparison data points for field galaxies at $z = 0.24$ from Lah et al. (2007) and the linear correlation between H α and radio luminosity for local galaxies found by Sullivan et al. (2001).

SFRs, to the field line emitters, their radio BB detection rate is a factor of >5 higher. However, for all emission line sources with radio detections, the H α luminosity correlates with the amount of radio emission.

4 DISCUSSION

Despite being extremely massive ($M_{200} \sim 2 \times 10^{15} M_{\odot}$; Dawson et al. 2015; Jee et al. 2015) and hot ($T = 6\text{--}12$ keV; O’Graeán et al. 2013), the ‘Sausage’ merging cluster hosts numerous massive, H α emission line galaxies displaying elevated levels of SF and AGN activity, outflows, high metallicities and low electron densities compared to galaxies in the field (Sobral et al. 2015; Stroe et al. 2015). In the present study, we find that the emission line cluster galaxies have similar H I neutral gas fraction as the field galaxies. The data reveal linear correlations of the H I , H α line emission and radio BB luminosity (Figs 7–8). By combining tracers of SF on different time-scales, H I , H α and radio BB data, we can understand the circumstances under which the elevated activity can be triggered and also the possible future evolution of the SF properties in the cluster galaxies.

4.1 H I and H α – tracing the gas that fuels future star formation episodes

We make a clear detection of H I for the emission line cluster galaxies giving an average mass of $(2.50 \pm 0.62) \times 10^9 M_\odot$, while the average H I mass for the field galaxies is $(1.86 \pm 1.20) \times 10^9 M_\odot$. Stroe et al. (2015) and Sobral et al. (2015) find that the stellar masses of H α cluster galaxies are on average higher than their field counterparts. For the samples used in the H I stacks, the cluster galaxies are about 1.5 times more massive than the field line emitters (see Table 6).

Note that the cluster and field line emitters are selected in the same way and that the spectroscopic samples are representative of their parent samples (see Sobral et al. 2015). Therefore the cluster and field emission line galaxy samples within the ‘Sausage’ field are fully comparable.

As mentioned in Section 1, previous studies of the H I content in cluster galaxies find that star-forming galaxies become increasingly H I deficient towards cluster cores, when controlling for stellar mass or optical disc size (e.g. Solanes et al. 2001; Lah et al. 2007; Verheijen et al. 2007). Contrary to previous work in the field, we find that our emission line cluster galaxies are as gas rich as their field counterparts, despite the two samples having similar H α luminosities and hence similar SFR (see Table 5 and Fig. 7).

The studies of Lah et al. (2007, 2009) indicate that a cluster at $z \sim 0.37$ and a blank field at $z \sim 0.24$ follow a similar relationship between SFR and H I mass to local field galaxies (Doyle & Drinkwater 2006). Even though our galaxies show evidence for a correlation between the amount of H α emission (or the SFR) and the H I mass, both the emission line and passive galaxies do not follow the Doyle & Drinkwater (2006) relationship (see Fig. 8). The H I masses of our sample are $>5\sigma$ away from the masses predicted by the relationship at the same SFR. This could be entirely driven by the different SF tracers used in the different studies [Doyle & Drinkwater (2006) use IR data, Lah et al. (2009) use [O II] and Lah et al. (2007) use the H α emission line] or the spatial or velocity range over which the H I signal was integrated over.

Given the massive cluster galaxies may reside in massive dark matter haloes, they could have retained their cold H I gas more easily during the cluster merger. Interestingly, while spiral galaxies in the Virgo Cluster are highly H I deficient, they are not deficient in molecular gas (Kenney & Young 1986, 1989; Stark et al. 1986). The authors attribute this to the preferential stripping of low-density gas located at the galaxy outskirts, therefore, not affecting dense molecular gas located towards the galaxy centre. Therefore, in the case of the ‘Sausage’ line emitters, with little to no ram pressure stripping of neutral and molecular gas, the larger reservoirs could fuel increased SFR in the cluster galaxies. If the cluster galaxies maintain their current average level of SF ($\sim 1.23 M_\odot \text{ yr}^{-1}$, see Table 6), and assuming 100 per cent efficiency in converting cold gas into stars, the H I reservoir would be depleted in ~ 2.0 Gyr. If we assume a molecular gas content equal to the H I mass, it would take about ~ 4.0 Gyr to consume the gas. However, as shown in Sobral et al. (2015), the cluster galaxies also lose gas through outflows. Assuming a maximal mass outflow rate similar to the SFR (Förster Schreiber et al. 2009; Genzel et al. 2011), the H I gas will have been used up in about ~ 1.0 Gyr (~ 2.0 Gyr if we include molecular gas). Assuming a more realistic outflow rate of about 0.1–0.5 SFR (as observed by Swinbank et al. 2015), the H I gas would be depleted in 1.35–1.85 Gyr, or 2.7–3.7 Gyr if molecular gas is considered. This is in line with calculations from Stroe et al. (2015) where the molecular gas content was esti-

mated using the total stellar mass, but atomic gas was not taken into account.

4.2 Radio broad data – tracing the SN emission

As shown by Sullivan et al. (2001), synchrotron radio emission in star-forming galaxies is generated in supernova remnants (SNRs). Given the time required for a $7\text{--}8 M_\odot$ star to evolve to the red giant phase and undergo core-collapse, SNRs are good tracers of SF episodes happening ~ 100 Myr ago (Condon 1992).

In the case of H α luminous cluster galaxies, undergoing strong SF and optical AGN activity, the H α luminosity correlates well with the radio BB continuum (see Fig. 9 and Table A1 in Appendix A). The values fall on the same correlation as the large sample (~ 350) of typical $z = 0.24$, H α luminous field galaxies studied by Lah et al. (2007) and follow the tight relationship found by Sullivan et al. (2001) for local field galaxies. For both samples, the galaxies classified as purely star forming or as hosting an optical AGNs follow the H α –radio correlation. However, the AGN-dominated cluster galaxies are expected to possess reasonable amounts of SF, as indicated by the spiral arm patterns in many of their hosts (for images see Sobral et al. 2015). Therefore, the sample includes photoionized broad line and narrow line regions (Sobral et al. 2015) hosted by spiral galaxies, constituting examples of Seyfert galaxies. Even though some of the cluster galaxies are currently dominated by AGN, they have undergone significant SF activity in the past.

Despite their similar average H α luminosities, the fraction of H α luminous cluster members with radio counterparts is a factor of >5 higher than their field counterparts, down to a similar sensitivity limit (see Table 6 and Fig. 9). This is consistent with results from Sobral et al. (2015) where they found evidence for strong outflows, probably driven by SN, only in the cluster galaxies and not in their field counterparts. Therefore, cluster galaxies have been undergoing SF for at least 100 Myr, while field galaxies have been relatively inactive or less active in the past. Increased SF episodes tens to a few hundred Myr ago triggered by the cluster merger and its associated shock would also lead to higher SN rates. Field galaxies have not undergone any interaction with the shock front, hence in their case, there was no trigger for SN.

Overall, line emission galaxies both inside and outside the ‘Sausage’ cluster follow the local relationship between H α and radio emission. Field emission line galaxies from a larger sample at $z \sim 0.24$ (~ 150 field galaxies; Lah et al. 2007) fall on the same relation. This indicates that the relationship between H α and BB radio emission does not evolve from $z \sim 0.2$ to the present and that it does not depend on environment. The stellar populations in all galaxies that have been undergoing SF for longer periods (~ 100 Myr), irrespective of redshift or environment, seem to evolve similarly from the massive, short-lived stars which are responsible for producing the bulk of H α emission to the slightly less massive stars whose explosions dominate the SN population. These results indicate that the star formation history for the ‘Sausage’ cluster galaxies is relatively constant, without any strong recent (~ 10 Myr) bursts of star formation or in the past 100 Myr.

A little bit over 10 per cent of the passive galaxies have a radio BB counterpart, a similar rate to field emission line galaxies, much four times lower than the cluster line emitters. The passive cluster members that have radio counterparts are giant ellipticals hosting radio jets and tails as indicated by the radio morphologies and luminosities. By contrast to the emission line galaxies, where the radio emission is most likely produced by SNR, the radio emission

in the elliptical galaxies traces shock-accelerated electrons in the jets and their back-flow (see also Appendix A).

4.3 Relationship to cluster merger state and shocks

As the radio BB and H α data indicate, the ‘Sausage’ cluster galaxies have been undergoing intense SF and AGN activity for at least 100 Myr and this is likely to last a further ~ 1 Gyr, given the large reservoirs of neutral hydrogen. The SF can last for another ~ 2 Gyr if comparable amounts of molecular gas are present. However, in the field galaxies around the cluster, we only find evidence of very recent SF episodes (~ 10 Myr). Despite the comparable amounts of H α , the cluster galaxies therefore underwent a significant event triggering SF about ~ 100 Myr ago, evolving differently than their field counterparts. The most significant difference between the cluster and field line emitters is the cluster merger and the passage of the merger-induced shock waves only affected the cluster members.

The ‘Sausage’ cluster is a result of a massive merger about 0.5 Gyr ago (e.g. van Weeren et al. 2011), which produced shocks travelling through the ICM at about 2500 km s $^{-1}$ (Stroe et al. 2014c). As the H α data indicate, the massive cluster members seem to have retained most of their neutral gas during the merger. Given their travelling speed, we expect the shocks to have traversed most cluster galaxies about 100–300 Myr ago. The SF time-scale imposed by the radio and optical tracers therefore matches well with the cluster merger time line. Our results fully support the interpretation previously proposed by Stroe et al. (2015) and Sobral et al. (2015), where the cluster merger induced shocks trigger gas collapse as they traverse the gas-rich cluster galaxies. This interpretation is also supported by simulations (Roediger et al. 2014).

5 CONCLUSIONS

We presented deep H α observations combined with H α and BB radio data to study the past, present and future SF activity in the ‘Sausage’ merging cluster. Our main results are the following.

(i) The cluster H α emission line galaxies (star forming and radio-quiet broad and narrow line AGN), selected down to the same SFR limit, have as much H α gas ($(2.50 \pm 0.62) \times 10^9 M_\odot$) as the field counterparts around the cluster ($(1.84 \pm 1.20) \times 10^9 M_\odot$), when accounting for the different stellar masses of the two samples. This indicates the massive cluster line emitters retained their gas during the cluster merger.

(ii) A stringent upper limit is placed on the average H α content of the passive galaxies in the ‘Sausage’ cluster: $M_{H\alpha} = (0.21 \pm 0.15) \times 10^9 M_\odot$. The ratio of H α to stellar mass for the passive galaxies is almost 40 times less than for cluster line emitters (significant at 4σ level).

(iii) If the present SF and outflow rate is maintained in the emission line cluster galaxies, their H α reservoirs will be depleted in ~ 0.75 –1.0 Gyr.

(iv) A large fraction of the emission line cluster galaxies have radio BB detections, indicating the presence of SNR. These sources have been therefore undergoing vigorous SF for at least 100 Myr.

(v) The relationship between H α and radio continuum emission shows no evolution from $z \sim 0.2$ to the present and also no dependence on environment.

Our H α observations represent an important milestone in the study of the ‘Sausage’ cluster SF history. The member galaxies are gas rich (gas to stellar mass ratio of ~ 0.34) and thus capable

of sustaining the increased SF and AGN activity measured in the cluster.

ACKNOWLEDGEMENTS

We would like to thank the referee for comments which greatly improved the clarity of the paper. We also thank Leah Morabito for useful discussions. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research has made use of NASA’s Astrophysics Data System. AS and HJAR acknowledge financial support from an NWO top subsidy (614.001.006). Part of this work performed under the auspices of the US DOE by LLNL under Contract DE-AC52-07NA27344. DS acknowledges financial support from the Netherlands Organisation for Scientific Research (NWO) through a Veni fellowship, from FCT through a FCT Investigator Starting Grant and Start-up Grant (IF/01154/2012/CP0189/CT0010) and from FCT grant PEst-OE/FIS/UI2751/2014. RvW is supported by NASA through the Einstein Postdoctoral grant number PF2-130104 awarded by the Chandra X-ray Center, which is operated by the Smithsonian Astrophysical Observatory for NASA under contract NAS8-03060. The Westerbork Synthesis Radio Telescope is operated by the Netherlands Institute for Radio Astronomy (ASTRON) with support from the Netherlands Foundation for Scientific Research (NWO). The Isaac Newton and William Herschel telescopes are operated on the island of La Palma by the Isaac Newton Group in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. Some of the data presented herein were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation. Based in part on data collected at Subaru Telescope, which is operated by the National Astronomical Observatory of Japan. Based in part on observations from the Karl G. Jansky Very Large Array, operated by the National Radio Astronomy Observatory, a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

REFERENCES

- Akamatsu H., Kawahara H., 2013, PASJ, 65, 16
- Andersen V., Owen F. N., 1995, AJ, 109, 1582
- Balogh M. L., Schade D., Morris S. L., Yee H. K. C., Carlberg R. G., Ellingson E., 1998, ApJ, 504, L75
- Bolatto A. D., Wolfire M., Leroy A. K., 2013, ARA&A, 51, 207
- Briggs D. S., 1995, PhD thesis, New Mexico Tech, USA
- Carilli C. L., Walter F., 2013, ARA&A, 51, 105
- Cayatte V., van Gorkom J. H., Balkowski C., Kotanyi C., 1990, AJ, 100, 604
- Cedrés B., Iglesias-Páramo J., Vilchez J. M., Reverte D., Petropoulou V., Hernández-Fernández J., 2009, AJ, 138, 873
- Chabrier G., 2003, PASP, 115, 763
- Chengalur J. N., Braun R., Wieringa M., 2001, A&A, 372, 768
- Chung A., van Gorkom J. H., Kenney J. D. P., Crowl H., Vollmer B., 2009, AJ, 138, 1741
- Chung S. M., Gonzalez A. H., Clowe D., Markevitch M., Zaritsky D., 2010, ApJ, 725, 1536
- Condon J. J., 1992, ARA&A, 30, 575

Cornwell T. J., Golap K., Bhatnagar S., 2005, in Shopbell P., Britton M., Ebert R., eds, ASP Conf. Ser. Vol. 347, Astronomical Data Analysis Software and Systems XIV. Astron. Soc. Pac., San Francisco, p. 86

Cornwell T. J., Golap K., Bhatnagar S., 2008, IEEE J. Selected Topics Signal Processing, 2, 647

Darvish B., Sobral D., Mobasher B., Scoville N. Z., Best P., Sales L. V., Smail I., 2014, *ApJ*, 796, 51

Dawson W. A. et al., 2015, *ApJ*, 805, 143

Doyle M. T., Drinkwater M. J., 2006, *MNRAS*, 372, 977

Dressler A., 1980, *ApJ*, 236, 351

Ensslin T. A., Biermann P. L., Klein U., Kohle S., 1998, *A&A*, 332, 395

Ferrari C., Benoist C., Maurogordato S., Cappi A., Slezak E., 2005, *A&A*, 430, 19

Finn R. A. et al., 2005, *ApJ*, 630, 206

Förster Schreiber N. M. et al., 2009, *ApJ*, 706, 1364

Fumagalli M., Fossati M., Hau G. K. T., Gavazzi G., Bower R., Sun M., Boselli A., 2014, *MNRAS*, 445, 4335

Gavazzi G., Jaffe W., 1985, *ApJ*, 294, L89

Gavazzi G., Catinella B., Carrasco L., Boselli A., Contursi A., 1998, *AJ*, 115, 1745

Genzel R. et al., 2011, *ApJ*, 733, 101

Goto T., Yamauchi C., Fujita Y., Okamura S., Sekiguchi M., Smail I., Bernardi M., Gomez P. L., 2003, *MNRAS*, 346, 601

Gunn J. E., Gott J. R., III, 1972, *ApJ*, 176, 1

Hudson D. S., Mittal R., Reiprich T. H., Nulsen P. E. J., Andernach H., Sarazin C. L., 2010, *A&A*, 513, A37

Hwang H. S., Lee M. G., 2009, *MNRAS*, 397, 2111

Jee M. J. et al., 2015, *ApJ*, 802, 46

Johnston-Hollitt M., Sato M., Gill J. A., Fleenor M. C., Brick A.-M., 2008, *MNRAS*, 390, 289

Katayama H., Hayashida K., Takahara F., Fujita Y., 2003, *ApJ*, 585, 687

Kenney J. D., Young J. S., 1986, *ApJ*, 301, L13

Kenney J. D. P., Young J. S., 1989, *ApJ*, 344, 171

Kennicutt R. C., Jr, 1998, *ARA&A*, 36, 189

Kocevski D. D., Ebeling H., Mullis C. R., Tully R. B., 2007, *ApJ*, 662, 224

Koyama Y. et al., 2013, *MNRAS*, 434, 423

Lah P. et al., 2007, *MNRAS*, 376, 1357

Lah P. et al., 2009, *MNRAS*, 399, 1447

Larson R. B., Tinsley B. M., Caldwell C. N., 1980, *ApJ*, 237, 692

Leroy A. K., Walter F., Brinks E., Bigiel F., de Blok W. J. G., Madore B., Thornley M. D., 2008, *AJ*, 136, 2782

Magri C., Haynes M. P., Forman W., Jones C., Giovanelli R., 1988, *ApJ*, 333, 136

Mann A. W., Ebeling H., 2012, *MNRAS*, 420, 2120

Miller N. A., Owen F. N., 2003, *AJ*, 125, 2427

Moore B., Katz N., Lake G., Dressler A., Oemler A., 1996, *Nature*, 379, 613

Morabito L. K. et al., 2014, *ApJ*, 795, L33

Morrison G. E., 1999, PhD thesis, Univ. New Mexico

Morrison G. E., Owen F. N., 2003, *AJ*, 125, 506

Offringa A. R., de Bruyn A. G., Biehl M., Zaroubi S., Bernardi G., Pandey V. N., 2010, *MNRAS*, 405, 155

Ogreal G. A., Brüggen M., Röttgering H., Simionescu A., Croston J. H., van Weeren R., Hoeft M., 2013, *MNRAS*, 429, 2617

Ogreal G. A., Brüggen M., van Weeren R., Röttgering H., Simionescu A., Hoeft M., Croston J. H., 2014, *MNRAS*, 440, 3416

Oosterloo T., van Gorkom J., 2005, *A&A*, 437, L19

Owen F. N., Ledlow M. J., Keel W. C., Wang Q. D., Morrison G. E., 2005, *AJ*, 129, 31

Owers M. S., Couch W. J., Nulsen P. E. J., Randall S. W., 2012, *ApJ*, 750, L23

Padovani P., Miller N., Kellermann K. I., Mainieri V., Rosati P., Tozzi P., 2011, *ApJ*, 740, 20

Poggianti B. M., Bridges T. J., Komiyama Y., Yagi M., Carter D., Mobasher B., Okamura S., Kashikawa N., 2004, *ApJ*, 601, 197

Rau U., Cornwell T. J., 2011, *A&A*, 532, A71

Rawle T. D. et al., 2012, *ApJ*, 756, 106

Roberts M. S., 1962, *AJ*, 67, 437

Roediger E., Brüggen M., Owers M. S., Ebeling H., Sun M., 2014, *MNRAS*, 443, L114

Salim S. et al., 2007, *ApJS*, 173, 267

Sanderson A. J. R., Edge A. C., Smith G. P., 2009, *MNRAS*, 398, 1698

Sault R. J., 1994, *A&AS*, 107, 55

Sault R. J., Teuben P. J., Wright M. C. H., 1995, in Shaw R. A., Payne H. E., Hayes J. J. E., eds, ASP Conf. Ser. Vol. 77, Astronomical Data Analysis Software and Systems IV. Astron. Soc. Pac., San Francisco, p. 433

Savitzky A., Golay M. J. E., 1964, *Analytical Chem.*, 36, 1627

Scott T. C., Cortese L., Brinks E., Bravo-Alfar H., Auld R., Minchin R., 2012, *MNRAS*, 419, L19

Sobral D., Best P. N., Smail I., Geach J. E., Cirasuolo M., Garn T., Dalton G. B., 2011, *MNRAS*, 411, 675

Sobral D., Best P. N., Matsuda Y., Smail I., Geach J. E., Cirasuolo M., 2012, *MNRAS*, 420, 1926

Sobral D., Stroe A., Dawson W. A., Wittman D., Jee M. J., Röttgering H., van Weeren R. J., Brüggen M., 2015, *MNRAS*, 450, 630

Solanes J. M., Manrique A., García-Gómez C., González-Casado G., Giovanelli R., Haynes M. P., 2001, *ApJ*, 548, 97

Stark A. A., Knapp G. R., Bally J., Wilson R. W., Penzias A. A., Rowe H. E., 1986, *ApJ*, 310, 660

Stroe A., van Weeren R. J., Intema H. T., Röttgering H. J. A., Brüggen M., Hoeft M., 2013, *A&A*, 555, A110

Stroe A., Sobral D., Röttgering H. J. A., van Weeren R. J., 2014a, *MNRAS*, 438, 1377

Stroe A. et al., 2014b, *MNRAS*, 441, L41

Stroe A., Harwood J. J., Hardcastle M. J., Röttgering H. J. A., 2014c, *MNRAS*, 445, 1213

Stroe A. et al., 2015, *MNRAS*, 450, 646

Sullivan M., Mobasher B., Chan B., Cram L., Ellis R., Treyer M., Hopkins A., 2001, *ApJ*, 558, 72

Swinbank M. et al., 2015, *MNRAS*, submitted

Umeda K. et al., 2004, *ApJ*, 601, 805

van Weeren R. J., Röttgering H. J. A., Brüggen M., Hoeft M., 2010, *Science*, 330, 347

van Weeren R. J., Brüggen M., Röttgering H. J. A., Hoeft M., 2011, *MNRAS*, 418, 230

Verheijen M., van Gorkom J. H., Szomoru A., Dwarakanath K. S., Poggianti B. M., Schiminovich D., 2007, *ApJ*, 668, L9

Wegner G. A., Chu D. S., Hwang H. S., 2015, *MNRAS*, 447, 1126

Wentzel D. G., van Woerden H., 1959, *Bull. Astron. Inst. Netherlands*, 14, 335

APPENDIX A: RADIO FLUXES AND MORPHOLOGIES

We tabulate the flux values of the 1.5 GHz VLA counterparts to the spectroscopic sources in Table A1. We also describe the morphology of the radio sources. Passive galaxies are hosts to jetted radio AGN, mainly pushed in wide or narrow angle tail morphologies by the interaction with the ICM. Emission line galaxies have mostly disc or spiral-like morphologies, indicating the radio emission is coming from star formation.

Table A1. The radio fluxes, errors and morphologies of the VLA 1.5 GHz counterparts to the spectroscopic sources. The same sources tabulated here are plotted in Fig. 9.

RA ($^{\circ}$)	Dec. ($^{\circ}$)	Flux (mJy)	Error (mJy)	Morphology
Passive, cluster				
340.6978	53.0939	15.034	0.022	NAT
340.7687	53.1234	0.023	0.006	Unresolved
340.6048	52.9719	6.304	0.033	WAT
340.7073	53.0401	0.034	0.007	Unresolved
340.8293	53.1228	12.420	0.094	WAT
340.7072	53.0081	5.109	0.125	Unresolved
340.7051	53.0920	0.160	0.013	Unresolved
340.7191	53.0806	90.106	0.090	Tailed
340.7132	53.0139	8.084	0.053	WAT
340.7385	53.1344	0.124	0.008	Unresolved
340.7720	52.9552	0.103	0.010	Unresolved
Emission line, field				
340.3917	53.0163	0.194	0.014	Unresolved
340.6917	52.8561	0.404	0.025	Unresolved
340.9183	52.8727	0.097	0.013	Unresolved
Emission line, cluster				
340.4575	53.0954	0.138	0.011	Unresolved
340.5930	53.0083	0.029	0.006	Unresolved
340.6151	52.9683	0.185	0.011	Unresolved
340.6518	53.0904	0.073	0.007	Unresolved
340.6598	53.0138	0.100	0.010	Unresolved
340.6642	52.9674	0.361	0.012	Unresolved
340.6711	52.9746	0.817	0.026	Disc
340.7136	52.9061	0.133	0.015	Disc
340.7225	52.9517	0.080	0.009	Unresolved
340.7369	52.9439	0.439	0.015	Disc
340.7401	53.0874	0.336	0.022	Disc
340.7475	53.0889	0.179	0.017	Disc
340.7502	53.0439	0.139	0.010	Unresolved
340.7796	52.9257	0.122	0.013	Disc
340.7870	53.0902	1.207	0.018	Unresolved
340.7941	53.0716	0.135	0.011	Disc
340.7985	53.0764	0.055	0.008	Unresolved
340.8037	53.0029	0.285	0.016	Disc
340.9204	53.0778	0.344	0.016	Disc

Notes. NAT: narrow angle tailed galaxy; WAT: wide angle tailed galaxy;
disc: galaxies with a disc or spiral-like morphology.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.