

The accretion history of dark matter haloes – II. The connections with the mass power spectrum and the density profile

Camila A. Correa,¹★ J. Stuart B. Wyithe,¹ Joop Schaye² and Alan R. Duffy^{1,3}

¹*School of Physics, University of Melbourne, Parkville, Victoria 3010, Australia*

²*Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, the Netherlands*

³*Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Melbourne, Victoria 3122, Australia*

Accepted 2015 March 26. Received 2015 March 23; in original form 2015 January 20

ABSTRACT

We explore the relation between the structure and mass accretion histories of dark matter haloes using a suite of cosmological simulations. We confirm that the formation time, defined as the time when the virial mass of the main progenitor equals the mass enclosed within the scale radius, correlates strongly with concentration. We provide a semi-analytic model for halo mass history that combines analytic relations with fits to simulations. This model has the functional form, $M(z) = M_0(1+z)^\alpha e^{\beta z}$, where the parameters α and β are directly correlated with concentration. We then combine this model for the halo mass history with the analytic relations between α , β and the linear power spectrum derived by Correa et al. to establish the physical link between halo concentration and the initial density perturbation field. Finally, we provide fitting formulae for the halo mass history as well as numerical routines. We derive the accretion rate as a function of halo mass, and demonstrate how the halo mass history depends on cosmology and the adopted definition of halo mass.

Key words: methods: numerical – galaxies: haloes – cosmology: theory.

1 INTRODUCTION

Dark matter haloes provide the potential wells inside which galaxies form. As a result, understanding their basic properties, including their formation history and internal structure, is an important step for understanding galaxy evolution. It is generally believed that the halo mass accretion history determines dark matter halo properties, such as their ‘universal’ density profile (Navarro, Frenk & White 1996, hereafter **NFW**). The argument is as follows. During hierarchical growth, haloes form through mergers with smaller structures and accretion from the intergalactic medium. Most mergers are minor (with smaller satellite haloes) and do not alter the structure of the inner halo. Major mergers (mergers between haloes of comparable mass) can bring material to the centre, but they are found not to play a pivotal role in modifying the internal mass distribution (Wang & White 2009). Halo formation can therefore be described as an ‘inside out’ process, where a strongly bound core collapses, followed by the gradual addition of material at the cosmological accretion rate. Through this process, haloes acquire a nearly universal density profile that can be described by a simple formula known as the ‘**NFW** profile’ (**NFW**).

The origin of this universal density profile is not fully understood. One possibility is that the **NFW** profile results from a relaxation

mechanism that produces equilibrium and is largely independent of the initial conditions and merger history (Navarro, Frenk & White 1997). However, another popular explanation, originally proposed by Syer & White (1998), is that the **NFW** profile is determined by the halo mass history, and it is then expected that haloes should also follow a universal mass history profile (Dekel, Devor & Hatzroni 2003; Manrique et al. 2003; Sheth & Tormen 2004; Dalal, Lithwick & Kuhlen 2010; Giocoli, Tormen & Sheth 2012; Salvador-Solé et al. 2012). This universal accretion history was recently illustrated by Ludlow et al. (2013), who showed that the halo mass histories, if scaled to certain values, follow the **NFW** profile. This was done by comparing the mass accretion history, expressed in terms of the critical density of the Universe, $M(\rho_{\text{crit}}(z))$, with the **NFW** density profile, expressed in units of enclosed mass and mean density within r , $M(\langle\rho\rangle(<r))$ at $z = 0$, in a mass-density plane.

In this work, we aim to provide a model that links the halo mass history with the halo *concentration*, a parameter that fully describes the internal structure of dark matter haloes. By doing so, we will gain insight into the origin of the **NFW** profile and its connection with the halo mass history. We also aim to find a physical explanation for the known correlation between the linear *rms* fluctuation of the density field, σ , and halo concentration.

This paper is organized as follows. We briefly introduce our simulations in Section 2, where we explain how we calculated the merger history trees and discuss the necessary numerical convergence conditions. Then we provide a model for the halo mass history, which

* E-mail: correac@student.unimelb.edu.au

Table 1. List of simulations. From left-to-right the columns show: simulation identifier; comoving box size; number of dark matter particles (there are equally many baryonic particles); initial baryonic particle mass; dark matter particle mass; comoving (Plummer-equivalent) gravitational softening; maximum physical softening; final redshift.

Simulation	L (h^{-1} Mpc)	N	m_b ($h^{-1} M_\odot$)	m_{dm} ($h^{-1} M_\odot$)	ϵ_{com} (h^{-1} kpc)	ϵ_{prop} (h^{-1} kpc)	z_{end}
REF_L100N512	100	512 ³	8.7×10^7	4.1×10^8	7.81	2.00	0
REF_L100N256	100	256 ³	6.9×10^8	3.2×10^9	15.62	4.00	0
REF_L100N128	100	128 ³	5.5×10^9	2.6×10^{10}	31.25	8.00	0
REF_L050N512	50	512 ³	1.1×10^7	5.1×10^7	3.91	1.00	0
REF_L025N512	25	512 ³	1.4×10^6	6.3×10^6	1.95	0.50	2
REF_L025N256	25	256 ³	1.1×10^7	5.1×10^7	3.91	1.00	2
REF_L025N128	25	128 ³	8.7×10^7	4.1×10^8	7.81	2.00	2
DMONLY_WMAP5_L400N512	400	512 ³	–	3.4×10^{10}	31.25	8.00	0
DMONLY_WMAP5_L200N512	200	512 ³	–	3.2×10^9	15.62	4.00	0
DMONLY_WMAP5_L100N512	100	512 ³	–	5.3×10^8	7.81	2.00	0
DMONLY_WMAP5_L050N512	50	512 ³	–	6.1×10^7	3.91	1.00	0
DMONLY_WMAP5_L025N512	25	512 ³	–	8.3×10^6	2.00	0.50	0

we refer to as the *semi-analytic model*. This semi-analytic model is described in Section 3, along with an analysis of the formation time definition. For this model we use the empirical McBride, Fakhouri & Ma (2009) formula. This functional form was motivated by EPS theory in a companion paper (Correa et al. 2015a, hereafter Paper I), and we calibrate the correlation between its two parameters (α and β) using numerical simulations. As a result, the semi-analytic model combines analytic relations with fits to simulations, to relate halo structure to the mass accretion history. In Section 3.6 we show how the semi-analytic model for the halo mass history depends on cosmology and the adopted definition for halo mass. In Section 4 we provide a detailed comparison between the semi-analytic halo mass history model provided in this work, and the analytic model presented in Paper I. The parameters in this analytic model depend on the linear power spectrum and halo mass, whereas in the semi-analytic model the parameters depend on concentration and halo mass. We therefore combine the two models to establish the physical relation between the linear power spectrum and halo concentration. We will expand on this in a forthcoming paper (Correa et al. 2015b, hereafter Paper III), where we predict the evolution of the concentration–mass relation and its dependence on cosmology. Finally, we provide a summary of formulae and discuss our main findings in Section 5.

2 SIMULATIONS

In this work we use the set of cosmological hydrodynamical simulations (the REF model) along with a set of dark matter only (DMONLY) simulations from the OWLS project (Schaye et al. 2010). These simulations were run with a significantly extended version of the N-Body Tree-PM, smoothed particle hydrodynamics (SPH) code GADGET3 (last described in Springel 2005). In order to assess the effects of the finite resolution and box size on our results, most simulations were run using the same physical model (DMONLY or REF) but different box sizes (ranging from $25 h^{-1}$ Mpc to $400 h^{-1}$ Mpc) and particle numbers (ranging from 128^3 to 512^3). The main numerical parameters of the runs are listed in Table 1. The simulation names contain strings of the form LxxxNyyy, where xxx is the simulation box size in comoving h^{-1} Mpc, and yyy is the cube root of the number of particles per species (dark matter or baryonic). For more details on the simulations we refer the reader to Appendix A.

Table 2. Cosmological parameters.

Simulation	Ω_m	Ω_Λ	h	σ_8	n_s
DMONLY_WMAP1	0.25	0.75	0.73	0.90	1.000
DMONLY_WMAP3	0.238	0.762	0.73	0.74	0.951
DMONLY_WMAP5	0.258	0.742	0.72	0.796	0.963
DMONLY_WMAP9	0.282	0.718	0.70	0.817	0.964
DMONLY_Planck1	0.317	0.683	0.67	0.834	0.962

Our DMONLY simulations assume the *WMAP5* cosmology, whereas the REF simulations assume *WMAP3*. To investigate the dependence on the adopted cosmological parameters, we include an extra set of five dark matter only simulations ($100 h^{-1}$ Mpc box size and 512^3 dark matter particles) which assume values for the cosmological parameters derived from the different releases of *Wilkinson Microwave Anisotropy Probe (WMAP)* and the *Planck* mission. Table 2 lists the sets of cosmological parameters adopted in the different simulations.

Halo mass histories are obtained from the simulation outputs by building halo merger trees. We define the halo mass history as the mass of the most massive halo (main progenitor) along the main branch of the merger tree. The method used to create the merger trees is described in detail in Appendix A1. While analysing the merger trees from the simulations, we look for a numerical resolution criterion under which mass accretion histories converge numerically. We begin by investigating the minimum number of particles haloes must contain so that the merger trees lead to accurate numerical convergence. We find a necessary minimum limit of 300 dark matter particles, which corresponds to a minimum dark matter halo mass of $M_{\text{halo}} \sim 2.3 \times 10^{11} M_\odot$ in the $100 h^{-1}$ Mpc box, $M_{\text{halo}} \sim 2.6 \times 10^{10} M_\odot$ in the $50 h^{-1}$ Mpc box, and $M_{\text{halo}} \sim 3.4 \times 10^9 M_\odot$ in the $25 h^{-1}$ Mpc box.

In a merger tree, when a progenitor halo contains less than 300 dark matter particles, it is considered unresolved and discarded from the analysis. As a result, the number of haloes in the sample that contribute to the median value of the mass history decreases with increasing redshift. Removing unresolved haloes from the merger tree can introduce a bias. When the number of haloes that are discarded drops to more than 50 per cent of the original sample, a spurious upturn in the median mass history occurs. To avoid this bias, the median mass history curve is only built out to the redshift




Figure 1. Median halo mass history as a function of redshift from simulations DMONLY (left panel) and REF (right panel) for haloes in 11 and seven different mass bins, respectively. The curves show the median value, and the 1σ error bars are determined by bootstrap resampling the haloes from the merger tree at a given output redshift. The different colour lines show the mass histories of haloes from different simulations. We find that a necessary condition for a halo to be defined, and mass histories to converge, is that haloes should have a minimum of 300 dark matter particles. The horizontal dashed dotted lines show the $300 \times m_{\text{dm}}$ limit for the simulation that matches the colour, where m_{dm} is the respective dark matter particle mass. When following a merger tree from a given halo sample, some haloes are discarded when unresolved. This introduces a bias and so an upturn in the median mass history. Therefore, mass history curves are stopped once fewer than 50 per cent of the original sample of haloes are considered. Simulations in the REF model with $25 h^{-1}$ Mpc comoving box size have a final redshift of $z = 2$, therefore the haloes mass histories begin at this redshift.

at which less than 50 per cent of the original number of haloes contribute to the median mass value.

Fig. 1 shows the effects of changing the resolution for the dark matter only and reference simulations. We first vary the box size while keeping the number of particles fixed (left panel). Then we vary the number of particles while keeping the box size fixed (right panel). The left panel (right panel) of Fig. 1 shows the mass history as a function of redshift for haloes in 11 (seven) different mass bins for the DMONLY (REF) simulation. All halo masses are binned in equally spaced logarithmic bins of size $\Delta \log_{10} M = 0.5$. The mass histories are computed by calculating the median value of the halo masses from the merger tree at a given output redshift; the error bars correspond to 1σ confidence intervals. The different coloured lines indicate the different simulations from which the halo mass histories were calculated. The horizontal dash-dotted lines in the panels show the $300 \times m_{\text{dm}}$ limit for the simulation that matches the colour. Haloes in the simulation with masses lower than this value are unresolved, and hence their mass histories are not considered. The mass histories from haloes whose main progenitors have masses lower than $10^{12} M_{\odot}$ at $z = 0$ were computed from simulations with $50 h^{-1}$ Mpc and $25 h^{-1}$ Mpc comoving box sizes. In the right-hand panel, where the mass histories from the REF model are shown, all mass history curves obtained from the REF simulation with a $25 h^{-1}$ Mpc comoving box size have a final redshift of $z = 2$. Therefore, these halo mass histories begin at this redshift.

3 SEMI-ANALYTIC MODEL FOR THE HALO MASS HISTORY

In the following subsections we study dark matter halo properties and provide a semi-analytic model that relates halo structure to the

mass accretion history. We begin with the NFW density profile and derive an analytic expression for the mean inner density, $\langle \rho \rangle (< r_{-2})$, within the scale radius, r_{-2} . We then define the formation redshift, and use the simulations to find the relation between $\langle \rho \rangle (< r_{-2})$ and the critical density of the universe at the formation redshift. We discuss the universality of the mass history curve and show how we can obtain a semi-analytic model for the mass history that depends on only one parameter (as expected from our EPS analysis presented in a companion paper). We then calibrate this single parameter fit using our numerical simulations. Finally, we show how the semi-analytic model for halo mass history depends on cosmology and halo mass definition.

3.1 Density profile

An important property of a population of haloes is their spherically averaged density profile. Based on N -body simulations, Navarro et al. (1997) found that the density profiles of CDM haloes can be approximated by a two-parameter profile,

$$\rho(r) = \frac{\rho_{\text{crit}} \delta_c}{(r/r_{-2})(1 + r/r_{-2})^2}, \quad (1)$$

where r is the radius, r_{-2} is the characteristic radius at which the logarithmic density slope is -2 , $\rho_{\text{crit}}(z) = 3H^2(z)/8\pi G$ is the critical density of the universe and δ_c is a dimensionless parameter related to the concentration c by

$$\delta_c = \frac{200}{3} \frac{c^3}{[\ln(1+c) - c/(1+c)]}, \quad (2)$$

which applies at fixed virial mass and where c is defined as $c = r_{200}/r_{-2}$, and r_{200} is the virial radius. A halo is often defined

Table 3. Notation reference. Unless specified otherwise, quantities are evaluated at $z = 0$.

Notation	Definition
M_{200}	$M_r(r_{200})$, total halo mass
r_{200}	Virial radius
r_{-2}	NFW scale radius
c	NFW concentration
M_z	$M(z)$, total halo mass at redshift z
$M_r(r)$	$M(<r)$, mass enclosed within r
x	r/r_{200}
$\langle\rho\rangle(<r_{-2})$	Mean density within r_{-2}
$M_r(r_{-2})$	$M(<r_{-2})$, enclosed mass within r_{-2}
z_{-2}	Formation redshift, when M_z equals $M_r(r_{-2})$
$\rho_{\text{crit},0}$	Critical density
$\rho_{\text{crit}}(z)$	Critical density at redshift z
$\rho_m(z)$	Mean background density at redshift z

so that the mean density $\langle\rho\rangle(<r)$ within the halo virial radius r_Δ is a factor Δ times the critical density of the universe at redshift z . Unfortunately, not all authors adopt the same definition, and readers should be aware of the difference in halo formation history and internal structure when different mass definitions are adopted (see Duffy et al. 2008; Diemer, More & Kravtsov 2013). We explore this in Section 3.6.2 to which the reader is referred to for further details. Throughout this work we use $\Delta = 200$. We denote $M_z \equiv M_{200}(z)$ as the halo mass as a function of redshift, $M_r \equiv M(<r)$ as the halo mass profile within radius r at $z = 0$, r_{200} as the virial radius at $z = 0$ and c as the concentration at $z = 0$. Note that the halo mass is defined as all matter within the radius r_{200} (see Table 3 for reference).

The NFW profile is characterized by a logarithmic slope that steepens gradually from the centre outwards, and can be fully specified by the concentration parameter and halo mass. Simulations have shown that these two parameters are correlated, with the average concentration of a halo being a weakly decreasing function of mass (e.g. NFW; Bullock et al. 2001; Eke, Navarro & Steinmetz 2001; Shaw et al. 2006; Neto et al. 2007; Duffy et al. 2008; Macciò, Dutton & van den Bosch 2008; Diemer & Kravtsov 2015; Dutton & Macciò 2014). Therefore, the NFW density profile can be described by a single free parameter, the concentration, which can be related to virial mass. The following relation was found by Duffy et al. (2008) from a large set of N -body simulations with the WMAP5 cosmology,

$$c = 6.67(M_{200}/2 \times 10^{12} h^{-1} M_\odot)^{-0.092}, \quad (3)$$

for haloes in equilibrium (relaxed).

The NFW profile can be expressed in terms of the mean internal density

$$\langle\rho\rangle(<r) = \frac{M_r(r)}{(4\pi/3)r^3} = \frac{200}{x^3} \frac{Y(cx)}{Y(c)} \rho_{\text{crit}}, \quad (4)$$

where x is defined as $x = r/r_{200}$ and $Y(u) = \ln(1+u) - u/(1+u)$. From this last equation we can verify that at $r = r_{200}$, $x = 1$ and $\langle\rho\rangle(<r_{200}) = 200\rho_{\text{crit}}$.

Evaluating $\langle\rho\rangle(<r)$ at $r = r_{-2}$, we obtain

$$\langle\rho\rangle(<r_{-2}) = \frac{M_r(r_{-2})}{(4\pi/3)r_{-2}^3} = 200c^3 \frac{Y(1)}{Y(c)} \rho_{\text{crit}}. \quad (5)$$

From this last expression we see that for a fixed redshift the mean inner density $\langle\rho\rangle(<r_{-2})$ can be written in terms of c . By substituting equation (3) into (5), we can obtain $\langle\rho\rangle(<r_{-2})$ as a function of virial

mass. Finally, we can compute the mass enclosed within r_{-2} . From equation (5) we obtain

$$M_r(r_{-2}) = M_{200} \frac{Y(1)}{Y(c)}, \quad (6)$$

where we used $M_{200} = (4\pi/3)r_{200}^3 200\rho_{\text{crit}}$.

Although the NFW profile is widely used and generally describes halo density profiles with high accuracy, it is worth noting that high-resolution numerical simulations have shown that the spherically averaged density profiles of dark matter haloes have small but systematic deviations from the NFW form (e.g. Navarro et al. 2004; Hayashi & White 2008; Navarro et al. 2010; Ludlow et al. 2010; Diemer & Kravtsov 2014). While there is no clear understanding of what breaks the structural similarity among haloes, an alternative parametrization is sometimes used (the Einasto profile), which assumes the logarithmic slope to be a simple power law of radius, $d \ln \rho / d \ln r \propto (r/r_{-2})^\alpha$ (Einasto 1965). Recently, Ludlow et al. (2013) investigated the relation between the accretion history and mass profile of cold dark matter haloes. They found that haloes whose mass profiles deviate from NFW and are better approximated by Einasto profiles also have accretion histories that deviate from the NFW shape in a similar way. However, they found the residuals from the systematic deviations from the NFW shape to be smaller than 10 per cent. We therefore only consider the NFW halo density profile in this work.

3.2 Formation redshift

Navarro et al. (1997) showed that the characteristic overdensity (δ_c) is closely related to formation time (z_f), which they defined as the time when half the mass of the main progenitor was first contained in progenitors larger than some fraction f of the mass of the halo at $z = 0$. They found that the ‘natural’ relation $\delta_c \propto \Omega_m(1+z_f)^3$ describes how the overdensity of haloes varies with their formation redshift. Subsequent investigations have used N -body simulations and empirical models to explore the relation between concentration and formation history in more detail (Wechsler et al. 2002; Zhao et al. 2003, 2009). A good definition of formation time that relates concentration to halo mass history was found to be the time when the main progenitor switches from a period of fast growth to one of slow growth. This is based on the observation that haloes that have experienced a recent major merger typically have relatively low concentrations, while haloes that have experienced a longer phase of relatively quiescent growth have larger concentrations. Moreover, Zhao et al. (2009) argue that halo concentration can be very well determined at the time the main progenitor of the halo has 4 per cent of its final mass.

The various formation time definitions each provide accurate fits to the simulations on which they are based and, at a given halo mass, show reasonably small scatter. However, our goal is to adopt a formation time definition that has a natural justification without invoking arbitrary mass fractions. To this end, we go back to the idea that haloes are formed ‘inside out’, and consider the formation time to be defined as the time when the initial bound core forms. We follow Ludlow et al. (2013) and define the formation redshift as the time at which the mass of the main progenitor equals the mass enclosed within the scale radius at $z = 0$, yielding

$$z_{-2} = z[M_z = M_r(r_{-2})]. \quad (7)$$

From now on we denote the formation redshift by z_{-2} . Interestingly, Ludlow et al. (2013) found that at this formation redshift, the critical density of the universe is directly proportional to the mean density

within the scale radius of haloes at $z = 0$: $\rho_{\text{crit}}(z_{-2}) \propto \langle \rho \rangle(< r_{-2})$. A possible interpretation of this relation is that the central structure of a dark matter halo (contained within r_{-2}) is established through collapse and later accretion and mergers increase the mass and size of the halo without adding much material to its inner regions, thus increasing the halo virial radius while leaving the scale radius and its inner density ($\langle \rho \rangle(< r_{-2})$) almost unchanged (Huss, Jain & Steinmetz 1999; Wang & White 2009).

3.3 Relation between halo formation time and concentration from numerical simulations

We now study the relation between $\rho_{\text{crit}}(z_{-2})$ and $\langle \rho \rangle(< r_{-2})$ using a set of DMONLY cosmological simulations from the OWLS project that adopt the WMAP-5 cosmology. We begin by considering two samples of haloes. Our complete sample contains all haloes that satisfy our resolution criteria while our ‘relaxed’ sample retains only those haloes for which the separation between the most bound particle and the centre of mass of the Friends-of-Friends (FoF) halo is smaller than $0.07R_{\text{vir}}$, where R_{vir} is the radius for which the mean internal density is Δ (as given by Bryan & Norman 1998) times the critical density. Neto et al. (2007) found that this simple criterion resulted in the removal of the vast majority of unrelaxed haloes and as such we do not use their additional criteria. At $z = 0$, our complete sample contains 2831 haloes, while our relaxed sample is reduced to 2387 (84 per cent).

To compute the mean inner density within the scale radius, $\langle \rho \rangle(< r_{-2})$, we need to fit the NFW density profile to each individual halo. We begin by fitting NFW profiles to all haloes at $z = 0$ that contain at least 10^4 dark matter particles within the virial radius. For each halo, all particles in the range $-1.25 \leq \log_{10}(r/r_{200}) \leq 0$, where r_{200} is the virial radius, are binned radially in equally spaced logarithmic bins of size $\Delta \log_{10} r = 0.078$. The density profile is then fit to these bins by performing a least square minimization of the difference between the logarithmic densities of the model and the data, using equal weighting. The corresponding mean enclosed mass, $M_r(r_{-2})$, and mean inner density at r_{-2} , $\langle \rho \rangle(< r_{-2})$, are found by interpolating along the cumulative mass and density profiles (measured while fitting the NFW profile) from $r = 0$ to $r_{-2} = r_{200}/c$, where c is the concentration from the NFW fit. Then we follow the mass history of these haloes through the snapshots, and interpolate to determine the redshift z_{-2} at which $M_z = M_r(r_{-2})$.

We perform a least-square minimization of the quantity $\Delta^2 = \frac{1}{N} \sum_{i=1}^N [(\rho_i)(\rho_{\text{crit},i}) - f(\rho_{\text{crit},i}, A)]^2$ to obtain the constant of proportionality, A . We find $\langle \rho \rangle(< r_{-2}) = (900 \pm 50)\rho_{\text{crit}}(z_{-2})$ for the relaxed sample, and $\langle \rho \rangle(< r_{-2}) = (854 \pm 47)\rho_{\text{crit}}(z_{-2})$ for the complete sample. The 1σ error was obtained from the least-squares fit. For comparison, Ludlow et al. (2014) found a constant value of 853 for their relaxed sample of haloes and the WMAP-1 cosmology. Fig. 2 shows the relation between the mean inner density at $z = 0$ and the critical density of the universe at redshift z_{-2} for various DMONLY simulations. Each dot in this panel corresponds to an individual halo from the complete sample in the DMONLY_WMAP5 simulations that have box sizes of $400 h^{-1}$ Mpc, $200 h^{-1}$ Mpc, $100 h^{-1}$ Mpc, $50 h^{-1}$ Mpc and $25 h^{-1}$ Mpc. The $\langle \rho \rangle(< r_{-2}) - \rho_{\text{crit}}(z_{-2})$ values are coloured by mass according to the colour bar at the top of the plot. The black (red) star symbols show the mean values of the complete (relaxed) sample in logarithmic mass bins of width $\delta \log_{10} M = 0.2$. As expected when unrelaxed haloes are discarded (e.g. Duffy et al. 2008), the relaxed sample contains on average slightly higher concentrations (by a factor of 1.16) and so higher formation times (by a factor of 1.1).




Figure 2. Relation between the mean density within the NFW scale radius at $z = 0$ and the critical density of the universe at the halo formation redshift, z_{-2} , for DMONLY simulations from the OWLS project. The simulations assume the WMAP-5 cosmological parameters and have box sizes of $400 h^{-1}$ Mpc, $200 h^{-1}$ Mpc, $100 h^{-1}$ Mpc, $50 h^{-1}$ Mpc and $25 h^{-1}$ Mpc. The black solid line indicates the relation shown in equation (8), which only depends on cosmology through the mass–concentration relation. The black (red) star symbols show the mean values of the complete (relaxed) sample in logarithmic mass bins of width $\delta \log_{10} M = 0.2$. The black dashed and solid lines show the relations found by fitting the data of the complete and relaxed samples, respectively. The filled circles correspond to values of individual haloes and are coloured by mass according to the colour bar at the top of the plot.

In Fig. 2 the best fit to the data points from the relaxed sample is shown by the solid line, while the dashed line is the fit to the complete sample. The $\rho_{\text{crit}}(z_{-2}) - \langle \rho \rangle(< r_{-2})$ correlation clearly shows that haloes that collapsed earlier have denser cores at $z = 0$. Using the mean inner density–critical density relation for the relaxed sample,

$$\langle \rho \rangle(< r_{-2}) = (900 \pm 50)\rho_{\text{crit}}(z_{-2}). \quad (8)$$

We replace $\langle \rho \rangle(< r_{-2})$ by equation (5) and calculate the dependence of formation redshift on concentration,

$$(1 + z_{-2})^3 = \frac{200}{900} \frac{c^3}{\Omega_m} \frac{Y(1)}{Y(c)} - \frac{\Omega_\Lambda}{\Omega_m}. \quad (9)$$

This last expression is tested in Fig. 3 (left panel) where we plot the median formation redshift as a function of concentration using different symbols for different sets of simulations from the OWLS project. The symbols correspond to the median values of the relaxed sample, and the error bars indicate 1σ confidence limits. The grey solid line shows the $z_{-2}-c$ relation given by equation (9), whereas the grey dashed line shows the same relation assuming a constant of 854 instead of 900 (as obtained for the complete sample). Similarly, using the Duffy et al. (2008) concentration–mass relation we obtain the formation redshift as a function of halo mass at $z = 0$ (right panel of Fig. 3). It is important to note that the $z_{-2}-c$ and $z_{-2}-M$ relations are valid in the halo mass range $10^{11}-10^{15} h^{-1} M_\odot$, at lower masses the concentration–mass relation begins to deviate from power laws (Ludlow et al. 2014).




Figure 3. Relation between formation redshift, z_{-2} , and halo concentration, c (left panel), and between formation redshift and $z = 0$ halo mass, M_{200} (right panel). The different symbols correspond to the median values of the relaxed sample and the error bars to 1σ confidence limits. The solid line in the left panel is not a fit but a prediction of the z_{-2} - c relation for relaxed haloes given by equation (9). Similarly, the dashed line is the prediction for the complete sample of haloes, assuming a constant of proportionality between $\langle\rho\rangle(<r_{-2})$ and ρ_{crit} of 854, rather than the value of 900 used for the relaxed sample. The grey area shows the scatter in z_{-2} plotted in Fig. B1 (right panel). Similarly, the solid line in the right panel is a prediction of the z_{-2} - M_{200} relation given by equations (9) and (3). The dashed line also shows the z_{-2} - M_{200} relation assuming $\langle\rho\rangle(<r_{-2}) = 854\rho_{\text{crit}}$ and the concentration–mass relation calculated using the complete sample.

In Appendix B we analyse the scatter in the formation time–mass relation and show that it correlates with the scatter in the concentration–mass relation. Thus concluding that the scatter in formation time determines the scatter in the concentration. Also, we investigate how the scatter in halo mass history drives the scatter in formation time.

3.4 The mass history

Fig. 4 shows the mass accretion history of haloes in different mass bins as a function of the mean background density. The mass histories are scaled to $M_r(r_{-2})$ and the mean background densities are scaled to $\rho_m(z_{-2}) = \rho_{\text{crit},0}\Omega_m(1+z_{-2})^3$. The figure shows that all halo mass histories look alike. This is in agreement with Ludlow et al. (2013), who found that the mass accretion history, expressed in terms of the critical density of the Universe, $M(\rho_{\text{crit}}(z))$, resembles that of the enclosed NFW density profile, $M(\rho)(<r)$. The similarity in the shapes between $M(\rho_{\text{crit}}(z))$ and $M(\rho)(<r)$ is still not clear, but it suggests that the physically motivated form $M(z) = M_0(1+z)^\alpha e^{\beta z}$, which is a result of rapid growth in the matter-dominated epoch followed by a slow growth in the dark energy epoch, produces the double power law of the NFW profile (see e.g. Lu et al. 2006). We use this feature to find a functional form that describes this unique universal curve in order to obtain an empirical expression for the mass accretion history at all redshifts and halo masses.

We are motivated by the extended Press–Schechter analysis of halo mass histories presented in Paper I. In that work, we showed through analytic calculations that when halo mass histories are described by a power-law times an exponential,

$$M(z) = M_0(1+z)^\alpha e^{\beta z}, \quad (10)$$

and that the parameters α and β are connected via the power spectrum of density fluctuations. In this work, however, we aim to relate halo structure to the mass accretion history. We therefore first deter-




Figure 4. Mass histories of haloes, obtained from different DMONLY_WMAP5 simulations, as indicated by the colours. The bottom left legends indicate the halo mass range at $z = 0$, selected from each simulation. For example, we selected haloes between 10^9 and $10^{11} M_\odot$ from the DMONLY_WMAP5_L025N512 simulation, divided them in equally spaced logarithmic bins of size $\Delta\log_{10}M = 0.2$, and calculated the median mass histories. The different curves show the median mass history of the main progenitors, normalized to the median enclosed mass of the main progenitors at $z = 0$, $M_r(r_{-2})$. The mass histories are plotted as a function of the mean background density of the universe, scaled to the mean background density at z_{-2} . The blue dashed line is a fit of expression (18) to the different mass history curves. The median value of the only adjustable parameter, γ , is indicated in the top-right part of the plot.

mine the correlation between the parameters α and β and concentration. To this end, we first find the α - β relation that results from the formation redshift definition discussed in the previous section. Thus, we evaluate the halo mass at z_{-2} ,

$$M_z(z_{-2}) = M_r(r_{-2}) = M_z(z=0)(1+z_{-2})^\alpha e^{\beta z_{-2}}. \quad (11)$$

Taking the natural logarithm, we obtain,

$$\ln\left(\frac{M_r(r_{-2})}{M_z(z=0)}\right) = \alpha \ln(1+z_{-2}) + \beta z_{-2}, \quad (12)$$

and hence

$$\alpha = \frac{\ln\left(\frac{M_r(r_{-2})}{M_z(z=0)}\right) - \beta z_{-2}}{\ln(1+z_{-2})}. \quad (13)$$

From this last equation we see that α can be written as a function of β , $M_r(r_{-2})$, $M_z(z=0)$ and z_{-2} . However, as $M_r(r_{-2})$ is a function of concentration and virial mass (see equation 6), we can write α in terms of β , concentration and z_{-2} ,

$$\alpha = \frac{\ln(Y(1)/Y(c)) - \beta z_{-2}}{\ln(1+z_{-2})}. \quad (14)$$

The next step is to find an expression for β . We find $\beta(z_{-2})$ by fitting equation (10) to all the data points plotted in Fig. 4. We now need to express $M(z)$ (equation 10) as a function of the mean background density. To do this, we replace $(1+z)$ by $(\rho_m(z)/\rho_{\text{crit},0}/\Omega_m)^{1/3}$ and divide both sides of equation (10) by $M_r(r_{-2})$, yielding

$$\begin{aligned} \frac{M_z(z)}{M_r(r_{-2})} &= \frac{M_z(z=0)}{M_r(r_{-2})} \left(\frac{\rho_m(z)}{\Omega_m \rho_{\text{crit},0}}\right)^{\alpha/3} \\ &\times \exp\left(\beta \left[\left(\frac{\rho_m(z)}{\Omega_m \rho_{\text{crit},0}}\right)^{1/3} - 1\right]\right). \end{aligned} \quad (15)$$

Multiplying both denominators and numerators by $\rho_m(z_{-2})$, we get, after rearranging,

$$\begin{aligned} \frac{M_z(z)}{M_r(r_{-2})} &= \frac{M_z(z=0)}{M_r(r_{-2})} \left(\frac{\rho_m(z_{-2})}{\Omega_m \rho_{\text{crit},0}}\right)^{\alpha/3} \left(\frac{\rho_m(z)}{\rho_m(z_{-2})}\right)^{\alpha/3} \\ &\times \exp\left(\beta \left[\left(\frac{\rho_m(z_{-2})}{\Omega_m \rho_{\text{crit},0}}\right)^{1/3} - 1\right]\right) \\ &\times \exp\left(\gamma \left[\left(\frac{\rho_m(z)}{\rho_m(z_{-2})}\right)^{1/3} - 1\right]\right), \end{aligned} \quad (16)$$

where we have defined $\gamma \equiv \beta(\rho_m(z_{-2})/\Omega_m/\rho_{\text{crit},0})^{1/3} = \beta(1+z_{-2})$. The term $\frac{M_z(z=0)}{M_r(r_{-2})} \left(\frac{\rho_m(z_{-2})}{\Omega_m \rho_{\text{crit},0}}\right)^{\alpha/3} \exp(\beta[\left(\frac{\rho_m(z_{-2})}{\Omega_m \rho_{\text{crit},0}}\right)^{1/3} - 1])$ in equation (16) is equal to unity, which can be seen by replacing $\rho_m(z_{-2})/\Omega_m \rho_{\text{crit},0} = (1+z_{-2})^3$ and comparing with equation (11). Hence equation (16) becomes

$$\begin{aligned} \frac{M_z(z)}{M_r(r_{-2})} &= \left(\frac{\rho_m(z)}{\rho_m(z_{-2})}\right)^{\alpha/3} \\ &\times \exp\left(\gamma \left[\left(\frac{\rho_m(z)}{\rho_m(z_{-2})}\right)^{1/3} - 1\right]\right). \end{aligned} \quad (17)$$

Thus, based on equation (17), the functional form to fit the mass accretion histories from the simulations can be written as

$$f(\tilde{z}, \gamma) = \alpha(z_{-2}, c, \gamma)\tilde{z}/3 + \gamma(e^{\tilde{z}/3} - 1), \quad (18)$$

where $f(\tilde{z}, \gamma) = \ln\left(\frac{M_z(z)}{M_r(r_{-2})}\right)$ and $\tilde{z} = \ln\left[\frac{\rho_m(z)}{\rho_m(z_{-2})}\right]$. From equation (14) we see that the parameter α is a function of z_{-2} , c and γ ,

$$\alpha = \frac{\ln(Y(1)/Y(c)) - \gamma z_{-2}/(1+z_{-2})}{\ln(1+z_{-2})}. \quad (19)$$

Therefore, γ is now the only adjustable parameter. We perform a χ^2 -like minimization of the quantity

$$\Delta^2 = \frac{1}{N} \sum_{i=1}^N [\log_{10}(M_z(z_i)/M_r(r_{-2})) - f(\tilde{z}_i, \gamma)]^2, \quad (20)$$

and find the value of γ that best fits all halo accretion histories. The sum in the χ^2 -like minimization is over the N available simulation output redshifts at $z_i (i = 1, N)$, with $\tilde{z}_i = \ln\left[\frac{\rho_{\text{crit},0}\Omega_m(1+z_i)^3}{\rho_m(z_{-2})}\right]$.

Fig. 4 shows halo mass histories (with $M_z(z)$ scaled to $M_r(r_{-2})$) for our complete halo sample as a function of the mean background density [$\rho_m(z)$ scaled to $\rho_m(z_{-2})$]. The blue dashed line is the fit of expression (18) to all the mass history curves included in the figure. Here, the only adjustable parameter is γ . We obtained $\gamma = -3.01 \pm 0.08$, yielding

$$\beta = -3(\rho_m(z_{-2})/\Omega_m/\rho_{\text{crit},0})^{-1/3} = -3/(1+z_{-2}). \quad (21)$$

Fig. 4 shows that the halo mass histories have a characteristic shape consisting of a rapid growth at early times, followed by a slower growth at late times. The change from rapid to slow accretion corresponds to the transition between the mass and dark energy dominated eras (see Paper I), and depends on the parameter β in the exponential (as can be seen from equation 10). The dependence of β on the formation redshift is given in equation (21), which shows that a more recent formation time, and hence a larger halo mass, results in a larger value of β , and so a steeper halo mass history curve. This last point can be seen in Fig. 5 from the mass histories of haloes in different mass bins (coloured lines shown in the panels). The panel on the left shows the mass history curves from the DMONLY simulation outputs (coloured lines in the background), and the mass histories predicted by equations (10), (14) and (21) (red dashed lines). From these panels we see that, (i) the mass history formula works remarkably well when compared with the simulation, and (ii) the larger the mass of a halo at $z=0$, the steeper the mass history curve at early times. In contrast, the mass history of low-mass haloes is essentially governed by the power law at late times.

The halo mass histories plotted in Fig. 5 come from the complete sample of haloes (relaxed and unrelaxed). We found no significant difference in mass growth when only relaxed haloes are considered. We therefore conclude that the fact that a halo is unrelaxed at a particular redshift does not affect its halo mass history, provided the concentration–mass relation fit from the relaxed halo sample is used. This is an interesting result because while deriving the semi-analytic model of halo mass history, we assumed that the halo density profile is described by the NFW profile at all times. Therefore, while the NFW is not a good fit for the density profile of unrelaxed haloes (Neto et al. 2007), our semi-analytic model (based on NFW profiles) is a good fit all haloes (relaxed and unrelaxed).

The right panel of Fig. 5 shows halo mass histories from the REF hydrodynamical simulations. We compute the halo mass as the total mass (gas and dark matter) within the virial radius (r_{200}). We find that the inclusion of baryons steepens the mass histories at high redshift; therefore the best description of $M(z)$ is given by equations (10), (14), (21), and the concentration–mass relation from the complete sample of haloes, $c = 5.74(M/2 \times 10^{12} h^{-1} M_\odot)^{-0.097}$.

3.5 The mass accretion rate

The accretion of gas and dark matter from the intergalactic medium is a fundamental driver of both the evolution of dark matter haloes and the formation of galaxies within them. For that reason, developing a theoretical model for the mass accretion rate is the basis




Figure 5. Mass histories of all haloes from simulations DMONLY_WMAP5 (left panel) and REF (right panel). Halo masses are binned in equally spaced logarithmic bins of size $\Delta \log_{10} M = 0.5$. The mass histories are computed by calculating the median value of the halo masses from the merger tree at a given output redshift; the error bars correspond to 1σ . The different colour lines show the mass histories of haloes from different simulations as indicated in the legends, while the red dashed curves correspond to the mass histories predicted by equations (10), (19) and (21).

for analytic and semi-analytic models that study galaxy formation and evolution. In this section we look for a suitable expression for the mean accretion rate of dark matter haloes. To achieve this, we take the derivative of the semi-analytic mass history model, $M(z)$, given by equation (10) with respect to time and replace dz/dt by $-H_0[\Omega_m(1+z)^5 + \Omega_\Lambda(1+z)^2]^{1/2}$, yielding

$$\frac{dM}{dt} = 71.6 M_\odot \text{ yr}^{-1} M_{12} h_{0.7} [-\alpha - \beta(1+z)] \times [\Omega_m(1+z)^3 + \Omega_\Lambda]^{1/2}, \quad (22)$$

where $h_{0.7} = h/0.7$, $M_{12} = M/10^{12} M_\odot$ and α and β are given by equations (14) and (21), respectively. As shown in the previous section, the parameters α and β depend on halo mass (through the formation time dependence). We find that this mass dependence is crucial for obtaining an accurate description for the mass history (as shown in Fig. 5). However, the factor of 2 (3) change for α (β) between halo masses of 10^8 and $10^{14} M_\odot$ is not significant when calculating the accretion rate. Therefore, we provide an approximation for the *mean* mass accretion rate as a function of redshift and halo mass, by averaging α and β over halo mass, yielding $\langle \alpha \rangle = 0.24$, $\langle \beta \rangle = -0.75$, and

$$\left\langle \frac{dM}{dt} \right\rangle = 71.6 M_\odot \text{ yr}^{-1} M_{12} h_{0.7} \times [-0.24 + 0.75(1+z)][\Omega_m(1+z)^3 + \Omega_\Lambda]^{1/2}. \quad (23)$$

Fig. 6 (top panel) compares the median dark matter accretion rate for different halo masses as a function of redshift (solid lines) to the mean accretion rate given by equation (23) (grey dashed lines). From the merger trees of the main haloes, we compute the mass growth rate of a halo of a given mass. We do this by following the main branch of the tree and computing $dM/dt = (M(z_1) - M(z_2))/\Delta t$, where $z_1 < z_2$, $M(z_1)$ is the descendant halo mass at time t and $M(z_2)$ is the most massive progenitor at time $t - \Delta t$. The median value of dM/dt for the complete set of resolved haloes is then plotted for different constant halo masses. We find very good agreement between the simulation outputs and the analytic estimate

given by equation (23). As expected, the larger the halo mass, the larger the dark matter accretion rate.

3.5.1 Baryonic accretion

Next, we estimate the gas accretion rate and compare our model with similar fitting formulae proposed by Fakhouri, Ma & Boylan-Kolchin (2010) and Dekel & Krumholz (2013). The bottom panel of Fig. 6 shows the gas accretion rate as a function redshift for a range of halo masses ($\log_{10} M/M_\odot = 11.2-12.8$). The grey circles correspond to the gas accretion rate measured in REF_L100N512. In this case we compute the total mass growth ($M = M_{\text{gas}} + M_{\text{DM}}$) from the merger trees, and then estimate the gas accretion rate by multiplying the total accretion rate by the universal baryon fraction $f_b = \Omega_b/\Omega_m$. The green solid line corresponds to our gas accretion rate model (given by Ω_b/Ω_m times equation 23). The blue dot-dashed line is the gas accretion rate proposed by Dekel & Krumholz (2013) ($dM_b/dt = 30 M_\odot \text{ yr}^{-1} f_b M_{12} (1+z)^{5/2}$), who derived the baryonic inflow on to a halo dM_b/dt from the averaged growth rate of halo mass through mergers and smooth accretion based on the EPS theory of gravitational clustering (Neistein, van den Bosch & Dekel 2006; Neistein & Dekel 2008). Lastly, we compare our model with the accretion rate formula from Fakhouri et al. (2010) [$dM_b/dt = 46.1 M_\odot \text{ yr}^{-1} f_b M_{12}^{1.1} (1 + 1.11z)(\Omega_m(1+z)^3 + \Omega_\Lambda)^{1/2}$], plotted as the purple dashed line. Fakhouri et al. (2010) constructed merger trees of dark matter haloes and quantified their merger rates and mass growth rate using the Millennium and Millennium II simulations. They defined the halo mass as the sum of the masses of all subhaloes within a FoF halo. We see that our accretion rate model is in excellent agreement with the formulae from Fakhouri et al. (2010) and Dekel & Krumholz (2013). We find that the Fakhouri et al. (2010) formula generally overpredicts the gas accretion rate in the low-redshift regime (e.g. it overpredicts it by a factor of 1.4 at $z = 0$ for a $10^{12} M_\odot$ mass halo). The Dekel & Krumholz (2013) formula underpredicts (overpredicts) the gas accretion rate in the low- (high-) redshift regime for haloes with masses larger (lower) than $10^{12} M_\odot$.




Figure 6. Mean accretion rate of dark matter (top panel) and gas (bottom panel) as a function of redshift for different halo masses. Top panel: the accretion rate obtained from the simulation outputs up to the redshift where the halo mass histories are converged. Grey dashed lines show the accretion rate estimated using equation (23). Bottom panel: gas accretion rate obtained from the REF_L100N512 simulation (grey circles), from Ω_b/Ω_m times equation (23) (green solid line), and from various fitting formulae taken from the literature.

3.6 Dependence on cosmology and mass definition

We have developed a semi-analytic model that relates the inner structure of a halo at redshift zero to its mass history. The model adopts the NFW profile, computes the mean inner density within the scale radius, and relates this to the critical density of the universe at the redshift where the halo virial mass equals the mass enclosed within r_{-2} . This relation enables us to find the formation redshift–halo mass dependence and to derive a one-parameter model for the halo mass history. In this section we consider the effects of cosmology and mass definition on the semi-analytic model.

3.6.1 Cosmology dependence

The adopted cosmological parameters affect the mean inner halo densities, concentrations, formation redshifts and halo mass histories. To investigate the dependence of halo mass histories on cosmology, we have run a set of dark matter only simulations with different cosmologies. Table 2 lists the sets of cosmological parameters adopted by the different simulations. Specifically, we assume

values for the cosmological parameters derived from measurements of the cosmic microwave background by the *WMAP* and the *Planck* missions (Spergel et al. 2003, 2007; Komatsu et al. 2009; Hinshaw et al. 2013; Planck Collaboration XVI et al. 2014).

It has been shown that haloes that formed earlier are more concentrated (Navarro et al. 1997; Bullock et al. 2001; Eke et al. 2001; Kuhlen et al. 2005; Macciò et al. 2007; Neto et al. 2007). Macciò et al. (2008) explored the dependence of halo concentration on the adopted cosmological model for field galaxies. They found that dwarf-scale field haloes are more concentrated by a factor of 1.55 in *WMAP1* compared to *WMAP3*, and by a factor of 1.29 for cluster-sized haloes. This reflects the fact that haloes of a fixed $z = 0$ mass assemble earlier in a universe with higher Ω_m , higher σ_8 and/or higher n_s .

The halo formation redshift can be related to the power at the corresponding mass scale, and therefore depends on both σ_8 and n_s . The parameter σ_8 sets the power at a scale of $8 h^{-1}$ Mpc, which corresponds to a mass of about $1.53 \times 10^{14} h^{-1} M_\odot (\Omega_m/\Omega_{m,WMAP5})$, and a wavenumber of k_8 . This last quantity is given by the relation $M = (4\pi\rho_m/3)(2\pi/k)^3$. For a power-law spectrum $P(k) \propto k^n$, the variance can be written as $\sigma^2(k)/\sigma_8^2 = (k/k_8)^{n+3}$. Therefore, the change in σ between *WMAP5* and *WMAP1* for a given halo mass that corresponds to a wavenumber k is

$$\frac{\sigma_{WMAP1}(k)}{\sigma_{WMAP5}(k)} = \frac{\sigma_{8,WMAP1}}{\sigma_{8,WMAP5}} \left(\frac{k}{k_8} \right)^{(n_s,WMAP1 - n_s,WMAP5)/2}. \quad (24)$$

A halo mass of $10^{12} M_\odot$ corresponds to a wavenumber of $k_{1.3} \sim 6k_8$. The total change in the mean power spectrum at this mass scale is $\frac{\sigma_{WMAP1}(k_{1.3})}{\sigma_{WMAP5}(k_{1.3})} = 1.27$. This is proportional to the change of the formation redshift,

$$(1 + z_{f,WMAP1}) = 1.27(1 + z_{f,WMAP5}). \quad (25)$$

Next, we test how this change affects the halo mass history. We showed that the mass history profile is well described by the expression $M(z) = M_z(z=0)(1+z)^\alpha e^{\beta z}$, where α and β both depend on the formation redshift. In the mass history model presented in Section 4 there are two best-fitting parameters that can be cosmology dependent. One is the constant value $A = 900$ in equation (8) that relates the mean inner density to the critical density at z_{-2} , and the other is the constant value $\gamma = -3$ in equation (21) that defines the β parameter.

To investigate the cosmology dependence of A , we analyse the $\langle \rho \rangle(<r_{-2}) - \rho_{crit}(z_{-2})$ relation in the simulations with different cosmologies. We do the same as in Section 4.3. First we fit the NFW profile to dark matter haloes to obtain c and r_{-2} , and calculate the cumulative mass, M_{-2} , and density, $\langle \rho \rangle(<r_{-2})$, from $r = 0$ to $r = r_{-2}$. Then we follow the halo mass histories through the snapshots and interpolate to calculate z_{-2} , the redshift for which $M(z)$ is equal to M_{-2} . Finally, we obtain the best-fitting $\langle \rho \rangle(<r_{-2}) - \rho_{crit}(z_{-2})$ relation. We find that the parameter A_{cosmo} , where cosmo is *WMAP1*, *WMAP3*, *WMAP5*, *WMAP9* or *Planck*, changes with cosmology. We show this in the top panel of Fig. 8. We find $A_{WMAP1} = 787 \pm 52.25$, $A_{WMAP3} = 850 \pm 39.60$, $A_{WMAP5} = 903 \pm 48.63$, $A_{WMAP9} = 820 \pm 51.03$ and $A_{Planck} = 798 \pm 43.73$. We do not find good agreement with Ludlow et al. (2013), who found $A_{WMAP1} = 853$ for *WMAP1* cosmology. This is due to the fact that we are only analysing the $\langle \rho \rangle(<r_{-2}) - \rho_{crit}(z_{-2})$ relation in the high-mass regime ($M = 10^{12.8} - 10^{13.8} M_\odot$), due to the limitations of the box size ($L = 100 h^{-1}$ Mpc). With a more complete halo population, we may obtain better agreement. We conclude that the A_{cosmo} parameter




Figure 7. Mass histories of haloes, obtained from simulations with the *WMAP1* cosmology (DMONLY_WMAP1_L100N512, blue solid lines) and the *WMAP5* cosmology (DMONLY_WMAP5_L100N512, green solid lines). The curves show the median mass history of the main progenitors, normalized to the median enclosed mass, $M_r(r_{-2})$, of the main progenitors at $z=0$. The mass histories are plotted as a function of the mean background density of the universe, scaled to the mean background density at z_{-2} . The blue dashed line is a fit of expression (18) to the different mass history curves. The median value of the only adjustable parameter, γ , is indicated in the top-right part of the plot. We find that γ is insensitive to cosmology.

depends on cosmology, at least for the halo mass ranges we are considering.

Next, we analyse how the change of the formation redshift due to cosmology affects β , defined as $\beta = -3/(1+z_f)$. We find from Fig. 7 that the constant value, -3 , is insensitive to cosmology. Fig. 7 shows the same analysis as Fig. 4, but for halo mass histories obtained from simulations with the *WMAP1* and *WMAP5* cosmology as indicated in the legends. We fit expression (18) to the mass history curves from different cosmologies and obtained the same adjustable parameter γ . We therefore conclude that $\gamma = -3$ is insensitive to cosmology.

If we then consider that the change in β between *WMAP1* and *WMAP5* is $\beta_{\text{WMAP1}} = -3/(1+z_{f,\text{WMAP1}}) = -3/1.27/(1+z_{f,\text{WMAP5}}) = \beta_{\text{WMAP5}}/1.27$, the change in the halo mass history between the *WMAP5* and *WMAP1* cosmologies, for a halo mass of $10^{12} M_\odot$ at $z=0$, corresponds to

$$\begin{aligned} \log_{10} \frac{M(z)_{\text{WMAP1}}}{M(z)_{\text{WMAP5}}} &= \log_{10} e^{(\beta_{\text{WMAP1}} - \beta_{\text{WMAP5}})z} \\ &\approx -0.12\beta_{\text{WMAP1}}z, \\ &\approx 0.1z. \end{aligned} \quad (26)$$

In the last step we replaced β_{WMAP1} by $-3/(1+z_{f,\text{WMAP1}}) = -0.75$ for a $10^{12} M_\odot$ halo. We obtained $z_{f,\text{WMAP1}}$ from the $\langle \rho \rangle(\langle r_{-2} \rangle) - \rho_{\text{crit}}(z_{-2})$ relation suitable for the *WMAP1* cosmology (see top panel of Fig. 8) and the concentration–mass relation from Neto et al. (2007).

Next, we test the change in halo mass history. For example, if a halo had a mass of $10^{11.4} M_\odot$ at $z=2$ in the *WMAP5* cosmology, it would have had a mass of $10^{11.6} M_\odot$ in the *WMAP1* cosmology. The value of σ_8 has a particularly large effect at high redshift, because structure formation proceeds faster in the *WMAP1* cosmology, as shown by the above expression. This last point can also be seen in the two panels of Fig. 8. The top panel is the same as the right panel of Fig. 3, and shows the formation redshift, z_{-2} , as a function of





Figure 8. Top panel: relation between formation redshift (z_{-2}) and halo mass at $z=0$ (M_0). The different symbols correspond to median values, the error bars to 1σ confidence limits and the grey area to the scatter. These were computed from the dark matter only simulations that assumed a *WMAP3* (light blue line), *WMAP5* (blue line), *WMAP1* (green line), *WMAP9* (purple line) and *Planck* cosmology (dark blue line). The solid lines are not fits, but predictions of the z_{-2} – M_{halo} relation given by equation (9). We also indicated the different values of the constant of proportionality A obtained by fitting the $\langle \rho_{-2} \rangle - \rho_{\text{crit}}(z_{-2})$ relation. Bottom panel: halo mass history of a halo of $10^{12} M_\odot$ at $z=0$ from DMONLY simulations with different cosmologies. The grey curves show that as long as a suitable concentration–mass relation is assumed for the cosmology under consideration, equations (37), (38) and (39) give a good estimate of the mass history curve.

halo mass (obtained from simulations with different cosmologies). We see that there are large differences between the *WMAP5* and *WMAP1* cosmologies due to the changes in σ_8 and n_s . Interestingly, there is only a small difference between the *Planck* and *WMAP1* cosmologies (in agreement with Dutton & Macciò 2014; Ludlow et al. 2014), and also between the *Planck* and *WMAP9* cosmologies for which we found $(1+z_f)_{\text{Planck}} = 1.01(1+z_f)_{\text{WMAP9}}$ for a $10^{12} M_\odot$ halo. The bottom panel of Fig. 8 shows the mass history of $10^{12} M_\odot$ haloes at $z=0$ from DMONLY simulations with

different cosmologies. As predicted, the change in mass history between the *WMAP5* and *WMAP1* cosmologies is $\log_{10} \frac{M(z)_{\text{WMAP1}}}{M(z)_{\text{WMAP5}}} \sim 0.1z$, while little difference is found between the *WMAP9* and *Planck* ($\Delta M(z) \sim 10^{-3}z$).

We found that as long as a suitable concentration–mass relation and value for the A parameter are assumed for the cosmology being considered, equations (37), (38) and (39) provide a good estimate of the mass history curve. This can be seen in the bottom panel of Fig. 8 by comparing the different halo mass histories. For the *WMAP5* and *WMAP3* cosmologies we assumed the concentration–mass relation found by Duffy et al. (2008), whereas we used the relation from Neto et al. (2007) for the other cosmologies. For a step-by-step description of how to use the mass history models (analytic and semi-analytic) that were presented in Sections 2 and 4.4, respectively, see Appendix C.

3.6.2 Mass definition dependence

So far our calculations have been based on a halo mass defined as the mass of all matter within the radius r_{200} at which the mean internal density $\langle \rho \rangle(<r_{200})$ is a factor of $\Delta = 200$ times the critical density of the universe, ρ_{crit} (from now on we denote this halo mass by M_{200}). In the literature a number of values have been used for Δ . Some authors opt to use $\Delta = 200$ (e.g. Jenkins et al. 2001) or $\Delta = 200\Omega_m(z)$ (e.g. NFW), while others (e.g. Bullock et al. 2001) choose $\Delta = \Delta_{\text{vir}}$ according to the spherical virialization criterion of Bryan & Norman (1998). These definitions can lead to sizeable differences in c for a given halo and, as discussed, the differences are also cosmology-dependent.

In this section we study how the structural properties and mass accretion histories depend on the adopted mass definition. We analyse halo mass histories of relaxed haloes using three different halo mass definitions. First, we use M_{200} . Secondly, we use M_{mean} , which is the mass within the radius r_{mean} for which the mean internal density is 200 times the mean background density. Finally, M_{vir} is the mass within the radius r_{vir} for which the mean internal density is Δ_{vir} times the critical density as determined by Bryan & Norman (1998). Note that halo masses and radii are determined using a spherical overdensity routine within the SUBFIND algorithm (Springel, White & Hernquist 2001) centred on the main subhalo of the FoF haloes (Davis et al. 1985). We perform all calculations for the three different halo definitions, taking the halo centre to be the location of the particle in the FoF group for which the gravitational potential is minimum.

Equation (8) shows that the formation redshift is directly proportional to the mean density within the scale radius ($(1+z_{-2})^3 \propto \langle \rho \rangle(<r_{-2})$), but the constant of proportionality depends on the mass definition that is adopted. Therefore, a change in the mass definition changes the formation time as

$$\frac{(1+z_f)_{\Delta_1}}{(1+z_f)_{\Delta_2}} \approx \left(\frac{\langle \rho \rangle(<r_{-2})_{\Delta_1}}{\langle \rho \rangle(<r_{-2})_{\Delta_2}} \right)^{1/3}, \quad (27)$$

where $\Delta_{1,2}$ refers to different overdensity criteria. That is, from equation (5), $\langle \rho \rangle(<r_{-2})$ changes according to the mass definition as $\langle \rho \rangle(<r_{-2})_{\Delta} = \Delta \times \rho_{\text{crit},0} c_{\Delta}^3 Y(1)/Y(c_{\Delta})$. Then $\langle \rho \rangle(<r_{-2})_{\Delta=200}$ refers to the mean internal density within r_{-2} , obtained by defining the mean internal density at the virial radius to be $200\rho_{\text{crit},0}$. If we consider $\Delta_1 = 200$ and $\Delta_2 = \text{mean} = 200\Omega_m$, and that there is a factor of 0.55 difference between the concentrations c_{200} and c_{mean} for a $10^{12} M_{\odot}$ halo, then we obtain (using equation 27) the relation $(1+z_f)_{200} \approx (0.255\Omega_m^{-1})^{1/3}(1+z_f)_{\text{mean}}$, where we have used the




Figure 9. Mass history of a $10^{12} M_{\odot}$ halo as a function of redshift. The different coloured lines show the change in the mass history when different halo mass definitions are used. The green line shows the mass history of a halo of $M_{200} = 10^{12} M_{\odot}$ at $z = 0$, whereas the dark blue (purple) line shows the mass history of a halo of $M_{\text{mean}} = 10^{12} M_{\odot}$ ($M_{\text{vir}} = 10^{12} M_{\odot}$) at $z = 0$. The dashed lines show the mass history predicted by equations (37), (38) and (39). The difference lies in the formation redshift definition which is affected by the change in the mean inner density (see equation 8). The value of $\langle \rho \rangle(<r_{-2})$ changes with the value of Δ we used in the definition of halo mass. The c – M relation corresponds to the mass definition under consideration.

fact that $\langle \rho \rangle(<r_{-2})_{\text{mean}} \sim (\Omega_m/0.255)\langle \rho \rangle(<r_{-2})_{200}$.¹ This implies that the change in the halo mass history due to different halo mass definitions is

$$\log_{10} \frac{M(z)_{200}}{M(z)_{\text{mean}}} = \log_{10}(1+z)^{\alpha_{200} - \alpha_{\text{mean}}} + \log_{10} e^{(\beta_{200} - \beta_{\text{mean}})z} \quad (28)$$

$$\approx 0.956\alpha_{200} \log_{10}(1+z) + [1 - (0.255/\Omega_m)^{1/3}] \log_{10}(e)\beta_{200}z \quad (29)$$

$$\approx 0.0543z, \quad (30)$$

where in step (29) we replaced $\alpha_{200} - \alpha_{\text{mean}} \approx 0.956\alpha_{200}$, which is valid for a $10^{12} M_{\odot}$ halo, and $\alpha_{200} = 0.2501$ and $\beta_{200} = -0.8147$.

The difference in mass history given by equation (30) can be seen in Fig. 9, which shows how the halo mass history is affected by the halo mass definition. The green line in Fig. 9 shows the mass history assuming the M_{200} mass definition. The purple line shows the M_{vir} definition, and the dark blue line shows the M_{mean} definition. The different dashed lines correspond to the mass histories $M(z) = M(z=0)(1+z)^{\alpha} e^{\beta z}$, where the difference lies in the mass definition that changes the mean inner density and the concentration–mass relation (for a relaxed halo sample). Duffy et al. (2008) studied how the halo mass definition changes the concentration–mass relation, and provided the parameters of the different c – M relations. They found that the concentration of a relaxed M_{mean} halo is 80 per cent larger than the concentration of a relaxed M_{200} halo. We adopt those fits in our calculations of the $M(z)$

¹ In this last step we used the approximation that $\langle \rho \rangle(<r_{-2})_{\Delta} = \Delta \times c^3 Y(1)/Y(c)\rho_{\text{crit},0} \approx \Delta \times 0.643c^{2.28}\rho_{\text{crit},0}$.




Figure 10. The top panel shows a comparison between the semi-analytic (solid lines) and the analytic model (dashed lines) for halo mass history. The bottom panel shows the residuals between the models. The different coloured lines correspond to the models for various halo masses in range $\log_{10}[M_0/M_{\odot}] = 5-15$ in steps of 2.

estimate and conclude that, as long as we use a concentration–mass relation that is consistent with the adopted halo mass definition, the expressions (37), (38) and (39) accurately reproduce the halo mass history.

The analytic estimate given by equation (30) predicts that the difference in mass history due to the change in the mass definition (M_{200} versus M_{mean}) is $\Delta \log_{10} M(z) \approx 0.0543z$. This can be seen in Fig. 9, where $\Delta \log_{10} M(z) = 0.1086$ at $z = 2$ [$M_{\text{mean}}(z = 2) = 10^{11.3035} M_{\odot}$ and $M_{200}(z = 2) = 10^{11.4117} M_{\odot}$].

4 COMPARISON BETWEEN SEMI-ANALYTIC AND ANALYTIC MODELS

In this section we compare the semi-analytic model derived in Section 3.4, with the analytic model for halo mass history derived in Paper I. Note that while the semi-analytic model is obtained through fits to simulations, the analytic model was based on the extended Press–Schechter (EPS) theory without calibration against simulations.

Fig. 10 shows a comparison between the models for various halo masses ($\log_{10}[M_0/M_{\odot}] = 5-15$). As can be seen from the figure, the models mostly agree on the mass histories of haloes with final masses between 10^{10} and $10^{14} M_{\odot}$. However, there are a few factors of difference in the mass histories of larger and smaller haloes, and the difference increases towards high redshift. We find that in the analytic model the halo mass decreases quite abruptly at high redshift for haloes with final masses $> 10^{14} M_{\odot}$. For instance, there is a factor of 9 difference at $z = 5$ between the models for a $10^{15} M_{\odot}$ halo. This difference is probably due to the progenitor definition. In the analytic model, the progenitor is defined as the halo with mass a factor of q lower ($q \sim 4$ for $M_0 > 10^{14} M_{\odot}$) at redshift z_f ($z_f \sim 0.9$), whereas in the semi-analytic model, the

progenitor is the halo that contains most of the 25 most bound dark matter particles in the following snapshot.

Fig. 10 also shows that the semi-analytic model overpredicts the mass histories of haloes with final masses $< 10^9 M_{\odot}$. This is expected because the parameters α and β in the semi-analytic model depend on the concentration–mass relation adopted. In this case we are using Duffy et al. (2008) relation, which is calibrated in the mass range $10^{10} - 10^{14} M_{\odot}$, for lower masses the concentration–mass relation deviates from a simple power law (Ludlow et al. 2014).

In Paper I we developed an analytic model of halo mass history, based on EPS theory, that only depends on the power spectrum of the primordial density perturbations. We found very good agreement between the halo mass histories predicted by our analytic model and published fits to simulation results (van den Bosch 2002; McBride et al. 2009; van den Bosch et al. 2014). In this work we have developed a semi-analytic model that uses the functional form for the mass history motivated by EPS theory, and linked the mass history to halo structure through empirical relations obtained from simulations.

We now combine these two models (semi-analytic and analytic) to establish the physical link between a halo concentration and the linear rms fluctuation of the primordial density field. From Fig. 10 we have found that there are a few factors of difference between the models. We now focus on the mass range $10^{11} - 10^{15} M_{\odot}$, where the factor of difference is less than 1.5. We set the mass history curve to be the same in the two models, that is

$$M(z)_{\text{Analytic}} = M(z)_{\text{Semi-analytic}} \quad (31)$$

for all redshifts. We then evaluate this equality at redshift 1 and obtain

$$f(M_0) \left(0.92 \frac{dD}{dz} \Big|_{z=0} - 0.3 \right) = \alpha_S(c) \ln(2) + \beta_S(c). \quad (32)$$

In this last equation, $\alpha_S(c)$ and $\beta_S(c)$ are given by equations (19) and (21), respectively, and depend on concentration, D is the linear growth factor and $f(M_0)$ depends on the rms of the primordial density field, σ . We approximate various terms in equation (32), including $f(M_0) \sim 1.155(\sigma(M_0)^2)^{0.277}$ and $Y(1)/Y(c) \sim 0.643c^{-0.71}$, and obtain

$$c = 3\sigma^{0.946} + 2.3, \quad (33)$$

which is suitable for a *WMAP5* cosmology. Note that equation (33) is not a fit to any simulation data; it has been derived from equation (32). Fig. 11 shows the c – σ relation at $z = 0$. In this figure we compare the predicted relation (solid line), as given by equation (33) (obtained by equalling the analytic and semi-analytic models), with the simulation outputs (coloured symbols). The different symbols correspond to the median values of the relaxed sample of haloes and the error bars to 1σ confidence limits. The good agreement between the analytic prediction and the simulation outputs clearly shows that the halo mass accretion history is the physical connection in the c – σ correlation.

5 SUMMARY AND CONCLUSION

In this work we have demonstrated that there is an intrinsic relation between halo assembly history and inner halo structure, and that the mass history is the physical connection between the inner halo structure and the power spectrum of initial density fluctuations.

We examined the density profiles and mass growth histories of a large sample of haloes and their progenitors within the OWLS




Figure 11. Comparison between the σ - c relation at $z = 0$ predicted by the combination of the mass history models (solid line), and the simulation outputs (coloured symbols).

simulations. We separated our halo sample into a ‘relaxed’ sample, and a ‘complete’ sample that includes both relaxed and unrelaxed haloes. We confirmed the finding of Ludlow et al. (2013) that for relaxed haloes the mean enclosed density within the NFW scale radius (r_{-2}), $\langle \rho \rangle (< r_{-2})$, is directly proportional to the critical density of the Universe at the formation redshift, z_{-2} , defined as the time at which the mass of the main progenitor equals the mass enclosed within the scale radius at $z = 0$,

$$\langle \rho \rangle (< r_{-2}) = 900 \rho_{\text{crit}}(z_{-2}). \quad (34)$$

In the above relation, the value, 900 ± 50 , is obtained through fits to the simulation data (suitable for WMAP5 cosmology). We showed in Section 3.6.1 that expression (34) provides a straightforward relation between formation time and concentration,

$$(1 + z_{-2}) = [(200/900)c^3 Y(1)/Y(c) - \Omega_{\Lambda}]^{1/3} \Omega_{\text{m}}^{-1/3}, \quad (35)$$

where $Y(c) = \ln(1+c) - c/(1+c)$. The overall trend of decreasing formation redshift with decreasing concentration is evident for the main branches of the descendant haloes. This implies that haloes which assemble earlier have a higher concentration, because the density of the Universe at the formation time was larger. To relate the formation time to halo mass, we used the following concentration–mass relation for relaxed haloes

$$c = 6.67(M/2 \times 10^{12} h^{-1} M_{\odot})^{-0.092}, \quad (36)$$

obtained by Duffy et al. (2008) by fitting to the simulation data and suitable for the WMAP5 cosmology. We found that, on average, halo concentrations differ by a factor of 1.16 between the relaxed and complete samples. The lower individual concentrations of unrelaxed haloes (due to spurious subhaloes or ongoing mergers that do not result in an accurate fit for an NFW density profile) produce incorrect enclosed halo masses and therefore lower formation times (by a factor of 1.1). However, on average, the formation time–concentration relation does not change, thus indicating that the halo mass history is not affected by the fact that a halo is out of equilibrium at a particular redshift. We used these findings to show that scatter in the halo mass history leads to scatter in the formation time (δz_{-2}), and hence to scatter in the concentration–mass relation (δc).

We found that formation time decreases with increasing mass (at a non-linear rate). This means that high-mass haloes are still accreting mass rapidly in the present epoch, while low-mass haloes typically accreted their mass early. Thus, the formation time–concentration relation provides the physical link between the halo mass history and internal structure. This result led us to provide a semi-analytic model for the halo mass history, which uses a direct, analytic correlation between the parameters α and β in the mass history (equation 37) and concentration,

$$M(z) = M_0(1+z)^{\alpha} e^{\beta z}, \quad (37)$$

$$\alpha = [\ln(Y(1)/Y(c)) - \beta z_{-2}] / \ln(1 + z_{-2}), \quad (38)$$

$$\beta = -3/(1 + z_{-2}), \quad (39)$$

where we obtained the constant value, -3.01 ± 0.08 , in the last relation (equation 39) by fitting the halo mass history model (equation 37) to the simulation data. Then we obtained a semi-analytic model for the mass accretion history, that adopts the functional form, equation (37), and the parameters α and β are now given by analytic relations that include numbers obtained from fits to simulation results. It is important to note that the semi-analytic model was derived assuming that the density profile of all haloes is NFW. Interestingly, we found that the semi-analytic model describes with high accuracy the mass histories of both relaxed and unrelaxed halo sample, even though the NFW profile is not a good fit to the density profile of unrelaxed haloes.

We investigated how cosmology affects the semi-analytic model. We found that as long as a suitable concentration–mass relation and the value for the best-fitting parameter in the $\langle \rho \rangle (< r_{-2}) - \rho_{\text{crit}}(z_{-2})$ relation are assumed for the cosmology being considered, the semi-analytic models describe the mass histories with high accuracy. In addition, we investigated how different mass definitions change the halo mass histories and we found that as long as we use a concentration–mass relation that is consistent with the adopted halo mass definition, the semi-analytic model accurately reproduces the halo mass history.

In Paper I, we presented an analytic model for the halo mass history, based on extended Press–Schechter theory and not calibrated against simulations data. We compared the analytic model of Paper I with the semi-analytic model presented here and found very good agreement in the mass range $10^9 - 10^{14} M_{\odot}$. However, we found that the analytic model predicts larger masses at high redshift for haloes with final masses $> 10^{14} M_{\odot}$, whereas the semi-analytic model overpredicts the mass history of low-mass haloes (haloes with final masses $< 10^9 M_{\odot}$). This is expected because the semi-analytic model depends on the adopted concentration–mass relation, which deviates from the assumed power law at low masses. The reader may find a step-by-step guide on how to implement the semi-analytic model in Appendix C, as well as numerical routines online.²

Interestingly, by combining these two models (semi-analytic and analytic) we established the physical link between a halo concentration and the initial density perturbation field, which explains the correlation between concentration and rms fluctuation of the primordial density field, σ (Fig. 11).

² Available at <https://bitbucket.org/astroduff/commah>.

Finally, by differentiating equation (37) we obtained the dark matter accretion rate,

$$\frac{dM}{dt} = 71.6 M_{\odot} \text{yr}^{-1} M_{12} h_{0.7} \times [-\alpha - \beta(1+z)][\Omega_m(1+z)^3 + \Omega_{\Lambda}]^{1/2}. \quad (40)$$

As the change in the α and β parameters over halo masses is not significant when calculating accretion rates, we provided a *mean* accretion rate, obtained by averaging the parameters α and β over halo mass,

$$\left\langle \frac{dM}{dt} \right\rangle = 71.6 M_{\odot} \text{yr}^{-1} M_{12} h_{0.7} \times [-0.24 + 0.75(1+z)][\Omega_m(1+z)^3 + \Omega_{\Lambda}]^{1/2}. \quad (41)$$

We then concluded that in order to predict halo mass growth, the concentration–mass relation from a relaxed sample should be used.

Putting the pieces together, we addressed the question of how the structure of haloes depends on the primordial density perturbation field. We found that concentration is the link between the halo mass profile and the halo mass history (and that one can be determined from the other). We also found that the ‘shape’ of the halo mass history is given by the linear growth factor and linear power spectrum of density fluctuations. Therefore, we concluded that halo concentrations are directly connected to the initial density perturbation field.

In a forthcoming paper (Paper III) we will combine the analytic model presented in Paper I and semi-analytic model presented here to predict the concentration–mass relation. We will investigate its evolution. We will show that extrapolations to low masses of power-law fits to simulation results are highly inadequate, and will investigate whether linear $\langle \rho \rangle (< r_{-2}) - \rho_{\text{crit}}(z_{-2})$ relation holds at redshifts other than 0.

ACKNOWLEDGEMENTS

We are grateful to the referee Aaron Ludlow for fruitful comments that substantially improved the original manuscript. CAC acknowledges the support of the 2013 John Hodgson Scholarship and the hospitality of Leiden Observatory. JSBW is supported by an Australian Research Council Laureate Fellowship. JS acknowledges support by the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement 278594-GasAroundGalaxies. We are grateful to the OWLS team for their help with the simulations.

REFERENCES

- Bryan G. L., Norman M. L., 1998, *ApJ*, 495, 80
 Bullock J. S., Kolatt T. S., Sigad Y., Somerville R. S., Kravtsov A. V., Klypin A. A., Primack J. R., Dekel A., 2001, *MNRAS*, 321, 559
 Correa C. A., Wyithe J. S. B., Schaye J., Duffy A. R., 2015a, *MNRAS*, 450, 1514 (Paper I)
 Correa C. A., Wyithe J. S. B., Schaye J., Duffy A. R., 2015b, preprint (arXiv:1502.00391) (Paper III)
 Dalal N., Lithwick Y., Kuhlen M., 2010, preprint (arXiv:1010.2539)
 Dalla Vecchia C., Schaye J., 2008, *MNRAS*, 387, 1431
 Davis M., Efstathiou G., Frenk C. S., White S. D. M., 1985, *ApJ*, 292, 371
 Dekel A., Krumholz M. R., 2013, *MNRAS*, 432, 455
 Dekel A., Devor J., Hetzroni G., 2003, *MNRAS*, 341, 326
 Diemer B., Kravtsov A. V., 2014, *ApJ*, 789, 1

- Diemer B., Kravtsov A. V., 2015, *ApJ*, 799, 108
 Diemer B., More S., Kravtsov A. V., 2013, *ApJ*, 766, 25
 Dolag K., Borgani S., Murante G., Springel V., 2009, *MNRAS*, 399, 497
 Duffy A. R., Schaye J., Kay S. T., Dalla Vecchia C., 2008, *MNRAS*, 390, L64
 Dutton A. A., Macciò A. V., 2014, *MNRAS*, 441, 3359
 Einasto J., 1965, *Trudy Astrofizicheskogo Instituta Alma-Ata*, 5, 87
 Eke V. R., Navarro J. F., Steinmetz M., 2001, *ApJ*, 554, 114
 Fakhouri O., Ma C.-P., Boylan-Kolchin M., 2010, *MNRAS*, 406, 2267
 Giocoli C., Tormen G., Sheth R. K., 2012, *MNRAS*, 422, 185
 Hayashi E., White S. D. M., 2008, *MNRAS*, 388, 2
 Hinshaw G. et al., 2013, *ApJS*, 208, 19
 Huss A., Jain B., Steinmetz M., 1999, *ApJ*, 517, 64
 Jenkins A., Frenk C. S., White S. D. M., Colberg J. M., Cole S., Evrard A. E., Couchman H. M. P., Yoshida N., 2001, *MNRAS*, 321, 372
 Klypin A. A., Trujillo-Gomez S., Primack J., 2011, *ApJ*, 740, 102
 Komatsu E. et al., 2009, *ApJS*, 180, 330
 Kuhlen M., Strigari L. E., Zentner A. R., Bullock J. S., Primack J. R., 2005, *MNRAS*, 357, 387
 Lacerna I., Padilla N., 2011, *MNRAS*, 412, 1283
 Lu Y., Mo H. J., Katz N., Weinberg M. D., 2006, *MNRAS*, 368, 1931
 Ludlow A. D., Navarro J. F., Springel V., Vogelsberger M., Wang J., White S. D. M., Jenkins A., Frenk C. S., 2010, *MNRAS*, 406, 137
 Ludlow A. D., Navarro J. F., Li M., Angulo R. E., Boylan-Kolchin M., Bett P. E., 2012, *MNRAS*, 427, 1322
 Ludlow A. D. et al., 2013, *MNRAS*, 432, 1103
 Ludlow A. D., Navarro J. F., Angulo R. E., Boylan-Kolchin M., Springel V., Frenk C. S., White S. D. M., 2014, *MNRAS*, 441, 378
 McBride J., Fakhouri O., Ma C.-P., 2009, *MNRAS*, 398, 1858
 Macciò A. V., Dutton A. A., van den Bosch F. C., Moore B., Potter D., Stadel J., 2007, *MNRAS*, 378, 55
 Macciò A. V., Dutton A. A., van den Bosch F. C., 2008, *MNRAS*, 391, 1940
 Manrique A., Raig A., Salvador-Solé E., Sanchis T., Solanes J. M., 2003, *ApJ*, 593, 26
 Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563 (NFW)
 Navarro J. F., Frenk C. S., White S. D. M., 1997, *ApJ*, 490, 493
 Navarro J. F. et al., 2004, *MNRAS*, 349, 1039
 Navarro J. F. et al., 2010, *MNRAS*, 402, 21
 Neistein E., Dekel A., 2008, *MNRAS*, 388, 1792
 Neistein E., van den Bosch F. C., Dekel A., 2006, *MNRAS*, 372, 933
 Neto A. F. et al., 2007, *MNRAS*, 381, 1450
 Planck Collaboration XVI et al., 2014, *A&A*, 571, A16
 Prada F., Klypin A. A., Cuesta A. J., Betancort-Rijo J. E., Primack J., 2012, *MNRAS*, 423, 3018
 Salvador-Solé E., Viñas J., Manrique A., Serra S., 2012, *MNRAS*, 423, 2190
 Schaye J., Dalla Vecchia C., 2008, *MNRAS*, 383, 1210
 Schaye J. et al., 2010, *MNRAS*, 402, 1536
 Shaw L. D., Weller J., Ostriker J. P., Bode P., 2006, *ApJ*, 646, 815
 Sheth R. K., Tormen G., 2004, *MNRAS*, 349, 1464
 Spergel D. N. et al., 2003, *ApJS*, 148, 175
 Spergel D. N. et al., 2007, *ApJS*, 170, 377
 Springel V., 2005, *MNRAS*, 364, 1105
 Springel V., White M., Hernquist L., 2001, *ApJ*, 549, 681
 Syer D., White S. D. M., 1998, *MNRAS*, 293, 337
 van de Voort F., Schaye J., Booth C. M., Haas M. R., Dalla Vecchia C., 2011, *MNRAS*, 414, 2458
 van den Bosch F. C., 2002, *MNRAS*, 331, 98
 van den Bosch F. C., Jiang F., Hearin A., Campbell D., Watson D., Padmanabhan N., 2014, *MNRAS*, 445, 1713
 Wang J., White S. D. M., 2009, *MNRAS*, 396, 709
 Wang H., Mo H. J., Jing Y. P., 2009, *MNRAS*, 396, 2249
 Wechsler R. H., Bullock J. S., Primack J. R., Kravtsov A. V., Dekel A., 2002, *ApJ*, 568, 52
 Wiersma R. P. C., Schaye J., Theuns T., Dalla Vecchia C., Tornatore L., 2009a, *MNRAS*, 399, 574
 Wiersma R. P. C., Schaye J., Smith B. D., 2009b, *MNRAS*, 393, 99
 Zhao D. H., Mo H. J., Jing Y. P., Börner G., 2003, *MNRAS*, 339, 12
 Zhao D. H., Jing Y. P., Mo H. J., Börner G., 2009, *ApJ*, 707, 354

APPENDIX A: SIMULATIONS

The DMONLY simulations contain only dark matter particles, which provide us with a useful baseline model for testing the impact of baryonic physics on halo mass histories and mass profiles, when comparing it with the REF simulation. The REF simulation includes sub-grid recipes for star formation (Schaye & Dalla Vecchia 2008), radiative (metal-line) cooling and heating (Wiersma, Schaye & Smith 2009b), stellar evolution, mass loss from massive stars and chemical enrichment (Wiersma et al. 2009a), and a kinetic prescription for supernova feedback (Dalla Vecchia & Schaye 2008). In this work we use hydrodynamical simulations (along with DMONLY) to test the effect of baryons on halo-mass histories and compute the gas accretion rate. We do not test other feedback schemes because it was shown by van de Voort et al. (2011) that the halo mass accretion is robust to variations in feedback.

A1 Construction of halo merger trees

The first step towards studying the mass assembly history of haloes is to identify gravitationally bound structures and build halo merger trees. We begin by selecting the largest halo in each FoF group (Davis et al. 1985) (i.e. the main subhalo of FoF groups that is not embedded inside larger haloes). Halo virial masses and radii are determined using a spherical overdensity routine within the SUBFIND algorithm (Springel et al. 2001; Dolag et al. 2009) centred on the main subhalo of FoF haloes. Therefore, we define halo masses as all matter within the radius r_{200} for which the mean internal density is 200 times the critical density. Throughout this work, we study the accretion history of the largest haloes in each FoF group. Subhaloes, defined as bound structures that reside within the virial radius of the largest ‘host’ halo, show distinct mass histories. The structures of subhaloes are strongly affected by the potential of their host haloes, as seen for example in the cessation of mass accretion on to subhaloes residing in dense environments (see Wang, Mo & Jing 2009; Lacerna & Padilla 2011). Consequently, the masses of subhaloes do not follow the mass history of their host haloes.

The merger trees of the largest haloes are then built as follows. First, at each output redshift (snapshot), we select ‘resolved’ haloes that contain more than 300 dark matter particles. We refer to these resolved haloes as ‘descendants’. We then link each descendant with a unique ‘progenitor’ at the previous output redshift. This is non-trivial due to halo fragmentation: subhaloes of a progenitor halo may have descendants that reside in more than one halo. The fragmentation can be spurious or due to a physical unbinding event. To correct this, we link the descendant to the progenitor that contains the majority of the descendant’s 25 most bound dark matter particles. Therefore, the main progenitor of a given dark matter halo is found by tracing backwards in time to the most massive halo along the main branch of its merger tree. The different mass histories are calculated by following the merger trees of a given sample of haloes. At each redshift the mass histories are computed by calculating the median mass value, determined by bootstrap resampling the haloes, from the merger tree. Along with the median value, the 1σ confidence interval is recorded.

APPENDIX B: ANALYSIS OF SCATTER

B1 Scatter in formation times and concentrations

We now analyse the scatter in the formation time–mass relation and show it relates to the scatter in the concentration–mass relation.

So far, we have shown that the formation time is related to halo concentration through the mean inner density ($\langle\rho\rangle(<r_{-2})$)–critical density ($\rho_{\text{crit}}(z_{-2})$) relation plotted in Fig. 2. Through first-order error propagation, we look for the corresponding scatter in the formation time,

$$900\rho_{\text{crit}}(z_{-2}) = \langle\rho\rangle(<r_{-2}), \quad (\text{B1})$$

$$900\delta(\rho_{\text{crit}}(z_{-2})) = \delta(\langle\rho\rangle(<r_{-2})), \quad (\text{B2})$$

$$\frac{900}{200}\delta[\Omega_{\text{m}}(1+z_{-2})^3 + \Omega_{\Lambda}] = \delta\left(\frac{c^3 Y(1)}{Y(c)}\right), \quad (\text{B3})$$

$$3\Omega_{\text{m}}(1+z_{-2})^2 z_{-2} \left(\frac{\delta z_{-2}}{z_{-2}}\right) = \frac{\langle\rho\rangle(<r_{-2})}{900\rho_0} \left(\frac{\delta c}{c}\right) \times \left(3 - \frac{c^2}{(1+c)^2} \frac{1}{Y(c)}\right), \quad (\text{B4})$$

where we used $\delta(c^3/Y(c)) = 3c^2\delta c/Y(c) - c^4\delta c/[Y(c)^2(1+c)^2]$. Equation (B4) relates the scatter in formation time ($|\delta z_{-2}|/z_{-2}$) to the scatter in the concentration ($|\delta c|/c$).

The grey shaded areas in the panels in Fig. 3 show the scatter in z_{-2} , while the panels in Fig. B1 show the scatter in the c – M relation (left panel) and in the z_{-2} – M relation (right panel). The grey contours in Fig. B1 enclose 68 per cent of the distribution while the individual points show the remaining 32 per cent. The black (green) solid line shows the mean scatter in the formation time per halo mass bin for the relaxed (complete) sample. The presence of unrelaxed haloes does not have any significant effect on either the scatter in the formation time or the mass histories. The average scatter in formation time is $\langle|\delta z_{-2}|/z_{-2}\rangle = 0.324$ ($\langle|\delta z_{-2}|/z_{-2}\rangle = 0.356$) for the relaxed (complete) sample.

The left panel of Fig. B1 shows that the average scatter in the concentration–mass relation is $\langle|\delta c|/c\rangle = 0.257$ for the full sample and $\langle|\delta c|/c\rangle = 0.218$ for the relaxed sample. In agreement with previous work (see e.g. Neto et al. 2007), the scatter in the concentration of the relaxed halo sample is lower than the scatter of the full sample. The extra scatter in the full sample is produced by the deviation of the density profiles from the NFW form for haloes experiencing ongoing mergers and for artificially linked haloes.

Assuming $\langle|\delta z_{-2}|/z_{-2}\rangle = 0.324$, $\langle|\delta c|/c\rangle$ can be obtained as a function of halo mass by applying equation (B4). This analytic estimate is plotted in the left panel for the relaxed sample (red dashed line). We find very good agreement between the scatter in concentration from equation (B4), and the median value plotted in black for the relaxed sample, and in green for the complete sample. Therefore, we conclude that the scatter in formation time determines the scatter in the concentration. However, at higher masses and redshifts the fraction of relaxed haloes decreases (e.g. Ludlow et al. 2012) and it has been found that the concentration–mass relation of the complete halo sample exhibits a strong flattening and upturn (Klypin, Trujillo-Gomez & Primack 2011; Prada et al. 2012). As a result, at high masses and redshifts the scatter in the concentration will probably depend on other variables besides the scatter in formation time.

In the following subsection we find that the scatter in the accretion history determines the scatter in the formation time.




Figure B1. Scatter in the concentration–mass relation (left panel) and the formation redshift–mass relation (right panel). Left panel: Y-axis shows the difference between the concentration predicted by Duffy et al. (2008) (see equation 3) and the actual concentration from the simulation output. This difference is divided by the concentration from the simulation output and plotted against virial halo mass. The different colours of the points indicate that the concentration outputs were obtained from DMONLY_WMAP5 simulations with different box sizes. The grey contours enclose 68 per cent of the distribution while the individual points show the remaining 32 per cent. The black (green) solid line shows the mean relative scatter in the concentration–mass relation per halo mass bin of the relaxed (complete) sample. The red dashed line is an analytic estimate of the scatter obtained by propagating the scatter in the formation redshift–mass relation to the concentration–mass relation (equation B4). Right panel: same as the left panel but the scatter is obtained from the difference in the formation redshift predicted from equation (9) and the simulation output. The black (green) solid line shows the median value in the scatter as a function of mass of the relaxed (complete) sample.

B2 Scatter in halo mass histories

In this section we analyse the scatter found when computing mass histories from the simulation outputs. We analytically estimate the scatter in the mass history due to both the scatter in the concentration and formation times (estimated in Section B1). We then compare this to the scatter obtained from the simulation outputs.

To compute the scatter in the mass history we perform a first-order error propagation in $M(z)$ (equation 10),

$$\delta M(z)/M(z) = \delta\alpha \ln(1+z) + z\delta\beta, \quad (\text{B5})$$

where $\delta\alpha$ ($\delta\beta$) is the scatter in α (β) due to the scatter in c and z_{-2} . From equation (14) we first compute $\delta\alpha$ due to the scatter in z_{-2} ,

$$\begin{aligned} \delta\alpha &= \delta(-\beta z_{-2}/\ln(1+z_{-2})) \\ &= \frac{-z_{-2}\delta\beta - \beta\delta z_{-2}}{\ln(1+z_{-2})} - \beta z_{-2}\delta[\ln(1+z_{-2})]^{-1} \\ &= -\frac{z_{-2}\beta(1+z_{-2}\ln(1+z_{-2}))}{(1+z_{-2})\ln(1+z_{-2})} \left(\frac{\delta z_{-2}}{z_{-2}} \right), \end{aligned} \quad (\text{B6})$$

where in the last line we used $\delta\beta = \frac{-\beta z_{-2}}{(1+z_{-2})} \left(\frac{\delta z_{-2}}{z_{-2}} \right)$, which follows from equation (21). Similarly, we calculate the scatter in α due to the scatter in c ,

$$\begin{aligned} \delta\alpha &= \delta(\ln(Y(1)/Y(c))/\ln(1+z_{-2})) \\ &= -\frac{\delta Y(c)}{Y(c)} \frac{1}{\ln(1+z_{-2})} \\ &= -\frac{c^2}{(1+c)^2 Y(c) \ln(1+z_{-2})} \left(\frac{\delta c}{c} \right), \end{aligned} \quad (\text{B7})$$

where in the last step we used $\delta Y(c) = \frac{c^2}{(1+c)^2} \left(\frac{\delta c}{c} \right)$. Finally, we substitute $\delta\beta = \frac{-\beta z_{-2}}{(1+z_{-2})} \left(\frac{\delta z_{-2}}{z_{-2}} \right)$ and equations (B6) and (B7) into equation (B5). From Section B1 we adopt the average scatter in z_{-2} and c for the complete sample, that is $\langle |\delta c|/c \rangle = 0.25$ and




Figure B2. Mean scatter in the halo mass history against redshift for different halo masses M_{200} . The y-axis shows the mean value of the difference between the mass history (M_{fit}) predicted by equations (10), (14) and (21), and the mass history (M_{sim}) from the simulation output. The difference is divided by the M_{sim} from the simulation output. The different coloured lines correspond to the median values of the scatter from simulations with different box sizes. The grey lines show the analytic estimates of the mass history curves given by equations (B5), (B6) and (B7).

$\langle \delta z_{-2}/z_{-2} \rangle = 0.35$, and compute $\langle |\delta M(z)/M(z)| \rangle$ as a function of redshift. Fig. B2 shows the scatter measured from the simulation outputs (coloured lines) and from the analytic estimates (grey dashed lines). The different coloured lines correspond to the median values of the scatter from simulations with different box sizes. We calculate these by averaging the difference between the mass history predicted by equations (10), (14) and (21) (for a given halo with mass M_{200} at $z = 0$), and each $M(z)$ given by the merger trees from the complete halo sample. We find very good agreement indicating that the scatter in the concentration comes from the scatter in the

formation time, which in turn comes from the scatter in the halo mass history.

APPENDIX C: STEP-BY-STEP GUIDE TO COMPUTE HALO MASS HISTORIES

C1 Semi-analytical model

This Appendix provides a step-by-step procedure that outlines how to calculate the halo mass histories using the numerical model presented in Section 4.4.

(i) First assume a cosmology and choose a concentration–mass relation from the literature. For instance, the Duffy et al. (2008) relation, $c = 6.67(M_0/2 \times 10^{12} h^{-1} M_\odot)^{-0.092}$, is suitable for the *WMAP5* cosmology, whereas Neto et al. (2007) is suitable for *WMAP1*.

(ii) Calculate the formation time,

$$z_{-2} = \left(\frac{200}{A_{\text{cosmo}}} \frac{c(M_0)^3 Y(1)}{\Omega_m Y(c(M_0))} - \frac{\Omega_\Lambda}{\Omega_m} \right)^{1/3} - 1. \quad (\text{C1})$$

Note that the value of A_{cosmo} in the above equation is cosmology dependent. $A_{\text{WMAP5}} = 900$ is suitable for the *WMAP5* cosmology. In

this work we obtained $A_{\text{WMAP1}} = 787$, $A_{\text{WMAP3}} = 850$, $A_{\text{WMAP9}} = 820$ and $A_{\text{Planck}} = 798$.

(iii) Calculate the parameters α and β ,

$$\alpha = [\ln(Y(1)/Y(c)) - \beta z_{-2}] / \ln(1 + z_{-2}), \quad (\text{C2})$$

$$\beta = -3/(1 + z_{-2}). \quad (\text{C3})$$

(iv) Finally, the mass history can be calculated as follows,

$$M(z) = M_0(1 + z)^\alpha e^{\beta z}. \quad (\text{C4})$$

The above model is suitable for any cosmology (as long as the concentration–mass relation and the value of A_{cosmo} correspond to the desired cosmology) and is valid over the halo mass range for which the concentration–mass and the z_{-2} – M_0 relations, obtained from simulations, are valid (e.g. 10^{10} – $10^{14} M_\odot$ for Duffy et al. 2008).

This paper has been typeset from a \LaTeX file prepared by the author.