



Universiteit
Leiden
The Netherlands

Diagnosis and treatment of obese children with insulin resistance

Aa, M.P. van der

Citation

Aa, M. P. van der. (2016, December 13). *Diagnosis and treatment of obese children with insulin resistance*. Retrieved from <https://hdl.handle.net/1887/44921>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/44921>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/44921> holds various files of this Leiden University dissertation

Author: Aa, Marloes van der

Title: Diagnosis and treatment of obese children with insulin resistance

Issue Date: 2016-12-13



Chapter 1

General introduction on diagnosis and treatment of obese children with insulin resistance

General introduction on diagnosis and treatment of obese children with insulin resistance

Nowadays, worldwide more people are overweight or obese than underweight [1]. In 2014, worldwide 10.8% (9.7–12.0) of adult men and 14.9% (13.6–16.1) of adult women were obese, whereas the prevalence of underweight was respectively 8.8% (7.4–10.3) in men and 9.7% (8.3–11.1) in women [1].

Obesity is a condition which is defined as abnormal or excessive body fat accumulation. This condition may impair health, and is of specific concern in view of the increasing prevalence in children [2, 3]. To classify overweight, and thereby obesity, the body mass index (BMI) is used. The BMI is calculated as weight (kg) divided by (height (m))², with adult cut-off values for overweight and obesity of respectively ≥ 25 kg/m² and ≥ 30 kg/m² [2]. In children normal growth results in an initial decrease of BMI until the age of 4–5 years, followed by an increase in BMI. Consequently, fixed cut-off values for BMI to classify obesity in children cannot be used, and therefore standard deviation scores (SDS), z-scores, or percentiles are used to define overweight and obesity [4–6]. These scores are based on the number of standard deviations below or above the median BMI for age and sex of an international population of six large, nationally representative growth studies [4, 5]. Cut-off values used in the Netherlands are BMI-SDS > 1.1 (BMI > p85) for overweight and BMI-SDS > 2.3 (BMI > p95) for obesity [7].

In the 1960's, childhood obesity was very uncommon, with prevalence rates for overweight (BMI > p85) and obesity (BMI > p95) of 4.2–4.6% for children 6–19 years old in the United States [8]. Since that moment, prevalence of childhood obesity is rising. Although prevalence rates from 2007–2012 suggest that the rising prevalence in childhood obesity may have reached a plateau since 2003–2004 [9–11], in 2013–2014 the prevalence was rising again. In 2013–2014 the prevalence rates in children and adolescents 2–17 years old in the US for overweight (BMI > p85) and obesity (BMI > p95) were 33.4 (95%CI 30.9–35.9) % and 17.4 (95%CI 15.2–19.6)%, respectively [3]. In the Netherlands, prevalence of obesity was 0.3% in native boys and 0.5% in native girls aged 2–21 years in 1980. In 2010 however, these prevalence rates for obesity were 1.8% and 2.2% in native boys and girls, respectively, and higher in Moroccan and Turkish descent (6.0% and 8.4% in boys and 7.5% and 8.4% in girls, respectively). These figures should be seen in the context of a prevalence of overweight of 13.3 and 14.9% in native boys and girls, up to 32.5 and 31.7% in boys and girls of Turkish descent [12].

Obesity is most frequently caused by an energy imbalance between intake of calories and calories burned. During the last decades, energy intake shifts towards high-caloric foods containing few vitamins, minerals and other healthy nutrients. At the same time, the energy expended is reduced because of physical activity is reduced

and a sedentary lifestyle is more common [2]. The energy not expended is stored as fat mass. However, there is evidence that more factors contribute to obesity, such as parental obesity, social economic state, maternal nutrition and glucose metabolism during pregnancy and psychological health [13-15]. In 2-4% of the obese children, an endocrine cause (such as hypothyroidism, growth hormone deficiency, Cushing syndrome or hypothalamic obesity), genetic cause (for example leptin deficiency, mutations in POMC or MC4R deficiency) or genetic syndrome (for example Prader-Willi syndrome, Bardet-Biedl and Alstrom syndrome) is found [13, 14, 16-18]. Finally, use of medication, for example anti-epileptic and antipsychotic drugs, can contribute to the development of obesity [18, 19].

Prognosis and consequences of childhood obesity

Childhood obesity is a strong predictor for obesity in adulthood. Whitaker et al. described in 1997 that 79% of children who were obese at age 10-14 years old, were still obese as young adults (21-29 years) [20]. A systematic review by Singh et al, described rates from 47-83% of obese children becoming obese adults [21]. Odds ratios for obese children to become obese adults varied from OR 1.3 for obese children aged 1-2 years to an OR of 22.3 for obese children aged 10-14 years [20, 22]. In adults who were obese during childhood (age 14-19 years) relative risk for all-cause mortality after 31.5 years of follow up was 1.82 (95% CI 1.48–2.43) in men and 2.03 (95% CI 1.51–2.72) in women [23].

There are multiple consequences affecting almost all organ tracts, of which many will be listed here. Consequences of childhood obesity are both psychosocial and somatic. Psychosocial consequences have a high burden on the quality of life on short-term and long-term. Short-term psychosocial consequences are for example poor self-esteem, being bullied, depression and eating disorders [24, 25]. Low self-esteem was found in 34% of obese girls vs 8% in non-obese girls [26]. Long-term psychosocial effects of obesity include higher risk of depression, oppositional defiant disorder and lower incomes [27-29].

Cardiovascular and metabolic consequences are very common, both on short-term and long-term. The cardiovascular short-term consequences are hypertension, dyslipidemia and endothelial dysfunction [16, 24]. In a cohort of 886 obese children, 42% had dyslipidemia and 32% had hypertension [30]. These short-term consequences, which are risk factors for cardiovascular events such as myocardial infarction and stroke, persist in adulthood when obese children become obese adults [31, 32]. BMI during adolescence was associated with death from cardiovascular disease in a follow up of up to 40 years [33]. The metabolic consequences of childhood obesity are insulin resistance (IR), impaired glucose tolerance, and, in a minority of the obese children, type 2 diabetes mellitus (T2DM). In adulthood, developing T2DM is more common.

Pulmonary short-term and long-term consequences are sleep apnoea, asthma and exercise intolerance [34-37]. Regarding the digestive tract, short-term and long-term consequences are equal, these include gall-stones [38, 39], constipation [40-42], hepatic steatosis or non-alcoholic fatty liver disease (NAFLD) [38, 43-45] and gastro-oesophageal reflux [46, 47]. Short-term musculoskeletal problems are flat feet [48-50], lower limb pain [51-53], malalignment of knees and fractures of fore-arm, humerus and femur [54-56]. Fifty to seventy percent of the patients with slipped femoral capital epiphysis was obese [56, 57]. Long-term consequences are arthrosis of knee and hip in early adulthood [58]. Childhood obesity has consequences for the urogenital tract in boys as hypogonadism occurs and in girls by the occurrence of PCOS, which may cause an irregular menstrual cycle, and on long-term subfertility [16, 24, 59]. PCOS is associated with IR [59-61].

Insulin resistance in children with obesity

IR is a state in which increased levels of insulin are measured in absence of diabetes mellitus, which results from peripheral tissues being less sensitive to insulin [62]. IR in obesity develops as a result of the excess fat mass. Fat is an endocrine active tissue, which excretes adipocytokines, such as tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6), and free fatty acids. These adipocytokines induce a chronic inflammatory state [63]. This inflammatory state and the free fatty acid release reduce the muscle glucose uptake, and more insulin is needed to maintain normoglycemia. As a result of this insulin resistance, a compensatory hyperinsulinemia arises. Hyperinsulinemia is correlated with NAFLD [64], hypertriglyceridemia [65, 66], hypertension [66] and T2DM [67-69].

Therefore, IR is described to play a key role in the development of metabolic and cardiovascular consequences in obesity [63, 70-72]. Although IR is related to obesity, not all obese children are insulin resistant, and not all insulin resistant children are obese. IR levels increase with the level of overweight [73, 74]. Other factors influencing IR are for example puberty, in which a physiological decrease of ~25-50% in insulin sensitivity was observed, and gender and ethnicity [75-77]. Black, African-American and Mexican American children have higher levels of both fasted insulin and post-glucose load insulin levels than white children, irrespective of their pubertal state [74, 78-81].

As IR is a precursor of T2DM, it can be considered an early marker for those who are at high risk of T2DM. Although the other precursors of T2DM, impaired fasting glucose (IFG) and impaired glucose tolerance (IGT) are clearly defined by the ADA criteria [82], for IR there is no uniform method and cut off value. As a result, the prevalence of IR reported within the same obese population varies between 40.5% and 80.1% in obese children and adolescents (6-18 years), depending on the definition used [83].

In another study the prevalence of IR in obese children aged 8-10 years was 27.3% or 58.4% depending on the definition used [84].

Treatment of childhood obesity

Lifestyle intervention is the cornerstone of the treatment of obesity, since lifestyle interventions aim to restore the balance between calories ingested and calories burned. The effectivity of lifestyle interventions for the treatment of obesity in children is studied frequently. A Cochrane review and meta-analysis showed a BMI reduction of -3.04 (95% confidence interval (CI) -3.14 to -2.94) and -3.37 (95% CI -3.38 to -3.17) after 6 and 12 months, respectively in children of 12 years and above. The change in BMI-SDS after 6 and 12 months was respectively -0.14 (95%CI -0.17 to -0.12) and -0.14 (-0.18 to -0.10). In younger children (<12 years) after 6 months a small decrease in BMI-SDS was achieved (-0.06 (95%CI -0.12 to -0.01), which declined after 12 months (-0.04 (-0.12 to 0.04) [85]. However, there is large variation in lifestyle programmes, and drop-out rates are high. The effectivity of lifestyle interventions is largely influenced by the motivation of parents. Parent-only interventions for children with obesity showed a reduction in child BMI [86-88]. Finally, to improve the effect of lifestyle interventions for childhood obesity, the use of smartphones and internet-based programmes was studied. Most studies showed an improved compliance and response, and lower drop-out rates. However, no difference in body weight was found between groups using the smartphones or internet-based programmes and the groups receiving standard care [89-91].

In addition to lifestyle intervention, treatment with pharmacological agents, has been explored in obese children and adolescents. The following pharmacological agents have been evaluated in pediatrics [92, 93]: orlistat which is FDA approved for the treatment of obesity, sibutramine (withdrawn in 2010 because of safety reasons [94]), exenatide, and metformin, registered for treatment of T2DM from the age of ten years. According to a Cochrane systematic review, orlistat has shown an additional reducing effect on the absolute BMI in children and adolescents, yet medication related adverse effects such as gastro-intestinal tract symptoms were observed [85]. A review and meta-analysis on the effect of metformin in obese children and adolescents without T2DM concluded that metformin is moderately effective in reducing BMI and IR in hyperinsulinemic obese children and adolescents on short term use (6 months or less): a reduction in mean BMI of 1.42 kg/m² and homeostasis model assessment of insulin resistance (HOMA-IR) score by 2.01 [95]. Long-term data are limited to one study of 48 weeks, in which a reduction in BMI of 0.9 kg/m² was reported in participants receiving metformin versus an increase of 0.2 kg/m² in the placebo group [96].

Finally, the effect of bariatric surgery for severe obese adolescents (BMI of > 40 kg/m² or > 35 kg/m² with associated co-morbidities) has been studied. In three trials, chil-

dren who failed to achieve weight loss with non-invasive treatments were included. Bariatric surgery (most frequently laparoscopic sleeve gastrectomy) resulted in this selected populations in significant changes in BMI compared to lifestyle intervention groups, with a follow up of two to four years. Complications occurred in 4.1-4.4% of the patients [97-99]. Three-year follow up of adolescents who underwent bariatric surgery, showed improvement in weight, cardiometabolic health and weight-related quality of life. Complications included deficiencies in micronutrients [99].

Objectives of this thesis

Childhood obesity is an increasing problem, with IR as an important consequence. The role of IR in the development of T2DM is clear. Given the lack of a generally accepted definition of IR in children, the exact prevalence and incidence of IR are unclear. Therefore, in **chapter 2**, a literature review on the epidemiology of IR in population based studies in pediatric populations is presented.

In **chapter 3**, the variety in definitions for IR in paediatrics is further investigated. The aim of this study is to review all published definitions for IR in children and to apply these definitions to a population of patients with obesity from a pediatric outpatient clinic. The application of all definitions in this population demonstrates the large heterogeneity in definitions.

The clinical application of IR as a screening measure for children at risk of T2DM is investigated in **chapter 4**. In this study the recommended screening for T2DM using fasted plasma glucose (FPG) is compared to a screening combining FPG with a IR measurement, to investigate whether IR is useful as an additional screening to identify more children at risk for T2DM.

As the recommended screening interval for children at risk for T2DM is 3 years, in **chapter 5**, a follow up study is performed in children at risk for T2DM, to evaluate weight, insulin sensitivity, and progression to T2DM approximately 3 years after being diagnosed with overweight/obesity and IR.

In the second part of this thesis, the effect of long-term treatment with metformin in obese children with IR is presented. **Chapter 6a** presents the study protocol of the randomized controlled double-blind trial (RCT) in which the effect of long-term treatment metformin on BMI and IR was studied. In **Chapter 6b** the results of this RCT are presented.

Since treatment effects from clinical trials might differ from the effects in daily clinical practice, the results of the RCT described in **chapter 6**, are compared to the effects of metformin on BMI in daily clinical practice. The aim of **chapter 7** was to compare the effects of metformin (in addition to a lifestyle intervention programme) on change in

BMI between obese adolescents treated with metformin in daily clinical practice and patients who participated in the above mentioned RCT.

A summary of the conclusions of chapter 2-7 is presented in **chapter 8**. Finally, the future perspectives are discussed.

References

1. NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. 2016 Apr 2;387(10026):1377-96.
2. Factsheet no 311: Overweight and obesity [Internet].; 2015. Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/>.
3. Skinner AC, Perrin EM, Skelton JA. Prevalence of obesity and severe obesity in US children, 1999-2014. *Obesity* (Silver Spring). 2016 May;24(5):1116-23.
4. Waterlow JC, Buzina R, Keller W, Lane JM, Nichaman MZ, Tanner JM. The presentation and use of height and weight data for comparing the nutritional status of groups of children under the age of 10 years. *Bull World Health Organ*. 1977;55(4):489-98.
5. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000 May 6;320(7244):1240-3.
6. Cole TJ, Flegal KM, Nicholls D, Jackson AA. Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ*. 2007 Jul 28;335(7612):194.
7. van der Baan-Slootweg OH. Centrumbehandeling van obesitas. *Praktische Pediatie*. 2009(2):124-8.
8. Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents, 1999-2000. *JAMA*. 2002 Oct 9;288(14):1728-32.
9. Lakshman R, Elks CE, Ong KK. Childhood obesity. *Circulation*. 2012 Oct 2;126(14):1770-9.
10. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA*. 2014 Feb 26;311(8):806-14.
11. de Wilde JA, Verkerk PH, Middelkoop BJ. Declining and stabilising trends in prevalence of overweight and obesity in Dutch, Turkish, Moroccan and South Asian children 3-16 years of age between 1999 and 2011 in the Netherlands. *Arch Dis Child*. 2014 Jan;99(1):46-51.
12. Factsheet Resultaten Vijfde Landelijke Groeistudie TNO, 10 juni 2010 [Internet].; 2011 [].
13. Skelton JA, Irby MB, Grzywacz JG, Miller G. Etiologies of obesity in children: nature and nurture. *Pediatr Clin North Am*. 2011 Dec;58(6):1333,54, ix.
14. Lee WW. An overview of pediatric obesity. *Pediatr Diabetes*. 2007 Dec;8 Suppl 9:76-87.
15. Hardy LR, Harrell JS, Bell RA. Overweight in children: definitions, measurements, confounding factors, and health consequences. *J Pediatr Nurs*. 2004 Dec;19(6):376-84.
16. Garver WS, Newman SB, Gonzales-Pacheco DM, Castillo JJ, Jelinek D, Heidenreich RA, et al. The genetics of childhood obesity and interaction with dietary macronutrients. *Genes Nutr*. 2013 May;8(3):271-87.
17. Crino A, Greggio NA, Beccaria L, Schiaffini R, Pietrobelli A, Maffei C. Diagnosis and differential diagnosis of obesity in childhood. *Minerva Pediatr*. 2003 Oct;55(5):461-70.
18. Crocker MK, Yanovski JA. Pediatric obesity: etiology and treatment. *Endocrinol Metab Clin North Am*. 2009 Sep;38(3):525-48.
19. Correll CU, Lencz T, Malhotra AK. Antipsychotic drugs and obesity. *Trends Mol Med*. 2011 Feb;17(2):97-107.
20. Whitaker RC, Wright JA, Pepe MS, Seidel KD, Dietz WH. Predicting obesity in young adulthood from childhood and parental obesity. *N Engl J Med*. 1997 Sep 25;337(13):869-73.
21. Singh AS, Mulder C, Twisk JW, van Mechelen W, Chinapaw MJ. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes Rev*. 2008 Sep;9(5):474-88.

22. Guo SS, Roche AF, Chumlea WC, Gardner JD, Siervogel RM. The predictive value of childhood body mass index values for overweight at age 35 y. *Am J Clin Nutr.* 1994 Apr;59(4):810-9.
23. Engeland A, Bjorge T, Sogaard AJ, Tverdal A. Body mass index in adolescence in relation to total mortality: 32-year follow-up of 227,000 Norwegian boys and girls. *Am J Epidemiol.* 2003 Mar 15;157(6):517-23.
24. Daniels SR. Complications of obesity in children and adolescents. *Int J Obes (Lond).* 2009 Apr;33 Suppl 1:S60-5.
25. Morrison KM, Shin S, Tarnopolsky M, Taylor VH. Association of depression & health related quality of life with body composition in children and youth with obesity. *J Affect Disord.* 2015 Feb 1;172:18-23.
26. Strauss RS. Childhood obesity and self-esteem. *Pediatrics.* 2000 Jan;105(1):e15.
27. Mustillo S, Worthman C, Erkanli A, Keeler G, Angold A, Costello EJ. Obesity and psychiatric disorder: developmental trajectories. *Pediatrics.* 2003 Apr;111(4 Pt 1):851-9.
28. Halfon N, Larson K, Slusser W. Associations between obesity and comorbid mental health, developmental, and physical health conditions in a nationally representative sample of US children aged 10 to 17. *Acad Pediatr.* 2013 Jan-Feb;13(1):6-13.
29. Gortmaker SL, Must A, Perrin JM, Sobol AM, Dietz WH. Social and economic consequences of overweight in adolescence and young adulthood. *N Engl J Med.* 1993 Sep 30;329(14):1008-12.
30. Moniz M, Marques T, Cabral M, Nizarali Z, Coelho R, Monteiro A, et al. Cardiovascular risk factors and childhood obesity. *Acta Med Port.* 2011 Dec;24 Suppl 2:327-32.
31. Srinivasan SR, Frontini MG, Xu J, Berenson GS. Utility of childhood non-high-density lipoprotein cholesterol levels in predicting adult dyslipidemia and other cardiovascular risks: the Bogalusa Heart Study. *Pediatrics.* 2006 Jul;118(1):201-6.
32. Srinivasan SR, Myers L, Berenson GS. Changes in metabolic syndrome variables since childhood in prehypertensive and hypertensive subjects: the Bogalusa Heart Study. *Hypertension.* 2006 Jul;48(1):33-9.
33. Twig G, Yaniv G, Levine H, Leiba A, Goldberger N, Derazne E, et al. Body-Mass Index in 2.3 Million Adolescents and Cardiovascular Death in Adulthood. *N Engl J Med.* 2016 Apr 13.
34. Eisenmann JC, Arnall DA, Kanuho V, Interpretter C, Coast JR. Obesity and pulmonary function in Navajo and Hopi children. *Ethn Dis.* 2007 Winter;17(1):14-8.
35. Lazarus R, Colditz G, Berkey CS, Speizer FE. Effects of body fat on ventilatory function in children and adolescents: cross-sectional findings from a random population sample of school children. *Pediatr Pulmonol.* 1997 Sep;24(3):187-94.
36. Raj D, Kabra SK, Lodha R. Childhood obesity and risk of allergy or asthma. *Immunol Allergy Clin North Am.* 2014 Nov;34(4):753-65.
37. Alonso-Alvarez ML, Cordero-Guevara JA, Teran-Santos J, Gonzalez-Martinez M, Jurado-Luque MJ, Corral-Penafiel J, et al. Obstructive sleep apnea in obese community-dwelling children: the NANOS study. *Sleep.* 2014 May 1;37(5):943-9.
38. Nunes MM, Medeiros CC, Silva LR. Cholelithiasis in obese adolescents treated at an outpatient clinic. *J Pediatr (Rio J).* 2014 Mar-Apr;90(2):203-8.
39. Koebnick C, Getahun D, Smith N, Porter AH, Der-Sarkissian JK, Jacobsen SJ. Extreme childhood obesity is associated with increased risk for gastroesophageal reflux disease in a large population-based study. *Int J Pediatr Obes.* 2011 Jun;6(2-2):e257-63.

40. vd Baan-Slootweg OH, Liem O, Bekkali N, van Aalderen WM, Rijcken TH, Di Lorenzo C, et al. Constipation and colonic transit times in children with morbid obesity. *J Pediatr Gastroenterol Nutr.* 2011 Apr;52(4):442-5.
41. Phatak UP, Pashankar DS. Prevalence of functional gastrointestinal disorders in obese and overweight children. *Int J Obes (Lond).* 2014 Oct;38(10):1324-7.
42. Misra S, Lee A, Gensel K. Chronic constipation in overweight children. *JPEN J Parenter Enteral Nutr.* 2006 Mar-Apr;30(2):81-4.
43. Malespin M, Slesman B, Lau A, Wong SS, Cotler SJ. Prevalence and correlates of suspected nonalcoholic fatty liver disease in Chinese American children. *J Clin Gastroenterol.* 2015 Apr;49(4):345-9.
44. Zhang X, Wan Y, Zhang S, Lu L, Chen Z, Liu H, et al. Nonalcoholic fatty liver disease prevalence in urban school-aged children and adolescents from the Yangtze River delta region: a cross-sectional study. *Asia Pac J Clin Nutr.* 2015;24(2):281-8.
45. AlKhater SA. Paediatric non-alcoholic fatty liver disease: an overview. *Obes Rev.* 2015 May;16(5):393-405.
46. Malaty HM, Fraley JK, Abudayyeh S, Fairly KW, Javed US, Aboul-Fotouh H, et al. Obesity and gastroesophageal reflux disease and gastroesophageal reflux symptoms in children. *Clin Exp Gastroenterol.* 2009;2:31-6.
47. Pashankar DS, Corbin Z, Shah SK, Caprio S. Increased prevalence of gastroesophageal reflux symptoms in obese children evaluated in an academic medical center. *J Clin Gastroenterol.* 2009 May-Jun;43(5):410-3.
48. Mueller S, Carlsohn A, Mueller J, Baur H, Mayer F. Influence of Obesity on Foot Loading Characteristics in Gait for Children Aged 1 to 12 Years. *PLoS One.* 2016 Feb 25;11(2):e0149924.
49. Sadeghi-Demneh E, Azadinia F, Jafarian F, Shamsi F, Melvin JM, Jafarpishe M, et al. Flatfoot and obesity in school-age children: a cross-sectional study. *Clin Obes.* 2016 Feb;6(1):42-50.
50. Stolzman S, Irby MB, Callahan AB, Skelton JA. Pes planus and paediatric obesity: a systematic review of the literature. *Clin Obes.* 2015 Apr;5(2):52-9.
51. Tsiros MD, Buckley JD, Howe PR, Walkley J, Hills AP, Coates AM. Musculoskeletal pain in obese compared with healthy-weight children. *Clin J Pain.* 2014 Jul;30(7):583-8.
52. Paulis WD, Silva S, Koes BW, van Middelkoop M. Overweight and obesity are associated with musculoskeletal complaints as early as childhood: a systematic review. *Obes Rev.* 2014 Jan;15(1):52-67.
53. Adams AL, Kessler JI, Deramerian K, Smith N, Black MH, Porter AH, et al. Associations between childhood obesity and upper and lower extremity injuries. *Inj Prev.* 2013 Jun;19(3):191-7.
54. Benson EC, Miller M, Bosch P, Szalay EA. A new look at the incidence of slipped capital femoral epiphysis in new Mexico. *J Pediatr Orthop.* 2008 Jul-Aug;28(5):529-33.
55. Fornari ED, Suszter M, Roocroft J, Bastrom T, Edmonds EW, Schlechter J. Childhood obesity as a risk factor for lateral condyle fractures over supracondylar humerus fractures. *Clin Orthop Relat Res.* 2013 Apr;471(4):1193-8.
56. Taylor ED, Theim KR, Mirch MC, Ghorbani S, Tanofsky-Kraff M, Adler-Wailes DC, et al. Orthopedic complications of overweight in children and adolescents. *Pediatrics.* 2006 Jun;117(6):2167-74.
57. Must A, Strauss RS. Risks and consequences of childhood and adolescent obesity. *Int J Obes Relat Metab Disord.* 1999 Mar;23 Suppl 2:S2-11.

58. Schonau E. Childhood and adolescent obesity--consequences for the locomotor system and treatment options. *Musculoskeletal complications of overweight children and adolescents. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 2013 Apr;56(4):528-31.
59. Anderson AD, Solorzano CM, McCartney CR. Childhood obesity and its impact on the development of adolescent PCOS. *Semin Reprod Med.* 2014 May;32(3):202-13.
60. DeUgarte CM, Bartolucci AA, Azziz R. Prevalence of insulin resistance in the polycystic ovary syndrome using the homeostasis model assessment. *Fertil Steril.* 2005 May;83(5):1454-60.
61. Huang J, Ni R, Chen X, Huang L, Mo Y, Yang D. Metabolic abnormalities in adolescents with polycystic ovary syndrome in south China. *Reprod Biol Endocrinol.* 2010 Nov 17;8:142,7827-8-142.
62. Kim G, Caprio S. Diabetes and insulin resistance in pediatric obesity. *Pediatr Clin North Am.* 2011 Dec;58(6):1355,61, ix.
63. Cruz ML, Shaibi GQ, Weigensberg MJ, Spruijt-Metz D, Ball GD, Goran MI. Pediatric obesity and insulin resistance: chronic disease risk and implications for treatment and prevention beyond body weight modification. *Annu Rev Nutr.* 2005;25:435-68.
64. Khan UI, McGinn AP, Isasi CR, Groisman-Perelstein A, Diamantis PM, Ginsberg M, et al. Differences in Cardiometabolic Risk between Insulin-Sensitive and Insulin-Resistant Overweight and Obese Children. *Child Obes.* 2015 Jun;11(3):289-96.
65. Asato Y, Katsuren K, Ohshiro T, Kikawa K, Shimabukuro T, Ohta T. Relationship between lipid abnormalities and insulin resistance in Japanese school children. *Arterioscler Thromb Vasc Biol.* 2006 Dec;26(12):2781-6.
66. Bao W, Srinivasan SR, Berenson GS. Persistent elevation of plasma insulin levels is associated with increased cardiovascular risk in children and young adults. The Bogalusa Heart Study. *Circulation.* 1996 Jan 1;93(1):54-9.
67. Dubinina IA, Chistiakov DA, Eremina IA, Brovkin AN, Zilberman LI, Nikitin AG, et al. Studying progression from glucose intolerance to type 2 diabetes in obese children. *Diabetes Metab Syndr.* 2014 Jul-Sep;8(3):133-7.
68. Morrison JA, Glueck CJ, Umar M, Daniels S, Dolan LM, Wang P. Hyperinsulinemia and metabolic syndrome at mean age of 10 years in black and white schoolgirls and development of impaired fasting glucose and type 2 diabetes mellitus by mean age of 24 years. *Metabolism.* 2011 Jan;60(1):24-31.
69. Nguyen QM, Srinivasan SR, Xu JH, Chen W, Berenson GS. Changes in risk variables of metabolic syndrome since childhood in pre-diabetic and type 2 diabetic subjects: the Bogalusa Heart Study. *Diabetes Care.* 2008 Oct;31(10):2044-9.
70. Levy-Marchal C, Arslanian S, Cutfield W, Sinaiko A, Druet C, Marcovecchio ML, et al. Insulin resistance in children: consensus, perspective, and future directions. *J Clin Endocrinol Metab.* 2010 Dec;95(12):5189-98.
71. Lee JM. Insulin resistance in children and adolescents. *Rev Endocr Metab Disord.* 2006 Sep;7(3):141-7.
72. Chiarelli F, Marcovecchio ML. Insulin resistance and obesity in childhood. *Eur J Endocrinol.* 2008 Dec;159 Suppl 1:S67-74.
73. Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics.* 1999 Jun;103(6 Pt 1):1175-82.

74. Lee JM, Okumura MJ, Davis MM, Herman WH, Gurney JG. Prevalence and determinants of insulin resistance among U.S. adolescents: a population-based study. *Diabetes Care*. 2006 Nov;29(11):2427-32.
75. Goran MI, Gower BA. Longitudinal study on pubertal insulin resistance. *Diabetes*. 2001 Nov;50(11):2444-50.
76. Jeffery AN, Metcalf BS, Hosking J, Streeter AJ, Voss LD, Wilkin TJ. Age before stage: insulin resistance rises before the onset of puberty: a 9-year longitudinal study (EarlyBird 26). *Diabetes Care*. 2012 Mar;35(3):536-41.
77. Nightingale CM, Rudnicka AR, Owen CG, Wells JC, Sattar N, Cook DG, et al. Influence of adiposity on insulin resistance and glycemia markers among U.K. Children of South Asian, black African-Caribbean, and white European origin: child heart and health study in England. *Diabetes Care*. 2013 Jun;36(6):1712-9.
78. Gungor N, Saad R, Janosky J, Arslanian S. Validation of surrogate estimates of insulin sensitivity and insulin secretion in children and adolescents. *J Pediatr*. 2004 Jan;144(1):47-55.
79. Arslanian S, Suprasongsin C, Janosky JE. Insulin secretion and sensitivity in black versus white prepubertal healthy children. *J Clin Endocrinol Metab*. 1997 Jun;82(6):1923-7.
80. Arslanian S, Suprasongsin C. Differences in the in vivo insulin secretion and sensitivity of healthy black versus white adolescents. *J Pediatr*. 1996 Sep;129(3):440-3.
81. Svec F, Nastasi K, Hilton C, Bao W, Srinivasan SR, Berenson GS. Black-white contrasts in insulin levels during pubertal development. The Bogalusa Heart Study. *Diabetes*. 1992 Mar;41(3):313-7.
82. American Diabetes Association. 2. Classification and Diagnosis of Diabetes. *Diabetes Care*. 2016 Jan;39 Suppl 1:S13-22.
83. Yin J, Li M, Xu L, Wang Y, Cheng H, Zhao X, et al. Insulin resistance determined by Homeostasis Model Assessment (HOMA) and associations with metabolic syndrome among Chinese children and teenagers. *Diabetol Metab Syndr*. 2013 Nov 15;5(1):71,5996-5-71.
84. Iamopas O, Chongviriyaphan N, Suthutvoravut U. Metabolic syndrome in obese Thai children and adolescents. *J Med Assoc Thai*. 2011 Aug;94 Suppl 3:S126-32.
85. Oude Luttikhuis H, Baur L, Jansen H, Shrewsbury VA, O'Malley C, Stolk RP, et al. Interventions for treating obesity in children. *Cochrane Database Syst Rev*. 2009 Jan 21;(1):CD001872. doi(1):CD001872.
86. Loveman E, Al-Khudairy L, Johnson RE, Robertson W, Colquitt JL, Mead EL, et al. Parent-only interventions for childhood overweight or obesity in children aged 5 to 11 years. *Cochrane Database Syst Rev*. 2015 Dec 21;(12):CD012008. doi(12):CD012008.
87. Ewald H, Kirby J, Rees K, Robertson W. Parent-only interventions in the treatment of childhood obesity: a systematic review of randomized controlled trials. *J Public Health (Oxf)*. 2014 Sep;36(3):476-89.
88. Jansen E, Mulkens S, Jansen A. Tackling childhood overweight: treating parents exclusively is effective. *Int J Obes (Lond)*. 2011 Apr;35(4):501-9.
89. Abraham AA, Chow WC, So HK, Yip BH, Li AM, Kumta SM, et al. Lifestyle intervention using an internet-based curriculum with cell phone reminders for obese Chinese teens: a randomized controlled study. *PLoS One*. 2015 May 6;10(5):e0125673.
90. Chaplais E, Naughton G, Thivel D, Courteix D, Greene D. Smartphone Interventions for Weight Treatment and Behavioral Change in Pediatric Obesity: A Systematic Review. *Telemed J E Health*. 2015 Oct;21(10):822-30.

91. Turner T, Spruijt-Metz D, Wen CK, Hingle MD. Prevention and treatment of pediatric obesity using mobile and wireless technologies: a systematic review. *Pediatr Obes*. 2015 Dec;10(6):403-9.
92. Wald AB, Uli NK. Pharmacotherapy in pediatric obesity: current agents and future directions. *Rev Endocr Metab Disord*. 2009 Sep;10(3):205-14.
93. Boland CL, Harris JB, Harris KB. Pharmacological management of obesity in pediatric patients. *Ann Pharmacother*. 2015 Feb;49(2):220-32.
94. Williams G. Withdrawal of sibutramine in Europe. *BMJ*. 2010 Feb 9;340:c824.
95. Park MH, Kinra S, Ward KJ, White B, Viner RM. Metformin for obesity in children and adolescents: a systematic review. *Diabetes Care*. 2009 Sep;32(9):1743-5.
96. Wilson DM, Abrams SH, Aye T, Lee PD, Lenders C, Lustig RH, et al. Metformin extended release treatment of adolescent obesity: a 48-week randomized, double-blind, placebo-controlled trial with 48-week follow-up. *Arch Pediatr Adolesc Med*. 2010 Feb;164(2):116-23.
97. Al-Sabah SK, Almazeedi SM, Dashti SA, Al-Mulla AY, Ali DA, Jumaa TH. The efficacy of laparoscopic sleeve gastrectomy in treating adolescent obesity. *Obes Surg*. 2015 Jan;25(1):50-4.
98. Alqahtani AR, Elahmedi MO. Pediatric bariatric surgery: the clinical pathway. *Obes Surg*. 2015 May;25(5):910-21.
99. Inge TH, Courcoulas AP, Jenkins TM, Michalsky MP, Helmrath MA, Brandt ML, et al. Weight Loss and Health Status 3 Years after Bariatric Surgery in Adolescents. *N Engl J Med*. 2016 Jan 14;374(2):113-23.

