Cover Page



Universiteit Leiden



The handle <u>http://hdl.handle.net/1887/43361</u> holds various files of this Leiden University dissertation

Author: Samimi-Namin, Kaveh Title: Diversity and distribution of octocorals and scleractinians in the Persian Gulf region Issue Date: 2016-09-28

Chapter 10

Detecting coral bleaching, using QuickBird multi-temporal data: A feasibility study at Kish Island, the Persian Gulf

Keivan Kabiri, Biswajeet Pradhan, Kaveh Samimi-Namin, Masoud Moradi

This chapter has been previously published and re-used here with permission of the publisher. For original publication please refer to: Estuarine, Coastal and Shelf Science 117: 273-281. doi: 10.1016/j.ecss.2012.12.006(http://www.sciencedirect.com/science/article/pii/S0272771412004568)

Abstract

Coral bleaching events have become more frequent and intense worldwide and speculated to be a severe threat for coral survival in future. The Persian Gulf, as one of the warmest seas, has also experienced coral mortalities and bleaching events. Historically, bleaching events are known to occur south of the Persian Gulf, such information is scarce in the northern side. Perhaps remoteness and inaccessibility to Iran main coral communities which have developed on offshore islands can explain such lack of data. To address this issue, the feasibility of using multi-temporal satellite images for detecting past bleaching events were investigated. Two QuickBird images (2005, 2008) were selected to detect 2007 bleaching event at Kish Island, Iran, and the accuracy of results were compared to in situ observations. Current study might represent "algae-challenged" scenario in terms of having 7 months' time lapse between bleaching event and post-bleaching satellite image. As a result of this, we had algae-covered corals instead of white bleached corals. In the proposed procedure pre and post-bleaching images were classified, and changes in reflectance values within coral classes were interpreted as bleaching areas. By using this method, we could eliminate the effect of miss-classification between bleached corals and sand; as well as algae covered corals and live corals. Furthermore, it is not necessary to have a post-bleaching image acquired during bleaching events, although having such image will increase the accuracy. The proposed technique detected ~28% of bleached corals and the results support the idea that coral bleaching can be distinguished by detecting the changes in reflectance values in pre and post-bleaching images. Understanding the occurrence, severity, and extent of past bleaching events may help us understand the population dynamics of Iran corals and reveal coral connectivity patterns in the Persian Gulf.

Keywords: coral reefs, bleaching, multi-temporal satellite images, remote sensing, Persian Gulf, Kish Island

1. Introduction

Coral reefs are under degradation due to anthropogenic stress and natural disturbances worldwide (Wilkinson 2008). As a result of temperature rise in the world oceans, the frequency and the scale of bleaching occurrences have increased since the early 1980s (Glynn 1993; Hoegh-Guldberg 1999; Hoegh-Guldberg et al. 2007; Baker et al. 2008). Climate change scenarios suggest an increase in tropical ocean temperature (IPCC 2007), potentially increasing bleaching events and killing many coral reefs around the world.

The Persian Gulf is known as the warmest sea in the world with high environmental fluctuations and temperature, well above the thermal tolerance for corals elsewhere (Wilkinson and Buddemeier 1994; Sheppard et al. 2010; Riegl et al. 2011). Although, the Persian Gulf corals and their algal symbionts acclimatized and adapted to elevated temperatures (Baker et al. 2004; Obura 2009), they could not escape temperature anomalies and have experienced bleaching events and mass mortalities (Coles and Fadlallah 1991; George and John 1999, 2000; Riegl, 2002, 2003; Riegl et al. 2011). Bleaching is the loss of coral symbiotic algae in response to environmental stresses (Hoegh-Guldberg et al. 2007; van Oppen and Lough 2009). In particular, temperature anomalies (Berkelmans and Willis 1999) will be a severe threat to coral survival for the next 30e50 years (Baker et al. 2008). Corals are not likely to withstand and recover from these events unless they acquire at least 0.2e 0.3 C thermal tolerance per decade (Donner et al. 2005). The Persian Gulf corals already cope with temperatures projected for corals elsewhere in future (Riegl et al. 2011). These corals are close to their tipping point, and further bleaching and mass mortalities will be inevitable in future. The Persian Gulf corals can serve as a good case study to investigate bleaching as well as implementing and testing different techniques to assess such events.

The Iranian coastline of the Persian Gulf is over 1300 km long, and main coral communities are located around offshore islands. Access to most of these areas is difficult, and establishing regular monitoring programs has not been successful so far. One of the current problems of coral studies in Iran is insufficient historical data on corals, their environments, and disturbances in particular bleaching events. This explains why few bleaching reports are available from the northern Persian Gulf compare with the south, where such data have been collected since 80's (Baker et al. 2008). Although, historical surface temperature anomalies can be acquired from satellite data, not all anomalies lead to bleaching. While in situ bleaching assessment is impossible, remote sensing techniques might be beneficial. Remote sensing has proved to be a useful tool for assessing impacts of environmental factors, ecological and geomorphological changes on coral reef communities (Palandro et al. 2003; Andréfouët and Riegl 2004; Mumby et al. 2004); depth estimation (Lyzenga 1978; Stumpf et al. 2003; Mishra et al. 2004), habitat mapping (Mumby et al. 1998; Andréfouët et al. 2003; Purkis and Pasterkamp 2004; Green et al. 2005; Mishra et al. 2006); and bleaching detection across coral communities (Clark et al. 2000; Elvidge et al. 2004; Rowlands et al. 2008). The latter, needs high spatial resolution sensors (Andréfouët et al. 2002), and also using multitemporal images can improve the accuracy of bleaching detection (Yamano and Tamura 2004), especially when suitable satellite image is not available at the time of bleaching incident. Change detection and multi-temporal approach has been popular in land-based surveys and used in marine realms to detect coastline changes, estimation of reef topography, and changes in habitats (Yamano 2007; Scopélitis et al. 2009; Lyons et al. 2011; Al Fugura et al. 2011). Elvidge et al. (2004) provided satisfactory account of multi-temporal images using pseudo-invariant features (PIF) and differential imaging (DI) based approach to detect coral bleaching in the Great Barrier Reef. Rowlands et al. (2008) used similar approach based on spectral radiance for detecting bleaching in Honduras and showed by having adequate habitat knowledge; this method can be valuable for detecting bleaching. Therefore, the multitemporal approach might be beneficial in detecting past bleaching events and may be a costeffective tool to assist future in situ observations and monitoring plans along the Iranian coastline. The current study is one of the first attempts to use multi-temporal images to detect bleaching and to assess its application for acquiring information of the past bleaching events in Iran, and in the Persian Gulf.

Here, we tested the potential and feasibility of using multi-temporal high resolution QuickBird imagery as a source of reference information on presence/absence, frequency, severity, and extent of coral bleaching events. We also examined the potential value and accuracy of such technique to acquire bleaching historical data, especially when there are no field observations available. Finally, we discussed the implications of this study for coral monitoring and management in remote places along Iranian coastline.

2. Materials and methods

2.1. Study area

The investigated area is a shallow subtidal sea bed, sloping gently offshore (~2%) to depth of ~10 m on the southeast side of the Kish Island. The island is a salt diapir and part of the southern Zagros Mountains, with fossil corals dating from the Holocene and upper Pleistocene geological periods (Preusser et al. 2003). It is located at northern side of the Persian Gulf (26° 320' N; 53° 580' E), about 17 km far from Iran mainland, with 90 km² area (Fig. 1). The tidal regime of the area is mixed, with a range of 1.8 m above the lowest astronomical tide datum. Water temperature usually ranges from 22 to 33 °C. Climate is very dry semi-equatorial with ~60% humidity, and air temperature ranges from 15 to 48 °C. The average annual rainfall is ~150 mm, which usually occurs during winter and fall. The site is one of the main diving destinations and therefore receives high anthropogenic impacts. Corals can be found all around the island on hard grounds, however, most developed and dense corals were located at southeast of the island, hence it was chosen as our study area (Fig. 1).

2.2. Field observations and in situ data collection

The distribution and community structure of corals in the study area has been under investigation since 2005 (Samimi-Namin, unpublished data). Only parts of the available data sets were used as training data to improve and verify the results. Corals extend parallel to shoreline in a belt between 2 and 12 m depth (Fig. 1), with variable density, diversity, and surface cover. The area was composed of three main coral assemblages, currently under study: type a, massive dead or alive Porites colonies with, 10–30% cover; type b, non-frame-building *Acropora* colonies, with 40–80% cover; and type c, mixture of Poritids, Faviids, and Siderastreids in deeper areas, with 10–30% cover. It is not known whether the area represents true framework building corals and perhaps can be best classified as non-frame-building coral communities *sensu* Riegl and Piller (1999). For the purpose of the study we consider the mean coral cover of about 50% for the whole coral area.

The bleaching occurred at the study site from 10 to 28 August 2007 (Wilkinson 2008) with

~85% (84.9 \pm 6.3) of total coral colonies being partially or completely bleached during the first two weeks of the incident. During the first few days of bleaching, most of the *Acropora* colonies were bleached while other corals such as Faviids and Poritids, managed to survive. Two weeks later, all *Acropora* colonies were bleached, and even the other persistent genera could not escape the bleaching. Six months later, the coral skeletons were covered by algae and were under deterioration processes by fouling organisms. The extent of bleaching and its severity was overlaid on priori distribution map. This information allowed us to identify bleached areas, which was then compared with satellite images.

2.3. Satellite data

Two QuickBird satellite images were acquired for multi-temporal analysis; pre-bleaching (September 11, 2005); and post bleaching (March 28, 2008). The spatial resolution of the multispectral images of QuickBird satellite equals to 2.44 m with central values of 485 nm, 560 nm, and 660 nm in red, blue, and green bands, respectively. The water level was 1.21 m and 0.51 m above the mean sea level (MSL) at time of acquisition in 2005 and 2008. Only 2005 image showed favourable low cloud cover and water turbidity conditions, though mild waves were apparent. The other image had partial cloud cover over the study area (11%) and high turbidity due to tide. We favoured using only blue and green bands for image interpretation due to low penetration of red band through water (Elvidge et al. 2004; Mishra et al. 2005).

2.4. Image processing and data analysis

A summary of the methods used in the current study is represented in Fig. 2. Pre-processing procedures, i.e., geometric, radiometric, and atmospheric corrections were applied on the satellite images. Several ground control points (GCPs) were selected from an existing vector map (scale = 1:2000) to perform geometric corrections and geo-referencing in the images. For radiometric corrections, ENVI ver. 4.8 software was used to convert the digital number (DN) values to at-sensor radiance L (μ W cm⁻² nm⁻¹ sr⁻¹) (Krause 2005). For atmospheric corrections, the Fast Line-of-Site Atmospheric Analysis of Spectral Hypercubes (FLAASH) module in the same software was utilized to convert the primary radiance to scaled reflectance (Adler-Golden et al. 1999). In addition, features such as terrestrial areas, boats, and jetties, were masked to avoid unpleasant effects on classification (Fig. 3c, d).

The most critical parameter influencing benthic habitat mapping is water column, which attenuates the substrate reflectance values; and this attenuation changes with depth and wavelength of the light passing through water (Lyzenga 1978, 1981). To minimize this effect, Eq. (1) was applied following Bierwirth et al. (1993) and Purkis and Pasterkamp (2004),

$$R_{b} = \left(\frac{1}{0.54}R_{rs}(z=a) - (1 - e^{-2kz})R_{w}\right) \div e^{-2kz}$$
(1)

where Rw is water column reflectance of optically deep water; Rb is wet substrate reflectance; Rrs(z = a) is remotely sensed effective reflectance just above the water/air boundary; and k, is effective diffuse attenuation coefficient of water body. The k values were calculated using the depth of penetration zone approach proposed by Jupp (1988) and used by Mumby and Edwards (2000) (Table 1). The depth values were extracted from a bathymetric map (scale = 1:2000; provided by Kish Free Zone Organization, No. KLS100D0037). The average reflectance values of an area deeper than 15 m on images were considered as Rw. The optical properties of the study area were assumed to be homogenous (Holden and LeDrew 2008).



Fig. 1. Study area at SE of Kish Island, Iran, representing coral communities along the shoreline.

The maximum likelihood classifier was applied to classify main habitat types of the study area (Fig. 3e, f), using over 200 training points. The purpose of classification in our study is to reduce the interferences between sands in detecting bleached corals. Overall accuracy was calculated using testing points which were different than our training data (Table 2). Results were optimized by post-processing operations, such as image filtering (majority filter), which eliminated the salt and pepper effect on classified maps (Eastman et al. 1995).

The differences in reflectance values between two images within the coral classes (ΔRc) were used to detect bleached corals. The average bottom reflectance values of a homogenous shallow sandy area (Rs) and mean value of corrected remotely sensed reflectance values of a deep area (Rw) in both images were computed (Table 1), where $\Delta Rs > \Delta Rc$ $>\Delta Rw$ due to the differences in light absorption through water column. The adjusted reflectance values for coral areas (Ric(adj)), were calculated using Eq. (2), to minimize the systematic errors (e.g., differences in sunlight angle, surf, waves),

where i and j refer to the year of image acquisition (j is the image with greater values). $R_c^{i(adj)} = 0.5 \times \left(\Delta R_s^{j,i} + \Delta R_w^{j,i}\right) + R_c^i$ (2)

The Rc(adj) in both images were plotted against each other to distinguish bleaching areas (Fig. 4, a and b). The Rc(adj) values in both images were multiplied to enhance the differences in reflectance values simultaneously (Fig. 4, c). The live corals will have lower reflectance values compared with bleached corals in each of the blue and green bands (Holden and LeDrew 1998; Yamano and Tamura 2004). Consequently, the pixels representing potential bleaching areas were recognized at a 95% confidence interval. Because of seven months time-lapse between the bleaching event (August 2007) and the second satellite image acquisition date (March 2008), all of the white bleached corals were interpreted as bleaching areas later on.

3. Results

3.1. Multi-temporal processing

The study area can be categorized between Jerlov oceanic water type III and coastal water type 1 (Jerlov 1976), indicating high turbidity and, therefore, high attenuation of light (Table 1). The Rs and Rw showed different values in each image (Table 1). These differences were due to systematic errors caused by the variation of parameters such as sunlight angle, water quality, and amount of sea waves in each image. The pre-processing and water column corrections enhanced the image quality (Fig. 3c, d). The corals were classified into two groups: 1) sparse corals, dominant by type a, and c; 2) dense corals, dominant by type b (Fig. 3). A total of 253 and 249 points (not considered as training data) were checked on 2005 and 2008 during extensive field observations. The overall accuracies of the classifications in 2005 and 2008 were 82.2% and 75.8%, respectively. The atmospheric Fig. 2. Concise methodology schema adopted in the current condition of the 2008 image (slightly study. reduced cloudy) the accuracy,

particularly in coral classes with an



average of 80.7% in 2005, and 66% in 2008 images (Table 2).

The processed QuickBird satellite images showed an increment in reflectance values of some pixels within coral classes in 2008 in comparison to 2005 in blue and green bands (Fig. 4, a and b). This increase was more distinguishable when multiplied values of two bands were considered (Fig. 4, c). Due to the fact that this anomaly occurred within the coral classes (dense and sparse), it should be the result of bleaching in the coral communities and were identified in 2008 (Fig. 5). Eliminating these points (with reflectance values >2s) improved the normality of the remained values (Fig. 4, d, e). As mentioned before, the accuracy of classification for coral classes was w70%; hence, about 30% of the areas with anomalous reflectance values may contain errors. Because there was no actual bleaching event in 2005, therefore the pixels categorized as bleaching areas in 2005 (8 pixels) were considered as error.

3.2. Remotely sensed data versus in situ observations

Priori knowledge of the study area indicated ~85% bleaching in the study area occurred on August 2007. The proposed image processing method was able to detect ~12% bleaching in

the area (mixture of coral and non-coral pixels) (Fig. 5). Considering 50% average coral cover for the whole study area, it can be concluded that ~28% of the total bleached corals were detected (Fig. 5).

4. Discussion

This paper discusses the application of multi-temporal satellite images to detect coral bleaching as a potential tool to acquire information of past bleaching events. Accordingly, the multispectral satellite data for the pre- and post-bleaching events in 2007 were used to assess the capability and accuracy of using this approach compare to in situ observations at one of the Iranian islands of the Persian Gulf.

The application of remote sensing for detecting bleaching and habitat mapping is limited by sensor characteristics (e.g. spatial, spectral, and temporal resolutions), and environmental complexities (e.g. water quality, depth, and selective absorption) (Holden and LeDrew 1998; Andréfouët et al. 2001; Yamano and Tamura 2004). It is known that mapping aquatic vegetation, coral reefs, and general bottom characteristics requires high spatial resolution multi-spectral/hyperspectral imagery (Holden and LeDrew 1999; Mumby and Edwards 2002; Mishra et al. 2006, 2007). On the other hand, clarity of the water column is an important factor influencing accuracy and quality of the results (Yamano and Tamura 2004). One of the drawbacks of using remote sensing in the Persian Gulf is turbidity of the water. An analysis of k among different bands in images, the Persian Gulf waters classifies in between oceanic water type III and coastal water type 1 (Jerlov 1976), indicating high turbidity and, therefore, high attenuation of light (Table 1). Turbidity decreases the water clarity and increases the water column attenuation effects, consequently diminishes the accuracy of results. To solve the turbidity difficulties, an integration of acoustic systems (e.g. side scan sonar and single/multibeam echosounders), satellite remote sensing, and in situ observations will provide more accurate results (Brown et al. 2011). This approach has been investigated in the Persian Gulf and has proved to be correct (Riegl et al. 2001; Riegl and Purkis 2005; Purkis and Riegl 2005). Although, such combined method leads to higher accuracy, it is still limited by extensive field operations and available resources (equipment, technicians, budget, etc.), therefore it might not be cost-effective, especially in remote places along the Iranian coastline.

A further complication is that analyses of a single image might not correctly identify pixels containing both healthy and bleached corals. Identification of bleached corals from carbonate rock and sand is difficult because of their similarity in reflectance (Clark et al. 2000; Rowlands et al. 2008), while, live and dead corals (covered by algae) can be easily separated from sands. Most of the misclassifications occurred between corals (classes 2 & 3) and rubble/ rock (class 4), and minimum interference observed between corals and sand (Table 2). This is in agreement with the previous studies which indicated that due to similarities in optical properties of corals and rocks, most interferences during classification occur between these two classes and not between corals and sandy areas (Riegl and Purkis 2005; Mishra et al. 2006). In the proposed method, the sand class was not considered in bleaching detection, therefore, the interference between bleached corals and sand is at minimum level (Table 2).

Yamano and Tamura (2004) showed that it is difficult to detect bleached corals if the amount of bleaching within an image pixel of the satellite is less than 23%. Andréfouët et al. (2002) showed that underestimation in bleaching detection increases when the resolution decreases. For example, at 2 m resolution aerial photographs, they observed 0.2e12.5% underestimation



Fig. 3. Image processing from raw to classified image for 2005 and 2008; (a-b) false colour composite images (FCC,421) of 2005 and 2008 (FCC,421), (c-d) bottom reflectance values of the 2005 and 2008 images (after radiometric corrections, eliminating land and non-desired objects, and water column corrections), (e-f) classified substrates extracted from in 2005 and 2008.

Table 1. The estimated diffuse attenuation
values (k), reflectance values of optically deep
water (R_{μ}) , and reflectance values of shallow
sandy areas (R) for three visible bands in
2005 and 2008 satellite images.
e

Year	k (m ⁻¹))		$R_{\rm w}$ (\times	10 ²)		$R_{\rm s}(\times 10^2)$				
	Red	Green	Blue	Red	Green	Blue	Red	Green	Blue		
2005	0.481	0.169	0.133	3.99	4.48	4.72	22.8	34.2	28.1		
2008	0.462	0.156	0.129	7.8	7.96	8.1	34.9	41.6	35.9		

Table 2. Accuracy assessment of classifications and confusion matrix in 2005 and 2008.1 = Sand, 2 = dense coral, 3 = sparse coral, 4 = rock and rubble, 5 = deep area. The bold numbers represent the number of correctly classified pixels.

2005								2008							
Class	1	2	3	4	5	Total	Producer accuracy (%)	Class	1	2	3	4	5	Total	Producer accuracy (%)
1	44	0	1	4	2	51	86.3	1	40	1	2	5	1	49	81.6
2	3	36	6	4	0	49	73.5	2	2	31	11	5	1	50	62
3	0	4	39	6	1	50	78	3	1	6	33	5	0	45	73.3
4	3	5	2	41	1	52	78.9	4	2	7	4	38	2	53	71.7
5	1	0	0	3	49	53	92.5	5	1	2	0	2	47	52	90.4
Total	51	45	48	58	53	253		Total	46	47	50	55	51	249	
User accuracy (%)	86.3	80.0	81.3	70.7	92.5			User accuracy (%)	87.0	66.0	66.0	69.1	92.2		
Overall accuracy 82.2%, $\kappa = 78.2\%$						Overall accuracy 75.8%, $\kappa = 69.9\%$									

compare to in situ data. Therefore, to obtain the most accurate results, the resolution should be close to colony average size; and without large bleached colonies or contiguous colony structures bleaching event cannot be detected at a resolution coarser than 2 m (Andréfouët et al. 2002). On the other hand, susceptibilities to bleaching are also different among coral taxa (Marshall and Baird 2000), therefore the bleaching estimation varies according to size, growth form, and susceptibility of the dominant species (Andréfouët et al. 2002). This suggests that the amount of bleaching recognized in satellite image varies across different coral assemblages and therefore comparison of *in situ* data to satellite data might not be useful unless the boundaries and coverage of each coral assemblage are known. As most of the study area comprises sparse corals and most of these corals have less than 0.5 m diameter size, unless they show high coverage they cannot be detected with high accuracy. Hence, most of the detected bleaching pixels are located in dense coral class where dominated by high cover and large size Acropora colonies (Figs. 3e-f, 5). Rowlands et al. (2008) also found that on high-density coral cover the image showed noticeably higher at-sensor radiance than reference image, however, when bleaching was 20% of total cover the radiance did not show any changes.

The PIF approach used by Elvidge et al. (2004) requires the availability of an image acquired during a bleaching event. Our approach is not limited by availability of such image, though having such image will increase the precession and accuracy of final results. Rowlands et al. (2008) followed the same approach, but did not provide any comparison between their tested approach and in situ observations.

Comparison of multi-temporal images allows more accurate assessment of the detection limits for bleached corals in a pixel, because an increase in reflectance will be recorded where bleaching has taken place (Yamano and Tamura 2004). The patterns of increased radiance associated with bleached coral in bands 1 and 2 in this study support finding elsewhere (Holden and LeDrew 1998; Elvidge et al. 2004; Rowlands et al. 2008) and when both of the images were considered at the same time this increase is more distinguishable (Fig. 4c). By suing the known spatial distribution of coral habitat avoided the sand/coral interface, and prevented misclassifying bleached coral as sand (Rowlands et al. 2008).

Healthy corals show low reflectance, while bleached corals show higher reflectance (Holden and LeDrew 1998). If corals do not recover from bleaching, their white carbonate skeleton will be covered by algae in few weeks or months (Clark et al. 2000). The same phenomena happened in our study area. Because of time difference between bleaching event (August 2007) and the post-bleaching image (March 2008), the bleached coral spectral signals were

Chapter 10



Fig. 4. Relationship between reflectance values. (a) Scatter chart of the corrected bottom reflectance values (Rb) of green band, (b) scatter chart of the corrected bottom reflectance values of blue band, (c) Scatter chart of the multiplied corrected bottom reflectance values in green and blue bands. Pixels more than $2 \times \sigma$ from the average value are indicated by black squares, (d) The multiplied values and normalized values before eliminating bleached corals, (e) Multiplied values and normalized values after eliminating bleached corals.



Fig. 5. Comparison of multi-temporal analysis and in situ bleaching extent from field observations.

masked by algae. Therefore, we dealt with bleached corals covered with algae and not any white bleached colony in our study (algae-challenged scenario). The similarity of algae colour range to live corals leads to complication in distinction between live and dead corals (Clark et al. 2000), however, the proposed approach will eliminate such problems (Fig. 4 c, d). The accuracy of bleaching detection potentially can be increased by decreasing the time laps between post-bleaching image and bleaching event. Although, there were no white bleached corals on 2008 image and all of bleached corals were covered by algae, the proposed method was able to detect 28% bleaching within coral boundaries (Fig. 5). This, suggest that although algae have lower reflectance than bleached corals, it is still higher than the reflectance of live corals at the study area (Fig. 4). Whether this much accuracy is enough to say bleaching

happened without any field observations is still subject to more tests and is a matter of user choice.

4.1. Implications for management

As a result of counter clock wise water circulation pattern in the Persian Gulf, and better environmental conditions along the Iranian coastline, coral species richness is higher compare to the southern part (Samimi-Namin and van Ofwegen 2009, 2012; Riegl et al. 2012). Main Iran coral communities have developed around the remote islands, where field monitoring and in situ observations are not always practical and in some cases even impossible. Our study provides a demonstration of how remote sensing can play a role in monitoring inaccessible and rarely visited coral areas along the Iranian coastline. The 2007 bleaching at northern side of the Persian Gulf caused severe coral mortalities at some of the Iranian islands, while virtually there was no bleaching report from the southern part at the same time (Baker et al. 2008). This suggests that the bleaching patterns and temperature anomalies are not constant throughout the Persian Gulf. Without field observation, the bleaching event demonstrated here could have been easily overlooked and perhaps it is the case for most of Iranian corals and explains the insufficient bleaching records along Iranian coastline. The proposed method can detect major changes in coral communities such as bleaching; even without field observations. Such information can be used by managers for planning further field operations and future data collection. Integrating multi-temporal approach together and SST data set will provide a powerful tool to track bleaching events in the past. Iranian corals are among the least known corals within the Persian Gulf and likely to be the main source of larvae for coral recruitment in the south part and therefore knowledge of bleaching

frequency, severity, and distribution is vital and might reveal the whole coral population dynamics and connectivity patterns within the Persian Gulf environment (Sheppard et al. 2010).

5. Conclusion

In this study, multi-temporal remotely sensed images and in situ data sets were combined to produce a spatial pattern of bleaching extent in coral communities at SE of Kish Island, Iran. This is the first time such an approach was tested along the Iranian coastline and the surrounding region. The priori distribution map and coral communities knowledge, made it possible to explore and compare the results with field observations. The study allowed us to explore and assess the usage of multi-temporal satellite images in monitoring bleaching events. The technique proved to be useful for recognizing bleaching, even by using an image acquired several months after the bleaching event, when all white bleached corals were covered by algae. Furthermore, distinguishing coral communities during classification, the interferences between sand and bleached corals; as well as healthy and dead corals was in minimum level. Therefore, by using more images and reducing the time lapses between image acquisition dates and bleaching events, more accurate results can be produced. Combining commercial satellite remote sensing observations with in situ data can produce accurate results and provide baseline information for future studies.

Author contributions

Conceived and designed the experiments: KK KSN. Performed the experiments: KSN KK. Analysed the data: KK KSN. Contributed reagents/materials/analysis tools: KK KSN BP MM. Wrote the paper: KSN KK BP.

Acknowledgements

H. Rezai, H. Alizadeh, and V. Chegini (Iranian National Institute for Oceanography) are acknowledged for their support and for facilitating the field surveys at Kish Island. M. Mohammadi (Kish Free Zone Organization) and A. Mirshahidi (Nautilus Diving Center, Kish) are appreciated for support during the field survey and facilitating diving activities. The research at NCB Naturalis and partial field work was supported by Schure-BeijerinckPoppingfonds (KNAW), Alida Buitendijkfonds, Jan Joost ter Pelkwijkfonds, and Martin-Fellowship (NCB Naturalis). The Alfred P. Sloan Foundation and the Census of Marine Life are gratefully acknowledged for the research grant provided to KSN; in this regard, N. D'Adamo (UNESCO, IOC, Perth), J.H. Ausubel (Rockefeller University), and P. Miloslavich (Universidad Simón Bolívar) are greatly appreciated for their continued support and encouragement. This research was supported by UPM University Research Grant (05-01-11-1283RU) to Stimulate Research (RUGS scheme, vote number 9199892) to BP. The authors are grateful to anonymous reviewers for providing constructive comments.

References

- Adler-Golden SM, Matthew MW, Bernstein LS, Levine RY, Berk A, Richtsmeier SC, Acharya PK, Anderson GP, Felde G, Gardner J, Hoke M, Jeong LS, Pukall B, Ratkowski A, Burke HH (1999) Atmospheric correction for Short-wave spectral imagery based on MODTRAN4. SPIE Proceedings on Imaging Spectrometry 3753, 61–69.
- Al Fugura A, Billa L, Pradhan B (2011) Semi-automated procedures for shoreline extraction using single RADARSAT-1 SAR image. Estuarine, Coastal and Shelf Science 95, 395–400.
- Andréfouët S, Muller-Karger FE, Hochberg EJ, Hu C, Carder KL (2001) Change detection in shallow coral reef environments using Landsat 7 ETM+ data. Remote Sensing of Environment 78, 150–162.
- Andréfouët S, Berkelmans R, Odriozola L, Done T, Oliver J, Müller-Karger F (2002) Choosing the appropriate spatial resolution for monitoring coral bleaching events using remote sensing. Coral Reefs 21, 147–154.
- Andréfouët S, Kramer P, Torres-Pulliza D, Joyce KE, Hochberg EJ, GarzaPérez R, Mumby PJ, Riegl B, Yamano H, White WH, Zubia M, Brock JC, Phinn SR, Naseer A, Hatcher BG, Muller-Karger FE (2003) Multi-site evaluation of IKONOS data for classification of tropical coral reef environments. Remote Sensing of Environment 88, 128–143.

- Andréfouët S, Riegl B (2004) Remote sensing: a key tool for interdisciplinary assessment of coral reef processes. Coral Reefs 23, 1e4.
- Baker AC, Starger CJ, McClanahan TR, Glynn PW (2004) Corals' adaptive response to climate change. Nature 430, 741.
- Baker AC, Glynn PW, Riegl B (2008) Climate change and coral reef bleaching: an ecological assessment of long-term impacts, recovery trends and future outlook. Estuarine, Coastal and Shelf Science 80, 435–471.
- Berkelmans R, Willis B (1999) Seasonal and local spatial patterns in the upper thermal limits of corals on the inshore Central Great Barrier Reef. Coral Reefs 18, 219–228.
- Bierwirth PN, Lee TJ, Burne RV (1993) Shallow sea floor reflectance and water depth derived by un mixing multispectral imagery. Photogrammetric Engineering and Remote Sensing 59, 331-338.
- Brown CJ, Smith SJ, Lawton P, Anderson JT (2011) Benthic habitat mapping: a review of progress towards improved understanding of the spatial ecology of the seafloor using acoustic techniques. Estuarine, Coastal and Shelf Science 92, 502–520.
- Clark CD, Mumby PJ, Chisholm JRM, Jaubert J, Andréfouët S (2000) Spectral discrimination of coral mortality states following a severe bleaching event. International Journal of Remote Sensing 21, 2321–2327.
- Coles SL, Fadlallah YH (1991) Reef coral survival and mortality at low temperatures in the Arabian Gulf new speciesspecific lower temperature limits. Coral Reefs 9 (4), 231–237.
- Donner SD, Skirving WJ, Little CM, Oppenheimer M, Hoegh-Guldberg O (2005) Global assessment of coral bleaching and required rates of adaptation under climate change. Global Change Biology 11, 2251e2265. http:// dx.doi.org/ 10.1111/j.1365e2486.2005.01073.x.
- Eastman JR, Jin W, Kyem PAK, Toledano J (1995) Raster procedures for multi-criteria/multi-objective decisions. Photogrammetric Engineering and Remote Sensing 61 (5), 539–547.
- Elvidge C, Dietz J, Berkelmans R, Andréfouët S, Skirving W, Strong A, Tuttle B (2004) Satellite observation of Keppel Islands (Great Barrier Reef) 2002 coral bleaching using IKONOS data. Coral Reefs 23 (3), 461–462.
- George JD, John DM (1999) High sea temperatures along the coast of Abu Dhabi (UAE), Arabian Gulf their impact upon corals and macroalgae. Reef Encounter 25, 21–23.
- George JD, John DM (2000) The effects of the recent prolonged high seawater temperatures on the coral reefs of Abu Dhabi (UAE). In: Proc International Symposium on Extent of Coral Bleaching, pp. 28–29.
- Glynn P (1993) Coral reef bleaching: ecological perspectives. Coral Reefs 12 (1),1–17.
- Green EP, Mumby PJ, Edwards AJ, Clark CD (2005) Remote Sensing Handbook for Tropical Coastal Management.
- Hoegh-Guldberg O (1999) Coral bleaching, climate change and the future of the world's coral reefs. Review. Marine and Freshwater Research 50, 839–866.
- Hoegh-Guldberg O, Anthony K, Berkelmans RWC, Dove S, Fabricius KE, Lough JM, Marshall PA, van Oppen MJH, Negri AP, Willis BL (2007) Vulnerability of Reef-building Corals on the Great Barrier Reef to Climate Change. pp. 271–307.
- Holden H, LeDrew E (1998) Spectral discrimination of healthy and non-healthy corals based on cluster analysis, principal components analysis and derivative spectroscopy. Remote Sensing of Environment 65, 217–224.
- Holden H, LeDrew E (1999) Hyperspectral identification of coral reef features. International Journal of Remote Sensing 20, 2545e2563.
- Holden H, LeDrew E (2008) An examination of variability in vertical radiometric profiles in a coral reef environment. Journal of Coastal Research 241, 224e231. IPCC, 2007. In: Pachauri, R., Reisinger, A. (Eds.), Climate Change Synthesis Report. Contribution of Working Groups I, II, and III to the Fourth Assessment of the Intergovernmental Panel on Climate Change, p. 104 (Geneva).
- Jerlov NG (1976) Marine Optics. Elsevier, Amsterdam, 231 pp.
- Jupp DLB (1988) Background and extensions to depth of penetration (DOP) mapping in shallow coastal waters. In: Proceedings of the Symposium on Remote Sensing of the Coastal Zone, Gold Coast, Queensland, pp. IV.2.1– IV.2.19.
- Krause K (2005) Radiometric Use of QuickBird Imagery. Technical Note, 2005–1107. Digital Globe Inc., Longmont, CO, US.
- Lyons M, Phinn S, Roelfsema C (2011) Integrating Quickbird multi-spectral satellite and field data: mapping bathymetry, seagrass cover, seagrass species and change in Moreton Bay, Australia in 2004 and 2007. Remote Sensing 3, 42–64.
- Lyzenga DR (1978) Passive remote sensing techniques for mapping water depth and bottom features. Applied Optics 17, 379–383.
- Lyzenga DR (1981) Remote sensing of bottom reflectance and water attenuation parameters in shallow water using aircraft and Landsat data. International Journal of Remote Sensing 2, 71–82.
- Marshall P, Baird A. (2000) Bleaching of corals on the Great Barrier Reef: differential susceptibilities among taxa. Coral Reefs 19, 155–163.

- Mishra, D.R., Narumalani, S., Lawson, M., Rundquist, D., 2004. Bathymetric mapping using IKONOS multispectral data. GIScience and Remote Sensing 41 (4), 301e321.
- Mishra DR, Narumalani S, Rundquist D, Lawson M (2005) Characterizing the vertical diffuse attenuation coefficient for down welling irradiance in coastal waters: implications for water penetration by high resolution satellite data. ISPRS Journal of Photogrammetry and Remote Sensing 60, 48–64.
- Mishra DR, Narumalani S, Rundquist D, Lawson M (2006) Benthic habitat mapping in tropical marine environments using QuickBird multispectral data. Photogrammetric Engineering and Remote Sensing 72 (9), 1037–1048.
- Mishra DR, Narumalani S, Rundquist D, Lawson M (2007) Enhancing the detection and classification of coral reef and associated benthic habitats: a hyperspectral remote sensing approach. Journal of Geophysical Research 112, C08014. http://dx.doi.org/10.1029/2006JC003892.
- Mumby PJ, Clark CD, Green EP, Edwards AJ (1998) Benefits of water column correction and contextual editing for mapping coral reefs. International Journal of Remote Sensing 19, 203–210.
- Mumby PJ, Edwards AJ (2000) Water column correction techniques. In: Green EP, Mumby PJ, Edwards AJ, Clark CD (Eds.), Remote Sensing Handbook for Tropical Coastal Management. Unesco, Paris, p. 316.
- Mumby PJ, Edwards AJ (2002) Mapping marine environments with IKONOS imagery: enhanced spatial resolution can deliver great thematic accuracy. Remote Sensing of Environment 82, 248–257.
- Mumby PJ, Skirving W, Strong AE, Hardy JT, LeDrew EF, Hochberg EJ, Stumpf RP, David LT (2004) Remote sensing of coral reefs and their physical environment. Marine Pollution Bulletin 48, 219–228.
- Obura DO (2009) Reef corals bleach to resist stress. Marine Pollution Bulletin 58, 206–212. http://dx.doi. org/10.1016/j.marpolbul.2008.10.002.
- Palandro D, Andréfouët S, Dustan P, Muller-Karger FE (2003) Change detection in coral reef communities using Ikonos satellite sensor imagery and historic aerial photographs. International Journal of Remote Sensing 24, 873–878.
- Preusser F, Radtke U, Fontugne M, Haghipour A, Hilgers A, Kasper HU, Nazari H, Pirazzoli PA (2003) ESR dating of raised coral reefs from Kish Island, Persian Gulf. Quaternary Science Reviews 22, 1317–1322.
- Purkis SJ, Pasterkamp R (2004) Integrating in situ reef-top reflectance spectra with Landsat TM imagery to aid shallow-tropical benthic habitat mapping. Coral Reefs 23, 5–20.
- Purkis SJ, Riegl BM (2005) Spatial and temporal dynamics of Arabian Gulf coral assemblages quantified from remotesensing and in situ monitoring data. Marne Ecology and Progress Series 287, 99–113.
- Riegl B, Piller WE (1999) Coral frame works revisited-reefs and coral carpets in the northern Red Sea. Coral Reefs 18, 241–253.
- Riegl BM, Korrubel JL, Martin C (2001) Mapping and monitoring of coral communities and their spatial patterns using a surface-based video method from a vessel. Bulletin of Marine Science 69 (2), 869–880.
- Riegl BM (2002) Effects of the 1996 and 1998 positive sea-surface temperature anomalies on corals, coral diseases and fish in the Arabian Gulf (Dubai, UAE). Marine Biology 140, 29–40.
- Riegl BM (2003) Climate change and coral reefs: different effects in two high latitude areas (Arabian Gulf, South Africa). Coral Reefs 22, 433-446.
- Riegl BM, Purkis SJ (2005) Detection of shallow subtidal corals from IKONOS satellite and QTC View (50, 200 kHz) single-beam sonar data (Arabian Gulf; Dubai, UAE). Remote Sensing of Environment 95, 96–114.
- Riegl BM, Purkis SJ, Al-Cibahy AS, Abdel-Moati MA, Hoegh-Guldberg O (2011) Present limits to heat-adaptability in corals and population-level responses to climate extremes. PLoS ONE 6 (9), e24802. http://dx.doi.org/10.1371/journal.pone.0024802.
- Riegl BM, Benzoni F, Samimi-Namin K, Sheppard C (2012) The hermatypic scleractinian (Hard) coral fauna of the gulf. In: Riegl, B.M., Purkis, S.J. (Eds.), Coral Reefs of the Gulf: Adaptation to Climatic Extremes. Springer, The Netherlands, pp. 187–224.
- Rowlands G, Purkis S, Riegl B (2008) The 2005 coral bleaching event Roatan (Honduras): use of pseudoinvariant features (PIFs) in satellite assessments. Journal of Spatial Science 53, 99–112.
- Samimi-Namin K, van Ofwegen LP (2009) Some shallow water octocorals (Coelenterata: Anthozoa) of the Persian Gulf. Zootaxa, 1–52.
- Samimi-Namin K, van Ofwegen L (2012) The octocoral fauna of the gulf. In: Riegl BM, Purkis SJ (Eds.), Coral Reefs of the Gulf: Adaptation to Climatic Extremes. Springer, Netherlands, pp. 225–252.
- Scopélitis J, Andréfouët S, Phinn S, Chabanet P, Naim O, Tourrand C, Done T (2009) Changes of coral communities over 35 years: integrating in situ and remote-sensing data on Saint-Leu Reef (la Réunion, Indian Ocean). Estuarine, Coastal and Shelf Science 84, 342–352.
- Sheppard C, Al-Husiani M, Al-Jamali F, Al-Yamani F, Baldwin R, Bishop J, Benzoni F, Dutrieux E, Dulvy NK, Durvasula SR, Jones DA, Loughland R, Medio D, Nithyanandan M, Pilling GM, Polikarpov I, Price AR, Purkis S, Riegl B, Saburova M, Samimi-Namin K, Taylor O, Wilson S, Zainal K (2010) The Gulf: a young sea in decline. Marine Pollution Bulletin 60, 13–38.

- Stumpf RP, Holderied K, Sinclair M (2003) Determination of water depth with high-resolution satellite imagery over variable bottom types. Limnology and Oceanography 48 (1, part 2), 547–556.
- van Oppen MJH, Lough JM (2009) Coral Bleaching: Patterns, Processes, Causes and Consequences. Springer, 178 pp.
- Wilkinson CR, Buddemeier RW (1994) Global Climate Change and Coral Reefs. Implications for People and Reefs. Report of the UNEP-IOC-ASPEI-IUCN Global Task Team on the implications of climate change on coral reefs. IUCN, Gland, Switzerland, 124 pp.
- Wilkinson CR (2008) Status of the Coral Reefs of the World: 2008. Status of Coral Reefs of the World: 2008. Global Coral Reef Monitoring Network and Reef and Rainforest Research Center, Townsville, Australia. 296pp.
- Yamano H, Tamura M (2004) Detection limits of coral reef bleaching by satellite remote sensing: simulation and data analysis. Remote Sensing of Environment 90, 86–103.
- Yamano H (2007) The use of multi-temporal satellite images to estimate intertidal reef-flat topography. Journal of Spatial Science 52, 73–79.