

Diversity and distribution of octocorals and scleractinians in the Persian Gulf region

Samimi-Namin, K.

Citation

Samimi-Namin, K. (2016, September 28). *Diversity and distribution of octocorals and scleractinians in the Persian Gulf region*. Retrieved from https://hdl.handle.net/1887/43361

Version:	Not Applicable (or Unknown)
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/43361

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <u>http://hdl.handle.net/1887/43361</u> holds various files of this Leiden University dissertation

Author: Samimi-Namin, Kaveh Title: Diversity and distribution of octocorals and scleractinians in the Persian Gulf region Issue Date: 2016-09-28

Appendix 2

The second observation of a live *Trimuricea* species (Octocorallia: Plexauridae)

Kaveh Samimi-Namin, Leen P. van Ofwegen

This appendix has been previously published and re-used here with permission of the publisher. For original publication please refer to: Coral Reefs 28: 517-517. doi: 10.1007/s00338-009-0471-2 (http://link.springer.com/article/10.1007/s00338-009-0471-2)

The genus *Trimuricea* (Octocorallia: Plexauridae) was established by Gordon (1926), with four species characterized by the presence of tripartite thornscales in the polyps (Fig. 1). The genus was not observed again until Grasshoff (1999) added a fifth species from New Caledonia, *T. caledonica*. Grasshoff and Bargibant (2001) were the first to show an underwater picture of a species of *Trimuricea*, that of *T. caledonica*. Here, for the second time, a live *Trimuricea* species is presented (Fig. 2). It was photographed and sampled from Larak

Island in the Persian Gulf (Fig. 3). The living tissue is yellowish white, polyps are magenta, and sclerites are colourless. The species is still under investigation; it shows affinities with *T. reticulata* (Thomson and Simpson, 1909), described from the Andaman Islands, Indian Ocean.

Acknowledgements

The first author wants to thank Dr. H. Rezai for encouragement and support during field trips. The funding for field trips was supported by the Iranian National Center for Oceanography (INCO) and Hormozgan University, specially Dr. V. Chegini and Dr. Kamrani are appreciated for their supports.

References

- Gordon I (1926) Notes on a number of muriceid genera (Alcyonaria, Gorgonaceae), with special reference to spiculation. Proceedings of the scientific meetings of the Zoological Society of London, pp 509–531.
- Grasshoff M (1999) The shallow water gorgonians of New Caledonia and adjacent islands (Coelenterata: Octocorallia). Senckenb Biol 78(1/2):1–245.
- Grasshoff M, Bargibant G (2001) Coral Reef Gorgonians of New Caledonia. Institut de recherche pour le developpement, Paris.
- Thomson JA, Simpson JJ (1909) An account of the alcyonarians collected by the Royal Indian Marine Survey Ship Investigator in the Indian Ocean; with a report on the species of *Dendronephthya* by Henderson WD II. The alcyonarians of the littoral area. The Indian Museum, Calcutta.



Fig. 3 Geographic location of Larak Island in the Strait of Hormuz



Fig. 2 A live colony of Trimuricea in Larak Island



Fig. 1 Scanning electron micrographs of characteristic tripartite sclerites of *Trimuricea*