Cover Page



Universiteit Leiden



The handle <u>http://hdl.handle.net/1887/43361</u> holds various files of this Leiden University dissertation

Author: Samimi-Namin, Kaveh Title: Diversity and distribution of octocorals and scleractinians in the Persian Gulf region Issue Date: 2016-09-28

Appendix 5

Raised reef on Larak suggests Acropora dominance on the Persian Gulf coral reefs since the Pleistocene

Samimi-Namin K, Riegl B

This appendix has been previously published and re-used here with permission of the publisher. For original publication please refer to: Coral Reefs 31: 763-763. doi:10.1007/s00338-012-0908-x (http://link.springer.com/article/10.1007/s00338-012-0908-x)

We show hitherto undescribed raised coral frameworks exposed at the NE coast of Larak (Iran), a salt diapir island, located in the Strait of Hormuz, Persian Gulf (Fig. 1a). The entire outcropping section is ~7 m thick. At least two fossil reef sequences were recognized, separated by clastic sediment layers: one near mean sea level (MSL) and the other one >4 m above MSL, the upper 2.5 m of which consist mostly of *Acropora* plates of the *clathrata/downingi* group (Fig. 1 b–d). Preliminary U/Th, 14C dates are problematic but suggest an upper Pleistocene age ranging from 30 to 150 ka. This is comparable to similar outcrops at nearby Hormoz and Qeshm (Bruthans et al. 2006). Outcrops suggest several consecutive layers of *Acropora*

dominance, comparable to the situation in the modern Gulf until recently, but not yet reported from other known raised deposits in the region (Pirazzoli et al. 2004; Bruthans et al. 2006).

The Larak outcrop allows us for the first time to clearly establish that Acropora has been a dominant member of Persian Gulf coral assemblages since the Pleistocene. While Acropora remain common in the northern Gulf, rapidly recurring bleaching events caused their abrupt decline in the southern Gulf (Riegl et al. 2011). Their dire situation in the southern Gulf may herald an interruption of long-term ecological persistence of Acropora dominance. This is eerily similar to the situation observed in the Caribbean (Pandolfi and Jackson 2006).

References

Bruthans J, Filippi M, Gers'l M, Zare

M, Melkova´ J, Pazdur A, Bosa´k



Pandolfi JM, Jackson JBC (2006) Ecological persistence interrupted in Caribbean coral reefs. Ecol Lett 9:818-826.

Pirazzoli PA, Reyss J-L, Funtugne M, Haghipour A, Hilgers A, Kasper HU, Nazari H, Preusser F, Radtke U (2004) Quaternary coral-reef terraces from Kish and Qeshm Islands, Persian Gulf: new radiometric ages and tectonic implications. Quaternary Int 120:15–27.

Riegl B, Purkis S, Al-Cibahy A, Abdel-Moati M, Hoegh-Guldberg O (2011) Present limits to heat- adaptability in corals and population-level responses to climate extremes.PLoS One 6/9.e24802: 1–8.



Fig. 1 a Overview of the outcrop. $\mathbf{b}-\mathbf{c}$ Details of *Acropora* tables in growth position. **d** *Acropora* dominance continues to the very top of the section, where it becomes obscured by heavier diagenesis.