



Universiteit
Leiden
The Netherlands

Glucocorticoid signature in a neuronal genomic context

Polman, J.A.E.

Citation

Polman, J. A. E. (2016, May 10). *Glucocorticoid signature in a neuronal genomic context*. Retrieved from <https://hdl.handle.net/1887/39295>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/39295>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/39295> holds various files of this Leiden University dissertation

Author: Polman, J.A.E.

Title: Glucocorticoid signature in a neuronal genomic context

Issue Date: 2016-05-10

References

- Adams, M., Meijer, O.C., Wang, J., Bhargava, A., and Pearce, D. (2003). Homodimerization of the glucocorticoid receptor is not essential for response element binding: activation of the phenylethanolamine N-methyltransferase gene by dimerization-defective mutants. *Mol. Endocrinol.* *17*, 2583–2592.
- Akhavan, D., Cloughesy, T.F., and Mischel, P.S. (2010). mTOR signaling in glioblastoma: lessons learned from bench to bedside. *Neuro. Oncol.* *12*, 882–889.
- Alboni, S., Tascadda, F., Corsini, D., Benatti, C., Caggia, F., Capone, G., Barden, N., Blom, J.M., and Brunello, N. (2011). Stress induces altered CRE/CREB pathway activity and BDNF expression in the hippocampus of glucocorticoid receptor-impaired mice. *Neuropharmacology* *60*, 1337–1346.
- Alvarez, D.N., Joels, M., and Krugers, H.J. (2003). Chronic unpredictable stress impairs long-term potentiation in rat hippocampal CA1 area and dentate gyrus in vitro. *Eur. J. Neurosci.* *17*, 1928–1934.
- Allen, J.M., Martin, J.B., and Heinrich, G. (1987). Neuropeptide Y gene expression in PC12 cells and its regulation by nerve growth factor: a model for developmental regulation. *Brain Res.* *427*, 39–43.
- Altonsy, M.O., Sasse, S.K., Phang, T.L., and Gerber, A.N. (2014). Context-dependent cooperation between nuclear factor kappaB (NF-kappaB) and the glucocorticoid receptor at a TNFAIP3 intronic enhancer: a mechanism to maintain negative feedback control of inflammation. *J. Biol. Chem.* *289*, 8231–8239.
- Andrus, B.M., Blizinsky, K., Vedell, P.T., Dennis, K., Shukla, P.K., Schaffer, D.J., Radulovic, J., Churchill, G.A., and Redei, E.E. (2012). Gene expression patterns in the hippocampus and amygdala of endogenous depression and chronic stress models. *Mol. Psychiatry* *17*, 49–61.
- Antonow-Schlorke, I., Schwab, M., Li, C., and Nathanielsz, P.W. (2003). Glucocorticoid exposure at the dose used clinically alters cytoskeletal proteins and presynaptic terminals in the fetal baboon brain. *J. Physiol* *547*, 117–123.
- Araya-Callis, C., Hiemke, C., Abumaria, N., and Flugge, G. (2012). Chronic psychosocial stress and citalopram modulate the expression of the glial proteins GFAP and NDRG2 in the hippocampus. *Psychopharmacology (Berl)* *224*, 209–222.
- Ayalew, M., Le-Niculescu, H., Levey, D.F., Jain, N., Changala, B., Patel, S.D., Winiger, E., Breier, A., Shekhar, A., Amdur, R., Koller, D., Nurnberger, J.I., Corvin, A., Geyer, M., Tsuang, M.T., Salomon, D., Schork, N.J., Fanous, A.H., O'Donovan, M.C., and Niculescu, A.B. (2012). Convergent functional genomics of schizophrenia: from comprehensive understanding to genetic risk prediction. *Mol. Psychiatry* *17*, 887–905.
- Bagamasbad, P., Ziera, T., Borden, S.A., Bonett, R.M., Rozeboom, A.M., Seasholtz, A., and Denver, R.J. (2012). Molecular basis for glucocorticoid induction of the Kruppel-like factor 9 gene in hippocampal neurons. *Endocrinology* *153*, 5334–5345.
- Bailey, T.L. and Elkan, C. (1994). Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proc. Int. Conf. Intell. Syst. Mol. Biol.* *2*, 28–36.
- Bailey, T.L., Williams, N., Mischel, C., and Li, W.W. (2006). MEME: discovering and analyzing DNA and protein sequence motifs. *Nucleic Acids Res.* *34*, W369–W373.
- Balic, A., Adams, D., and Mina, M. (2009). Prx1 and Prx2 cooperatively regulate the morphogenesis of the medial region of the mandibular process. *Dev. Dyn.* *238*, 2599–2613.

- Bardwell,V.J. and Treisman,R. (1994). The POZ domain: a conserved protein-protein interaction motif. *Genes Dev.* 8, 1664–1677.
- Barsegyan,A., Mackenzie,S.M., Kurose,B.D., McGaugh,J.L., and Roozendaal,B. (2010). Glucocorticoids in the prefrontal cortex enhance memory consolidation and impair working memory by a common neural mechanism. *Proc. Natl. Acad. Sci. U. S. A* 107, 16655–16660.
- Barski,A., Cuddapah,S., Cui,K., Roh,T.Y., Schones,D.E., Wang,Z., Wei,G., Chepelev,I., and Zhao,K. (2007). High-resolution profiling of histone methylations in the human genome. *Cell* 129, 823–837.
- Beato,M., Chalepakis,G., Schauer,M., and Slater,E.P. (1989). DNA regulatory elements for steroid hormones. *J. Steroid Biochem.* 32, 737–747.
- Bekinschtein,P., Katche,C., Slipczuk,L.N., Igaz,L.M., Cammarota,M., Izquierdo,I., and Medina,J.H. (2007). mTOR signaling in the hippocampus is necessary for memory formation. *Neurobiol. Learn. Mem.* 87, 303–307.
- Biddie,S.C., John,S., and Hager,G.L. (2010). Genome-wide mechanisms of nuclear receptor action. *Trends Endocrinol. Metab* 21, 3–9.
- Biddie,S.C., John,S., Sabo,P.J., Thurman,R.E., Johnson,T.A., Schiltz,R.L., Miranda,T.B., Sung,M.H., Trump,S., Lightman,S.L., Vinson,C., Stamatoyannopoulos,J.A., and Hager,G.L. (2011). Transcription factor AP1 potentiates chromatin accessibility and glucocorticoid receptor binding. *Mol. Cell* 43, 145–155.
- Binder,E.B. (2009). The role of FKBP5, a co-chaperone of the glucocorticoid receptor in the pathogenesis and therapy of affective and anxiety disorders. *Psychoneuroendocrinology* 34 *Suppl 1*, S186–S195.
- Binder,E.B., Salyakina,D., Lichtner,P., Wochnik,G.M., Ising,M., Putz,B., Papiol,S., Seaman,S., Lucae,S., Kohli,M.A., Nickel,T., Kunzel,H.E., Fuchs,B., Majer,M., Pfennig,A., Kern,N., Brunner,J., Modell,S., Baghai,T., Deiml,T., Zill,P., Bondy,B., Rupprecht,R., Messer,T., Kohnlein,O., Dabitz,H., Bruckl,T., Muller,N., Pfister,H., Lieb,R., Mueller,J.C., Lohmusaar,E., Strom,T.M., Bettecken,T., Meitinger,T., Uhr,M., Rein,T., Holsboer,F., and Muller-Myhok,B. (2004). Polymorphisms in FKBP5 are associated with increased recurrence of depressive episodes and rapid response to antidepressant treatment. *Nat. Genet.* 36, 1319–1325.
- Biola,A., Andreau,K., David,M., Sturm,M., Haake,M., Bertoglio,J., and Pallardy,M. (2000). The glucocorticoid receptor and STAT6 physically and functionally interact in T-lymphocytes. *FEBS Lett.* 487, 229–233.
- Blankenberg,D., Von,K.G., Coraor,N., Ananda,G., Lazarus,R., Mangan,M., Nekrutenko,A., and Taylor,J. (2010). Galaxy: a web-based genome analysis tool for experimentalists. *Curr. Protoc. Mol. Biol. Chapter 19*, Unit-21.
- Bledsoe,R.K., Montana,V.G., Stanley,T.B., Delves,C.J., Apolito,C.J., McKee,D.D., Consler,T.G., Parks,D.J., Stewart,E.L., Willson,T.M., Lambert,M.H., Moore,J.T., Pearce,K.H., and Xu,H.E. (2002). Crystal structure of the glucocorticoid receptor ligand binding domain reveals a novel mode of receptor dimerization and coactivator recognition. *Cell* 110, 93–105.
- Bodnoff,S.R., Humphreys,A.G., Lehman,J.C., Diamond,D.M., Rose,G.M., and Meaney,M.J. (1995). Enduring effects of chronic corticosterone treatment on spatial learning, synaptic plasticity, and hippocampal neuropathology in young and mid-aged rats. *J. Neurosci.* 15, 61–69.

- Bosia, C., Osella, M., Baroudi, M.E., Cora, D., and Caselle, M. (2012). Gene autoregulation via intronic microRNAs and its functions. *BMC. Syst. Biol.* 6, 131.
- Boulikroun, S., Fay, M., Zennaro, M.C., Escoubet, B., Jaisser, F., Blot-Chabaud, M., Farman, N., and Courtois-Coutry, N. (2002). Characterization of rat NDRG2 (N-Myc downstream regulated gene 2), a novel early mineralocorticoid-specific induced gene. *J. Biol. Chem.* 277, 31506–31515.
- Bourgeron, T. (2009). A synaptic trek to autism. *Curr. Opin. Neurobiol.* 19, 231–234.
- Bourtchuladze, R., Frenguelli, B., Blendy, J., Cioffi, D., Schutz, G., and Silva, A.J. (1994). Deficient long-term memory in mice with a targeted mutation of the cAMP-responsive element-binding protein. *Cell* 79, 59–68.
- Brose, K., Bland, K.S., Wang, K.H., Arnott, D., Henzel, W., Goodman, C.S., Tessier-Lavigne, M., and Kidd, T. (1999). Slit proteins bind Robo receptors and have an evolutionarily conserved role in repulsive axon guidance. *Cell* 96, 795–806.
- Brudno, M., Do, C.B., Cooper, G.M., Kim, M.F., Davydov, E., Green, E.D., Sidow, A., and Batzoglou, S. (2003). LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic DNA. *Genome Res.* 13, 721–731.
- Bruna, A., Nicolas, M., Munoz, A., Kyriakis, J.M., and Caelles, C. (2003). Glucocorticoid receptor-JNK interaction mediates inhibition of the JNK pathway by glucocorticoids. *EMBO J.* 22, 6035–6044.
- Bruni, F., Polosa, P.L., Gadaleta, M.N., Cantatore, P., and Roberti, M. (2010). Nuclear respiratory factor 2 induces the expression of many but not all human proteins acting in mitochondrial DNA transcription and replication. *J. Biol. Chem.* 285, 3939–3948.
- Cannarile, L., Zollo, O., D'Adamio, F., Ayroldi, E., Marchetti, C., Tabilio, A., Bruscoli, S., and Riccardi, C. (2001). Cloning, chromosomal assignment and tissue distribution of human GILZ, a glucocorticoid hormone-induced gene. *Cell Death. Differ.* 8, 201–203.
- Carroll, J.S., Meyer, C.A., Song, J., Li, W., Geistlinger, T.R., Eeckhoute, J., Brodsky, A.S., Keeton, E.K., Fertuck, K.C., Hall, G.F., Wang, Q., Bekiranov, S., Sementchenko, V., Fox, E.A., Silver, P.A., Gingeras, T.R., Liu, X.S., and Brown, M. (2006). Genome-wide analysis of estrogen receptor binding sites. *Nat. Genet.* 38, 1289–1297.
- Cereseto, M., Reines, A., Ferrero, A., Sifonios, L., Rubio, M., and Wikinski, S. (2006). Chronic treatment with high doses of corticosterone decreases cytoskeletal proteins in the rat hippocampus. *Eur. J. Neurosci.* 24, 3354–3364.
- Champagne, D.L., Bagot, R.C., van, H.F., Ramakers, G., Meaney, M.J., de Kloet, E.R., Joels, M., and Krugers, H. (2008). Maternal care and hippocampal plasticity: evidence for experience-dependent structural plasticity, altered synaptic functioning, and differential responsiveness to glucocorticoids and stress. *J. Neurosci.* 28, 6037–6045.
- Chan, O., Inouye, K., Akirav, E.M., Park, E., Riddell, M.C., Matthews, S.G., and Vranic, M. (2005). Hyperglycemia does not increase basal hypothalamo-pituitary-adrenal activity in diabetes but it does impair the HPA response to insulin-induced hypoglycemia. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 289, R235–R246.
- Chandler, V.L., Maler, B.A., and Yamamoto, K.R. (1983). DNA sequences bound specifically by glucocorticoid receptor in vitro render a heterologous promoter hormone responsive in vivo. *Cell* 33, 489–499.
- Chang, C.M. and Chao, C.C. (2013). Protein kinase CK2 enhances Mcl-1 gene expression through the serum response factor-mediated pathway in the rat hippocampus. *J. Neurosci. Res.* 91, 808–817.

- Chang,T.J., Scher,B.M., Waxman,S., and Scher,W. (1993). Inhibition of mouse GATA-1 function by the glucocorticoid receptor: possible mechanism of steroid inhibition of erythroleukemia cell differentiation. *Mol. Endocrinol.* 7, 528–542.
- Chao,H.M., Choo,P.H., and McEwen,B.S. (1989). Glucocorticoid and mineralocorticoid receptor mRNA expression in rat brain. *Neuroendocrinology* 50, 365–371.
- Charmandari,E., Chrousos,G.P., Ichijo,T., Bhattacharyya,N., Vottero,A., Souvatzoglou,E., and Kino,T. (2005). The human glucocorticoid receptor (hGR) beta isoform suppresses the transcriptional activity of hGRalpha by interfering with formation of active coactivator complexes. *Mol. Endocrinol.* 19, 52–64.
- Chebotaev,D., Yemelyanov,A., and Budunova,I. (2007). The mechanisms of tumor suppressor effect of glucocorticoid receptor in skin. *Mol. Carcinog.* 46, 732–740.
- Chen,Y., Dube,C.M., Rice,C.J., and Baram,T.Z. (2008). Rapid loss of dendritic spines after stress involves derangement of spine dynamics by corticotropin-releasing hormone. *J. Neurosci.* 28, 2903–2911.
- Chen,Y.J., Tan,B.C., Cheng,Y.Y., Chen,J.S., and Lee,S.C. (2010). Differential regulation of CHOP translation by phosphorylated eIF4E under stress conditions. *Nucleic Acids Res.* 38, 764–777.
- Chinenov,Y., Gupte,R., Dobrovolna,J., Flammer,J.R., Liu,B., Michelassi,F.E., and Rogatsky,I. (2012). Role of transcriptional coregulator GRIP1 in the anti-inflammatory actions of glucocorticoids. *Proc. Natl. Acad. Sci. U. S. A* 109, 11776–11781.
- Choi,K.C., Kim,S.H., Ha,J.Y., Kim,S.T., and Son,J.H. (2010). A novel mTOR activating protein protects dopamine neurons against oxidative stress by repressing autophagy related cell death. *J. Neurochem.* 112, 366–376.
- Chong,Z.Z., Shang,Y.C., Wang,S., and Maiese,K. (2012). Shedding new light on neurodegenerative diseases through the mammalian target of rapamycin. *Prog. Neurobiol.* 99, 128–148.
- Chrousos,G.P. and Kino,T. (2009). Glucocorticoid signaling in the cell. Expanding clinical implications to complex human behavioral and somatic disorders. *Ann. N. Y. Acad. Sci.* 1179, 153–166.
- Cole,T.J., Blendy,J.A., Monaghan,A.P., Kriegstein,K., Schmid,W., Aguzzi,A., Fantuzzi,G., Hummler,E., Unsicker,K., and Schutz,G. (1995). Targeted disruption of the glucocorticoid receptor gene blocks adrenergic chromaffin cell development and severely retards lung maturation. *Genes Dev.* 9, 1608–1621.
- Conway-Campbell,B.L., Pooley,J.R., Hager,G.L., and Lightman,S.L. (2012). Molecular dynamics of ultradian glucocorticoid receptor action. *Mol. Cell Endocrinol.* 348, 383–393.
- Conway-Campbell,B.L., Sarabdjitsingh,R.A., McKenna,M.A., Pooley,J.R., Kershaw,Y.M., Meijer,O.C., de Kloet,E.R., and Lightman,S.L. (2010). Glucocorticoid ultradian rhythmicity directs cyclical gene pulsing of the clock gene period 1 in rat hippocampus. *J. Neuroendocrinol.* 22, 1093–1100.
- Corradetti,M.N., Inoki,K., and Guan,K.L. (2005). The stress-induced proteins RTP801 and RTP801L are negative regulators of the mammalian target of rapamycin pathway. *J. Biol. Chem.* 280, 9769–9772.
- Costeas,P.A. and Chinsky,J.M. (2000). Glucocorticoid regulation of branched-chain alpha-ketoacid dehydrogenase E2 subunit gene expression. *Biochem. J.* 347, 449–457.
- Costello,D.A., Claret,M., Al-Qassab,H., Plattner,F., Irvine,E.E., Choudhury,A.I., Giese,K.P., Withers,D.J., and Pedarzani,P. (2012). Brain deletion of insulin receptor substrate 2 disrupts hippocampal synaptic plasticity and metaplasticity. *PLoS. One.* 7, e31124.

- Dai,J., Wang,X., Chen,Y., Wang,X., Zhu,J., and Lu,L. (2009). Expression quantitative trait loci and genetic regulatory network analysis reveals that Gabraz is involved in stress responses in the mouse. *Stress*. 12, 499–506.
- Datson,N.A., Meijer,L., Steenbergen,P.J., Morsink,M.C., van der Laan,S., Meijer,O.C., and de Kloet,E.R. (2004). Expression profiling in laser-microdissected hippocampal subregions in rat brain reveals large subregion-specific differences in expression. *Eur. J. Neurosci*. 20, 2541–2554.
- Datson,N.A., Morsink,M.C., Meijer,O.C., and de Kloet,E.R. (2008). Central corticosteroid actions: Search for gene targets. *Eur. J. Pharmacol*. 583, 272–289.
- Datson,N.A., Polman,J.A., de Jonge,R.T., van Boheemen,P.T., van Maanen,E.M., Welten,J., McEwen,B.S., Meiland,H.C., and Meijer,O.C. (2011). Specific regulatory motifs predict glucocorticoid responsiveness of hippocampal gene expression. *Endocrinology* 152, 3749–3757.
- Datson,N.A., Speksnijder,N., Mayer,J.L., Steenbergen,P.J., Korobko,O., Goeman,J., de Kloet,E.R., Joels,M., and Lucassen,P.J. (2012). The transcriptional response to chronic stress and glucocorticoid receptor blockade in the hippocampal dentate gyrus. *Hippocampus* 22, 359–371.
- Datson,N.A., van den Oever,J.M., Korobko,O.B., Magarinos,A.M., de Kloet,E.R., and McEwen,B.S. (2013). Previous history of chronic stress changes the transcriptional response to glucocorticoid challenge in the dentate gyrus region of the male rat hippocampus. *Endocrinology* 154, 3261–3272.
- Datson,N.A., van der Perk,J., de Kloet,E.R., and Vreugdenhil,E. (2001a). Expression profile of 30.000 genes in rat hippocampus using SAGE. *Hippocampus* 11, 430–444.
- Datson,N.A., van der Perk,J., de Kloet,E.R., and Vreugdenhil,E. (2001b). Identification of corticosteroid-responsive genes in rat hippocampus using serial analysis of gene expression. *Eur. J. Neurosci*. 14, 675–689.
- De Bosscher K., Van,C.K., Meijer,O.C., and Haegeman,G. (2008). Selective transrepression versus transactivation mechanisms by glucocorticoid receptor modulators in stress and immune systems. *Eur. J. Pharmacol*. 583, 290–302.
- De Bosscher K., Vanden Berghe,W., and Haegeman,G. (2006). Cross-talk between nuclear receptors and nuclear factor kappaB. *Oncogene* 25, 6868–6886.
- de Kloet,E.R. (2014). From receptor balance to rational glucocorticoid therapy. *Endocrinology* 155, 2754–2769.
- de Kloet,E.R., Joels,M., and Holsboer,F. (2005). Stress and the brain: from adaptation to disease. *Nat. Rev. Neurosci*. 6, 463–475.
- de Kloet,E.R., Vreugdenhil,E., Oitzl,M.S., and Joels,M. (1998). Brain corticosteroid receptor balance in health and disease. *Endocr. Rev*. 19, 269–301.
- de Lange,P., Segeren,C.M., Koper,J.W., Wiemer,E., Sonneveld,P., Brinkmann,A.O., White,A., Brogan,I.J., de Jong,F.H., and Lamberts,S.W. (2001). Expression in hematological malignancies of a glucocorticoid receptor splice variant that augments glucocorticoid receptor-mediated effects in transfected cells. *Cancer Res*. 61, 3937–3941.
- De Rijk,R.H., Schaaf,M., Stam,F.J., de Jong,I.E., Swaab,D.F., Ravid,R., Vreugdenhil,E., Cidlowski,J.A., de Kloet,E.R., and Lucassen,P.J. (2003). Very low levels of the glucocorticoid receptor beta isoform in the human hippocampus as shown by Taqman RT-PCR and immunocytochemistry. *Brain Res. Mol. Brain Res*. 116, 17–26.

- Deweindt,C., Albagli,O., Bernardin,F., Dhordain,P., Quief,S., Lantoine,D., Kerckaert,J.P., and Leprince,D. (1995). The LAZ3/BCL6 oncogene encodes a sequence-specific transcriptional inhibitor: a novel function for the BTB/POZ domain as an autonomous repressing domain. *Cell Growth Differ.* 6, 1495–1503.
- Di Nardo A., Kramvis,I., Cho,N., Sadowski,A., Meikle,L., Kwiatkowski,D.J., and Sahin,M. (2009). Tuberous sclerosis complex activity is required to control neuronal stress responses in an mTOR-dependent manner. *J. Neurosci.* 29, 5926–5937.
- Djordjevic,J., Djordjevic,A., Adzic,M., Mitic,M., Lukic,I., and Radojcic,M.B. (2015). Alterations in the Nrf2-Keap1 signaling pathway and its downstream target genes in rat brain under stress. *Brain Res.* 1602, 20–31.
- Drouin,J., Sun,Y.L., Chamberland,M., Gauthier,Y., De,L.A., Nemer,M., and Schmidt,T.J. (1993). Novel glucocorticoid receptor complex with DNA element of the hormone-repressed POMC gene. *EMBO J.* 12, 145–156.
- Du,J., Wang,Y., Hunter,R., Wei,Y., Blumenthal,R., Falke,C., Khairova,R., Zhou,R., Yuan,P., Machado-Vieira,R., McEwen,B.S., and Manji,H.K. (2009). Dynamic regulation of mitochondrial function by glucocorticoids. *Proc. Natl. Acad. Sci. U. S. A* 106, 3543–3548.
- Dwyer,J.M., Lepack,A.E., and Duman,R.S. (2012). mTOR activation is required for the antidepressant effects of mGluR(2)/(3) blockade. *Int. J. Neuropsychopharmacol.* 15, 429–434.
- Ellisen,L.W., Ramsayer,K.D., Johannessen,C.M., Yang,A., Beppu,H., Minda,K., Oliner,J.D., McKeon,F., and Haber,D.A. (2002). REDD1, a developmentally regulated transcriptional target of p63 and p53, links p63 to regulation of reactive oxygen species. *Mol. Cell* 10, 995–1005.
- Euskirchen,G.M., Rozowsky,J.S., Wei,C.L., Lee,W.H., Zhang,Z.D., Hartman,S., Emanuelson,O., Stolc,V., Weissman,S., Gerstein,M.B., Ruan,Y., and Snyder,M. (2007). Mapping of transcription factor binding regions in mammalian cells by ChIP: comparison of array- and sequencing-based technologies. *Genome Res.* 17, 898–909.
- Fang,Z.H., Lee,C.H., Seo,M.K., Cho,H., Lee,J.G., Lee,B.J., Park,S.W., and Kim,Y.H. (2013). Effect of treadmill exercise on the BDNF-mediated pathway in the hippocampus of stressed rats. *Neurosci. Res.* 76, 187–194.
- Feldker,D.E., Morsink,M.C., Veenema,A.H., Datson,N.A., Proutski,V., Lathouwers,D., de Kloet,E.R., and Vreugdenhil,E. (2006). The effect of chronic exposure to highly aggressive mice on hippocampal gene expression of non-aggressive subordinates. *Brain Res.* 1089, 10–20.
- Fernandez-Chacon,R., Konigstorfer,A., Gerber,S.H., Garcia,J., Matos,M.F., Stevens,C.F., Brose,N., Rizo,J., Rosenmund,C., and Sudhof,T.C. (2001). Synaptotagmin I functions as a calcium regulator of release probability. *Nature* 410, 41–49.
- Ferreira,R., Ohneda,K., Yamamoto,M., and Philipsen,S. (2005). GATA1 function, a paradigm for transcription factors in hematopoiesis. *Mol. Cell Biol.* 25, 1215–1227.
- First,M., Gil-Ad,I., Taler,M., Tarasenko,I., Novak,N., and Weizman,A. (2011). The effects of fluoxetine treatment in a chronic mild stress rat model on depression-related behavior, brain neurotrophins and ERK expression. *J. Mol. Neurosci.* 45, 246–255.
- Gallihier-Beckley,A.J. and Cidlowski,J.A. (2009). Emerging roles of glucocorticoid receptor phosphorylation in modulating glucocorticoid hormone action in health and disease. *IUBMB. Life* 61, 979–986.

- Gauthier, J.M., Bourachot, B., Doucas, V., Yaniv, M., and Moreau-Gachelin, F. (1993). Functional interference between the Spi-1/PU.1 oncoprotein and steroid hormone or vitamin receptors. *EMBO J.* *12*, 5089–5096.
- Gemert Van, arvalho, D.M., arst, H., an der Laan, S., hang, M., eijer, O.C., ell, J.W., and oels, M. (2009). Dissociation between rat hippocampal CA1 and dentate gyrus cells in their response to corticosterone: effects on calcium channel protein and current. *Endocrinology* *150*, 4615–4624.
- Gemert van, N.G., Meijer, O.C., Morsink, M.C., and Joels, M. (2006). Effect of brief corticosterone administration on SGK1 and RGS4 mRNA expression in rat hippocampus. *Stress.* *9*, 165–170.
- Gertz, J., Savic, D., Varley, K.E., Partridge, E.C., Safi, A., Jain, P., Cooper, G.M., Reddy, T.E., Crawford, G.E., and Myers, R.M. (2013). Distinct properties of cell-type-specific and shared transcription factor binding sites. *Mol. Cell* *52*, 25–36.
- Gillespie, C.F., Phifer, J., Bradley, B., and Ressler, K.J. (2009). Risk and resilience: genetic and environmental influences on development of the stress response. *Depress. Anxiety.* *26*, 984–992.
- Glass, C.K. and Saijo, K. (2010). Nuclear receptor transrepression pathways that regulate inflammation in macrophages and T cells. *Nat. Rev. Immunol.* *10*, 365–376.
- Goecks, J., Nekrutenko, A., and Taylor, J. (2010). Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. *Genome Biol.* *11*, R86.
- Goshima, Y., Ito, T., Sasaki, Y., and Nakamura, F. (2002). Semaphorins as signals for cell repulsion and invasion. *J. Clin. Invest* *109*, 993–998.
- Gould, E., McEwen, B.S., Tanapat, P., Galea, L.A., and Fuchs, E. (1997). Neurogenesis in the dentate gyrus of the adult tree shrew is regulated by psychosocial stress and NMDA receptor activation. *J. Neurosci.* *17*, 2492–2498.
- Gourley, S.L., Wu, F.J., Kiraly, D.D., Ploski, J.E., Kedves, A.T., Duman, R.S., and Taylor, J.R. (2008). Regionally specific regulation of ERK MAP kinase in a model of antidepressant-sensitive chronic depression. *Biol. Psychiatry* *63*, 353–359.
- Gray, J.D., Rubin, T.G., Hunter, R.G., and McEwen, B.S. (2014). Hippocampal gene expression changes underlying stress sensitization and recovery. *Mol. Psychiatry* *19*, 1171–1178.
- Greene, J.G., Borges, K., and Dingledine, R. (2009). Quantitative transcriptional neuroanatomy of the rat hippocampus: evidence for wide-ranging, pathway-specific heterogeneity among three principal cell layers. *Hippocampus* *19*, 253–264.
- Greene, L.A. and Tischler, A.S. (1976). Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc. Natl. Acad. Sci. U. S. A* *73*, 2424–2428.
- Gromak, N. (2012). Intronic microRNAs: a crossroad in gene regulation. *Biochem. Soc. Trans.* *40*, 759–761.
- Gupta, S., Stamatoyannopoulos, J.A., Bailey, T.L., and Noble, W.S. (2007). Quantifying similarity between motifs. *Genome Biol.* *8*, R24.
- Gutsaeva, D.R., Suliman, H.B., Carraway, M.S., Demchenko, I.T., and Piantadosi, C.A. (2006). Oxygen-induced mitochondrial biogenesis in the rat hippocampus. *Neuroscience* *137*, 493–504.

- Harris,A.P., Holmes,M.C., de Kloet,E.R., Chapman,K.E., and Seckl,J.R. (2012). Mineralocorticoid and glucocorticoid receptor balance in control of HPA axis and behaviour. *Psychoneuroendocrinology*.
- He,X., Chatterjee,R., John,S., Bravo,H., Sathyanarayana,B.K., Biddie,S.C., FitzGerald,P.C., Stamatoyannopoulos,J.A., Hager,G.L., and Vinson,C. (2013). Contribution of nucleosome binding preferences and co-occurring DNA sequences to transcription factor binding. *BMC Genomics* 14, 428.
- Heine,V.M., Maslam,S., Zareno,J., Joels,M., and Lucassen,P.J. (2004). Suppressed proliferation and apoptotic changes in the rat dentate gyrus after acute and chronic stress are reversible. *Eur. J. Neurosci.* 19, 131–144.
- Hellman,L.M. and Fried,M.G. (2007). Electrophoretic mobility shift assay (EMSA) for detecting protein-nucleic acid interactions. *Nat. Protoc.* 2, 1849–1861.
- Her,S., Claycomb,R., Tai,T.C., and Wong,D.L. (2003). Regulation of the rat phenylethanolamine N-methyltransferase gene by transcription factors Sp1 and MAZ. *Mol. Pharmacol.* 64, 1180–1188.
- Herman,J.P. (2013). Neural control of chronic stress adaptation. *Front Behav. Neurosci.* 7, 61.
- Herrlich,P. and Ponta,H. (1994). Mutual cross-modulation of steroid/retinoic acid receptor and AP-1 transcription factor activities: a novel property with practical implications. *Trends Endocrinol. Metab* 5, 341–346.
- Hoeffler,C.A. and Klann,E. (2010). mTOR signaling: at the crossroads of plasticity, memory and disease. *Trends Neurosci.* 33, 67–75.
- Hoo,R.L., Chu,J.Y., Yuan,Y., Yeung,C.M., Chan,K.Y., and Chow,B.K. (2010). Functional identification of an intronic promoter of the human glucose-dependent insulinotropic polypeptide gene. *Gene* 463, 29–40.
- Horie-Inoue,K., Takayama,K., Bono,H.U., Ouchi,Y., Okazaki,Y., and Inoue,S. (2006). Identification of novel steroid target genes through the combination of bioinformatics and functional analysis of hormone response elements. *Biochem. Biophys. Res. Commun.* 339, 99–106.
- Hou,L. and Klann,E. (2004). Activation of the phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling pathway is required for metabotropic glutamate receptor-dependent long-term depression. *J. Neurosci.* 24, 6352–6361.
- Howell,J.J. and Manning,B.D. (2011). mTOR couples cellular nutrient sensing to organismal metabolic homeostasis. *Trends Endocrinol. Metab* 22, 94–102.
- Howell,K.R., Kutiyawalla,A., and Pillai,A. (2011). Long-term continuous corticosterone treatment decreases VEGF receptor-2 expression in frontal cortex. *PLoS One.* 6, e20198.
- Howland,J.G. and Wang,Y.T. (2008). Synaptic plasticity in learning and memory: stress effects in the hippocampus. *Prog. Brain Res.* 169, 145–158.
- Huang,d.W., Sherman,B.T., and Lempicki,R.A. (2009a). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 37, 1–13.
- Huang,d.W., Sherman,B.T., and Lempicki,R.A. (2009b). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* 4, 44–57.
- Huang,X., Zhang,H., Yang,J., Wu,J., McMahon,J., Lin,Y., Cao,Z., Gruenthal,M., and Huang,Y. (2010). Pharmacological inhibition of the mammalian target of rapamycin pathway suppresses acquired epilepsy. *Neurobiol. Dis.* 40, 193–199.

- Hudson, W.H., Youn, C., and Ortlund, E.A. (2013). The structural basis of direct glucocorticoid-mediated transrepression. *Nat. Struct. Mol. Biol.* 20, 53–58.
- Hunter, R.G., McCarthy, K.J., Milne, T.A., Pfaff, D.W., and McEwen, B.S. (2009). Regulation of hippocampal H₃ histone methylation by acute and chronic stress. *Proc. Natl. Acad. Sci. U. S. A* 106, 20912–20917.
- Hutchins, A.P., Poulain, S., and Miranda-Saavedra, D. (2012). Genome-wide analysis of STAT3 binding in vivo predicts effectors of the anti-inflammatory response in macrophages. *Blood* 119, e110–e119.
- Huynh, K.D. and Bardwell, V.J. (1998). The BCL-6 POZ domain and other POZ domains interact with the co-repressors N-CoR and SMRT. *Oncogene* 17, 2473–2484.
- Igreja, C. and Izaurralde, E. (2011). CUP promotes deadenylation and inhibits decapping of mRNA targets. *Genes Dev.* 25, 1955–1967.
- Imai, E., Miner, J.N., Mitchell, J.A., Yamamoto, K.R., and Granner, D.K. (1993). Glucocorticoid receptor-cAMP response element-binding protein interaction and the response of the phosphoenolpyruvate carboxykinase gene to glucocorticoids. *J. Biol. Chem.* 268, 5353–5356.
- Jernigan, C.S., Goswami, D.B., Austin, M.C., Iyo, A.H., Chandran, A., Stockmeier, C.A., and Karolewicz, B. (2011). The mTOR signaling pathway in the prefrontal cortex is compromised in major depressive disorder. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 35, 1774–1779.
- Ji, H., Jiang, H., Ma, W., Johnson, D.S., Myers, R.M., and Wong, W.H. (2008). An integrated software system for analyzing ChIP-chip and ChIP-seq data. *Nat. Biotechnol.* 26, 1293–1300.
- Jilg, A., Lesny, S., Peruzki, N., Schwegler, H., Selbach, O., Dehghani, F., and Stehle, J.H. (2010). Temporal dynamics of mouse hippocampal clock gene expression support memory processing. *Hippocampus* 20, 377–388.
- Joels, M., Velzing, E., Nair, S., Verkuyl, J.M., and Karst, H. (2003). Acute stress increases calcium current amplitude in rat hippocampus: temporal changes in physiology and gene expression. *Eur. J. Neurosci.* 18, 1315–1324.
- John, S., Johnson, T.A., Sung, M.H., Biddie, S.C., Trump, S., Koch-Paiz, C.A., Davis, S.R., Walker, R., Meltzer, P.S., and Hager, G.L. (2009). Kinetic complexity of the global response to glucocorticoid receptor action. *Endocrinology* 150, 1766–1774.
- John, S., Sabo, P.J., Thurman, R.E., Sung, M.H., Biddie, S.C., Johnson, T.A., Hager, G.L., and Stamatoyannopoulos, J.A. (2011). Chromatin accessibility pre-determines glucocorticoid receptor binding patterns. *Nat. Genet.* 43, 264–268.
- Johnson, D.S., Mortazavi, A., Myers, R.M., and Wold, B. (2007). Genome-wide mapping of in vivo protein-DNA interactions. *Science* 316, 1497–1502.
- Johnson, L.R., Farb, C., Morrison, J.H., McEwen, B.S., and LeDoux, J.E. (2005). Localization of glucocorticoid receptors at postsynaptic membranes in the lateral amygdala. *Neuroscience* 136, 289–299.
- Jonat, C., Rahmsdorf, H.J., Park, K.K., Cato, A.C., Gebel, S., Ponta, H., and Herrlich, P. (1990). Antitumor promotion and antiinflammation: down-modulation of AP-1 (Fos/Jun) activity by glucocorticoid hormone. *Cell* 62, 1189–1204.
- Kapranov, P. (2009). From transcription start site to cell biology. *Genome Biol.* 10, 217.
- Karatsoreos, I.N. and McEwen, B.S. (2013). Resilience and vulnerability: a neurobiological perspective. *Front. Rep.* 5, 13.

- Karst, H. and Joels, M. (2003). Effect of chronic stress on synaptic currents in rat hippocampal dentate gyrus neurons. *J. Neurophysiol.* 89, 625–633.
- Karst, H., Karten, Y.J., Reichardt, H.M., de Kloet, E.R., Schutz, G., and Joels, M. (2000). Corticosteroid actions in hippocampus require DNA binding of glucocorticoid receptor homodimers. *Nat. Neurosci.* 3, 977–978.
- Kassel, O. and Herrlich, P. (2007). Crosstalk between the glucocorticoid receptor and other transcription factors: molecular aspects. *Mol. Cell Endocrinol.* 275, 13–29.
- Kel, A.E., Gossling, E., Reuter, I., Cheremushkin, E., Kel-Margoulis, O.V., and Wingender, E. (2003). MATCH: A tool for searching transcription factor binding sites in DNA sequences. *Nucleic Acids Res.* 31, 3576–3579.
- Kelly, E.J., Sandgren, E.P., Brinster, R.L., and Palmiter, R.D. (1997). A pair of adjacent glucocorticoid response elements regulate expression of two mouse metallothionein genes. *Proc. Natl. Acad. Sci. U. S. A.* 94, 10045–10050.
- Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., and Haussler, D. (2002). The human genome browser at UCSC. *Genome Res.* 12, 996–1006.
- Kim, J. and Hoffman, D.A. (2008). Potassium channels: newly found players in synaptic plasticity. *Neuroscientist.* 14, 276–286.
- Kim, J.J., Song, E.Y., and Kosten, T.A. (2006). Stress effects in the hippocampus: synaptic plasticity and memory. *Stress.* 9, 1–11.
- Kinyamu, H.K., Collins, J.B., Grissom, S.F., Hebbar, P.B., and Archer, T.K. (2008). Genome wide transcriptional profiling in breast cancer cells reveals distinct changes in hormone receptor target genes and chromatin modifying enzymes after proteasome inhibition. *Mol. Carcinog.* 47, 845–885.
- Komatsuzaki, Y., Hatanaka, Y., Murakami, G., Mukai, H., Hojo, Y., Saito, M., Kimoto, T., and Kawato, S. (2012). Corticosterone induces rapid spinogenesis via synaptic glucocorticoid receptors and kinase networks in hippocampus. *PLoS. One.* 7, e34124.
- Kooi van der, Onuma, H., Oeser, J.K., Svitek, C.A., Allen, S.R., Vander Kooi, C.W., Chazin, W.J., and O'Brien, R.M. (2005). The glucose-6-phosphatase catalytic subunit gene promoter contains both positive and negative glucocorticoid response elements. *Mol. Endocrinol.* 19, 3001–3022.
- Krugers, H.J., Goltstein, P.M., van der Linden, S., and Joels, M. (2006). Blockade of glucocorticoid receptors rapidly restores hippocampal CA₁ synaptic plasticity after exposure to chronic stress. *Eur. J. Neurosci.* 23, 3051–3055.
- Krum, S.A., Miranda-Carboni, G.A., Lupien, M., Eeckhoutte, J., Carroll, J.S., and Brown, M. (2008). Unique ERalpha cisromes control cell type-specific gene regulation. *Mol. Endocrinol.* 22, 2393–2406.
- Kumar, S. (2011). Remote homologue identification of Drosophila GAGA factor in mouse. *Bioinformatics.* 7, 29–32.
- Kunarsow, G., Chia, N.Y., Jeyakani, J., Hwang, C., Lu, X., Chan, Y.S., Ng, H.H., and Bourque, G. (2010). Transposable elements have rewired the core regulatory network of human embryonic stem cells. *Nat. Genet.* 42, 631–634.
- Landt, S.G., Marinov, G.K., Kundaje, A., Kheradpour, P., Pauli, F., Batzoglou, S., Bernstein, B.E., Bickel, P., Brown, J.B., Cayting, P., Chen, Y., DeSalvo, G., Epstein, C., Fisher-Aylor, K.I., Euskirchen, G., Gerstein, M., Gertz, J., Hartemink, A.J., Hoffman, M.M., Iyer, V.R., Jung, Y.L., Karmakar, S., Kellis, M., Kharchenko, P.V., Li, Q., Liu, T., Liu, X.S., Ma, L., Milosavljevic, A., Myers, R.M., Park, P.J., Pazin, M.J., Perry, M.D., Raha, D., Reddy, T.E., Rozowsky, J., Shores, N.,

- Sidow,A., Slattery,M., Stamatoyannopoulos,J.A., Tolstorukov,M.Y., White,K.P., Xi,S., Farnham,P.J., Lieb,J.D., Wold,B.J., and Snyder,M. (2012). ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. *Genome Res.* 22, 1813–1831.
- Langlais,D., Couture,C., Balsalobre,A., and Drouin,J. (2012). The Stat3/GR interaction code: predictive value of direct/indirect DNA recruitment for transcription outcome. *Mol. Cell* 47, 38–49.
- Leal,G., Comprido,D., and Duarte,C.B. (2014). BDNF-induced local protein synthesis and synaptic plasticity. *Neuropharmacology* 76 Pt C, 639–656.
- Lee,P.R., Brady,D., and Koenig,J.I. (2003). Corticosterone alters N-methyl-D-aspartate receptor subunit mRNA expression before puberty. *Brain Res. Mol. Brain Res.* 115, 55–62.
- Lee,R.S., Tamashiro,K.L., Yang,X., Purcell,R.H., Harvey,A., Willour,V.L., Huo,Y., Rongione,M., Wand,G.S., and Potash,J.B. (2010). Chronic corticosterone exposure increases expression and decreases deoxyribonucleic acid methylation of Fkbp5 in mice. *Endocrinology* 151, 4332–4343.
- Lein,E.S., Hawrylycz,M.J., Ao,N., Ayres,M., Bensinger,A., Bernard,A., Boe,A.F., Boguski,M.S., Brockway,K.S., Byrnes,E.J., Chen,L., Chen,L., Chen,T.M., Chin,M.C., Chong,J., Crook,B.E., Czaplinska,A., Dang,C.N., Datta,S., Dee,N.R., Desaki,A.L., Desta,T., Diep,E., Dolbeare,T.A., Donelan,M.J., Dong,H.W., Dougherty,J.G., Duncan,B.J., Ebbert,A.J., Eichele,G., Estin,L.K., Faber,C., Facer,B.A., Fields,R., Fischer,S.R., Fliss,T.P., Frensley,C., Gates,S.N., Glattfelder,K.J., Halverson,K.R., Hart,M.R., Hohmann,J.G., Howell,M.P., Jeung,D.P., Johnson,R.A., Karr,P.T., Kawal,R., Kidney,J.M., Knapik,R.H., Kuan,C.L., Lake,J.H., Laramée,A.R., Larsen,K.D., Lau,C., Lemon,T.A., Liang,A.J., Liu,Y., Luong,L.T., Michaels,J., Morgan,J.J., Morgan,R.J., Mortrud,M.T., Mosqueda,N.F., Ng,L.L., Ng,R., Orta,G.J., Overly,C.C., Pak,T.H., Parry,S.E., Pathak,S.D., Pearson,O.C., Puchalski,R.B., Riley,Z.L., Rockett,H.R., Rowland,S.A., Royall,J.J., Ruiz,M.J., Sarno,N.R., Schaffnit,K., Shapovalova,N.V., Sivasay,T., Slaughterbeck,C.R., Smith,S.C., Smith,K.A., Smith,B.I., Sodt,A.J., Stewart,N.N., Stumpf,K.R., Sunkin,S.M., Sutram,M., Tam,A., Teemer,C.D., Thaller,C., Thompson,C.L., Varnam,L.R., Visel,A., Whitlock,R.M., Wohnoutka,P.E., Wolkey,C.K., Wong,V.Y., Wood,M., Yayaoglu,M.B., Young,R.C., Youngstrom,B.L., Yuan,X.F., Zhang,B., Zwingman,T.A., and Jones,A.R. (2007). Genome-wide atlas of gene expression in the adult mouse brain. *Nature* 445, 168–176.
- Lein,E.S., Zhao,X., and Gage,F.H. (2004). Defining a molecular atlas of the hippocampus using DNA microarrays and high-throughput in situ hybridization. *J. Neurosci.* 24, 3879–3889.
- Lemetre,C. and Zhang,Z.D. (2013). A brief introduction to tiling microarrays: principles, concepts, and applications. *Methods Mol. Biol.* 1067, 3–19.
- Li,H. and Durbin,R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics.* 25, 1754–1760.
- Li,N., Lee,B., Liu,R.J., Banasr,M., Dwyer,J.M., Iwata,M., Li,X.Y., Aghajanian,G., and Duman,R.S. (2010). mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science* 329, 959–964.
- Lim,J.H. (2014). Zinc finger and BTB domain-containing protein 3 is essential for the growth of cancer cells. *BMB. Rep.* 47, 405–410.
- Lipton,J.O. and Sahin,M. (2014). The neurology of mTOR. *Neuron* 84, 275–291.

- Lisowski,P., Juszczak,G.R., Goscik,J., Wieczorek,M., Zwierzchowski,L., and Swiergiel,A.H. (2011). Effect of chronic mild stress on hippocampal transcriptome in mice selected for high and low stress-induced analgesia and displaying different emotional behaviors. *Eur. Neuropsychopharmacol.* 21, 45–62.
- Liu,Q., Qiu,J., Liang,M., Golinski,J., van,L.K., Jung,J.E., You,Z., Lo,E.H., Degterev,A., and Whalen,M.J. (2014). Akt and mTOR mediate programmed necrosis in neurons. *Cell Death. Dis.* 5, e1084.
- Liu,Y., Wei,L., Batzoglou,S., Brutlag,D.L., Liu,J.S., and Liu,X.S. (2004). A suite of web-based programs to search for transcriptional regulatory motifs. *Nucleic Acids Res.* 32, W204–W207.
- Livak,K.J. and Schmittgen,T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C(T)}$ Method. *Methods* 25, 402–408.
- Lu,N.Z. and Cidlowski,J.A. (2005). Translational regulatory mechanisms generate N-terminal glucocorticoid receptor isoforms with unique transcriptional target genes. *Mol. Cell* 18, 331–342.
- Lu,Y., Wang,C., Xue,Z., Li,C., Zhang,J., Zhao,X., Liu,A., Wang,Q., and Zhou,W. (2015). PI3K/AKT/mTOR signaling-mediated neuropeptide VGF in the hippocampus of mice is involved in the rapid onset antidepressant-like effects of GLYX-13. *Int. J. Neuropsychopharmacol.* 18.
- Ma,Y., Wu,X., Li,X., Fu,J., Shen,J., Li,X., and Wang,H. (2012). Corticosterone regulates the expression of neuropeptide Y and reelin in MLO-Y4 cells. *Mol. Cells* 33, 611–616.
- Magarinos,A.M., McEwen,B.S., Flugge,G., and Fuchs,E. (1996). Chronic psychosocial stress causes apical dendritic atrophy of hippocampal CA3 pyramidal neurons in subordinate tree shrews. *J. Neurosci.* 16, 3534–3540.
- Malagelada,C., Jin,Z.H., and Greene,L.A. (2008). RTP801 is induced in Parkinson's disease and mediates neuron death by inhibiting Akt phosphorylation/activation. *J. Neurosci.* 28, 14363–14371.
- Mardis,E.R. (2007). CHIP-seq: welcome to the new frontier. *Nat. Methods* 4, 613–614.
- Masuno,K., Haldar,S.M., Jeyaraj,D., Mailloux,C.M., Huang,X., Panettieri,R.A., Jr., Jain,M.K., and Gerber,A.N. (2011). Expression profiling identifies *Klf15* as a glucocorticoid target that regulates airway hyperresponsiveness. *Am. J. Respir. Cell Mol. Biol.* 45, 642–649.
- McEwen, B. S. Allostasis and Allostatis Load. *Encyclopedia of Stress.* 1, 145–150. 2000. Academic Press. Ref Type: Generic
- McLaughlin,K.J., Gomez,J.L., Baran,S.E., and Conrad,C.D. (2007). The effects of chronic stress on hippocampal morphology and function: an evaluation of chronic restraint paradigms. *Brain Res.* 1161, 56–64.
- Meijsing,S.H., Pufall,M.A., So,A.Y., Bates,D.L., Chen,L., and Yamamoto,K.R. (2009). DNA binding site sequence directs glucocorticoid receptor structure and activity. *Science* 324, 407–410.
- Meyer,M.B., Goetsch,P.D., and Pike,J.W. (2010). Genome-wide analysis of the VDR/RXR cistrome in osteoblast cells provides new mechanistic insight into the actions of the vitamin D hormone. *J. Steroid Biochem. Mol. Biol.* 121, 136–141.
- Mikkelsen,T.S., Ku,M., Jaffe,D.B., Issac,B., Lieberman,E., Giannoukos,G., Alvarez,P., Brockman,W., Kim,T.K., Koche,R.P., Lee,W., Mendenhall,E., O'Donovan,A., Presser,A., Russ,C.,

- Xie,X., Meissner,A., Wernig,M., Jaenisch,R., Nusbaum,C., Lander,E.S., and Bernstein,B.E. (2007). Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature* 448, 553–560.
- Miller,J.A., Nathanson,J., Franjic,D., Shim,S., Dalley,R.A., Shapouri,S., Smith,K.A., Sunkin,S.M., Bernard,A., Bennett,J.L., Lee,C.K., Hawrylycz,M.J., Jones,A.R., Amaral,D.G., Sestan,N., Gage,F.H., and Lein,E.S. (2013). Conserved molecular signatures of neurogenesis in the hippocampal subgranular zone of rodents and primates. *Development* 140, 4633–4644.
- Miller,J.C., Jimenez,P., and Mathe,A.A. (2007). Restraint stress influences AP-1 and CREB DNA-binding activity induced by chronic lithium treatment in the rat frontal cortex and hippocampus. *Int. J. Neuropsychopharmacol.* 10, 609–619.
- Moghbelinejad,S., Nassiri-Asl,M., Farivar,T.N., Abbasi,E., Sheikhi,M., Taghilo,M., Farsad,F., Samimi,A., and Hajiali,F. (2014). Rutin activates the MAPK pathway and BDNF gene expression on beta-amyloid induced neurotoxicity in rats. *Toxicol. Lett.* 224, 108–113.
- Moisan,M.P., Minni,A.M., Dominguez,G., Helbling,J.C., Foury,A., Henkous,N., Dorey,R., and Beracochea,D. (2014). Role of corticosteroid binding globulin in the fast actions of glucocorticoids on the brain. *Steroids* 81, 109–115.
- Morimoto,M., Morita,N., Ozawa,H., Yokoyama,K., and Kawata,M. (1996). Distribution of glucocorticoid receptor immunoreactivity and mRNA in the rat brain: an immunohistochemical and in situ hybridization study. *Neurosci. Res.* 26, 235–269.
- Morsink,M.C., Joels,M., Sarabdjitsingh,R.A., Meijer,O.C., de Kloet,E.R., and Datson,N.A. (2006a). The dynamic pattern of glucocorticoid receptor-mediated transcriptional responses in neuronal PC12 cells. *J. Neurochem.* 99, 1282–1298.
- Morsink,M.C., Steenbergen,P.J., Vos,J.B., Karst,H., Joels,M., de Kloet,E.R., and Datson,N.A. (2006b). Acute activation of hippocampal glucocorticoid receptors results in different waves of gene expression throughout time. *J. Neuroendocrinol.* 18, 239–252.
- Mozaffari,M., Hoogveen-Westerveld,M., Kwiatkowski,D., Sampson,J., Ekong,R., Povey,S., den Dunnen,J.T., van den Ouweland,A., Halley,D., and Nellist,M. (2009). Identification of a region required for TSC1 stability by functional analysis of TSC1 missense mutations found in individuals with tuberous sclerosis complex. *BMC. Med. Genet.* 10, 88.
- Mundade,R., Ozer,H.G., Wei,H., Prabhu,L., and Lu,T. (2014). Role of ChIP-seq in the discovery of transcription factor binding sites, differential gene regulation mechanism, epigenetic marks and beyond. *Cell Cycle* 13, 2847–2852.
- Nelson,C.C., Hendy,S.C., Shukin,R.J., Cheng,H., Bruchovsky,N., Koop,B.F., and Rennie,P.S. (1999). Determinants of DNA sequence specificity of the androgen, progesterone, and glucocorticoid receptors: evidence for differential steroid receptor response elements. *Mol. Endocrinol.* 13, 2090–2107.
- Nelson,J.D., Denisenko,O., and Bomsztyk,K. (2006). Protocol for the fast chromatin immunoprecipitation (ChIP) method. *Nat. Protoc.* 1, 179–185.
- Nishi,M., Tanaka,M., Matsuda,K., Sunaguchi,M., and Kawata,M. (2004). Visualization of glucocorticoid receptor and mineralocorticoid receptor interactions in living cells with GFP-based fluorescence resonance energy transfer. *J. Neurosci.* 24, 4918–4927.
- Oakley,R.H. and Cidlowski,J.A. (2013). The biology of the glucocorticoid receptor: new signaling mechanisms in health and disease. *J. Allergy Clin. Immunol.* 132, 1033–1044.

- Oakley, R.H., Sar, M., and Cidlowski, J.A. (1996). The human glucocorticoid receptor beta isoform. Expression, biochemical properties, and putative function. *J. Biol. Chem.* *271*, 9550–9559.
- Oitzl, M.S., Reichardt, H.M., Joels, M., and de Kloet, E.R. (2001). Point mutation in the mouse glucocorticoid receptor preventing DNA binding impairs spatial memory. *Proc. Natl. Acad. Sci. U. S. A* *98*, 12790–12795.
- Okamoto, S., Sherman, K., Bai, G., and Lipton, S.A. (2002). Effect of the ubiquitous transcription factors, SP1 and MAZ, on NMDA receptor subunit type 1 (NR1) expression during neuronal differentiation. *Brain Res. Mol. Brain Res.* *107*, 89–96.
- Ongwijitwat, S., Liang, H.L., Graboyes, E.M., and Wong-Riley, M.T. (2006). Nuclear respiratory factor 2 senses changing cellular energy demands and its silencing down-regulates cytochrome oxidase and other target gene mRNAs. *Gene* *374*, 39–49.
- Ongwijitwat, S. and Wong-Riley, M.T. (2005). Is nuclear respiratory factor 2 a master transcriptional coordinator for all ten nuclear-encoded cytochrome c oxidase subunits in neurons? *Gene* *360*, 65–77.
- Orlovsky, M.A., Dosenko, V.E., Spiga, F., Skibo, G.G., and Lightman, S.L. (2014). Hippocampus remodeling by chronic stress accompanied by GR, proteasome and caspase-3 overexpression. *Brain Res.* *1593*, 83–94.
- Orsetti, M., Di, B.F., Canonico, P.L., Genazzani, A.A., and Ghi, P. (2008). Gene regulation in the frontal cortex of rats exposed to the chronic mild stress paradigm, an animal model of human depression. *Eur. J. Neurosci.* *27*, 2156–2164.
- Ott, C.J., Blackledge, N.P., Leir, S.H., and Harris, A. (2009). Novel regulatory mechanisms for the CFTR gene. *Biochem. Soc. Trans.* *37*, 843–848.
- Ovcharenko, I., Loots, G.G., Nobrega, M.A., Hardison, R.C., Miller, W., and Stubbs, L. (2005). Evolution and functional classification of vertebrate gene deserts. *Genome Res.* *15*, 137–145.
- Owen, D. and Matthews, S.G. (2003). Glucocorticoids and sex-dependent development of brain glucocorticoid and mineralocorticoid receptors. *Endocrinology* *144*, 2775–2784.
- Oyadomari, S. and Mori, M. (2004). Roles of CHOP/GADD153 in endoplasmic reticulum stress. *Cell Death. Differ.* *11*, 381–389.
- Paakinaho, V., Kaikkonen, S., Makkonen, H., Benes, V., and Palvimo, J.J. (2014). SUMOylation regulates the chromatin occupancy and anti-proliferative gene programs of glucocorticoid receptor. *Nucleic Acids Res.* *42*, 1575–1592.
- Paakinaho, V., Makkonen, H., Jaaskelainen, T., and Palvimo, J.J. (2010). Glucocorticoid receptor activates poised FKBP51 locus through long-distance interactions. *Mol. Endocrinol.* *24*, 511–525.
- Pan, D., Kocherginsky, M., and Conzen, S.D. (2011). Activation of the glucocorticoid receptor is associated with poor prognosis in estrogen receptor-negative breast cancer. *Cancer Res.* *71*, 6360–6370.
- Pardridge, W.M. and Mietus, L.J. (1979). Transport of steroid hormones through the rat blood-brain barrier. Primary role of albumin-bound hormone. *J. Clin. Invest* *64*, 145–154.
- Pascale, A., Amadio, M., Caffino, L., Racagni, G., Govoni, S., and Fumagalli, F. (2011). ELAV-GAP43 pathway activation following combined exposure to cocaine and stress. *Psychopharmacology (Berl)* *218*, 249–256.

- Payvar,F., DeFranco,D., Firestone,G.L., Edgar,B., Wrangle,O., Okret,S., Gustafsson,J.A., and Yamamoto,K.R. (1983). Sequence-specific binding of glucocorticoid receptor to MTV DNA at sites within and upstream of the transcribed region. *Cell* 35, 381–392.
- Pearce,D., Matsui,W., Miner,J.N., and Yamamoto,K.R. (1998). Glucocorticoid receptor transcriptional activity determined by spacing of receptor and nonreceptor DNA sites. *J. Biol. Chem.* 273, 30081–30085.
- Pei,H., Li,L., Fridley,B.L., Jenkins,G.D., Kalari,K.R., Lingle,W., Petersen,G., Lou,Z., and Wang,L. (2009). FKBP51 affects cancer cell response to chemotherapy by negatively regulating Akt. *Cancer Cell* 16, 259–266.
- Pei,H., Lou,Z., and Wang,L. (2010). Emerging role of FKBP51 in AKT kinase/protein kinase B signaling. *Cell Cycle* 9, 6–7.
- Pei,J.J. and Hugon,J. (2008). mTOR-dependent signalling in Alzheimer’s disease. *J. Cell Mol. Med.* 12, 2525–2532.
- Phuc Le P., Friedman,J.R., Schug,J., Brestelli,J.E., Parker,J.B., Bochkis,I.M., and Kaestner,K.H. (2005). Glucocorticoid receptor-dependent gene regulatory networks. *PLoS. Genet.* 1, e16.
- Pilar-Cuellar,F., Vidal,R., Diaz,A., Castro,E., dos,A.S., Pascual-Brazo,J., Linge,R., Vargas,V., Blanco,H., Martinez-Villayandre,B., Pazos,A., and Valdizan,E.M. (2013). Neural plasticity and proliferation in the generation of antidepressant effects: hippocampal implication. *Neural Plast.* 2013, 537265.
- Pillai,S. and Chellappan,S.P. (2009). ChIP on chip assays: genome-wide analysis of transcription factor binding and histone modifications. *Methods Mol. Biol.* 523, 341–366.
- Polman,J.A., Hunter,R.G., Speksnijder,N., van den Oever,J.M., Korobko,O.B., McEwen,B.S., de Kloet,E.R., and Datson,N.A. (2012a). Glucocorticoids modulate the mTOR pathway in the hippocampus: differential effects depending on stress history. *Endocrinology* 153, 4317–4327.
- Polman,J.A., Welten,J.E., Bosch,D.S., de Jonge,R.T., Balog,J., van der Maarel,S.M., de Kloet,E.R., and Datson,N.A. (2012b). A genome-wide signature of glucocorticoid receptor binding in neuronal PC12 cells. *BMC. Neurosci.* 13, 118.
- Pratt,W.B. (1990). Glucocorticoid receptor structure and the initial events in signal transduction. *Prog. Clin. Biol. Res.* 322, 119–132.
- Pratt,W.B. and Toft,D.O. (1997). Steroid receptor interactions with heat shock protein and immunophilin chaperones. *Endocr. Rev.* 18, 306–360.
- Qi,S., Mizuno,M., Yonezawa,K., Nawa,H., and Takei,N. (2010). Activation of mammalian target of rapamycin signaling in spatial learning. *Neurosci. Res.* 68, 88–93.
- Quinlan,A.R. and Hall,I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics.* 26, 841–842.
- Rani,C.S., Elango,N., Wang,S.S., Kobayashi,K., and Strong,R. (2009). Identification of an activator protein-1-like sequence as the glucocorticoid response element in the rat tyrosine hydroxylase gene. *Mol. Pharmacol.* 75, 589–598.
- Rao,N.A., McCalman,M.T., Moulos,P., Francoijs,K.J., Chatziioannou,A., Kolisis,F.N., Alexis,M.N., Mitsiou,D.J., and Stunnenberg,H.G. (2011). Coactivation of GR and NFκB alters the repertoire of their binding sites and target genes. *Genome Res.* 21, 1404–1416.
- Ravasi,T., Suzuki,H., Cannistraci,C.V., Katayama,S., Bajic,V.B., Tan,K., Akalin,A., Schmeier,S., Kanamori-Katayama,M., Bertin,N., Carninci,P., Daub,C.O., Forrest,A.R.,

- Gough,J., Grimmond,S., Han,J.H., Hashimoto,T., Hide,W., Hofmann,O., Kamburov,A., Kaur,M., Kawaji,H., Kubosaki,A., Lassmann,T., van,N.E., MacPherson,C.R., Ogawa,C., Radovanovic,A., Schwartz,A., Teasdale,R.D., Tegner,J., Lenhard,B., Teichmann,S.A., Arakawa,T., Ninomiya,N., Murakami,K., Tagami,M., Fukuda,S., Imamura,K., Kai,C., Ishihara,R., Kitazume,Y., Kawai,J., Hume,D.A., Ideker,T., and Hayashizaki,Y. (2010). An atlas of combinatorial transcriptional regulation in mouse and man. *Cell* 140, 744–752.
- Reddy,T.E., Gertz,J., Crawford,G.E., Garabedian,M.J., and Myers,R.M. (2012). The hypersensitive glucocorticoid response specifically regulates period 1 and expression of circadian genes. *Mol. Cell Biol.* 32, 3756–3767.
- Reddy,T.E., Pauli,F., Sprouse,R.O., Neff,N.F., Newberry,K.M., Garabedian,M.J., and Myers,R.M. (2009). Genomic determination of the glucocorticoid response reveals unexpected mechanisms of gene regulation. *Genome Res.* 19, 2163–2171.
- Reichardt,H.M., Kaestner,K.H., Tuckermann,J., Kretz,O., Wessely,O., Bock,R., Gass,P., Schmid,W., Herrlich,P., Angel,P., and Schutz,G. (1998). DNA binding of the glucocorticoid receptor is not essential for survival. *Cell* 93, 531–541.
- Reul,J.M. and de Kloet,E.R. (1985). Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation. *Endocrinology* 117, 2505–2511.
- Robertson,G., Hirst,M., Bainbridge,M., Bilenky,M., Zhao,Y., Zeng,T., Euskirchen,G., Bernier,B., Varhol,R., Delaney,A., Thiessen,N., Griffith,O.L., He,A., Marra,M., Snyder,M., and Jones,S. (2007). Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. *Nat. Methods* 4, 651–657.
- Ru,W., Peng,Y., Zhong,L., and Tang,S.J. (2012). A role of the mammalian target of rapamycin (mTOR) in glutamate-induced down-regulation of tuberous sclerosis complex proteins 2 (TSC2). *J. Mol. Neurosci.* 47, 340–345.
- Ruiz-Llorente,S., Carrillo Santa de,P.E., Sastre-Perona,A., Montero-Conde,C., Gomez-Lopez,G., Fagin,J.A., Valencia,A., Pisano,D.G., and Santisteban,P. (2012). Genome-wide analysis of Pax8 binding provides new insights into thyroid functions. *BMC. Genomics* 13, 147.
- Sakai,D.D., Helms,S., Carlstedt-Duke,J., Gustafsson,J.A., Rottman,F.M., and Yamamoto,K.R. (1988). Hormone-mediated repression: a negative glucocorticoid response element from the bovine prolactin gene. *Genes Dev.* 2, 1144–1154.
- Sapolsky,R.M., Romero,L.M., and Munck,A.U. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr. Rev.* 21, 55–89.
- Sarabdjitsingh,R.A., Isenia,S., Polman,A., Mijalkovic,J., Lachize,S., Datson,N., de Kloet,E.R., and Meijer,O.C. (2010a). Disrupted corticosterone pulsatile patterns attenuate responsiveness to glucocorticoid signaling in rat brain. *Endocrinology* 151, 1177–1186.
- Sarabdjitsingh,R.A., Meijer,O.C., and de Kloet,E.R. (2010b). Specificity of glucocorticoid receptor primary antibodies for analysis of receptor localization patterns in cultured cells and rat hippocampus. *Brain Res.* 1331, 1–11.
- Schaaf,M.J., de,J.J., de Kloet,E.R., and Vreugdenhil,E. (1998). Downregulation of BDNF mRNA and protein in the rat hippocampus by corticosterone. *Brain Res.* 813, 112–120.
- Schaaf,M.J., Lewis-Tuffin,L.J., and Cidlowski,J.A. (2005). Ligand-selective targeting of the glucocorticoid receptor to nuclear subdomains is associated with decreased receptor mobility. *Mol. Endocrinol.* 19, 1501–1515.

- Scheinman,R.I., Gualberto,A., Jewell,C.M., Cidlowski,J.A., and Baldwin,A.S., Jr. (1995). Characterization of mechanisms involved in transrepression of NF-kappa B by activated glucocorticoid receptors. *Mol. Cell Biol.* 15, 943-953.
- Schiene-Fischer,C. and Yu,C. (2001). Receptor accessory folding helper enzymes: the functional role of peptidyl prolyl cis/trans isomerases. *FEBS Lett.* 495, 1-6.
- Schmidt,M.V., Trumbach,D., Weber,P., Wagner,K., Scharf,S.H., Liebl,C., Datson,N., Namendorf,C., Gerlach,T., Kuhne,C., Uhr,M., Deussing,J.M., Wurst,W., Binder,E.B., Holsboer,F., and Muller,M.B. (2010). Individual stress vulnerability is predicted by short-term memory and AMPA receptor subunit ratio in the hippocampus. *J. Neurosci.* 30, 16949-16958.
- Schoneveld,O.J., Gaemers,I.C., and Lamers,W.H. (2004). Mechanisms of glucocorticoid signalling. *Biochim. Biophys. Acta* 1680, 114-128.
- Schule,R., Rangarajan,P., Kliewer,S., Ransone,L.J., Bolado,J., Yang,N., Verma,I.M., and Evans,R.M. (1990). Functional antagonism between oncoprotein c-Jun and the glucocorticoid receptor. *Cell* 62, 1217-1226.
- Shah,O.J., Anthony,J.C., Kimball,S.R., and Jefferson,L.S. (2000a). 4E-BP1 and S6K1: translational integration sites for nutritional and hormonal information in muscle. *Am. J. Physiol Endocrinol. Metab* 279, E715-E729.
- Shah,O.J., Anthony,J.C., Kimball,S.R., and Jefferson,L.S. (2000b). Glucocorticoids oppose translational control by leucine in skeletal muscle. *Am. J. Physiol Endocrinol. Metab* 279, E1185-E1190.
- Shah,O.J., Kimball,S.R., and Jefferson,L.S. (2000c). Acute attenuation of translation initiation and protein synthesis by glucocorticoids in skeletal muscle. *Am. J. Physiol Endocrinol. Metab* 278, E76-E82.
- Shimizu,H., Arima,H., Ozawa,Y., Watanabe,M., Banno,R., Sugimura,Y., Ozaki,N., Nagasaki,H., and Oiso,Y. (2010). Glucocorticoids increase NPY gene expression in the arcuate nucleus by inhibiting mTOR signaling in rat hypothalamic organotypic cultures. *Peptides* 31, 145-149.
- Shors,T.J., Chua,C., and Falduto,J. (2001). Sex differences and opposite effects of stress on dendritic spine density in the male versus female hippocampus. *J. Neurosci.* 21, 6292-6297.
- Sierra,A., Gottfried-Blackmore,A., Milner,T.A., McEwen,B.S., and Bulloch,K. (2008). Steroid hormone receptor expression and function in microglia. *Glia* 56, 659-674.
- Silverman,M.N. and Sternberg,E.M. (2012). Glucocorticoid regulation of inflammation and its functional correlates: from HPA axis to glucocorticoid receptor dysfunction. *Ann. N. Y. Acad. Sci.* 1261, 55-63.
- Simoes,D.C., Psarra,A.M., Mauad,T., Pantou,I., Roussos,C., Sekeris,C.E., and Gratiou,C. (2012). Glucocorticoid and estrogen receptors are reduced in mitochondria of lung epithelial cells in asthma. *PLoS. One.* 7, e39183.
- Simon,R., Lam,A., Li,M.C., Ngan,M., Menenzes,S., and Zhao,Y. (2007). Analysis of gene expression data using BRB-ArrayTools. *Cancer Inform.* 3, 11-17.
- Slipczuk,L., Bekinshtein,P., Katche,C., Cammarota,M., Izquierdo,I., and Medina,J.H. (2009). BDNF activates mTOR to regulate GluR1 expression required for memory formation. *PLoS. One.* 4, e6007.
- Smyth,G.K.Limma: linear models for microarray data. In: Gentleman,R., Carey,V., Dudoit,S., Irizarry,R., Huber,W., eds. *Bioinformatics and computational biology solutions using R and bioconductor.* 397-420. 2005. New York, Springer. Ref Type: Generic

- So,A.Y., Chaivorapol,C., Bolton,E.C., Li,H., and Yamamoto,K.R. (2007). Determinants of cell- and gene-specific transcriptional regulation by the glucocorticoid receptor. *PLoS. Genet.* 3, e94.
- So,A.Y., Cooper,S.B., Feldman,B.J., Manuchehri,M., and Yamamoto,K.R. (2008). Conservation analysis predicts in vivo occupancy of glucocorticoid receptor-binding sequences at glucocorticoid-induced genes. *Proc. Natl. Acad. Sci. U. S. A* 105, 5745-5749.
- Solomon,S.S., Majumdar,G., Martinez-Hernandez,A., and Raghow,R. (2008). A critical role of Sp1 transcription factor in regulating gene expression in response to insulin and other hormones. *Life Sci.* 83, 305-312.
- Song,C.Z., Tian,X., and Gelehrter,T.D. (1999). Glucocorticoid receptor inhibits transforming growth factor-beta signaling by directly targeting the transcriptional activation function of Smad3. *Proc. Natl. Acad. Sci. U. S. A* 96, 11776-11781.
- Song,J., Ugai,H., Nakata-Tsutsui,H., Kishikawa,S., Suzuki,E., Murata,T., and Yokoyama,K.K. (2003). Transcriptional regulation by zinc-finger proteins Sp1 and MAZ involves interactions with the same cis-elements. *Int. J. Mol. Med.* 11, 547-553.
- Song,J., Ugai,H., Ogawa,K., Wang,Y., Sarai,A., Obata,Y., Kanazawa,I., Sun,K., Itakura,K., and Yokoyama,K.K. (2001). Two consecutive zinc fingers in Sp1 and in MAZ are essential for interactions with cis-elements. *J. Biol. Chem.* 276, 30429-30434.
- Sotiropoulos,I., Catania,C., Riedemann,T., Fry,J.P., Breen,K.C., Michaelidis,T.M., and Almeida,O.F. (2008). Glucocorticoids trigger Alzheimer disease-like pathobiochemistry in rat neuronal cells expressing human tau. *J. Neurochem.* 107, 385-397.
- Sousa,N., Lukoyanov,N.V., Madeira,M.D., Almeida,O.F., and Paula-Barbosa,M.M. (2000). Reorganization of the morphology of hippocampal neurites and synapses after stress-induced damage correlates with behavioral improvement. *Neuroscience* 97, 253-266.
- Spiga,F. and Lightman,S.L. (2009). Dose-dependent effects of corticosterone on nuclear glucocorticoid receptors and their binding to DNA in the brain and pituitary of the rat. *Brain Res.* 1293, 101-107.
- Spilman,P., Podlutskaya,N., Hart,M.J., Debnath,J., Gorostiza,O., Bredesen,D., Richardson,A., Strong,R., and Galvan,V. (2010). Inhibition of mTOR by rapamycin abolishes cognitive deficits and reduces amyloid-beta levels in a mouse model of Alzheimer's disease. *PLoS. One.* 5, e9979.
- Stavreva,D.A., Wiench,M., John,S., Conway-Campbell,B.L., McKenna,M.A., Pooley,J.R., Johnson,T.A., Voss,T.C., Lightman,S.L., and Hager,G.L. (2009). Ultradian hormone stimulation induces glucocorticoid receptor-mediated pulses of gene transcription. *Nat. Cell Biol.* 11, 1093-1102.
- Stocklin,E., Wissler,M., Gouilleux,F., and Groner,B. (1996). Functional interactions between Stat5 and the glucocorticoid receptor. *Nature* 383, 726-728.
- Stormo,G.D. (2000). DNA binding sites: representation and discovery. *Bioinformatics.* 16, 16-23.
- Strahle,U., Klock,G., and Schutz,G. (1987). A DNA sequence of 15 base pairs is sufficient to mediate both glucocorticoid and progesterone induction of gene expression. *Proc. Natl. Acad. Sci. U. S. A* 84, 7871-7875.
- Surjit,M., Ganti,K.P., Mukherji,A., Ye,T., Hua,G., Metzger,D., Li,M., and Chambon,P. (2011). Widespread negative response elements mediate direct repression by agonist-liganded glucocorticoid receptor. *Cell* 145, 224-241.

- Susa,T., Ishikawa,A., Kato,T., Nakayama,M., and Kato,Y. (2009). Molecular cloning of paired related homeobox 2 (prx2) as a novel pituitary transcription factor. *J. Reprod. Dev.* 55, 502–511.
- Swiech,L., Perycz,M., Malik,A., and Jaworski,J. (2008). Role of mTOR in physiology and pathology of the nervous system. *Biochim. Biophys. Acta* 1784, 116–132.
- Tang,S.J., Reis,G., Kang,H., Gingras,A.C., Sonenberg,N., and Schuman,E.M. (2002). A rapamycin-sensitive signaling pathway contributes to long-term synaptic plasticity in the hippocampus. *Proc. Natl. Acad. Sci. U. S. A* 99, 467–472.
- Taupenot,L. (2007). Analysis of regulated secretion using PC12 cells. *Curr. Protoc. Cell Biol. Chapter 15*, Unit.
- Teyssier,J.R., Ragot,S., Chauvet-Gelinier,J.C., Trojak,B., and Bonin,B. (2011). Activation of a DeltaFOSB dependent gene expression pattern in the dorsolateral prefrontal cortex of patients with major depressive disorder. *J. Affect. Disord.* 133, 174–178.
- Thomas-Chollier,M., Watson,L.C., Cooper,S.B., Pufall,M.A., Liu,J.S., Borzym,K., Vingron,M., Yamamoto,K.R., and Meijings,S.H. (2013). A naturally occurring insertion of a single amino acid rewires transcriptional regulation by glucocorticoid receptor isoforms. *Proc. Natl. Acad. Sci. U. S. A* 110, 17826–17831.
- Trapp,T., Rupprecht,R., Castren,M., Reul,J.M., and Holsboer,F. (1994). Heterodimerization between mineralocorticoid and glucocorticoid receptor: a new principle of glucocorticoid action in the CNS. *Neuron* 13, 1457–1462.
- Tsolakidou,A., Trumbach,D., Panhuysen,M., Putz,B., Deussing,J., Wurst,W., Sillaber,I., Holsboer,F., and Rein,T. (2008). Acute stress regulation of neuroplasticity genes in mouse hippocampus CA3 area—possible novel signalling pathways. *Mol. Cell Neurosci.* 38, 444–452.
- Ueda,K., Fujiki,K., Shirahige,K., Gomez-Sanchez,C.E., Fujita,T., Nangaku,M., and Nagase,M. (2014). Genome-wide analysis of murine renal distal convoluted tubular cells for the target genes of mineralocorticoid receptor. *Biochem. Biophys. Res. Commun.* 445, 132–137.
- Unlap,T. and Jope,R.S. (1995). Inhibition of NFkB DNA binding activity by glucocorticoids in rat brain. *Neurosci. Lett.* 198, 41–44.
- van der Laan,S., Sarabdjitsingh,R.A., Van Batenburg,M.F., Lachize,S.B., Li,H., Dijkmans,T.F., Vreugdenhil,E., de Kloet,E.R., and Meijer,O.C. (2008). Chromatin immunoprecipitation scanning identifies glucocorticoid receptor binding regions in the proximal promoter of a ubiquitously expressed glucocorticoid target gene in brain. *J. Neurochem.* 106, 2515–2523.
- Van Eekelen,J.A., Kiss,J.Z., Westphal,H.M., and de Kloet,E.R. (1987). Immunocytochemical study on the intracellular localization of the type 2 glucocorticoid receptor in the rat brain. *Brain Res.* 436, 120–128.
- van Steensel B., van Binnendijk,E.P., Hornsby,C.D., van der Voort,H.T., Krozowski,Z.S., de Kloet,E.R., and van,D.R. (1996). Partial colocalization of glucocorticoid and mineralocorticoid receptors in discrete compartments in nuclei of rat hippocampus neurons. *J. Cell Sci.* 109 (Pt 4), 787–792.
- Vandevyver,S., Dejager,L., and Libert,C. (2012). On the trail of the glucocorticoid receptor: into the nucleus and back. *Traffic.* 13, 364–374.
- Varley,K.E., Gertz,J., Bowling,K.M., Parker,S.L., Reddy,T.E., Pauli-Behn,F., Cross,M.K., Williams,B.A., Stamatoyannopoulos,J.A., Crawford,G.E., Absher,D.M., Wold,B.J., and Myers,R.M. (2013). Dynamic DNA methylation across diverse human cell lines and tissues. *Genome Res.* 23, 555–567.

- Vazquez,B.N., Laguna,T., Notario,L., and Lauzurica,P. (2012). Evidence for an intronic cis-regulatory element within CD69 gene. *Genes Immun.* 13, 356–362.
- Vermeer,H., Hendriks-Stegeman,B.I., van der Burg,B., van Buul-Offers,S.C., and Jansen,M. (2003). Glucocorticoid-induced increase in lymphocytic FKBP51 messenger ribonucleic acid expression: a potential marker for glucocorticoid sensitivity, potency, and bioavailability. *J. Clin. Endocrinol. Metab* 88, 277–284.
- Vielkind,U., Walencewicz,A., Levine,J.M., and Bohn,M.C. (1990). Type II glucocorticoid receptors are expressed in oligodendrocytes and astrocytes. *J. Neurosci. Res.* 27, 360–373.
- Viguerie,N., Picard,F., Hul,G., Roussel,B., Barbe,P., Iacovoni,J.S., Valle,C., Langin,D., and Saris,W.H. (2012). Multiple effects of a short-term dexamethasone treatment in human skeletal muscle and adipose tissue. *Physiol Genomics* 44, 141–151.
- Vinckevicius,A. and Chakravarti,D. (2012). Chromatin immunoprecipitation: advancing analysis of nuclear hormone signaling. *J. Mol. Endocrinol.* 49, R113–R123.
- Vyas,A., Mitra,R., Shankaranarayana Rao,B.S., and Chattarji,S. (2002). Chronic stress induces contrasting patterns of dendritic remodeling in hippocampal and amygdaloid neurons. *J. Neurosci.* 22, 6810–6818.
- Wang,H., Kubica,N., Ellisen,L.W., Jefferson,L.S., and Kimball,S.R. (2006a). Dexamethasone represses signaling through the mammalian target of rapamycin in muscle cells by enhancing expression of REDD1. *J. Biol. Chem.* 281, 39128–39134.
- Wang,J., Rao,S., Chu,J., Shen,X., Lévasseur,D.N., Theunissen,T.W., and Orkin,S.H. (2006b). A protein interaction network for pluripotency of embryonic stem cells. *Nature* 444, 364–368.
- Wang,J.C., Derynck,M.K., Nonaka,D.F., Khodabakhsh,D.B., Haqq,C., and Yamamoto,K.R. (2004). Chromatin immunoprecipitation (ChIP) scanning identifies primary glucocorticoid receptor target genes. *Proc. Natl. Acad. Sci. U. S. A* 101, 15603–15608.
- Wang,Z., Malone,M.H., Thomenius,M.J., Zhong,F., Xu,F., and Distelhorst,C.W. (2003). Dexamethasone-induced gene 2 (dig2) is a novel pro-survival stress gene induced rapidly by diverse apoptotic signals. *J. Biol. Chem.* 278, 27053–27058.
- Wang,Z.X., Teh,C.H., Chan,C.M., Chu,C., Rossbach,M., Kunarso,G., Allapitchay,T.B., Wong,K.Y., and Stanton,L.W. (2008). The transcription factor Zfp281 controls embryonic stem cell pluripotency by direct activation and repression of target genes. *Stem Cells* 26, 2791–2799.
- Watanabe,Y., Gould,E., and McEwen,B.S. (1992). Stress induces atrophy of apical dendrites of hippocampal CA3 pyramidal neurons. *Brain Res.* 588, 341–345.
- Weinmann,A.S., Yan,P.S., Oberley,M.J., Huang,T.H., and Farnham,P.J. (2002). Isolating human transcription factor targets by coupling chromatin immunoprecipitation and CpG island microarray analysis. *Genes Dev.* 16, 235–244.
- Wieland,S., Dobbeling,U., and Rusconi,S. (1991). Interference and synergism of glucocorticoid receptor and octamer factors. *EMBO J.* 10, 2513–2521.
- Williams,A., Sarkar,S., Cuddon,P., Ttofi,E.K., Saiki,S., Siddiqi,F.H., Jahreiss,L., Fleming,A., Pask,D., Goldsmith,P., O’Kane,C.J., Floto,R.A., and Rubinsztein,D.C. (2008). Novel targets for Huntington’s disease in an mTOR-independent autophagy pathway. *Nat. Chem. Biol.* 4, 295–305.
- Wu,P., Jiang,C., Shen,Q., and Hu,Y. (2009). Systematic gene expression profile of hypothalamus in calorie-restricted mice implicates the involvement of mTOR signaling in neuroprotective activity. *Mech. Ageing Dev.* 130, 602–610.

- Xie, W., Barr, C.L., Kim, A., Yue, F., Lee, A.Y., Eubanks, J., Dempster, E.L., and Ren, B. (2012). Base-resolution analyses of sequence and parent-of-origin dependent DNA methylation in the mouse genome. *Cell* 148, 816–831.
- Yamamoto, K.R., Darimont, B.D., Wagner, R.L., and Iniguez-Lluhi, J.A. (1998). Building transcriptional regulatory complexes: signals and surfaces. *Cold Spring Harb. Symp. Quant. Biol.* 63, 587–598.
- Yan, H., Frost, P., Shi, Y., Hoang, B., Sharma, S., Fisher, M., Gera, J., and Lichtenstein, A. (2006). Mechanism by which mammalian target of rapamycin inhibitors sensitize multiple myeloma cells to dexamethasone-induced apoptosis. *Cancer Res.* 66, 2305–2313.
- Yang, P.C., Yang, C.H., Huang, C.C., and Hsu, K.S. (2008). Phosphatidylinositol 3-kinase activation is required for stress protocol-induced modification of hippocampal synaptic plasticity. *J. Biol. Chem.* 283, 2631–2643.
- Yang, T.T., Tsao, C.W., Li, J.S., Wu, H.T., Hsu, C.T., and Cheng, J.T. (2007). Changes of dopamine content and cell proliferation by dexamethasone via pituitary adenylate cyclase-activating polypeptide in PC12 cell. *Neurosci. Lett.* 426, 45–48.
- Yang-Yen, H.F., Chambard, J.C., Sun, Y.L., Smeal, T., Schmidt, T.J., Drouin, J., and Karin, M. (1990). Transcriptional interference between c-Jun and the glucocorticoid receptor: mutual inhibition of DNA binding due to direct protein-protein interaction. *Cell* 62, 1205–1215.
- Yehuda, R., Cai, G., Golier, J.A., Sarapas, C., Galea, S., Ising, M., Rein, T., Schmeidler, J., Muller-Myhsok, B., Holsboer, F., and Buxbaum, J.D. (2009). Gene expression patterns associated with posttraumatic stress disorder following exposure to the World Trade Center attacks. *Biol. Psychiatry* 66, 708–711.
- Yu, C.Y., Mayba, O., Lee, J.V., Tran, J., Harris, C., Speed, T.P., and Wang, J.C. (2010). Genome-wide analysis of glucocorticoid receptor binding regions in adipocytes reveal gene network involved in triglyceride homeostasis. *PLoS. One.* 5, e15188.
- Yudt, M.R. and Cidlowski, J.A. (2002). The glucocorticoid receptor: coding a diversity of proteins and responses through a single gene. *Mol. Endocrinol.* 16, 1719–1726.
- Yudt, M.R., Jewell, C.M., Bienstock, R.J., and Cidlowski, J.A. (2003). Molecular origins for the dominant negative function of human glucocorticoid receptor beta. *Mol. Cell Biol.* 23, 4319–4330.
- Zalachoras, I., Houtman, R., and Meijer, O.C. (2013). Understanding stress-effects in the brain via transcriptional signal transduction pathways. *Neuroscience* 242, 97–109.
- Zhang, L., Li, H., Hu, X., Li, X.X., Smerin, S., and Ursano, R. (2011). Glucocorticoid-induced p11 over-expression and chromatin remodeling: a novel molecular mechanism of traumatic stress? *Med. Hypotheses* 76, 774–777.
- Zhang, Y., Liu, T., Meyer, C.A., Eeckhoutte, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* 9, R137.

