



Universiteit
Leiden
The Netherlands

Rhizobial NodFE-dependent fatty acids which lack an α - β unsaturation are subject to differential transfer leading to novel phospholipids

Geiger, O.; Glushka, J.; Lugtenberg, E.J.J.; Spaink, H.P.; Thomas-Oates, J.

Citation

Geiger, O., Glushka, J., Lugtenberg, E. J. J., Spaink, H. P., & Thomas-Oates, J. (1998). Rhizobial NodFE-dependent fatty acids which lack an α - β unsaturation are subject to differential transfer leading to novel phospholipids. *Molecular Plant-Microbe Interactions*, 11(1), 33-44. doi:10.1094/MPMI.1998.11.1.33

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/64319>

Note: To cite this publication please use the final published version (if applicable).

NodFE-Dependent Fatty Acids That Lack an α - β Unsaturation Are Subject to Differential Transfer, Leading to Novel Phospholipids

Otto Geiger,¹ John Glushka,² Ben J. J. Lugtenberg,¹ Herman P. Spaink,¹ and Jane E. Thomas-Oates³

¹Institute of Molecular Plant Sciences, Leiden University, Leiden, The Netherlands; ²Complex Carbohydrate Research Center, 220 Riverbend Road, Athens, GA 30602, U.S.A.; ³Bijvoet Center for Biomolecular Research, Department of Mass Spectrometry, Utrecht University, Sorbonnelaan 16, 3584 CA Utrecht, The Netherlands

Accepted 23 September 1997.

In *Rhizobium leguminosarum*, the *nodABC* and *nodFEL* operons are involved in the production of lipo-chitin oligosaccharide signals that mediate host specificity. A *nodFE*-determined, highly unsaturated C18:4 fatty acid (*trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid) is essential for the ability of the signals to induce nodule meristems and pre-infection thread structures on the host plant *Vicia sativa*. Of the *nod* genes, induction of only *nodFE* is sufficient to modify fatty acid biosynthesis to yield *trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid, with an absorbance maximum of 303 nm. This unusual C18:4 fatty acid is not only found in the lipo-chitin oligosaccharides but is also associated with the phospholipids (O. Geiger, J. E. Thomas-Oates, J. Glushka, H. P. Spaink, and B. J. J. Lugtenberg, 1994, *J. Biol. Chem.* 269:11090-11097). Here we report that the phospholipids can contain other *nodFE*-derived fatty acids, a C18:3 *trans*-4, *trans*-6, *cis*-11-octadecatrienoic acid that has a characteristic absorption maximum at 225 nm, and a C18:2 octadecadienoic acid. Neither this C18:3 nor this C18:2 fatty acid has to date been observed attached to lipo-chitin oligosaccharides, suggesting that an as yet unknown acyl transferase (presumably NodA), responsible for the transfer of the fatty acyl chain to the glycan backbone of the lipo-chitin oligosaccharides, does not transfer all fatty acids synthesized by the action of NodFE to the lipo-chitin oligosaccharides. Rather, it must have a preference for α - β unsaturated fatty acids during transfer.

Rhizobium bacteria interact with leguminous plants in a host-specific manner and can cause the formation of nitrogen-fixing root nodules. Plant-specific flavonoids induce rhizobial

nod genes needed for nodulation. In *Rhizobium leguminosarum*, the *nodABC* and *nodFEL* operons are involved in the production of lipo-chitin oligosaccharide (LCO) signal molecules that mediate host specificity. A *nodL*-determined *O*-acetyl substituent and a *nodFE*-determined, highly unsaturated C18:4 fatty acid (*trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid) are both essential for the ability of the signals to induce nodule meristems (Spaink et al. 1991) and pre-infection thread structures (van Brussel et al. 1992) on the host plant *Vicia sativa*.

We now focus on the question as to how these LCO signals are synthesized in *Rhizobium*. For the synthesis of a basic LCO, active *nodABC* genes are required (Spaink et al. 1991). During biosynthesis the NodC protein seems to function as a glycosyl transferase producing chitin oligosaccharides (Germia et al. 1994; Kamst et al. 1995). Purified NodB protein shows *N*-deacetylase activity and removes an *N*-acetyl group from the *N*-acetyl glucosamine of the nonreducing terminal residue of chitin oligosaccharides (John et al. 1993). Finally, the NodA protein is involved in the transfer of a fatty acyl residue to the free amino group of the chito-oligosaccharide backbone (Atkinson et al. 1994; Röhrig et al. 1994). Recently, it has even been demonstrated that NodA, which was formerly considered to be the product of a common *nod* gene, can function as a host-specific determinant during the transfer of fatty acids in LCO biosynthesis (Debellé et al. 1996; Ritsema et al. 1996; Roche et al. 1996).

Here, we report the results of our study of the biosynthesis and transfer of the unusual *nodFE*-derived fatty acyl residues. Of the *nod* genes, induction of *nodFE* alone is sufficient for the synthesis of the host-specific 303 nm-absorbing *trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid (C18:4) (Geiger et al. 1994), which in *R. leguminosarum* bv. *viciae* is the only *nodFE*-derived fatty acid transferred in the formation of host-specific, mitogenic LCO signal molecules (Spaink et al. 1991). Here we show that two other *nodFE*-derived fatty acyl residues, a 225 nm-absorbing *trans*-4, *trans*-6, *cis*-11-octadecatrienoic acid (C18:3) and an octadecadienoic acid (C18:2), are formed, although they are not incorporated into LCOs in *R. leguminosarum* and can only be detected in membrane phospholipids.

Corresponding author: Otto Geiger
E-mail: geig1332@mailsz.zz.tu-berlin.de

Present address of O. Geiger: Technische Universität Berlin, Institut für Biotechnologie, FG Technische Biochemie, Seestraße 13, 13353 Berlin, Germany.

RESULTS

Lipid extracts contain several *nodFE*-dependent fatty acids.

During our search for metabolites produced by the *nodFE* gene products, we observed that NodFE are sufficient to redirect fatty acid biosynthesis in *Rhizobium* in order to obtain an unusual C18:4 fatty acid. This 303 nm-absorbing fatty acid was found to be an important structural element of the host-specific LCOs of *R. leguminosarum* bv. *viciae* (Spaink et al. 1991), but it was also detected linked to the *sn*-2 position of phosphatidylcholine (Geiger et al. 1994). We had also noticed that other phospholipids showed an absorbance maximum at 225 nm and that they were also formed only after the induction of the *nodFE* genes (Geiger et al. 1994). A 225 nm-absorbance maximum is consistent with species bearing two conjugated double bonds, suggesting to us that, in addition to the 303 nm-absorbing C18:4 fatty acid, in which three *trans* double bonds are in conjugation with the carbonyl group, the phospholipids may alternatively bear different *nodFE*-dependent fatty acyl groups.

We therefore analyzed the fatty acid composition of crude lipid extracts obtained from *R. leguminosarum* LPR5045.pMP280.pMP1255, a strain that is able to overexpress the genes *nodFE* in the presence of the flavonoid inducer naringenin (Geiger et al. 1994). Figure 1 shows gas chromatograms of methyl ester derivatives obtained from lipid extracts from a noninduced (Fig. 1A) and a naringenin-induced (Fig. 1B) bacterial cell suspension. The major fatty acid methyl ester present in the preparations obtained under both experimental conditions is that of *cis*-vaccenic acid, which is known to be the predominant fatty acid in *R. legumi-*




Fig. 1. Gas chromatographic separation of methyl esters obtained from (A) uninduced and (B) *nodFE*-induced lipid extracts from *Rhizobium leguminosarum* LPR5045.pMP280.pMP1255. Detection was performed with a flame ionization detector. In the *nodFE*-induced situation, two novel substances are indicated by their masses for the molecular ions at *m/z* 294 and 292, respectively, when analyzed in a mass spectrometer.

nosarum and which has a retention time of 31.4 min. In addition, two peaks with retention times of 30.9 and 32.6 min are only present when *nodFE* have been induced (Fig. 1B). If the gas chromatograph is linked to a mass spectrometer (with electron ionization) the two peaks with retention times of 30.9 and 32.6 min give M^+ molecular ions at *m/z* 294 and *m/z* 292, respectively. These masses are consistent with methyl esters of C18:2 and C18:3 fatty acids. A fragment ion is observed at *m/z* 74 in the spectrum of both compounds, corresponding to a McLafferty rearrangement ion (data not shown) indicative of fatty acid methyl esters. In the case of flame ionization detection, the peak area of the substance with a retention time of 30.9 min is substantially larger than that of the substance eluting at 32.6 min, suggesting that there is significantly more methyl ester of the C18:2 than of the C18:3 fatty acid present. Two-dimensional, thin-layer chromatographic separation into the individual phospholipid classes and their gas chromatographic (GC) analysis after transmethylation indicated that the C18:2 (retention time 30.9 min) and the C18:3 (retention time 32.6 min) fatty acids were associated with the different classes of phospholipids (data not shown). Minor peaks with retention times of 30.9 and 32.6 min, showing M^+ molecular ions at *m/z* 294 and *m/z* 292, respectively, were also detected in methanolized lipid extract of flavonoid-induced *R. leguminosarum* bv. *viciae* RBL5560 wild type (data not shown), suggesting that both *nodFE*-related substances are formed when the *nodFE* genes are expressed at wild-type levels. These results are surprising and seem to contradict our earlier finding that the *nodFE* genes are responsible for the synthesis of a C18:4 fatty acid that can be linked to the phospholipids (Geiger et al. 1994).

Purification of *nodFE*-dependent phospholipids.

In order to understand which fatty acids are synthesized by the action of NodFE from *R. leguminosarum* bv. *viciae*, we decided to purify several *nodFE*-dependent phospholipids and determine their covalent structures.

Chloroform phases of Bligh-Dyer extracts were chromatographed on DEAE-cellulose, which allowed separation of the neutral phospholipids (phosphatidylethanolamine, PE; monomethylphosphatidylethanolamine, MMPE; dimethylphosphatidylethanolamine, DMPE; and phosphatidylcholine, PC) from anionic phospholipids (phosphatidylglycerol, PG; and cardiolipin, CL). The neutral phospholipid fraction eluted with chloroform/methanol (9:1, vol/vol) whereas the anionic phospholipids were detected in the chloroform/methanol (4:1, vol/vol) eluate containing 50 mM ammonium acetate. Neutral and anionic fractions were further separated with normal phase high-pressure liquid chromatography (HPLC) on silica gel, which allows separation based mainly on the structures of the head groups. Under these conditions, phosphatidylcholine (in the neutral fraction) is bound strongly to the column, elutes at a retention time of 34 to 40 min, and can therefore be totally separated from all other phospholipid classes. The phosphatidylcholine fraction was further subjected to reverse phase HPLC, allowing base line separation of 303 nm-absorbing phosphatidylcholine (retention time of 131 to 148 min) from 225 nm-absorbing phosphatidylcholine (retention time of 150 to 170 min) (Fig. 2).

A second class of substances found in the neutral fraction, monomethylphosphatidylethanolamine (MMPE), elutes with a

retention time of 15 to 17 min during normal phase HPLC. In a final purification step (reverse phase HPLC) the 303 nm-absorbing MMPE (retention time of 124 to 132 min) and the 225 nm-absorbing MMPE (retention time of 135 to 152 min) could be separated, yielding nearly homogenous preparations.

Separation of the anionic fraction with normal phase HPLC yielded phosphatidylglycerol (PG) with a retention time of 8 to 9 min. In a subsequent purification step (reverse phase HPLC) the 303 nm-absorbing PG (retention time of 38 to 42 min) and the 225 nm-absorbing PG (retention time of 44 to 49 min) could be separated.

Purified 303 nm-absorbing phospholipids showed absorbance ratios $A_{303}/A_{206} = 5.0$ to 7.0 depending on the phospholipid species and the solvent used. Purified 225 nm-absorbing phospholipids showed absorbance ratios $A_{225}/A_{206} = 2.0$ to 2.4 depending on the phospholipid species and the solvent used. These data suggest that the molar absorptivity of 303 nm-absorbing phospholipids at 303 nm is at least twice that of 225 nm-absorbing phospholipids at 225 nm.

Chemical structure of 225 nm-absorbing phosphatidylcholine.

The purified 225 nm-absorbing phosphatidylcholine was analyzed by 2- and 4-sector mass spectrometric (MS) methods in both the positive and the negative ion modes.

The positive ion fast atom bombardment-mass spectrum (FAB-MS) contained an intense M^+ molecular ion at m/z 782 and two somewhat less intense M^+ molecular ions at m/z 784 and 758, which are accompanied by thioglycerol adduct ions at m/z 890, 892, and 866 consistent with the presence of unsaturated fatty acids (Fukuda et al. 1985). Positive collision-induced-dissociation tandem mass spectrometry (CID MS-MS) analysis of these three M^+ ions yielded spectra all containing the same, intense product ion at m/z 184, corresponding to the choline phosphate head group ($N^+(\text{CH}_3)_3\text{-CH}_2\text{CH}_2\text{-PO}_2(\text{OH})_2$), which ion is diagnostic for phosphatidylcholine phospholipids (Hiyashi et al. 1989). The species giving rise to the ion at m/z 782 thus corresponds to a phosphatidylcholine containing fatty acyl chains with a total of 36 carbon atoms and four double bonds, while that giving the ion at m/z 784 probably derives from a related impurity with one




Fig. 2. Separation of 225 nm-absorbing phosphatidylcholine (PC) from 303 nm-absorbing PC with high-pressure liquid chromatography (HPLC) on reverse phase silica gel. The PC fraction, as obtained after chromatography on silica gel, was separated into molecular species as detailed in Materials and Methods. Absorptions at 206 (solid line), 225 (dashed line), and 303 (dotted line) nm are indicated.

fewer double bond. The compound giving the ion at m/z 758 corresponds to a phosphatidylcholine with a total of 34 carbon atoms and two double bonds.

In order to determine the fatty acyl chains present and their sites of esterification to the glycerol, the same negative ion CID method was used as proved successful in making these assignments for the 303 nm-absorbing phosphatidylcholine (Geiger et al. 1994). This involves collisional activation of the negative ion generated on loss of the choline head group, and examination of the relative intensities of the resulting product ions. The ion generated on elimination of the *sn*-2 fatty acid is always more intense than that generated by the analogous *sn*-1 elimination (Huang et al. 1992). Consequently, the negative ion at m/z 695 (corresponding to $[\text{M} - \text{choline}]^-$ for the species giving M^+ at m/z 782) was subjected to CID MS-MS analysis. The resulting spectrum contained two sets of ions. Those at m/z 277 and 281 represent carboxylate anions for C18:3 and C18:1 fatty acids, respectively, while those at m/z 417 and 413 arise on elimination of a C18:3 and a C18:1 fatty acid from the parent ion, respectively. Since the ion at m/z 417 is significantly more intense than that at m/z 413, it can be concluded that the unusual C18:3 fatty acid is located at *sn*-2, while the *sn*-1 position bears the commonly occurring C18:1 fatty acid.

The ^1H nuclear magnetic resonance (NMR) spectra of the 225 nm-absorbing PC showed resonances and coupling patterns expected for acyl side chains containing double bonds. In particular, the two-dimensional, double-quantum-filtered proton-proton correlated (DQF-COSY) spectrum (Fig. 3) revealed four signals between δ 5.3 and δ 5.9 that correspond to methine protons from conjugated double bonds in positions 4 to 7 of one of the acyl chains. The figure indicates their connectivity to the upfield signals of the methylene protons (δ 2.4 to δ 1.2) in the fatty acid chain. The 225 nm-absorbing PC contains the C18:3 fatty acid, where the two conjugated double bonds are both in *trans* (E-E) configuration. This is confirmed by the proton coupling constants across the bonds of 14.5 Hz. These results are similar to those found for the acyl chain of the 303 nm-absorbing PC and the acyl chain of mitogenic LCOs from *R. leguminosarum* bv. *viciae* (Geiger et al. 1994; Spaink et al. 1991). However, the chemical shifts of protons from positions 2 and 3 (δ 2.30, δ 2.23) clearly indicate saturated methylene groups. Additional signals at δ 5.2 are due to the isolated double bonds in both chains at position 11 and 12, and resonances belonging to the phosphatidylcholine group are also indicated (c1 to c5 in Figure 3). The spectrum also shows signals due to some unidentified contaminants.

From the MS and the NMR spectroscopic studies we conclude that the 225 nm-absorbing phosphatidylcholine has the primary chemical structure presented in Figure 4.

Chemical structures of *nodFE*-dependent monomethylphosphatidyl ethanolamines and phosphatidylglycerols.

The purified 303 nm- and 225 nm-absorbing monomethylphosphatidyl ethanolamine and phosphatidylglycerol fractions were analyzed by 2- and 4-sector MS methods in both the positive and the negative ion modes. The results of the analyses and the tentative structural assignments are summarized in Table 1.

The fraction containing the 303 nm-absorbing MMPE contained three different MMPE molecular species having

$[M+H]^+$ pseudomolecular ions at m/z 704, m/z 730 (most intense), and m/z 752 that co-migrated through all purification procedures. These pseudomolecular ions were accompanied by a thioglycerol adduct ion at m/z 812, 838, and 860, respectively. These three molecular species were represented by corresponding $[M-H]^-$ pseudomolecular ions at m/z 702, 728, and 750 in the negative ion mode. Each $[M+H]^+$ ion was subjected to CID MS-MS in the positive ion mode. In each spectrum one major product ion was observed (Fig. 5), corresponding to the loss of 155 amu resulting from the elimination of the monomethylethanolamine phosphate head group (see Table 1).

Negative ion CID MS-MS of the $[M-H]^-$ pseudomolecular

ions was carried out in order to establish which fatty acids are present in each species, based on a consideration of the carboxylate anions produced. The $[M-H]^-$ precursor ion at m/z 702 yields a negative ion CID MS-MS spectrum containing ions corresponding to carboxylate anions at m/z 253 for a C16:1 fatty acid, and at m/z 255 for a C16:0 fatty acid. Similarly, the precursor $[M-H]^-$ ion at m/z 728 yields product ions at m/z 253 (C16:1) and 281 (C18:1), while the precursor with m/z 750 yields product ions at 275 and 281, indicating it corresponds to a MMPE bearing one C18:4 and one C18:1 fatty acyl chain (Fig. 5 insert), and is thus presumably responsible for the 303 nm-absorption maximum.




Fig. 3. Double-quantum-filtered proton-proton correlated (DQF-COSY) spectrum (600 MHz) of the 225 nm-absorbing phosphatidylcholine (PC). Connectivity indicated corresponds to the conjugated double bond system (peaks labeled 4 to 7) and to the rest of the acyl chain (peaks 2, 3, and 8 to 15).

The positive ion FAB-MS of the purified 225 nm-absorbing MMPE gave a major $[M+H]^+$ pseudomolecular ion at m/z 754 accompanied by a thioglycerol adduction at m/z 862. In the negative ion mode, an $[M-H]^-$ pseudomolecular ion for the same species was observed at m/z 752.

In order to test for the presence of MMPE, the $[M+H]^+$ ion at m/z 754 was subjected to CID MS-MS in the positive ion mode. One major product ion was observed (Fig. 6) at m/z 599, corresponding to the loss of the monomethylethanolamine phosphate head group $[M-PO_4CH_2CH_2N(CH_3)]^+$ (155 amu) as expected from MMPE.

CID MS-MS in the negative ion mode of the precursor ion at m/z 752 was carried out (Fig. 6 insert) in order to establish the fatty acids present. Two product ions were observed at m/z 277 and 281, corresponding to the carboxylate anions of a C18:3 and a C18:1 fatty acid.

The fraction corresponding to the 303 nm-absorption maximum for species assumed to correspond to phosphatidylglycerols failed to give pseudomolecular ions when analyzed by positive ion mode FAB-MS in a matrix of thioglycerol, but gave intense ions at m/z 769 and 791 when analyzed in a matrix of *m*-NBA. These ions correspond to $[M+H]^+$ and $[M+Na]^+$ pseudomolecular ions for a PG species bearing a total of 36 carbon atoms and five double bonds. Positive ion mode CID analysis of each of these two pseudomolecular ions yielded spectra containing an intense fragment ion at m/z 597, which corresponds to the loss of the glycerol phosphate head group from the $[M+H]^+$ ion (Fig. 7) and the loss of glycerol sodium phosphate from the $[M+Na]^+$ species, together with a fragment ion at m/z 195 for the glycerol sodium phosphate head group, demonstrating that these components do indeed correspond to PGs. In addition to the intense fragment ion at m/z 597 in the CID mass spectrum obtained from the 769 precursor, a second, much less intense fragment ion was observed at m/z 575, which can be rationalized as arising by loss of a glycerol sodium phosphate head group from a species having a total of 34 carbons and two double bonds, presumably from one C16:1 and one C18:1 fatty acyl chain. In other words, there is evidence for the presence of a small amount of a contaminating species present in this fraction that would not give rise to a 303 nm-absorption maximum, but that is incompletely fractionated during the chromatography.




Fig. 4. Structures of 225 nm-absorbing phosphatidylcholine (PC) and 303 nm-absorbing PC from *Rhizobium leguminosarum* bv. *viciae*.

The negative ion mode FAB-MS of this fraction, obtained in a matrix of thioglycerol, contains a pair of $[M-H]^-$ pseudomolecular ions at m/z 767, corresponding to a PG species with a total of 36 carbon atoms and five double bonds, and m/z 745, corresponding to the species having 34 carbon atoms and two double bonds. The $[M-H]^-$ ions are accompanied by thioglycerol adduct ions at m/z 853 and 875. The negative ion collision spectrum obtained from the ion at m/z 767 gives ions at m/z 275 and 281 (Fig. 7 insert), corresponding to carboxylate ions for its component fatty acids, a C18:4 (m/z 275) and a C18:1 (m/z 281) species.

The fraction eluting very close to that with the 303 nm-absorption maximum and having a 225 nm-absorption maximum was also analyzed by positive ion mode FAB-MS in a matrix of *m*-NBA. This gave a series of ions at m/z 769, 771, 773, 791, 793, and 795, the latter two being much the most intense, and corresponding to $[M+Na]^+$ pseudomolecular ions for species bearing a total of 36 carbon atoms and four and three double bonds, respectively. Positive ion CID MS-MS analysis of the precursor at m/z 793 yielded one fragment ion at m/z 599 corresponding to the loss of the glycerol sodium phosphate head group from an $[M+Na]^+$ ion for a PG bearing a total of 36 carbon atoms and four double bonds, together with an ion at m/z 621, for the loss of glycerol phosphate. In the same way, the positive ion CID MS-MS analysis of the

Table 1. Summary of fast atom bombardment-mass spectrometric data obtained from high-pressure liquid chromatography (HPLC)-purified phospholipids with structural assignments^a

| Fraction | Ionization mode | $[M+H]^+$ or $[M-H]^-$ | Fragments | Assignments ^b |
|----------|-----------------|------------------------|-----------|--------------------------|
| A | Pos | 704 | 549 | MMPE |
| | Neg | 702 | 253, 255 | C16:1, C16:0 |
| | Pos | 730 | 575 | MMPE |
| | Neg | 728 | 253, 281 | C16:1, C18:1 |
| | Pos | 752 | 597 | MMPE |
| | Neg | 750 | 275, 281 | C18:4, C18:1 |
| B | Pos | 754 | 599 | MMPE |
| | Neg | 752 | 277, 281 | C18:3, C18:1 |
| C | Pos | 780 | 597 | PC |
| | Neg | 778 | 275, 281 | C18:4, C18:1 |
| D | Pos | 782 | 184 | PC |
| | Neg | 780 | 277, 281 | C18:3, C18:1 |
| | Pos | 784 | 184 | PC |
| | Pos | 758 | 184 | PC |
| E | Pos | 769 | 597 | PG |
| | Neg | 767 | 275, 281 | C18:4, C18:1 |
| F | Pos | 771 | 599 | PG |
| | Neg | 769 | 277, 281 | C18:3, C18:1 |
| | Pos | 773 | 601 | PG |
| | Neg | 771 | 279, 281 | C18:2, C18:1 |

^a From the neutral fraction, phospholipids with a retention time on silica gel of 15 to 17 (A, B) or 34 to 40 (C, D) min were separated further on reverse phase HPLC. Reverse phase separation yielded a 303 nm-absorbing fraction with a retention time of 124 to 132 min (A) and a 225 nm-absorbing fraction with a retention time of 135 to 152 min (B), or in the alternative case a 303 nm-absorbing fraction with a retention time of 128 to 144 min (C), and a 225 nm-absorbing fraction with a retention time of 150 to 164 min (D). The anionic fraction that eluted with a retention time of 8 to 9 min from silica gel was separated further on reverse phase HPLC, yielding a 303 nm-absorbing fraction with a retention time of 38 to 42 min (E), and a 225 nm-absorbing fraction with a retention time of 44 to 49 min (F).

^b See Results for definitions of abbreviations.

precursor ion at m/z 795 yielded fragments, corresponding to the loss of the glycerol sodium phosphate head group (m/z 601) from a PG species bearing a total of 36 carbon atoms and three double bonds, and present as its $[M+Na]^+$ ion, together with a second fragment at m/z 623 produced by the loss of the glycerol phosphate head group. The negative ion mass spectrum obtained in a matrix of thioglycerol contains a pair of intense $[M-H]^-$ ions at m/z 769 and 771, together with thioglycerol adduct ions at m/z 877 and 879, corresponding to PGs bearing 36 carbon atoms and a total of four and three double bonds, respectively. Negative ion CID spectra of the $[M-H]^-$ ions contain product ions corresponding to carboxylate anions of the component fatty acyl chains—the precursor ion at m/z 769 generates such ions as m/z 277 (C18:3) and 281 (C18:1) (Fig. 8A), while that at m/z 771 yields product ions at m/z 279 (C18:2) and 281 (C18:1) (Fig. 8B).

***nodFE* from *R. leguminosarum* bv. *viciae* are involved in the synthesis of at least three polyunsaturated fatty acids.**

In an earlier paper we described the production of 303 nm-absorbing phospholipids containing a C18:4 fatty acid caused by the presence of *nodFE* from *R. leguminosarum* bv. *viciae* (Geiger et al. 1994). Separation and purification of phospholipid species based on their acyl chains allows baseline separation of phospholipids bearing C18:4 (303 nm absorbance) from phospholipids bearing C18:3 (225 nm absorbance) fatty acyl moieties. Phospholipids having a *nodFE*-derived C18:2 (no specific absorbance, but absorbance at 205 nm) fatty acyl chain are sometimes found as minor impurities in fractions containing phospholipids with a 225-nm absorbance. Peak areas of GC analysis of fatty acyl methyl esters (Fig. 1) indicate that *nodFE*-induced lipid extracts from *R. leguminosarum* bv. *viciae* contain more C18:2 than C18:3 fatty acids. Chromatographic separation of the 303 nm-absorbing from the 225 nm-absorbing species of individual phospholipid classes (Fig. 2) and their relative molar absorptivities indicate that in each case significantly more of the 225 nm-absorbing, C18:3 fatty acid-containing species is present than of the 303 nm-absorbing, C18:4 fatty acid-containing species.

ration of phospholipids bearing C18:4 (303 nm absorbance) from phospholipids bearing C18:3 (225 nm absorbance) fatty acyl moieties. Phospholipids having a *nodFE*-derived C18:2 (no specific absorbance, but absorbance at 205 nm) fatty acyl chain are sometimes found as minor impurities in fractions containing phospholipids with a 225-nm absorbance. Peak areas of GC analysis of fatty acyl methyl esters (Fig. 1) indicate that *nodFE*-induced lipid extracts from *R. leguminosarum* bv. *viciae* contain more C18:2 than C18:3 fatty acids. Chromatographic separation of the 303 nm-absorbing from the 225 nm-absorbing species of individual phospholipid classes (Fig. 2) and their relative molar absorptivities indicate that in each case significantly more of the 225 nm-absorbing, C18:3 fatty acid-containing species is present than of the 303 nm-absorbing, C18:4 fatty acid-containing species.

DISCUSSION

Three *nodFE*-related polyunsaturated fatty acids are produced in *R. leguminosarum* bv. *viciae*.

GC-MS analysis of methanolized Bligh-Dyer extracts of whole cells from *R. leguminosarum* bv. *viciae* revealed the methyl esters of two novel *nodFE*-induced fatty acids with the masses 294, corresponding to a C18:2 fatty acid, and 292, rep-




Fig. 5. Mass spectrometry of 303 nm-absorbing monomethylphosphatidylethanolamine (MMPE). Positive ion collision-induced-dissociation tandem mass spectrum of the 303 nm-absorbing MMPE with partial negative ion collision spectrum as insert.

representing a C18:3 fatty acid. Authentic, chemically synthesized *trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid (C18:4) methyl ester, with a mass of 290, migrates on GC with the same retention time as other major fatty acid methyl esters (data not shown) and therefore the *trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid, which is formed in lesser amounts than the *nodFE*-dependent C18:2 or C18:3 fatty acids, escaped detection in whole-cell Bligh-Dyer extracts as its methyl ester. Diode array analysis of HPLC elution profiles of phospholipids allowed detection of either one of two *nodFE*-induced absorbances. If a phospholipid showed a 225-nm absorbance, it contained a C18:3 fatty acid (mass of fatty acid methyl ester = 292; presumably *trans*-4, *trans*-6, *cis*-11-octadecatrienoic acid), and if a phospholipid showed an absorbance maximum at 303 nm it contained a C18:4 fatty acid, identified as *trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid (mass of fatty acid methyl ester = 290) (Geiger et al. 1994), the latter being identical to the *nodFE*-derived polyunsaturated fatty acid, conferring host specificity on the mitogenic LCOs of *R. leguminosarum* bv. *viciae* (Spaink et al. 1991). We therefore conclude that *nodFE* are responsible for the synthesis of three polyunsaturated fatty acids: a C18:2 fatty acid (mass of methyl ester = 294, probably *trans*-6, *cis*-11-octadecadienoic acid); a 225 nm-absorbing C18:3 fatty acid (mass of methyl ester = 292; *trans*-4, *trans*-6, *cis*-11-

octadecatrienoic acid); and the 303 nm-absorbing C18:4 fatty acid (mass of methyl ester = 290), identified previously as *trans*-2, *trans*-4, *trans*-6, *cis*-11-octadecatetraenoic acid (Fig. 9). It is interesting to note that when a plasmid carrying the *nodFE* genes of *R. leguminosarum* bv. *viciae* was introduced into *R. meliloti nodFE*- and *nodFEG*-deleted strains, LCOs with polyunsaturated C18 fatty acids (C18:2, C18:3, and C18:4) were detected (Demont et al. 1993). However, the C18:3 and the C18:2 fatty acyl groups found on LCOs of *R. meliloti* harboring the *nodFE* genes of *R. leguminosarum* bv. *viciae* are strikingly different from the C18:3 and the C18:2 fatty acyl groups we have found in the phospholipids of *R. leguminosarum* bv. *viciae* and which we describe here. The C18:3 and the C18:2 fatty acyl groups found on LCOs of *R. meliloti* harboring the *nodFE* genes of *R. leguminosarum* bv. *viciae* are described as *trans*-2, *trans*-4, *cis*-11-octadecatrienoic acid (absorbance maximum at 270 nm) and *trans*-2, *cis*-11-octadecadienoic acid (absorbance maximum around 220 nm), respectively, and are therefore both α - β unsaturated fatty acids. These latter two fatty acids have not been described in *R. leguminosarum* bv. *viciae* and their occurrence in *R. meliloti* might be due to the presence of an extra enoyl reductase in *R. meliloti* (Demont et al. 1993).

From the GC elution profile of methanolized Bligh-Dyer extracts of whole cells from *Rhizobium leguminosarum* we




Fig. 6. Mass spectrometry of 225 nm-absorbing monomethylphosphatidylethanolamine (MMPE). Positive ion collision-induced-dissociation tandem mass spectrum of the 225 nm-absorbing MMPE with partial negative ion collision spectrum as insert.

conclude that more C18:2 fatty acid than C18:3 fatty acid is produced. From the observation that the 225-nm absorbance in a given phospholipid class is always higher than the 303-nm absorbance, and considering that for a less conjugated system (C18:3) the molar extinction coefficients are lower than for a more conjugated system (C18:4), we also conclude that the C18:3 fatty acid is more abundant than the C18:4 fatty acid.

Purification of phospholipids containing *nodFE*-dependent fatty acids.

Anionic phospholipids (PG and CL) were separated from neutral phospholipids by chromatography on DEAE-cellulose 52. Purification into classes according to head group was achieved with normal phase HPLC. Further purification by reverse phase HPLC (Spherisorb ODS-2) allows a separation on the basis of the fatty acyl residues, thereby allowing the isolation of individual phospholipid species. The following *nodFE*-dependent phospholipids have been purified and extensively characterized by MS and MS-MS in the positive and negative ion modes. The results allow a determination of the polar head groups (in the positive mode) as well as the identification (and for PCs the determination of the sites of attachment) of the phospholipid-linked fatty acids (in the negative mode): (i) the 303 nm-absorbing PC contains a C18:4 at *sn*-2 position and a C18:1 fatty acid at *sn*-1 position (Fig. 4); (ii)

the 225 nm-absorbing PC contains a C18:3 at *sn*-2 position and a C18:1 fatty acid at *sn*-1 position (Fig. 4); (iii) the 303 nm-absorbing MMPE contains one C18:4 and one C18:1 fatty acid; (iv) the 225 nm-absorbing MMPE contains one C18:3 and one C18:1 fatty acid; (v) the 303 nm-absorbing PG contains one C18:4 and one C18:1 fatty acid; (vi) the 225 nm-absorbing PG contains one C18:3 and one C18:1 fatty acid; and (vii) as a minor impurity in the 225 nm-absorbing PG we find a PG that contains a C18:2 and a C18:1 fatty acid.

In conclusion, NodFE are involved in the synthesis of three polyunsaturated fatty acids (C18:2, C18:3, and C18:4) and, of all the *nod* genes, only the *nodFE* genes are needed for their synthesis. All three fatty acids are incorporated into the phospholipids (amounts: C18:2 > C18:3 > C18:4) and they show a characteristic absorbance maximum at 303 (C18:4) or 225 (C18:3) nm, or show only absorbance at 206 nm but no specific absorbance (C18:2).

Biosynthesis and acyl transfer of *nodFE*-dependent fatty acids.

Surprisingly, the mitogenic, host-specific LCOs from *R. leguminosarum* bv. *viciae* contain only one sort of *nodFE*-derived fatty acid, the α - β unsaturated C18:4 fatty acid. Therefore, we report here for the first time on novel fatty acids synthesized by the biochemical action of NodFE but which




Fig. 7. Mass spectrometry of 303 nm-absorbing phosphatidylglycerol (PG). Positive ion collision-induced-dissociation tandem mass spectrum of the 303 nm-absorbing PG with partial negative ion collision spectrum as insert.

are not used in LCO biosynthesis. These observations have yielded an insight into LCO biosynthesis, especially as to how the acyltransferase might work. Only the C18:4 fatty acid is selectively transferred to the LCOs. There are two possible explanations for this effect: Either (i) the acyltransferase (presumably NodA) that transfers fatty acyl residues to de-*N*-acetylated chitooligomers has a high degree of selectivity for α - β unsaturated fatty acids, or (ii) the LCO acyltransferase is selective for acyl-NodF intermediates and fatty acids lacking an α - β unsaturation (C18:3 and C18:2) might not be linked to NodF. A model on the biosynthesis of the different *nodFE*-derived fatty acids in *R. leguminosarum* bv. *viciae* (Fig. 10) explains how they can be linked to different acyl carrier proteins, either the constitutive AcpP or the nodulation protein NodF. It is currently thought that the acyl carrier protein NodF and the presumable condensing enzyme NodE modify normal fatty acid biosynthesis in such a way that when α - β unsaturated fatty acyl residues are linked to NodF they do not undergo the second reduction step of a normal fatty acid synthesis elongation cycle (Geiger et al. 1994). In the following

round of fatty acid elongation an C14:2 α - β unsaturated acyl-NodF could be condensed either with malonyl-NodF or malonyl-AcpP, leading to the formation of C16:3 acyl-NodF and C16:3 acyl-AcpP derivatives with two *trans* double bonds in conjugation to the carbonyl group, respectively. The C16:3 acyl-AcpP, which is not protected against the second reduction step of the fatty acid elongation cycle, would be reduced in the α - β position of the acyl residue, leading to C16:2 acyl-AcpP. As *R. leguminosarum* predominantly synthesizes fatty acids with a chain length of 18 carbons (Spaink et al. 1991), it is to be expected that the C16:2 acyl-AcpP is converted in a subsequent complete elongation cycle to the C18:2 octadecadienoyl-AcpP. The C16:3 acyl-NodF, harboring two *trans* double bonds in conjugation to the carbonyl group, would be condensed in another round of fatty acid elongation with malonyl-NodF or malonyl-AcpP, leading to the formation of C18:4 acyl-NodF and C18:4 acyl-AcpP derivatives with three *trans* double bonds in conjugation to the carbonyl group, respectively. The unprotected C18:4 acyl-AcpP would be subject to a second reduction, leading to an α - β saturated C18:3




Fig. 8. Fatty acyl residues in *nodFE*-derived phosphatidylglycerols (PGs). Partial collision-induced-dissociation tandem mass spectra in the negative ion mode of (A) the 225 nm-absorbing PG and (B) the C18:2 fatty acyl-containing PG.

octadecatrienoyl-AcpP. The C18:4 octadecatetraenoyl-NodF remains unaltered until acyl transfer reactions occur. Therefore, of the *nodFE*-derived fatty acids only the C18:4 would be linked to NodF and transferred to LCOs and phospholipids, whereas the others (C18:3 and C18:2) would be linked to the constitutive acyl carrier protein AcpP and would not be transferred to LCOs, but only to phospholipids (Fig. 10). Such a mechanism would lead to the introduction of only the C18:4 fatty acid into LCOs and to a significant discrimination against the other *nodFE*-derived polyunsaturated fatty acids, the C18:3 and the C18:2.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions.

Strains and plasmids are listed in Table 2. Broad host range plasmids were mobilized from *Escherichia coli* K12 strain KMBL1164 to *R. leguminosarum* with pRK2013 as a helper plasmid (Ditta et al. 1980) as previously described (Spaink et al. 1987).

Cultures of *R. leguminosarum* were grown on medium B (van Brussel et al. 1977) at 30°C on a gyratory shaker. If strains harbored broad host range plasmids, 0.5 mg of streptomycin per ml was added to maintain IncQ plasmids and 2 µg of tetracycline per ml was added to maintain IncP plasmids. For induction, naringenin (1.5 µM final concentration) was added at a cell density of 5×10^7 per ml. Cells were usually harvested after 3 generations of further growth.

Extraction of lipids.

Lipids were extracted by a modified Bligh and Dyer (Bligh and Dyer 1959) procedure. Wet cell paste was made up to 1 volume with water, and to the suspension 3.75 volumes of methanol/chloroform (2:1, vol/vol) was added. The mixture

was gently stirred for 1 h at room temperature. After centrifugation, the supernatant extract was decanted into a glass tube and the pellet was reextracted with 4.75 volumes methanol/chloroform/water (2:1:0.8, vol/vol) and centrifuged. To the combined supernatant extracts were added 2.5 volumes each of chloroform and water, and the mixture was centrifuged. The lower chloroform phase was withdrawn and brought to dryness in a rotary evaporator. The lipid residue was immediately redissolved in methanol/chloroform (1:1) and was stored under nitrogen at -20°C.

Methanolysis of lipids.

Dried lipid extracts were treated with 0.5 ml of hydrochloric methanol (0.6 N HCl in methanol) in sealed Teflon tubes under nitrogen at 65°C for 15 h. After addition of 0.5 ml of water the methyl esters were extracted twice with two volumes of 0.5 ml of methylene chloride.

Analysis of fatty acid methyl esters with GC and GC linked to MS.

The combined methylene chloride extracts were dried down and taken up in a total volume of 50 µl of methylene chloride. Usually, 1 µl of sample was injected and analysis was performed on a 60-m DB-1 column (J & W) with the following temperature program: the initial temperature of 75°C was raised at the rate of 10° per min and was held constant after the final temperature of 250°C was reached. The injector temperature was 250°C.

For quantification of fatty acid methyl esters a flame ionization detector (at 250°C) was used with a Hewlett-Packard 5890 instrument fitted with the DB-1 column.

When GC-MS was performed, a Hewlett-Packard 5995 instrument fitted with the DB-1 column was coupled to a HP 300 data system. MS temperatures were 300°C for the transfer line, 200°C for the ion source, and 230°C for the analyzer. Data were acquired in the electron impact mode (70 eV). The scanning range was 50 to 650 *m/z* at 1 s per decade.

Purification of phospholipids.

In order to obtain enough material to allow NMR spectroscopy on individual phospholipid species, 70 liters of rhizobial culture was grown to obtain 655 g of wet cells of naringenin-induced *R. leguminosarum* LPR5045.pMP280.pMP1255. Bligh-Dyer extraction yielded about 4.5 g of total lipids.

A crude separation of neutral and anionic phospholipids was achieved by chromatography on DEAE-cellulose (DE-52, Whatman) (Kates 1986). The sample was applied as a solution in 20 ml of chloroform (4.0 g of total lipids) to a 200-ml column of DEAE 52-cellulose in its acetate form, which had been equilibrated with chloroform. The column was then sequentially eluted with the following solvents: 5 volumes of chloroform, 9 volumes of chloroform-methanol (9:1, vol/vol), 5 volumes of chloroform-methanol (1:1, vol/vol), and 10 volumes of chloroform-methanol (4:1, vol/vol) containing 50 mM ammonium acetate. The individual fractions were brought to dryness in a rotary evaporator and they were immediately dissolved in methanol/chloroform (1:1, vol/vol) and stored under nitrogen at -20°C.

All major phospholipid classes were separated by normal-phase silica gel HPLC (Patton and Robbins 1990), either on an analytical Hypersil column 4.6 × 160 mm (Shandon, Pitts-




Fig. 9. Proposed structures for *nodFE*-derived fatty acids in *Rhizobium leguminosarum* bv. *viciae*.

burgh, PA) with a flow rate of 1 ml per min, or on a Hyperprep 120 Silica 12U column 4.6 × 250 mm (Alltech, Deerfield, IL) with a flow rate of 2 ml per min. The mobile phase consisted of hexane/2-propanol/ethanol/25 mM potassium phosphate (pH 7.0)/acetic acid (485:376:100:56:0.275, vol/vol). Diode array spectroscopic detection was performed with a Pharmacia RSD 2140 optical unit.

The purified phospholipid classes were separated into molecular species on a 4 × 250 mm Superpac Spherisorb ODS-2 column (Pharmacia) (Patton et al. 1982). They were eluted with methanol/water/acetonitrile (90.5:7:2.5, vol/vol) containing 20 mM choline chloride at a flow rate of 0.5 ml per min.

Table 2. Bacterial strains and plasmids

| Strain or plasmid | Relevant characteristics | Source or reference |
|--------------------------------|--|---------------------|
| <i>Rhizobium leguminosarum</i> | | |
| LPR5045 | <i>R. leguminosarum</i> bv. <i>trifolii</i> Ri ^F cured of Sym plasmid | Zaat et al. 1987 |
| RBL5560 | LPR5045 with Sym plasmid pRL1JI | Zaat et al. 1987 |
| Plasmids | | |
| pMP280 | IncP carrying pr. <i>nodD-nodD</i> | Spaink et al. 1987 |
| pMP604 | IncP carrying FITA-type <i>nodD604</i> | Spaink et al. 1989 |
| pMP1255 | IncQ carrying <i>nodFE</i> | Geiger et al. 1991 |




Fig. 10. Model for the biosynthesis and acyl transfer of the different *nodFE*-derived fatty acids.

MS.

Positive and negative ion mode mass spectra were obtained from the purified 303 nm- and 225 nm-absorbing phospholipids. Samples redissolved in chloroform/methanol (1:1, vol/vol) were analyzed by loading 1 to 3 μ l of sample solution into a matrix of monothioglycerol (positive ion mode) or *meta*-nitrobenzyl alcohol (*m*-NBA) (negative ion mode). Mass spectra were obtained with MS1 of a JEOL JMS-SX/SX102A tandem mass spectrometer operated at 10 kV accelerating voltage. The FAB gun was operated at 6 kV, with an emission current of 10 mA, and with xenon as the bombarding gas. Spectra were scanned at a speed of 30 s for the full mass range specified by the accelerating voltage used, and were recorded and averaged with JEOL COMPLEMENT software run on a Hewlett-Packard 9000 series data system. Tandem mass spectra were obtained from the same samples following CID in the third field free region, with the same instrument in its 4-sector mode under similar conditions, and with helium as the collision gas at a pressure sufficient to reduce the parent ion to one third of its original intensity.

¹H NMR analysis.

The PC sample with an absorption maximum at 225 nm ($E_{225} = 32/\text{ml}$) was dissolved in 0.6 ml of CD₃OD/CDCl₃ (1:5, vol/vol). Spectra were acquired on a Bruker AMX600 at 25°C. Two-dimensional DQF-COSY experiments (Piantini et al. 1982) were performed in phase-sensitive mode by the TPPI method (Marion and Wüthrich 1983).

ACKNOWLEDGMENTS

This work was funded in part by the European Communities BIO-TECH Programme 1993-1996, as part of the Project of Technical Priority, Contract No. BIO2 CT93 0400 (to B.J.J.L.), and The Netherlands Organization for Scientific Research (J.E.T.-O. and H.P.S.).

LITERATURE CITED

Atkinson, E. M., Palcic, M. M., Hindsgaul, O., and Long, S. R. 1994. Biosynthesis of *Rhizobium meliloti* lipooligosaccharide Nod factors: NodA is required for an *N*-acyltransferase activity. Proc. Natl. Acad. Sci. USA 91:8418-8422.

Bligh, E. G., and Dyer, W. J. 1959. A rapid method of total lipid extraction and purification. Can. J. Biochem. Physiol. 37:911-917.

Debellé, F., Plazanet, C., Roche, P., Pujol, C., Savagnac, A., Rosenberg, C., Promé, J. C., and Dénarié, J. 1996. The NodA proteins of *Rhizobium meliloti* and *Rhizobium tropici* specify the *N*-acylation of Nod factors by different fatty acids. Mol. Microbiol. 22:303-314.

Demont, N., Debellé, F., Aurelle, H., Dénarié, J., and Promé, J. C. 1993. Role of the *Rhizobium meliloti* *nodF* and *nodE* genes in the biosynthesis of lipo-oligosaccharidic nodulation factors. J. Biol. Chem. 268:20134-20142.

Ditta, G., Stanfield, S., Corbin, D., and Helinski, D. R. 1980. Broad host-range DNA cloning system for gram-negative bacteria: Construction of a gene bank of *Rhizobium meliloti*. Proc. Natl. Acad. Sci. USA 77:7347-7351.

Fukuda, M., Dell, A., Oates, J. E., Wu, P., Klock, J. C., and Fukuda, M. 1985. Structure of glycosphingolipids isolated from human granulocytes. J. Biol. Chem. 260:1067-1082.

Geiger, O., Spaink, H. P., and Kennedy, E. P. 1991. Isolation of the *Rhizobium leguminosarum* NodF Nodulation Protein: NodF carries a 4'-phosphopantetheine prosthetic group. J. Bacteriol. 173:2872-2878.

Geiger, O., Thomas-Oates, J. E., Glushka, J., Spaink, H. P., and Lugtenberg, B. J. J. 1994. Phospholipids of *Rhizobium* contain *nodE*-determined highly unsaturated fatty acid moieties. J. Biol. Chem. 269:11090-11097.

Geremia, R. A., Mergaert, P., Geelen, D., van Montagu, M., and Holsters, M. 1994. The NodC protein of *Azorhizobium caulinodans* is an *N*-acetylglucosaminyltransferase. Proc. Natl. Acad. Sci. USA 91:2669-2673.

Hiyashi, A., Matsubara, T., Morita, M., Kinoshita, T., and Nakamura, T. 1989. Structural analysis of choline phospholipids by fast atom bombardment mass spectrometry and tandem mass spectrometry. J. Biochem. (Tokyo) 106:264-269.

Huang, Z.-H., Gage, D. A., and Sweeley, C. C. 1992. Characterization of diacylglycerolphosphocholine molecular species by FAB-CAD-MS/MS: A general method not sensitive to the nature of the fatty acyl groups. J. Am. Soc. Mass Spectrom. 3:71-78.

John, M., Röhrig, H., Schmidt, J., Wieneke, U., and Schell, J. 1993. *Rhizobium* NodB protein involved in nodulation signal synthesis is a chitooligosaccharide deacetylase. Proc. Natl. Acad. Sci. USA 90:625-629.

Kamst, E., van der Drift, K. M. G. M., Thomas-Oates, J. E., Lugtenberg, B. J. J., and Spaink, H. P. 1995. Mass spectrometric analysis of chitin oligosaccharides produced by *Rhizobium* NodC protein in *Escherichia coli*. J. Bacteriol. 177:6282-6285.

Kates, M. 1986. Techniques of Lipidology. 2nd ed. Elsevier Science Publishers, Amsterdam.

Marion, D., and Wüthrich, K. 1983. Application of phase sensitive two-dimensional correlated spectroscopy (COSY) for measurements of 1H-1H spin-spin coupling constants in proteins. Biochem. Biophys. Res. Commun. 113:967-974.

Patton, G. M., Fasulo, J. M., and Robbins, S. J. 1982. Separation of phospholipids and individual molecular species of phospholipids by high-performance liquid chromatography. J. Lipid Res. 23:190-196.

Patton, G. M., and Robbins, S. J. 1990. Extraction of phospholipids and analysis of phospholipid molecular species. Methods Enzymol. 187:195-215.

Piantini, U., Sorensen, O., and Ernst, R. R. 1982. Multiple quantum filters for elucidating NMR coupling networks. J. Am. Chem. Soc. 104:6800-6801.

Ritsema, T., Wijffjes, A. H. M., Lugtenberg, B. J. J., and Spaink, H. P. 1996. *Rhizobium* nodulation protein NodA is a host-specific determinant of the transfer of fatty acids in Nod factor biosynthesis. Mol. Gen. Genet. 251:44-51.

Roche, P., Maillat, F., Plazanet, C., Debellé, F., Ferro, M., Truchet, G., Promé, J.-C., and Dénarié, J. 1996. The common *nodABC* genes of *Rhizobium meliloti* are host-range determinants. Proc. Natl. Acad. Sci. USA 93:15305-15310.

Röhrig, H., Schmidt, J., Wieneke, U., Kondorosi, E., Barlier, I., Schell, J., and John, M. 1994. Biosynthesis of lipooligosaccharide nodulation factors: *Rhizobium* NodA protein is involved in *N*-acylation of the chitooligosaccharide backbone. Proc. Natl. Acad. Sci. USA 91:3122-3126.

Spaink, H. P., Okker, R. J. H., Wijffelman, C. A., Pees, E., and Lugtenberg, B. J. J. 1987. Promoters in the nodulation region of the *Rhizobium leguminosarum* sym plasmid pRL1J1. Plant Mol. Biol. 9:27-37.

Spaink, H. P., Sheeley, D. M., van Brussel, A. A. N., Glushka, J., York, W. S., Tak, T., Geiger, O., Kennedy, E. P., Reinhold, V. N., and Lugtenberg, B. J. J. 1991. A novel highly unsaturated fatty acid moiety of lipo-oligosaccharide signals determines host specificity of *Rhizobium*. Nature (London) 354:125-130.

Spaink, H. P., Wijffelman, C. A., Okker, R. J. H., and Lugtenberg, B. J. J. 1989. Localization of functional regions of the *Rhizobium nodD* product using hybrid *nodD* genes. Plant Mol. Biol. 12:59-73.

Spaink, H. P., Wijffelman, C. A., Pees, E., Okker, R. J. H., and Lugtenberg, B. J. J. 1987. *Rhizobium* nodulation gene *nodD* as a determinant of host specificity. Nature (London) 328:337-340.

van Brussel, A. A. N., Bakhuizen, R., van Spronsen, P. C., Spaink, H. P., Tak, T., Lugtenberg, B. J. J., and Kijne, J. W. 1992. Induction of pre-infection thread structures in the leguminous host plant by mitogenic lipo-oligosaccharides of *Rhizobium*. Science 257:70-72.

van Brussel, A. A. N., Planqué, K., and Quispel, A. 1977. The wall of *Rhizobium leguminosarum* in bacteroid and free-living forms. J. Gen. Microbiol. 101:51-56.

Zaat, S. A. J., Wijffelman, C. A., Spaink, H. P., van Brussel, A. A. N., Okker, R. J. H., and Lugtenberg, B. J. J. 1987. Induction of the *nodA* promoter of *Rhizobium leguminosarum* Sym plasmid pRL1J1 by plant flavanones and flavones. J. Bacteriol. 169:198-204.