

A site-specific recombinase is required for competitive root colonization by *Pseudomonas fluorescens* WCS365

LINDA C. DEKKERS, CLAARTJE C. PHOELICH, LESLIE VAN DER FITS, AND BEN J. J. LUGTENBERG*

Leiden University, Institute of Molecular Plant Sciences, Clusius Laboratory, Wassenaarseweg 64, 2333AL Leiden, The Netherlands

Communicated by R. James Cook, Agricultural Research Center, Pullman, WA, March 18, 1998 (received for review April 1, 1997)

ABSTRACT A colonization mutant of the efficient root-colonizing biocontrol strain *Pseudomonas fluorescens* WCS365 is described that is impaired in competitive root-tip colonization of gnotobiotically grown potato, radish, wheat, and tomato, indicating a broad host range mutation. The colonization of the mutant is also impaired when studied in potting soil, suggesting that the defective gene also plays a role under more natural conditions. A DNA fragment that is able to complement the mutation for colonization revealed a multicistronic transcription unit composed of at least six ORFs with similarity to *lppL*, *lysA*, *dapF*, *orf235/233*, *xerC/sss*, and the largely incomplete *orf238*. The transposon insertion in PCL1233 appeared to be present in the *orf235/233* homologue, designated *orf240*. Introduction of a mutation in the *xerC/sss* homologue revealed that the *xerC/sss* gene homologue rather than *orf240* is crucial for colonization. *xerC* in *Escherichia coli* and *sss* in *Pseudomonas aeruginosa* encode proteins that belong to the λ integrase family of site-specific recombinases, which play a role in phase variation caused by DNA rearrangements. The function of the *xerC/sss* homologue in colonization is discussed in terms of genetic rearrangements involved in the generation of different phenotypes, thereby allowing a bacterial population to occupy various habitats. Mutant PCL1233 is assumed to be locked in a phenotype that is not well suited to compete for colonization in the rhizosphere. Thus we show the importance of phase variation in microbe–plant interactions.

The use of microorganisms, including fluorescent *Pseudomonas* spp., to protect plants against soil-borne diseases is an alternative for the use of chemical pesticides. The biocontrol activity of these strains usually results from the production of one or more antifungal factors. The application of fluorescent *Pseudomonas* spp. and other plant-growth-promoting rhizobacteria is hampered by inconsistency of performance in the field (1, 2). Although the mechanisms underlying biocontrol are complex and diverse, the need to bring the plant-growth-promoting rhizobacteria cells and their antifungal factors to the right sites at the right time is universal. The importance of this process, designated as root colonization, is underscored in two studies. Schippers *et al.* (1) showed that inadequate colonization leads to decreased biocontrol activity, and Bull *et al.* (3) reported an inverse relation between the number of bacteria present on the wheat root and the number of take-all lesions seen on the plant. For these and other reasons, root colonization is often considered the limiting factor for biocontrol in the rhizosphere (1, 2).

Two approaches were used in our laboratory to identify traits involved in root colonization. The first approach is to guess which traits are involved in colonization, isolate mutants in these traits, and then test these mutants for colonization in

competition with the parental strain. With this approach, motility (4) and synthesis of the O-antigen of lipopolysaccharide (LPS) (5) were shown to be essential for colonization. Moreover, mutants auxotrophic for amino acids or vitamin B₁ (6) and mutants with a slightly increased generation time (7) also appeared to be reduced in colonization. Our second approach involves random transposon mutagenesis, using *Tn5lacZ* (8), of the efficient root colonizer *Pseudomonas fluorescens* WCS365 (9). Individual mutants were tested for their colonization ability in competition with the parental strain in a gnotobiotic system as described by Simons *et al.* (7). This approach enables us to obtain knowledge of bacterial colonization traits. In this paper, we describe such a mutant strain, PCL1233, and the results indicate the involvement of a site-specific recombinase in root colonization.

MATERIALS AND METHODS

Bacterial Strains and Culture Conditions. The characteristics of all bacterial strains and plasmids used in this work are described in Table 1. Wild-type *P. fluorescens* WCS365 cells and derivatives of this strain were grown overnight at 28°C on solidified King B medium (18) or in liquid King B or standard succinate medium (SSM) (19) with vigorous shaking.

Various *Escherichia coli* strains were grown overnight at 37°C in liquid or on solidified Luria–Bertani medium (LBM) (20). When appropriate a final concentration of the following antibiotics was added to these media: nalidixic acid, 15 μ g/ml; tetracycline, 40 μ g/ml; streptomycin, 500 μ g/ml; spectinomycin, 200 μ g/ml; kanamycin 50 μ g/ml; and carbenicillin acid, 100 μ g/ml. To distinguish between the wild-type *P. fluorescens* WCS365 cells and its *Tn5lacZ* derivatives, 5-bromo-4-chloro-3-indolyl β -D-galactoside was added to the medium to 40 μ g/ml.

The ability to produce siderophores such as pyoverdine was tested on solidified LBM supplemented with 2.5 or 5 mM ZnSO₄ (13). Addition of the salt was followed by adjustment of the pH of the medium.

Plasmid Constructions. To isolate and study the DNA fragment from mutant PCL1233, which contains the *Tn5lacZ* (8) insertion, we isolated a flanking region of the transposon by using *SalI* to digest the chromosomal DNA of mutant PCL1233. Subsequently, ligation of this digested chromosomal DNA into pIC20H (17) and transformation to *E. coli* resulted in plasmid pMP5206 (Fig. 1 and Table 1). Plasmid pMP5206 was subsequently used to construct pMP5211 (Fig. 1 and Table 1). After introduction of pMP5211 into *P. fluorescens* WCS365 and selection for a single homologous recombination event, total DNA was isolated and digested with *Bam*HI. Self-ligation and selection for kanamycin resistance (21) resulted in plasmid pMP5209, which contains an 8.5-kb *Eco*RI fragment of *P. fluorescens* WCS365 spanning the *Tn5lacZ* insertion (Table 1).

Abbreviations: LPS, lipopolysaccharide; IHF, integration host factor. Data deposition: The sequence reported in this paper has been deposited in the GenBank database (accession no. Y12268).

*To whom reprint requests should be addressed. e-mail: Lugtenberg@ruibim.leidenuniv.nl.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1998 by The National Academy of Sciences 0027-8424/98/957051-6\$2.00/0
PNAS is available online at <http://www.pnas.org>.

Table 1. Bacterial strains and plasmids used in this study

Strains and plasmids	Characteristic(s)
<i>P. fluorescens</i>	
WCS365	Biocontrol strain in a <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> -tomato system (L.C.D., unpublished results), causes systemic acquired resistance in <i>Arabidopsis thaliana</i> ecotype <i>Columbia</i> (H. Gerrits, personal communication, efficient colonizer of the potato root (9)
PCL1500	Tn5 <i>lacZ</i> (8) derivative of WCS365 that behaves as wild type in colonization assays (10)
PCL1233	A Tn5 <i>lacZ</i> derivative of WCS365 that was isolated as a colonization-impaired mutant (11)
PCL1224	PCL1233 harboring plasmid pWTT2081*
PCL1228	PCL1233 harboring pMP5213, used for homogenotization*
PCL1231	Wild-type WCS365 obtained by homogenotization*
PCL1234	PCL1233 harboring pMP5215, used for complementation of the mutation*
PCL1239	WCS365 harboring pWTT2081*
PCL1250	PCL1233 harboring pMP5244*
<i>P. aeruginosa</i>	
7NSK ₂	Plant-growth-promoting <i>P. aeruginosa</i> strain (12)
SSS	Derivative of <i>P. aeruginosa</i> 7NSK ₂ mutated in the <i>sss</i> gene, a homologue of <i>xerC</i> (13, 14).
<i>E. coli</i>	
XL1-Blue	<i>SupE44 hsdR17 recA1 endA1 hlyA46 thi relA1 lac-F'</i> [<i>proAB</i> + <i>lacI^q lacZ</i> <i>M15</i> Tn10(<i>ter</i> ⁻)], used for transformation and propagation of pBluescript phagemids (Stratagene)
DH5 α	<i>EndA1 gyrSA96 hrdR17(rK- mK-) supE44 recA1</i> , used for propagation of plasmids (15)
Plasmids	
pME3049	Suicide plasmid, used for picking up wild-type genes (16)
pIC20H/R	Used for cloning and subcloning of fragments (17)
pMP2740	Plasmid used for homogenotization (17)
pWTT2081	Plasmid stably maintained in the rhizosphere; used for complementation of PCL1233 (10)
PBluescript	Vector used for single-stranded DNA sequencing (10)
pMP5206	pIC20H containing a <i>SalI</i> fragment of the chromosomal region of PCL1233 flanked by Tn5 <i>lacZ</i> (Fig. 1)*
pMP5211	pME3049, which contains a 1.8-kb <i>EcoRI</i> fragment from pMP5206 used to isolate a larger DNA fragment from WCS365 by homologous recombination* (Fig. 1)
pMP5209	pME3049 containing an 8.5-kb fragment of WCS365, obtained by homologous recombination and spanning the Tn5 <i>lacZ</i> insertion*
pMP5230/31	A 5-kb <i>HindIII</i> fragment derived from pMP5209 cloned into pBluescript used for single-stranded sequencing* (Fig. 1)
pMP5213	The 8.5-kb <i>EcoRI</i> fragment of pMP5209 in pMP2740, used for homogenotization*
pMP5215	Complementing 5-kb <i>HindIII</i> fragment, derived from pMP5209 and cloned into pWTT2081*
pMP5244	pWTT2081 containing the 5-kb fragment of pMP5215 with a small insertion in <i>sss</i> * (Fig. 1)

*This paper.

Various subclones of pMP5209 were generated (Fig. 1 and Table 1). Plasmid pMP5215 was used for complementation studies. Plasmid pMP5244 was constructed after digestion of plasmid pMP5215 with *Bgl*III (Fig. 1 and Table 1). To create a blunt end, the *Bgl*III site present in the *xerC/sss* homologue was subsequently filled in by using the Klenow fragment of DNA polymerase I, and the blunt ends were religated.

Plasmid pMP5213, harboring the total 8.5-kb *EcoRI* fragment isolated from WCS365 (Table 1), was introduced into mutant strain PCL1233. A selection was made for a double homologous recombination, which results in white kanamycin-sensitive colonies. One of these was designated PCL1231.

Triparental mating and electroporation were used to transfer the various plasmids to *P. fluorescens* WCS365. Electroporations were performed according to the manufacturer's guidelines using a Bio-Rad Gene Pulser.

DNA Modifications. Standard molecular techniques were performed as described (20). Single-strand sequencing of pMP5230/31, which contain a subclone of 5 kb of the originally isolated fragment, was performed by using the dideoxynucleotide chain-termination method (23). Computer analysis of the obtained sequence results was performed with the GCG Wisconsin software. The site of the Tn5*lacZ* insertion was determined by sequencing pMP5206 with a specific Tn5 primer constructed for the ends of the Tn5*lacZ*. To determine whether the integration host factor (IHF) was actually binding to the putative IHF binding site, an IHF binding study was performed (24).

Methods Used for Isolation and Characterization of Mutant PCL1233. The method of transposon mutagenesis used to obtain colonization-impaired mutants has been described (7). Individual mutants were screened by inoculation of potato stem cuttings with a 1:1 (OD₆₂₀ = 0.1) mixture of cells of the parental strain and one mutant in the compact gnotobiotic system (6, 7, 21). To determine the numbers of mutant and wild type cells on the root tip in a competitive root colonization, a time course study was performed by sampling and estimating the numbers of each strain on 10 plants approximately every other day starting at day 5. Competitive root tip




FIG. 1. Restriction map of the DNA fragment from *P. fluorescens* WCS365, which includes the 5-kb *HindIII* fragment that complements the colonization defect. (A) The 5-kb *HindIII* fragment in which arrows indicate the direction and the size of the six ORFs. (B) The fragments used for complementation or single strand-DNA sequencing. B, *BamHI*; C, *ClaI*; E, *EcoRI*; Ev, *EcoRV*; H3 *HindIII*; K, *KpnI*; S, *SmaI*; Sa, *SalI*.

colonization on tomato, radish, and wheat was determined in the gnotobiotic system described by Simons *et al.* (7), referred to herein as the standard system.

To determine colonization abilities in unsterile potting soil (Jongking Grond B.V., Aalsmeer, The Netherlands), germinated tomato seeds were bacterized with a 1:1 mixture ($OD_{620} = 0.1$) of mutant and parental cells and contained in plastic pots. The plants were grown for 10 days in a climate-controlled growth chamber. Root samples were taken, the adhering soil and bacteria were removed by shaking, and soil suspensions were plated on King B plates supplemented with nalidixic acid, cycloheximide (100 $\mu\text{g/ml}$), and 5-bromo-4-chloro-3-indolyl β -D-galactoside (80 $\mu\text{g/ml}$).

To determine whether mutant PCL1233 is actually a derivative of *P. fluorescens* WCS365 and whether differences exist in the ability of the mutant and the wild type to oxidize 95 different C-sources, the Biolog system (22) was used according to the manufacturer's guidelines. Isolation of cell envelopes and visualization of cell-envelope protein patterns was performed according to Lugtenberg *et al.* (25). Analysis of LPS ladder patterns was performed according to de Weger *et al.* (26). To measure motility, semisolid (0.35% agar) agar plates containing King B medium diluted 1:20 were used (4). Growth in competition in SSM was carried out as described by Dekkers *et al.* (27).

RESULTS AND DISCUSSION

Mutant Strain PCL1233 Is a Colonization-Impaired Mutant. After screening on potato roots and preliminary identification as a competitive colonization mutant (11, 28), mutant strain PCL1233 was selected and tested for previously described colonization traits. The parent strain *P. fluorescens* WCS365 and the mutant strain PCL1233 were indistinguishable with respect to motility, the production of the O-antigen of LPS, amino acids, and vitamin B₁. The parent and mutant were also indistinguishable with respect to growth rate in SSM

Table 2. Populations of mutant strain PCL1233 with the wild-type (WCS365) on root tips with different plant species and rooting media

Colonization conditions	Mean $\log_{10}(\text{CFU} + 1)/$ cm of root tip	
	WCS365	PCL1233
Gnotobiotic compact sand column in competition on potato		
Exp. 1	5.5 ^a	0.3 ^b
Exp. 2	5.4 ^a	1.4 ^b
Strains alone in the gnotobiotic compact sand column		
Potato	—	5.6 ^a
	5.7 ^a	—
Tomato	—	4.3 ^a
	4.4 ^a	—
Wheat	—	4.1 ^a
	5.0 ^a	—
Standard gnotobiotic sand column in competition		
Potato	3.6 ^a	1.1 ^b
Tomato	4.2 ^a	1.6 ^b
Wheat	4.7 ^a	3.8 ^b
Radish	4.3 ^a	2.9 ^b
Potting soil in competition		
Tomato	4.1 ^a	1.4 ^b

Inoculation was carried out on sterile stem cuttings of potato and germinated seeds of radish, tomato, and wheat (7). In every experiment, 10 plants were inoculated. When values from the same experiment are followed by a different letter, they are significantly different at $P = 0.05$ according to the Wilcoxon Mann-Whitney test (29).

Table 3. Populations of wild-type (WCS365 or PCL1239), complemented mutant (PCL1234), and mutant complemented with a deleted *xerC* (PCL1250), on potato root tips 14 days after inoculation of potato stem cuttings with a 1:1 mixture

Colonization test	Mean $\log_{10}(\text{CFU} + 1)/$ cm of root tip	
	Wild type	Various <i>xerC</i> derivatives
WCS365 vs. PCL1233	4.6 ^a	2.9 ^b
	6.1 ^a	3.1 ^b
PCL1239 vs. PCL1234	4.6 ^a	4.6 ^a
	5.1 ^a	5.5 ^a
PCL1239 vs. PCL1250	5.6 ^a	3.8 ^b
	5.0 ^a	3.4 ^b
7NSK2 vs. SSS	5.3 ^a	1.6 ^b
	5.6 ^a	4.5 ^b

Statistical analysis was performed as described in Table 2.

in competition, in their cell envelope proteins, and in the oxidation of 95 different carbon sources in Biolog plates (22). We conclude that mutant PCL1233 does not differ from the wild-type strain *P. fluorescens* WCS365 in any of the established colonization traits. Mutant PCL1233 was therefore selected for further molecular analysis.

Root Colonization Characteristics of Mutant PCL1233.

Root colonization experiments in which cells of mutant PCL1233 and parental strain *P. fluorescens* WCS365 were coinoculated on sterile potato plantlets showed consistent and statistically significant, at least 50-fold, less colonized root tips for mutant PCL1233 compared with its wild-type parent (Tables 2 and 3). However, inoculation of mutant PCL1233 alone on potato, tomato, and wheat resulted in bacterial numbers at the root tips comparable to those measured for the wild type (Table 2), indicating that mutant strain PCL1233 is not colonization-defective because of supersensitivity to compounds exuded by the roots.

The results in competition with the parent (Table 2) show that the mutation also exhibits an inability to colonize roots of tomato, radish, and wheat, indicating that the colonization trait impaired in mutant PCL1233 has a broad host range. Also in potting soil, mutant PCL1233 appeared to be defective in tomato root tip colonization when coinoculated with the parental strain (Table 2). Similar results in potting soil were described by Glandorf (11) for the colonization of the lower parts of potato roots. These results in soil indicate that the colonization trait impaired in mutant PCL1233 is likely to also play a role under more natural conditions. A time course experiment in which PCL1233 was coinoculated 1:1 with the parental strain on the potato root showed that the wild-type numbers isolated from the root tip remained constant or decreased slowly from 6.7 $\log_{10}(\text{CFU} + 1)/\text{cm}$ of root (where CFU is colony-forming units) at day 5 to 6.4 at day 14. The numbers of mutant cells at the growing root tip decreased in time: values of 5.9, 5.4, 5.1, 5.1, and 3.8 $\log_{10}(\text{CFU} + 1)/\text{cm}$ of root were found on days 5, 7, 10, 12, and 14, respectively.

Identification of the DNA Fragment with the Colonization Gene.

Two approaches were followed to test whether the *Tn5lacZ* insertion is responsible for the colonization defect. (i) Strain PCL1234 was tested in competition with PCL1239, to rule out the possibility of a genetic and/or physiological burden for the complemented mutant in the rhizosphere (10). Introduction of pMP5215 in strain PCL1233 caused complementation of the colonization defect in PCL1233 (Table 3). (ii) The wild-type fragment present in pMP5213 was exchanged for the mutated fragment present in mutant PCL1233. After selection, strain PCL1231, a white derivative of PCL1233, was isolated that behaved indistinguishably from *P. fluorescens* WCS365 in a competitive colonization assay (results not

shown), indicating that the colonization defect of PCL1233 is linked to the Tn5*lacZ* insertion.

Homologues of ORF1–6. Sequence analysis of 4,525 bp of pMP5230/31 revealed the presence of six ORFs, designated ORF1–6, of which ORF6 is incomplete (Fig. 1A). The average G+C content of the predicted ORFs is 63.7%, which is consistent with values found in other *Pseudomonas* species (30). In view of the close proximity of the six ORFs and the fact that no putative promoter sequences were found in the intergenic regions between the six ORFs, they probably form one multicistronic transcriptional unit. All ORFs, except ORF5, were preceded by a putative Shine–Dalgarno sequence that is correctly spaced from the various start codons. Lack of a Shine–Dalgarno sequence in front of ORF5 suggests that ORF4 and ORF5 are translationally coupled.

The first gene in the operon (Fig. 1A) shows an ORF of 171 bp; the deduced amino acid sequence indicates a 57-amino acid protein with a calculated molecular weight of 6,073. The predicted product shows 53% identity and 66.7% similarity at the amino acid level to LppL, a lipopeptide described for *Pseudomonas aeruginosa* (31).

ORF2 (Fig. 1A; 1,248 bp) could encode a protein of 416 amino acids with a calculated molecular weight of 45,222. The putative amino acid sequence shows homology with LysA from various bacteria (32–35). The best identity is found with LysA of *P. aeruginosa* (32), of which 84% of the amino acids are identical and 92.5% are similar. *lysA* encodes *meso*-diaminopimelate decarboxylase, the last enzyme in L-lysine biosynthesis.

ORF3 (Fig. 1A; 828 bp) could encode a protein of 276 amino acids with a calculated molecular weight of 29,920 and substantial amino acid sequence similarity (71.5%) and identity (54%) with diaminopimelate epimerase or DapF of *E. coli* K12 (36). This enzyme is involved in the formation of *meso*-diaminopimelate, an essential component of peptidoglycan in rod-shaped bacteria and the direct precursor of L-lysine. Part of this *dapF* gene was also found in the biocontrol strain *P. aeruginosa* 7NSK2 (13). Sequence homology of ORF3 with the *dapF* is even more striking, with 80% identity and 90% similarity over 108 amino acids.

The site of the Tn5*lacZ* insertion was shown to be in ORF4 (Fig. 1A; 720 bp), which encodes a putative protein of 240 amino acids, designated Orf240, with a molecular weight of 26,820. The Tn5*lacZ* insertion is located 150 bp downstream of the *orf240* start codon. The function of this ORF, designated *orf235* in *E. coli* (14) (29% identity and 53% similarity at the amino acid level) and *orf233* in *P. aeruginosa* 7NSK2 (13) (67% identity and 81.5% similarity at the amino acid level), is unknown.

ORF5 contains 897 bp and encodes a putative protein of 299 amino acids with a calculated molecular weight of 33,772. The putative protein sequence of this ORF, designated Sss, shows a high similarity (Fig. 2) to the Sss protein of *P. aeruginosa* (71% identity and 83% similarity) (13) and the XerC protein of *E. coli* (48% identity and 68% similarity) (14). Both XerC and Sss are members of the λ integrase family of site-specific recombinases. Other well known but slightly less homologous members of this family are XerD, formerly known as XrpB (38), FimE, and FimB (37, 39) (see Fig. 2), and λ integrase (40).

Because ORF6 is not necessary for complementation, it was only partially sequenced. The deduced amino acid sequence of this ORF shows homology with Orf238 of *E. coli*, which, as in *P. fluorescens* WCS365, is also located downstream of *xerC* (14). The deduced amino acid sequence of the first 66 amino acids of ORF6 shows significant similarity (44%) and identity (34%) to Orf238, the function of which in *E. coli* is not known (14). A striking resemblance is observed for *E. coli*, *P. aeruginosa* 7NSK2, and *P. fluorescens* WCS365 in that the genetic organization of the *dapF*, *orf240/235/233* genes, and genes

	241			280
FimB	IHPHMLRHS	CGFALANMGI	DTRLIQDYLG	HRNIRHTVWY
FimE	THPHMLRHA	CGYELAERGA	DTRLIQDYLG	HRNIRHTVRY
SssPa	HLHPHMLRHS	FASHLLESSG	DLRAVQELLG	HADIATTQIY
Sssps	NLHPHMLRHS	FASHLLESSQ	DLRAVQELLG	HSDIKTTQIY
XerC	HVHPHKLRLHS	FATHMLESSG	DLRGVQELLG	HANLSTTQIY
XerD	KLSPHVLRLHA	FATHLLNHGA	DLRVQMLLG	HSDLSTTQIY
	*****	**_****_	***_*_*_*_*	*_*_*_*_*_*_*
	281		310	
FimB	TASNAGRFYG	IWDRARGRQR	HAVAL	
FimE	TASNAARFAG	LWERNNLINE	KLKREEV	
SssPa	THLDFQHLAS	VYDAPIPRAK	QGQRDGGNDP	
Sssps	THLDFQHLAT	VYDSAHPRAK	RIKGDSE	
XerC	THLDFQHLAS	VYDAAHPRAK	RGK	
XerD	THVATERLRQ	LHQQHHPRA		
	**	*****	*****	

Fig. 2. Comparison of the C-terminal part of the deduced amino acid sequence of the Sss protein of *P. fluorescens* WCS365 (boldface type) with that of other site-specific recombinases. Amino acids found in more than four sequences are marked by an asterisk. The conserved tyrosine residue is boxed. The protein is compared with FimB and FimE (37, 39) of *E. coli*, Sss of *P. aeruginosa* 7NSK2 (SssPa) (13), and XerC (14) and XerD (38) of *E. coli*.

located further downstream is identical for all three strains, although the genetic information preceding these four ORFs is different.

Regulation of the Promoter. A putative σ^{70} promoter sequence (–35 region TAGGCA 17 bp –10 region TATACT) was found in front of ORF1. A putative IHF binding site (41) was present between the σ^{70} binding site and the ATG start codon of the first ORF. However, when a DNA fragment containing this IHF binding site was incubated with purified *E. coli* IHF protein, no IHF was bound to the DNA (results not shown).

A 300-bp region upstream of the ATG start codon of the *lppL* gene of *P. aeruginosa* is 71% homologous at the DNA level to the region upstream of the *lppL* homologue in strain WCS365. Even the putative –10 and –35 sequences, including the spacing of –10 and –35, are completely homologous, indicating strong conservation of this particular promoter region. This latter observation suggests an important role for this upstream region, perhaps in binding of certain regulatory elements. In *P. aeruginosa*, *lppL* is also located upstream of the *lysA* gene. A location in the outer membrane is suggested for LppL but its function is still unknown (31). A role in the regulation of *lysA* mediated by an attenuation-type system of regulation was proposed by Jann *et al.* (31); this could explain the conservation of the promoter region upstream of the *lppL/lysA* genes in both *Pseudomonas* species.

Three global regulators have been described to affect switching induced by *fimB* and *fimE*, namely IHF, the histone-like protein H-NS, and the leucine-responsive element (Lrp) (37, 42–46). Lrp differentially regulates switching in various media and is involved in the stimulation of switching upon addition of aliphatic amino acids (47). Furthermore, the isolation of fimbriate *E. coli* is favored under certain growth conditions (ref. 46 and references therein). Gally *et al.* (47) showed further that the *fim* switch is differently regulated by environmental conditions such as temperature and medium (47). For XerC, the presence of L-arginine functioning as a corepressor of ArgR positively influences site-specific recombination at *cer* (48). As a result of these findings and the presence of the *lysA* and *dapF* genes upstream of the *sss* homologue in *P. fluorescens* WCS365, we tested a possible regulatory role for the amino acid L-lysine in the regulation of switching induced by Sss. However, L-lysine added at 1 mM to the gnotobiotic system had no effect on the competitive colonization behavior of wild type and mutant (results not shown). If L-lysine suppressed transcription of the operon, one should expect the parent to behave as the mutant.

The ORF5 Site-Specific Recombinase Functions in Competitive Colonization. Because the mutation is located in *orf240* and pMP5215 complements for colonization, *orf240* and the *xerC/sss* homologue are the only candidates that can be responsible for the colonization defect of mutant PCL1233. A small insertion was introduced in the *xerC/sss* gene present on the complementing plasmid (pMP5244). This mutation is located downstream of the actual transposon insertion in mutant PCL1233. Transfer of plasmid pMP5244 to PCL1233 resulted in strain PCL1250. In contrast with PCL1234, PCL1250 is not complemented for root tip colonization (Table 3). Therefore, *xerC/sss* is crucial for colonization, whereas a role of *orf240* cannot be excluded.

The *sss* mutant of *P. aeruginosa* 7NSK2 was reported to have a prolonged lag phase (13) and inoculation of this mutant on maize led to a 4-fold reduction in colonization in a system quite different from ours (13, 49). In contrast, we observed a normal logarithmic-growth phase for mutant PCL1233. When the colonizing ability of the *sss* mutant of *P. aeruginosa* 7NSK2 was tested in our gnotobiotic system, it appeared to be at least 10- to 1,000-fold impaired in its ability to colonize potato root tips after inoculation with a 1:1 mixture of mutant and parental cells (Table 3). These results confirm independently the role of Sss in colonization. Furthermore, *sss* mutants were described not to be able to produce the siderophore pyoverdine when grown in LBM supplemented with Zn^{2+} (13). Incubation of *P. fluorescens* WCS365 and mutant PCL1233 on LBM supplemented with Zn^{2+} showed that neither *P. fluorescens* WCS365 nor mutant PCL1233 produces a siderophore under these circumstances, whereas 7NSK2 does. Höfte *et al.* (49) attributed the observed slightly decreased root colonization capacity for the *sss* mutant of 7NSK2 to an altered ability of the mutant to induce pyoverdine and related outer membrane proteins (49). In contrast to their results, (i) the effect of an *sss* mutation on colonization is much larger in our more sensitive root tip assay and (ii) we did not find the siderophore effect described by Höfte *et al.* (12).

Members of the λ integrase family of site-specific recombinases such as XerC and XerD have been described to have various functions in *E. coli*. XerC and XerD act cooperatively in the process of monomerization of ColE1 plasmids and of the circular chromosome at, respectively, *cer* and *dif* sites (38). *xerC* mutants of *E. coli* are reported to have a slow recovery from stationary phase and to show a variable growth rate in the exponential phase when compared with a *xerC*⁺ strain (50). This is clearly not the case for mutant PCL1233, which also does not show cell division defects (results not shown).

The observed homologies indicate that the *xerC/sss* homologue of *P. fluorescens* WCS365 is a member of the λ integrase family of site-specific recombinases. These enzymes promote conservative reciprocal recombination (which does not require DNA synthesis) between two small (approximately 15 bp) homologous DNA sequences. The presence and orientation of two of these sequences will lead to inversion or excision of the DNA fragment situated between these small recognition sites (51). Comparison of all site-specific recombinases shows that homology is particularly striking in the C-terminal part (Fig. 2), in which a perfectly conserved tyrosine residue is present and thought to form a linkage with the DNA (40).

Most DNA rearrangements involved in phenotypic switching regulate expression of phase-variable cell surface antigens (52) such as fimbriae, flagella, LPS, and lipoprotein. FimE and FimB play a role in phase variation of type 1 fimbriae in *E. coli*. Both proteins are thought to be site-specific recombinases (37) acting on a specific 314-bp DNA element located in the promoter region of *fimA*, the main fimbrial subunit gene (39, 53). From the homology found with FimB and FimE (Fig. 2), we postulate that the Sss protein is involved in DNA rearrangements that lead to phenotypes with different colonization abilities. DNA rearrangements can also regulate the

production of two forms of LPS related to pathogenicity in the intracellular pathogen *Francisella tularensis* (54) and antigenic variation of surface lipoprotein antigens in *Mycoplasma bovis* (55). Also the regulation of the alternate expression of two flagellin genes in *Salmonella typhimurium* is regulated by an invertible DNA element modulated by recombination (56). The fact that most DNA rearrangements involved in phenotypic switching regulate expression of phase-variable cell surface antigens and our observation that mutant PCL1233 is only impaired in competitive colonization but not when inoculated alone lead us to speculate that PCL1233 is impaired in competition for nutrients or for sites on the root surface. Flagella, crucial for colonization (4), presumably play a role in competition for nutrients through chemotaxis. Moreover, the composition of the LPS may affect motility because O-antigen-negative mutants of *P. fluorescens* WCS365 are often less motile than their parental strain (L.C.D., unpublished results). Fimbriae, LPS, and cell-surface proteins could be involved in competition for sites on the root surface. We favor the hypothesis that a cell-surface-related molecule or trait is subject to phase variation by Sss and that, as a result, competition for nutrients or sites on the root surface is affected. This hypothesis is also consistent with the observation that the mutant is impaired only in competition. The discovery that Sss plays a role in root colonization shows the importance of phase variation in this plant-microbe interaction.

Colony sector formation is often associated with DNA rearrangements. Interestingly, we observed that old colonies of WCS365 contain morphologically distinct sectors. Sectoring has been observed less frequently in mutant strain PCL1233 than in WCS365 (L.C.D., unpublished results). Dybvig (52) postulated that bacteria are able to generate subpopulations in certain environments, by means of DNA rearrangements, that result in different abilities to adjust to sudden environmental changes. By this mechanism, such a bacterial population is at all times able to respond adequately to environmental changes even if only a few cells of a subpopulation are surviving. According to this notion, mutant PCL1233 is locked in a genetic configuration that is less rhizosphere competent when compared with cells of the parental strain.

We thank N. Goosen for supplying us with purified *E. coli* IHF and technical assistance with binding studies, H. Spaik and L. de Weger for useful discussions, and M. Höfte for *P. aeruginosa* 7NSK2 and its mutant. We thank I. Brand for isolation of mutant PCL1233. The investigations were partly funded by the European Union Biotech Program BIO2-CT93.0196. C.C.P. is funded by the Netherlands Organization for Scientific Research-Ministry of Agriculture, Nature Management and Fisheries Priority Program Crop Protection, Project 805.45.008.

1. Schippers, B., Bakker, A. W. & Bakker, P. A. H. M. (1987) *Annu. Rev. Phytopathol.* **25**, 339–358.
2. Weller, D. M. (1988) *Annu. Rev. Phytopathol.* **26**, 379–407.
3. Bull, C. T., Weller, D. M. & Thomashow, L. S. (1991) *Phytopathology* **81**, 954–959.
4. de Weger, L. A., van der Vlugt, C. I. M., Wijffjes, A. H. M., Bakker, P. A. H. M., Schippers, B. & Lugtenberg, B. J. J. (1987b) *J. Bacteriol.* **169**, 2769–2773.
5. de Weger, L. A., Bakker, P. A. H. M., Schippers, B., van Loosdrecht, M. C. M. & Lugtenberg, B. J. J. (1989) in *Signal Molecules in Plants and Plant-Microbe Interactions*, ed. Lugtenberg, B. J. J. (Springer, Berlin), pp. 197–202.
6. Simons, M., Permentier, H. P., de Weger, L. A., Wijffelman, C. A. & Lugtenberg, B. J. J. (1997) *Mol. Plant-Microbe Interact.* **10**, 102–106.
7. Simons, M., van der Bij, A. J., Brand, J., de Weger, L. A., Wijffelman, C. A. & Lugtenberg, B. J. J. (1996) *Mol. Plant-Microbe Interact.* **9**, 600–607.
8. Lam, S. T., Ellis, D. M. & Ligon, J. M. (1990) *Plant and Soil* **129**, 11–18.
9. Geels, F. P. & Schippers, B. (1983) *Phytopathol. Z.* **108**, 193–206.

10. van der Bij, A. J., de Weger, L. A., Tucker, W. T. & Lugtenberg, B. J. J. (1996) *Appl. Environ. Microbiol.* **62**, 1076–1080.
11. Glandorf, D. C. M. (1992) Ph.D. thesis (University of Utrecht, Utrecht, The Netherlands).
12. Höfte, M., Seong, K. Y., Jurkevitch, E. & Verstraete, H. (1991) *Plant and Soil* **130**, 249–257.
13. Höfte, M., Dong, Q., Kourambas, S., Krishnapillai, V., Sherratt, D. & Mergeay, M. (1994) *Mol. Microbiol.* **14**, 1011–1020.
14. Colloms, S. D., Sykora, P., Szatmari, G. & Sherratt, D. J. (1990) *J. Bacteriol.* **172**, 6973–6980.
15. Boyer, H. W. & Roulland-Dussoix, D. (1969) *J. Mol. Biol.* **41**, 459–472.
16. Voisard, C., Bull, C. T., Keel, C., Laville, J., Maurhofer, M., Schnider, U., Défago, G. & Haas, D. (1994) in *Molecular Ecology of Rhizosphere Microorganisms*, eds. O’Gara, F., Dowling, D. N. & Boesten, B. (VCH, New York), pp. 67–89.
17. Marsh, J. L., Erfle, M. & Wykes, E. J. (1984) *Gene* **32**, 481–485.
18. King, E. O., Ward, M. K. & Raney, D. E. (1954) *J. Lab. Clin. Med.* **44**, 301–307.
19. Meyer, J. M. & Abdallah, M. A. (1978) *J. Gen. Microbiol.* **107**, 319–328.
20. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) in *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY), 2nd Ed.
21. de Weger, L. A., Dekkers, L. C., van der Bij, A. J. & Lugtenberg, B. J. J. (1994) *Mol. Plant-Microbe Interact.* **7**, 32–38.
22. Bochner, B. R. (1989) *Nature (London)* **339**, 157–158.
23. Sanger, F., Nicklen, S. & Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 5463–5467.
24. Goosen, N., van Ulsen, P., Zulianello, L. & van de Putte, P. (1996) *Methods Enzymol.* **274**, 232–242.
25. Lugtenberg, B. J. J., Meyers, J., Peters, R., van der Hoek, P. & van Alphen, L. (1975) *FEBS Lett.* **58**, 254–258.
26. de Weger, L. A., Jann, B., Jann, K. & Lugtenberg, B. (1987) *J. Bacteriol.* **169**, 1441–1446.
27. Dekkers, L. C., Bloemendaal, C. P., de Weger, L. A., Wijffelman, C. A., Spaink, H. P., Lugtenberg, B. J. J. (1998) *Mol. Plant-Microbe Interact.* **11**, 45–56.
28. Lugtenberg, B. J. J. & de Weger, L. A. (1992) in *Pseudomonas: Molecular Biology and Biotechnology*, eds. Galli, E., Silver, S. & Witholt, B. (Am. Soc. Microbiol., Washington, DC), pp. 13–19.
29. Sokal, R. R. & Rohlf, F. J. (1981) *Biometry* (Freeman, San Francisco).
30. Palleroni, N. J. (1975) in *General Properties and Taxonomy of the Genus Pseudomonas*, eds. Clarke, P. H. & Richmond, M. A. (Wiley, London), pp. 1–36.
31. Jann, A., Cavard, D., Martin, C., Cami, B. & Patte, J.-C. (1990) *Mol. Microbiol.* **4**, 667–682.
32. Martin, C., Cami, B., Yeh, P., Stragier, P., Parsot, C. & Patte, J. (1988) *Mol. Biol. Evol.* **5**, 549–559.
33. Marcel, T., Archer, J. A. C., Mengin-Lecreux, D. & Sinskey, A. J. (1990) *Mol. Microbiol.* **4**, 1819–1830.
34. Yamamoto, J., Shimizu, M. & Yamane, K. (1991) *Agric. Biol. Chem.* **55**, 1615–1626.
35. Mills, D. A. & Flickinger, M. C. (1993) *Appl. Environ. Microbiol.* **59**, 2927–2937.
36. Richaud, C. & Printz, C. (1988) *Nucleic Acids Res.* **16**, 10367.
37. Dorman, C. J. & Higgins, C. F. (1987) *J. Bacteriol.* **169**, 3840–3843.
38. Blakely, G., May, G., McCulloch, R., Arciszewska, L. K., Burke, M., Lovett, S. T. & Sherratt, D. (1993) *Cell* **75**, 351–361.
39. Klemm, P. (1986) *EMBO J.* **5**, 1389–1393.
40. Argos, P., Landy, A., Abremski, K., Egan, J. B., Haggard-Ljungquist, E., Hoess, R. H., Kahn, M. L., Kalionis, B., Narayana, S. V. L., Pierson, L. S., III *et al.* (1986) *EMBO J.* **5**, 433–440.
41. Goosen, N. & van de Putte, P. (1995) *Mol. Microbiol.* **16**, 1–7.
42. Higgins, C. F., Dorman, C. J., Stirling, D. A., Waddell, L., Booth, I. R., May, G. & Bremer, E. (1988) *Cell* **52**, 569–584.
43. Eisenstein, B. I., Sweet, D. S., Vaughn, V. & Friedman, D. I. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 6506–6510.
44. Blomfield, I. C., Calie, P. J., Eberhardt, K. J., McClain, M. S. & Eisenstein, B. I. (1993) *J. Bacteriol.* **175**, 27–36.
45. Kawula, T. H. & Orndorff, P. E. (1991) *J. Bacteriol.* **173**, 4116–4123.
46. Spears, P. A., Schauer, D. & Orndorff, P. E. (1986) *J. Bacteriol.* **168**, 179–185.
47. Gally, D. L., Bogan, J. A., Eisenstein, B. I. & Blomfield, I. C. (1993) *J. Bacteriol.* **175**, 6186–6193.
48. Stirling, C. J., Szatmari, G., Stewart, G., Smith, M. C. M. & Sherratt, D. J. (1988) *EMBO J.* **7**, 4389–4395.
49. Höfte, M., Boelens, J. & Verstraete, W. (1992) *J. Plant Nutr.* **15**, 2253–2262.
50. Blakely, G., Colloms, S., May, G., Burke, M. & Sherratt, D. (1991) *New Biol.* **3**, 789–798.
51. Sadowski, P. (1986) *J. Bacteriol.* **165**, 341–347.
52. Dybvig, K. (1993) *Mol. Microbiol.* **10**, 465–471.
53. Abraham, J. M., Freitag, C. S., Clements, J. R. & Eisenstein, B. I. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 5724–5727.
54. Cowley, S. C., Myltseva, S. V. & Nano, F. E. (1996) *Mol. Microbiol.* **20**, 867–874.
55. Lysnyansky, I., Rosengarten, R. & Yogev, D. (1996) *J. Bacteriol.* **178**, 5395–5401.
56. Zeig, J., Silverman, H., Hilmen, H. & Simon, M. (1977) *Science* **196**, 170–175.