Time-delay matrix, midgap spectral peak, and thermopower of an Andreev billiard

M. Marciani,¹ P. W. Brouwer,² and C. W. J. Beenakker¹

¹*Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands*

²*Dahlem Center for Complex Quantum Systems and Fachbereich Physik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany*

(Received 13 May 2014; revised manuscript received 16 June 2014; published 9 July 2014)

We derive the statistics of the time-delay matrix (energy derivative of the scattering matrix) in an ensemble of superconducting quantum dots with chaotic scattering (Andreev billiards), coupled ballistically to M conducting modes (electron-hole modes in a normal metal or Majorana edge modes in a superconductor). As a first application we calculate the density of states ρ_0 at the Fermi level. The ensemble average $\langle \rho_0 \rangle = \delta_0^{-1} M [\max(0, M + 2\alpha/\beta)]^{-1}$ deviates from the bulk value $1/\delta_0$ by an amount depending on the Altland-Zirnbauer symmetry indices α, β . The divergent average for $M = 1, 2$ in symmetry class D ($\alpha = -1, \beta = 1$) originates from the midgap spectral peak of a closed quantum dot, but now no longer depends on the presence or absence of a Majorana zero mode. As a second application we calculate the probability distribution of the thermopower, contrasting the difference for paired and unpaired Majorana edge modes.

DOI: [10.1103/PhysRevB.90.045403](https://doi.org/10.1103/PhysRevB.90.045403)

PACS number(s): 73.23.-b, 74.78.Na, 73.63.-b

I. INTRODUCTION

A semiconductor quantum dot feels the proximity to a superconductor even when a magnetic field has closed the excitation gap that would open in zero magnetic field: The average density of states has either a peak or a dip [1],

$$\rho_{\pm}(E) = \delta_0^{-1} \pm \frac{\sin(2\pi E/\delta_0)}{2\pi E}, \quad (1)$$

see Fig. 1, within a mean level spacing δ_0 from the Fermi level at $E = 0$ (in the middle of the superconducting gap). The appearance of a midgap spectral peak or dip distinguishes the two symmetry classes C (dip, when spin-rotation symmetry is preserved) and D (peak, spin-rotation symmetry is broken by strong spin-orbit coupling). These Altland-Zirnbauer symmetry classes exist because of the $\pm E$ electron-hole symmetry in a superconductor, and are a late addition to the Wigner-Dyson symmetry classes conceived in the 1960s to describe universal properties of nonsuperconducting systems [2].

Electron-hole symmetry in the absence of spin-rotation symmetry allows for a nondegenerate level at $E = 0$, a so-called Majorana zero mode [3,4]. The class-D spectral peak is then converted into a dip, $\rho_+ \rightarrow \rho_- + \delta(E)$, such that the integrated density of states remains the same as without the zero mode [5,6]. The entire spectral weight of this Fermi-level anomaly is $1/2$, consistent with the notion that a Majorana zero mode is a half fermion [7].

Here we study what happens if the quantum dot is coupled to M conducting modes, so that the discrete spectrum of the closed system is broadened into a continuum. We focus on the strong-coupling limit, typically realized by a ballistic point contact, complementing earlier work on the limit of weak coupling by a tunnel barrier or a localized conductor [8–15]. The simplicity of the strong-coupling limit allows for an analytical calculation using random-matrix theory of the entire probability distribution of the Fermi-level density of states—not just the ensemble average. Using the same random-matrix approach we also calculate the probability distribution of the thermopower of the quantum dot, which is nonzero in spite of electron-hole symmetry when the superconductor contains gapless Majorana edge modes [16].

The key technical ingredient that makes these calculations possible is the joint probability distribution of the scattering matrix S and the time-delay matrix $Q = -i\hbar S^\dagger dS/dE$, in the limit $E \rightarrow 0$. This is known for the Wigner-Dyson ensembles [17], and here we extend that to the Altland-Zirnbauer ensembles. The Fermi-level density of states then follows directly from the trace of Q , while the thermopower requires also knowledge of the statistics of S . We find that these probability distributions depend on the symmetry class (C or D), and on the number M of conducting modes, but are the same irrespective of whether the quantum dot contains a Majorana zero mode or not. A previous calculation [14] had found that the density-of-states signature of a Majorana zero mode becomes less evident when the quantum dot is coupled by a tunnel barrier to the continuum. We conclude that ballistic coupling completely removes any trace of the Majorana zero mode in the density of states, as well as in the thermopower—but not, we hasten to add, in the Andreev conductance [18].

The outline of the paper is as follows. In the next section we present the geometry of an “Andreev billiard” [19], a semiconductor quantum dot with Andreev reflection from a superconductor and a point-contact coupling to a metallic conductor. (Systems of this type have been studied experimentally, for example in Refs. [20–22].) We derive a formula relating the thermopower to the scattering matrix S and time-delay matrix Q , in a form which is suitable for a random-matrix approach. The distribution of the transmission eigenvalues T_n of S was already derived in Ref. [23]; what we need additionally is the distribution of the eigenvalues D_n of Q (the delay times), which we present in Sec. III. The distributions of the Fermi-level density of states and thermopower are given in Secs. IV and V, respectively. We conclude in Sec. VI.

II. SCATTERING FORMULA FOR THE THERMOPOWER

We study the thermopower of a quantum dot connecting a two-dimensional topological superconductor and a semiconductor two-dimensional electron gas (see Fig. 2). In equilibrium the normal-metal contact and the superconducting contact have a common temperature T_0 and chemical potential E_F .

FIG. 1. (Color online) Ensemble-averaged density of states (1) of an Andreev billiard in symmetry class C (ρ_- , dashed curve), or class D without a Majorana zero mode (ρ_+ , solid curve). The class-D billiard with a Majorana zero mode has the smooth density of states ρ_- together with the delta-function contribution from the zero mode. In this paper we investigate how the midgap spectral peak or dip evolves when the billiard is opened via a ballistic point contact to a metallic reservoir. We find that the distinction between classes C and D remains, but the signature of the Majorana zero mode is lost.

Application of a temperature difference δT induces a voltage difference V at zero electrical current. The ratio $S = -V/\delta T$ is the thermopower or Seebeck coefficient.

In the low-temperature limit $\delta T \ll T_0 \rightarrow 0$ the thermopower is given by the Cutler-Mott formula [24],

$$S/S_0 = -\lim_{E \rightarrow 0} \frac{1}{G} \frac{dG}{dE}, \quad S_0 = \frac{\pi^2 k_B^2 T_0}{3e}, \quad (2)$$

in terms of the electrical conductance $G(E)$ near the Fermi level ($E = 0$). See Ref. [16] for a demonstration that this relationship, originally derived for normal metals, still holds

FIG. 2. (Color online) Andreev-billiard geometry to measure the thermopower S of a semiconductor quantum dot coupled to chiral Majorana modes at the edge of a topological superconductor. A temperature difference δT induces a voltage difference $V = -S\delta T$ under the condition that no electrical current flows between the contacts. For a random-matrix theory we assume that the Majorana modes are uniformly mixed with the modes in the point contact, by chaotic scattering events in the quantum dot.

when one of the contacts is superconducting and G is the Andreev conductance.

Without gapless Majorana modes in the superconductor the Andreev conductance is an even function of E , so the ratio S/S_0 vanishes in the low-temperature limit. For that reason, with some exceptions [25,26], most studies of the effect of a superconductor on thermoelectric transport take a three-terminal geometry, where the temperature difference is applied between two normal contacts and the conductance is not so constrained [27–36]. As pointed out by Hou, Shtengel, and Refael [16], Majorana edge modes break the $\pm E$ symmetry of the conductance allowing for thermoelectricity in a two-terminal geometry—even if they themselves carry only heat and no charge.

In a random-matrix formulation of the problem two matrices enter, the scattering matrix at the Fermi level $S_0 \equiv S(E = 0)$ and the Wigner-Smith time-delay matrix [37–39]

$$Q = -i\hbar \lim_{E \rightarrow 0} S^\dagger \frac{dS}{dE}. \quad (3)$$

Before proceeding to the random-matrix theory, we first express the thermopower in terms of these two matrices. The existing expressions in the literature [40,41] cannot be directly applied for this purpose, since they do not incorporate Andreev reflection processes.

The Andreev conductance is given by [42]

$$G(E)/G_0 = \frac{1}{2}N - \text{Tr} r_{ee}(E)r_{ee}^\dagger(E) + \text{Tr} r_{he}(E)r_{he}^\dagger(E), \quad (4)$$

in terms of the matrix of reflection amplitudes

$$r = \begin{pmatrix} r_{ee} & r_{eh} \\ r_{he} & r_{hh} \end{pmatrix} \quad (5)$$

for electrons and holes injected via a point contact into the quantum dot. The submatrix r_{ee} describes normal reflection (from electron back to electron), while r_{he} describes Andreev reflection (from electron to hole, induced by the proximity effect of the superconductor that interfaces with the quantum dot). The conductance quantum is $G_0 = e^2/h$ and N is the total number of modes in the point contact (counting spin and electron-hole degrees of freedom), so r has dimension $N \times N$.

Without edge modes in the superconductor, the reflection matrix r would be unitary at energies E below the superconducting gap. In that case one can simplify Eq. (4) as $G/G_0 = 2 \text{Tr} r_{he} r_{he}^\dagger$. Because of the gapless edge modes the more general formula (4) is needed, which does not assume unitarity of r .

Equivalently, Eq. (4) may be written in terms of the full unitary scattering matrix $S(E)$,

$$G(E)/G_0 = \frac{1}{2}N - \frac{1}{2}\text{Tr} \mathcal{P} \tau_z S(E) \mathcal{P} (1 + \tau_z) S^\dagger(E), \quad (6)$$

where the Pauli matrix τ_z acts on the electron-hole degree of freedom and $\mathcal{P}$ projects onto the modes at the point contact:

$$S = \begin{pmatrix} r & t' \\ t & r' \end{pmatrix}, \quad \mathcal{P} \tau_z = \begin{pmatrix} \tau_z & 0 \\ 0 & 0 \end{pmatrix}. \quad (7)$$

The off-diagonal matrix blocks t, t' couple the N' Majorana edge modes to the N electron-hole modes in the point contact, mediated by the quasibound states in the quantum dot. The

incoming and outgoing Majorana edge modes are coupled by the $N' \times N'$ submatrix r' .

Electron-hole symmetry in class D is most easily accounted for by first making a unitary transformation from S to

$$S' = \begin{pmatrix} U & 0 \\ 0 & U \end{pmatrix} S \begin{pmatrix} U^\dagger & 0 \\ 0 & U^\dagger \end{pmatrix}, \quad U = \sqrt{\frac{1}{2}} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix}. \quad (8)$$

In this so-called Majorana basis [43] the electron-hole symmetry relation reads

$$S'(E) = S'^*(-E). \quad (9)$$

The Pauli matrix τ_z transforms into τ_y , so the conductance is given in the Majorana basis by

$$G(E)/G_0 = \frac{1}{2}N - \frac{1}{2}\text{Tr} \mathcal{P} \tau_y S'(E) \mathcal{P} (1 + \tau_y) S'^\dagger(E). \quad (10)$$

In what follows we will omit the prime, for ease of notation.

To first order in E the energy dependence of the scattering matrix is given by

$$S(E) = S_0[1 + iE\hbar^{-1}Q + \mathcal{O}(E^2)]. \quad (11)$$

Unitarity and electron-hole symmetry together require that S_0 is real orthogonal and Q is real symmetric, both in the Majorana basis. The conductance, still to first order in E , then takes the form

$$\begin{aligned} G(E)/G_0 &= \frac{1}{2}N - \frac{1}{2}\text{Tr} \mathcal{P} \tau_y S_0 \mathcal{P} (1 + \tau_y) S_0^\dagger \\ &\quad - \frac{1}{2}iE\hbar^{-1} \text{Tr} \mathcal{P} \tau_y S_0 [Q\mathcal{P}(1 + \tau_y) - \mathcal{P}(1 + \tau_y)Q] S_0^\dagger. \end{aligned} \quad (12)$$

Since $\text{Tr} \mathcal{P} \tau_y X$ vanishes for any symmetric matrix X , we can immediately set some of the traces in Eq. (12) to zero:

$$\begin{aligned} G(E)/G_0 &= \frac{1}{2}N - \frac{1}{2}\text{Tr} \mathcal{P} \tau_y S_0 \mathcal{P} \tau_y S_0^\dagger \\ &\quad - \frac{1}{2}iE\hbar^{-1} \text{Tr} \mathcal{P} \tau_y S_0 (Q\mathcal{P} - \mathcal{P}Q) S_0^\dagger. \end{aligned} \quad (13)$$

The resulting thermopower is

$$S/S_0 = i\hbar^{-1} \frac{\text{Tr} \mathcal{P} \tau_y S_0 (Q\mathcal{P} - \mathcal{P}Q) S_0^\dagger}{N - \text{Tr} \mathcal{P} \tau_y S_0 \mathcal{P} \tau_y S_0^\dagger}, \quad (14)$$

in the Majorana basis. Equivalently, in the electron-hole basis one has

$$S/S_0 = i\hbar^{-1} \frac{\text{Tr} \mathcal{P} \tau_z S_0 (Q\mathcal{P} - \mathcal{P}Q) S_0^\dagger}{N - \text{Tr} \mathcal{P} \tau_z S_0 \mathcal{P} \tau_z S_0^\dagger}. \quad (15)$$

This scattering formula for the thermopower is a convenient starting point for a random-matrix calculation. Notice that the commutator of Q and $\mathcal{P}$ in the numerator ensures a vanishing thermopower in the absence of gapless modes in the superconductor, because then the projector $\mathcal{P}$ is just the identity.

III. DELAY-TIME DISTRIBUTION IN THE ALT LAND-ZIRNBAUER ENSEMBLES

Chaotic scattering in the quantum dot mixes the N' Majorana edge modes with the N electron-hole modes in the point contact. The assumption that the mixing uniformly covers the whole available phase space produces one of the

TABLE I. The two Altland-Zirnbauer symmetry classes that support chiral Majorana edge modes, with d -wave pairing (class C) or p -wave pairing (class D). The “canonical basis” is the basis in which the scattering matrix elements are quaternion (class C) or real (class D). The degeneracies d_T and d_E refer to transmission eigenvalues and energy eigenvalues, respectively. The α and β parameters determine the exponents in the probability distributions (17) and (18) of the transmission eigenvalues and inverse delay times.

Symmetry class	Class C	Class D
pair potential	spin-singlet d wave	spin-triplet p wave
canonical basis	electron-hole	Majorana
S -matrix elements	quaternion	real
S -matrix space	symplectic	orthogonal
circular ensemble	CQE	CRE
d_T	4	1
d_E	2	1
α	2	-1
β	4	1

circular ensembles of random-matrix theory, distinguished by fundamental symmetries that restrict the available phase space [44]. Two Altland-Zirnbauer symmetry classes support chiral Majorana modes at the edge of a two-dimensional superconductor [45–47], corresponding to spin-singlet d -wave pairing (symmetry class C) or spin-triplet p -wave pairing (symmetry class D). Time-reversal symmetry is broken in both; in class C there is electron-hole symmetry as well as spin-rotation symmetry, while in class D only electron-hole symmetry remains. (See Table I.)

The uniformity of the circular ensembles is expressed by the invariance

$$P[S(E)] = P[U \cdot S(E) \cdot U'] \quad (16)$$

of the distribution functional $P[S(E)]$ upon multiplication of the scattering matrix by a pair of energy-independent matrices U, U' , restricted by symmetry to a subset of the full unitary group: In class C they are quaternion symplectic [48] in the electron-hole basis (circular quaternion ensemble, CQE), while in class D they are real orthogonal in the Majorana basis (circular real ensemble, CRE).

The unitary invariance (16) of the Wigner-Dyson scattering matrix ensembles was postulated in Ref. [49] and derived from the corresponding Hamiltonian ensembles in Ref. [50]. We extend the derivation to the Altland-Zirnbauer ensembles in Appendix A 1. The key step in this extension is to ascertain that the class-D unitary invariance applies to U, U' in the full orthogonal group—without any restriction on the sign of the determinant.

For the thermopower statistics we need the joint distribution $P(S_0, Q)$ of Fermi-level scattering matrix and time-delay matrix. The invariance (16) implies $P(S_0, Q) = P(-1, Q)$ (take $U = -S_0^\dagger, U' = 1$), so Q is statistically independent of S_0 and the two matrices can be considered separately [17, 51].

The uniform distribution of S_0 in the symplectic group (CQE, class C) or orthogonal group (CRE, class D) directly gives the probability distribution of the transmission eigenvalues $T_n \in [0, 1]$ of quasiparticles from the normal metal into the superconductor. [These are the quantities that

determine the thermal conductance $\propto \sum_n T_n$, not the electrical conductance (4).] For a transmission matrix of dimension $N' \times N$ there are $N_{\min} = \min(N, N')$ nonzero transmission eigenvalues, fourfold degenerate ($d_T = 4$) in class C and nondegenerate ($d_T = 1$) in class D. The $N_{\min}/d_T$ distinct T_n 's have probability distribution [23]

$$P(\{T_n\}) \propto \prod_k T_k^{\beta|\delta N|/2} T_k^{-1+\beta/2} (1 - T_k)^{\alpha/2} \prod_{i < j} |T_i - T_j|^\beta, \quad (17)$$

with $\delta N = (N - N')/d_T$ and parameters α, β listed in Table I [52].

The Hermitian positive-definite matrix Q has dimension $\mathcal{M} \times \mathcal{M}$ with $\mathcal{M} = N + N'$. Its eigenvalues $D_n > 0$ are the delay times, and $\gamma_n \equiv 1/D_n$ are the corresponding rates. The degeneracy d_T of the D_n 's is the same as that of the T_n 's. The derivation of the distribution $P(\gamma_1, \gamma_2, \dots, \gamma_M)$ of the $M = \mathcal{M}/d_T$ distinct delay rates is given in Appendix A, for all four Altland-Zirnbauer symmetry classes: C, D without time-reversal symmetry and CI, DIII with time-reversal symmetry. The result is

$$P(\{\gamma_n\}) \propto \prod_k \Theta(\gamma_k) \gamma_k^{\alpha+M\beta/2} \exp\left(-\frac{1}{2}\beta t_0 \gamma_k\right) \prod_{i < j} |\gamma_i - \gamma_j|^\beta. \quad (18)$$

The unit step function $\Theta(\gamma)$ ensures that the probability vanishes if any γ_n is negative. The characteristic time t_0 is defined by

$$t_0 = \frac{d_E}{d_T} \frac{2\pi\hbar}{\delta_0}, \quad (19)$$

in terms of the average spacing δ_0 of d_E -fold degenerate energy levels in the isolated quantum dot [53]. For $\alpha = 0$ and $d_E = d_T$ we recover the result of Ref. [17] for the Wigner-Dyson ensembles.

The difference between the Altland-Zirnbauer and Wigner-Dyson ensembles manifests itself in a nonzero value of α and in a difference in the degeneracies d_E and d_T of energy and transmission eigenvalues (see Table I). One has $d_T = d_E$ in the absence of particle-hole symmetry or when the particle-hole conjugation operator $\mathcal{C}$ squares to $+1$; when $\mathcal{C}^2 = -1$ one has $d_T = 2d_E$ [54].

Already at this stage we can conclude that the thermopower distribution in the circular ensemble does not depend on the presence or absence of Majorana zero modes inside the quantum dot, for example, bound to the vortex core in a chiral p -wave superconductor [3,4]. The parity of the number n_M of Majorana zero modes fixes the sign of the determinant of the orthogonal class-D scattering matrix,

$$\text{Det } S_0 = (-1)^{n_M}. \quad (20)$$

The unitary invariance (16) of the CRE implies, on the one hand, that $P(S_0, Q)$ is unchanged under the transformation $S_0 \mapsto U S_0$, $U = \text{diag}(-1, 1, 1, \dots, 1)$, that inverts the sign of $\text{Det } S_0$. [Here we make essential use of the fact that Eq. (16) in class D applies to the full orthogonal group.] On the other hand, the same transformation leaves the thermopower (14) unaffected, provided we assign the first matrix element to a superconducting edge mode (so $\mathcal{P}\tau_y$ commutes with U).

IV. FERMI-LEVEL ANOMALY IN THE DENSITY OF STATES

A. Analytical calculation

A striking difference between the Wigner-Dyson and Altland-Zirnbauer ensembles appears when one considers the density of states at the Fermi level ρ_0 , related to the time-delay matrix by

$$\rho_0 = \frac{1}{2\pi\hbar} \frac{d_T}{d_E} \sum_{n=1}^M D_n. \quad (21)$$

(The factor d_T/d_E is needed because delay times and energy levels may have a different degeneracy. The density of states counts degenerate levels once.) In the Wigner-Dyson ensembles the average density of states equals exactly $1/\delta_0$, independently of the symmetry index β and of the number of channels M that couple the discrete spectrum inside the quantum dot to the continuum outside [17,56].

In the Altland-Zirnbauer ensembles, instead, we find from Eq. (18) that [57]

$$\begin{aligned} \delta_0 \langle \rho_0 \rangle &= \frac{1}{t_0} \left\langle \sum_{n=1}^M D_n \right\rangle = \frac{M}{\max(0, M + 2\alpha/\beta)} \\ &= \begin{cases} M/(M+1) & \text{in class C for any } M \geq 1, \\ M/(M-2) & \text{in class D for } M \geq 3, \\ \infty & \text{in class D for } M = 1, 2. \end{cases} \end{aligned} \quad (22)$$

It is known [1,6,8–15] that the tunneling density of states of a superconducting quantum dot with broken time-reversal symmetry, weakly coupled to the outside, has a Fermi-level anomaly consisting of a narrow dip in symmetry class C and a narrow peak in class D. Equation (22) shows the effect of level broadening upon coupling via M channels to the continuum. For $M \rightarrow \infty$ the normal-state result $1/\delta_0$ is recovered, but for small M the Fermi-level anomaly persists.

For $M = 1, 2$ the average density of states in class D diverges, because of a long tail in the probability distribution of $\kappa \equiv \delta_0 \rho_0$:

$$P_D(\kappa) = \begin{cases} (2\pi)^{-1/2} \kappa^{-3/2} e^{-(2\kappa)^{-1}} & \text{for } M = 1, \\ \kappa^{-3} (2 + \kappa) e^{-2/\kappa} & \text{for } M = 2. \end{cases} \quad (23)$$

See Fig. 3 for a plot and a comparison with the class-C distribution, that has a finite average for all M .

The result (23) holds irrespective of the sign of $\text{Det } S_0$; in other words, the statistics of the Fermi-level anomaly in the CRE does not depend on the presence or absence of an unpaired Majorana zero-mode in the quantum dot. As we remarked at the end of the previous section, in connection with the thermopower, this is a direct consequence of the unitary invariance (16) of the circular ensemble.

B. Numerical check

As a check on our analytical result we have calculated $P(\rho_0)$ numerically from the Gaussian ensemble of random Hamiltonians. We focus on symmetry class D, where we can test in particular for the effect of a Majorana zero mode.

FIG. 3. (Color online) Probability distributions of the Fermi-level density of states, for $M = 1$ and $M = 2$ modes coupling the quantum dot to the continuum, in symmetry classes C and D. The ensemble average diverges for class D; see Eq. (22).

The Hamiltonian H is related to the scattering matrix $S(E)$ by the Weidenmüller formula [58,59],

$$\begin{aligned} S(E) &= \frac{1 + i\pi W^\dagger (H - E)^{-1} W}{1 - i\pi W^\dagger (H - E)^{-1} W} \\ &= 1 + 2\pi i W^\dagger (H_{\text{eff}} - E)^{-1} W, \\ H_{\text{eff}} &= H - i\pi W W^\dagger. \end{aligned} \quad (24)$$

The $M_0 \times M$ matrix W couples the M_0 energy levels in the quantum dot to $M \ll M_0$ scattering channels. Ballistic coupling corresponds to

$$W_{nm} = \delta_{nm} \sqrt{M_0 \delta_0 / \pi}. \quad (25)$$

FIG. 4. (Color online) Histograms: Probability distributions of the Fermi-level density of states in symmetry class D for $M = 1$, $M_0 = 140, 141$ and $M = 2$, $M_0 = 200, 201$, calculated numerically from Eq. (27) by averaging the Hamiltonian over the Gaussian ensemble. For each dimensionality M of the scattering matrix we compare an even-dimensional Hamiltonian, without a Majorana zero mode, to an odd-dimensional Hamiltonian with a zero mode. The black curve is the analytical result (23) for the circular scattering matrix ensemble, predicting no effect from the Majorana zero mode for this case of ballistic coupling. Notice that there is no fit parameter in this comparison between numerics and analytics.

The density of states is determined by the scattering matrix via [60]

$$\rho(E) = -\frac{i}{2\pi} \frac{d}{dE} \ln \text{Det } S(E). \quad (26)$$

From Eqs. (24) and (26) we obtain an expression for the Fermi-level density of states in terms of the Hamiltonian,

$$\rho_0 = \text{Tr} \left([1 - 2\pi i W^\dagger (H_{\text{eff}}^\dagger)^{-1} W] W^\dagger H_{\text{eff}}^{-2} W \right). \quad (27)$$

In the Majorana basis the class-D Hamiltonian is purely imaginary, $H = iA$, with A a real antisymmetric matrix. The Gaussian ensemble has probability distribution [6,55]

$$P(A) \propto \prod_{n>m} \exp \left(-\frac{\pi^2 A_{nm}^2}{2M_0 \delta_0^2} \right). \quad (28)$$

The dimensionality of A is odd if the quantum dot contains an unpaired Majorana zero mode; otherwise it is even.

Numerical results for the probability distribution of ρ_0 for $M = 1, 2$ scattering channels are shown in Fig. 4. The agreement with the analytical distribution (23) is excellent, including the absence of any effect from the Majorana zero mode.

V. THERMOPOWER DISTRIBUTION

We apply the general thermopower formulas (14) and (15) to a single-channel point contact, with transmission probability T into the edge mode of the superconductor. There are two independent delay times D_1, D_2 in class C, each with a twofold spin degeneracy and a twofold electron-hole degeneracy ($d_T = 4$). Because of this degeneracy the class-C edge mode contains Kramers pairs of Majorana fermions. In class D the Majorana edge mode is unpaired and all delay times are nondegenerate ($d_T = 1$). The point contact contributes two and the edge mode one more, so class D has a total of three independent delay times D_1, D_2, D_3 .

Equations (14) and (15) can be expressed in terms of these quantities; see Appendix B. We denote the dimensionless thermopower by $p = (\hbar/t_0)S/S_0$ and add a subscript C, D to indicate the symmetry class. For class C we have

$$p_C = \frac{(D_2/t_0 - D_1/t_0)\xi \sqrt{T(1-T)}}{1 - (1-T)\cos 2\beta}. \quad (29)$$

The independent variables β, ξ enter via the eigenvectors of S_0 and Q , with distribution

$$P(\beta, \xi) = \frac{3}{4}(1 - \xi^2) \sin 2\beta, \quad |\xi| < 1, \quad 0 < \beta < \pi/2. \quad (30)$$

The class-D distribution p_D has a more lengthy expression, involving three delay times; see Appendix B. These are all averages in the grand-canonical ensemble, without including effects from the charging energy of the quantum dot (which could force a transition into the canonical ensemble) [61].

The resulting distributions, shown in Fig. 5, are qualitatively different, with a quadratic maximum in class C and a cusp in class D. The variance diverges in class D, while in class C

$$\langle p_C^2 \rangle = \frac{2}{15}(3 \ln 2 - 2) = 0.011. \quad (31)$$

FIG. 5. (Color online) Probability distribution of the dimensionless thermopower $p = \mathcal{S} \times \hbar / t_0 S_0$ in symmetry class C (black solid curve, bottom and left axes), and in class D (blue dashed curve, top and right axes). These are results for the quantum dot of Fig. 2 connecting a single-channel point contact to the unpaired Majorana edge mode of a chiral p -wave superconductor (class D), or to the paired Majorana mode of a chiral d -wave superconductor (class C).

VI. CONCLUSION

Perhaps the most remarkable conclusion of our analysis is that the density of states of a Majorana zero mode is not topologically protected in an open system.

Take a superconducting quantum dot with an unpaired Majorana zero mode and bring it into contact with a metallic contact, as in Fig. 6. Is something left of the spectral peak? The answer is “yes” for tunnel coupling [8–15], as it should be if the level broadening is less than the level spacing in the quantum dot. What we have found is that the answer is “no” for ballistic coupling, with level broadening comparable to level spacing.

As an intuitive explanation, one might argue that this is the ultimate consequence of the fact that the two average densities of states $\rho_+(E)$ and $\rho_-(E) + \delta(E)$ of a closed quantum dot without and with a Majorana zero mode are markedly different [5,6], see Eq. (1), and yet have the same integrated spectral weight of half a fermion. Still, we had not expected to find that the entire probability distribution

of the Fermi-level density of states becomes identical in the topologically trivial and nontrivial system, once the quantum dot is coupled ballistically to $M \geq 1$ conducting modes.

It would be a mistake to conclude that the whole notion of a topologically nontrivial superconductor applies only to a closed system. Indeed, the Andreev conductance remains sensitive to the presence or absence of a Majorana zero mode, even for ballistic coupling, when no trace is left in the density of states [18]. This can be seen most directly for the case $M = 2$ of a superconducting quantum dot coupled to a normal metal by a pair of spin-resolved electron-hole modes. The Andreev conductance is then given simply by

$$G = \frac{e^2}{h} (1 - \text{Det } S_0), \quad (32)$$

and so is in one-to-one relationship with the topological quantum number $\text{Det } S_0 = \pm 1$. In contrast, the Fermi-level density of states has the same probability distribution (23) regardless of the sign of $\text{Det } S_0$.

We have applied our results for the probability distribution of the time-delay matrix to a calculation of the thermopower induced by edge modes of a chiral p -wave or chiral d -wave superconductor [16]. The search for electrical edge conduction in such topological superconductors, notably Sr_2RuO_4 [62], has remained inconclusive [63], in part because of the charge neutrality of an unpaired Majorana mode at the Fermi level [64–67]. Figure 5 shows that both unpaired and paired Majorana edge modes can produce a nonzero thermopower—of random sign, with a magnitude of order $S_0/\delta_0 = (0.3 \text{ mV/K}) \times k_B T_0/\delta_0$. This is a small signal, but it has the attractive feature that it directly probes for the existence of propagating edge modes—irrespective of their charge neutrality.

ACKNOWLEDGMENTS

This research was supported by the Foundation for Fundamental Research on Matter (FOM), the Netherlands Organization for Scientific Research (NWO/OCW), the Alexander von Humboldt Foundation, and an ERC Synergy grant.

APPENDIX A: DERIVATION OF THE DELAY-TIME DISTRIBUTION FOR THE ALTLAND-ZIRNBAUER ENSEMBLES

Repeating the steps of Refs. [17] and [50] we extend the calculation of the joint distribution $P(S_0, Q)$ from the nonsuperconducting Wigner-Dyson ensembles to the superconducting Altland-Zirnbauer ensembles. We treat the two symmetry classes C, D without time-reversal symmetry, of relevance for the main text (see Table I), and for completeness also consider the time-reversal symmetric classes CI and DIII (see Table II).

1. Unitary invariance

Since the entire calculation relies on the unitary invariance (16) of the Altland-Zirnbauer circular ensembles, we demonstrate that first. Following Ref. [50] we construct the $\mathcal{M} \times \mathcal{M}$ energy-dependent unitary scattering matrix $S(E)$ in

FIG. 6. (Color online) Geometry to detect a Majorana zero mode by a measurement of the Andreev conductance of a ballistic point contact to a superconducting quantum dot. The probability distribution of the conductance depends on the presence or absence of the Majorana zero mode, while the distribution of the density of states does not.

TABLE II. The two Altland-Zirnbauer classes with time-reversal symmetry.

Symmetry class	Class CI	Class DIII
S-matrix space	symplectic & symmetric	orthogonal & self-dual
d_T	4	2
d_E	2	2
α	1	-1
β	2	2

terms of an $\mathcal{M}_0 \times \mathcal{M}_0$ energy-independent unitary matrix U ,

$$S(E) = \mathcal{P}U(e^{-2\pi i E/M_0\delta_0} + \mathcal{R}U)^{-1}\mathcal{P}^T. \quad (\text{A1})$$

The rectangular $\mathcal{M} \times \mathcal{M}_0$ matrix $\mathcal{P}$ has elements $\mathcal{P}_{nm} = \delta_{nm}$ and $\mathcal{R} = 1 - \mathcal{P}^T\mathcal{P}$. The eigenvalues $e^{i\phi_n}$ of U have the same degeneracy d_E as the energy eigenvalues, so there are $M_0 = \mathcal{M}_0/d_E$ distinct eigenvalues on the unit circle, arranged symmetrically around the real axis.

The $\mathcal{M}_0 \times \mathcal{M}_0$ Hermitian matrix H is related to U via a Cayley transform,

$$U = e^{2\pi i \epsilon/M_0\delta_0} \frac{\pi H/M_0\delta_0 + i}{\pi H/M_0\delta_0 - i} \quad (\text{A2})$$

$$\Leftrightarrow H = \frac{iM_0\delta_0}{\pi} \frac{U + e^{2\pi i \epsilon/M_0\delta_0}}{U - e^{2\pi i \epsilon/M_0\delta_0}}.$$

The factor $e^{2\pi i \epsilon/M_0\delta_0}$ with $\epsilon \rightarrow 0$ is introduced to regularize the singular inverse when U has an eigenvalue pinned at +1, as we will discuss in just a moment.

We can immediately observe that if we take a circular ensemble for U , with distribution function $P(U) = P(U'U) = P(UU')$, then the unitary invariance (16) of the distribution functional $P[S(E)]$ is manifestly true. So what we have to verify is that the construction (A1)–(A2) with U in the circular ensemble is, first, equivalent to the Weidenmüller formula (24), and second, produces a Gaussian ensemble for H . It is sufficient if the equivalence holds in the low-energy range $|E| \lesssim M_0\delta_0 \ll M_0\delta_0$.

First, substitution of Eq. (A2) into Eq. (A1) gives

$$S(E) = \frac{1 + i\mathcal{P} \frac{M_0\delta_0 - i\pi H \tan(\pi E^+/M_0\delta_0)}{\pi H - M_0\delta_0 \tan(\pi E^+/M_0\delta_0)} \mathcal{P}^T}{1 - i\mathcal{P} \frac{M_0\delta_0 - i\pi H \tan(\pi E^+/M_0\delta_0)}{\pi H - M_0\delta_0 \tan(\pi E^+/M_0\delta_0)} \mathcal{P}^T}$$

$$= \frac{1 + i\mathcal{P} \frac{M_0\delta_0}{\pi(H-E^+)} \mathcal{P}^T}{1 - i\mathcal{P} \frac{M_0\delta_0}{\pi(H-E^+)} \mathcal{P}^T} + \mathcal{O}(M/M_0), \quad (\text{A3})$$

with $E^+ = E + \epsilon$. This is the Weidenmüller formula (24), with the ballistic coupling matrix $W = \mathcal{P}^T(M_0\delta_0/\pi^2)^{1/2}$ from Eq. (25).

Second, the Cayley transform (A2) produces a Lorentzian instead of a Gaussian distribution for H , but in the low-energy range the two ensembles are equivalent [68]. One also readily checks that a uniform distribution with spacing $2\pi/M_0$ of the distinct eigenphases ϕ_n of U produces a mean spacing δ_0 of the distinct eigenvalues E_n of H , through the relation $(\pi/M_0\delta_0)E_n = \cotan(\phi_n/2) \approx (\pi - \phi_n)/2$ in the low-energy range.

The finite- ϵ regularization is irrelevant in the class C and CI circular ensembles, because there the U 's with an eigenvalue +1 are of measure zero. In the class D and DIII circular ensembles, in contrast, an eigenvalue may be pinned at unity and the regularization is essential. Let us analyze this for class D (the discussion in class DIII is similar). The matrix U in class D is real orthogonal, with determinant $\text{Det } U = (-1)^{n_M}$ fixed by the parity of the number of Majorana zero modes [cf. Eq. (20)]. This implies that U has an eigenvalue pinned at +1 if M_0 is even and n_M is odd, or if M_0 is odd and n_M is even. The Cayley transform (A2) then maps to an eigenvalue of H at infinity. This eigenvalue does not contribute to the low-energy scattering matrix (A3), so that it can be removed from the spectrum of H . Hence, whereas the dimension M_0 of the unitary matrix U can be arbitrary, the dimension of H is always even for even n_M and odd for odd n_M .

2. Broken time-reversal symmetry, classes C and D

We now proceed with the calculation of the distribution of the time-delay matrix, first in symmetry classes C and D. The starting point is the Weidenmüller formula (24) or (A3) for the energy-dependent scattering matrix. Differentiation gives the time-delay matrix defined in Eq. (3),

$$Q^{-1} = \frac{1}{2\pi\hbar} \lim_{\epsilon \rightarrow 0} [1 - i\pi W^\dagger(H - \epsilon)^{-1}W]$$

$$\times \frac{1}{W^\dagger(H - \epsilon)^{-2}W} [1 + i\pi W^\dagger(H - \epsilon)^{-1}W], \quad (\text{A4})$$

in terms of the Hamiltonian H of the closed quantum dot and the coupling matrix W to the scattering channels. The dimensionality of H is $d_E M_0 \times d_E M_0$ while the dimensionality of Q and S is $d_T M \times d_T M$ (and W has dimension $d_E M_0 \times d_T M$). The unitary invariance (16) implies $P(S_0, Q) = P(-1, Q)$, so we may restrict ourselves to the case that H has a zero eigenvalue with multiplicity $d_T M$ —since then $S_0 = \lim_{E \rightarrow 0} S(E) = -1$.

Restricting H to its $d_T M$ -dimensional null space we have, using the ballistic coupling matrix (25),

$$W^\dagger(H - \epsilon)^{-p}W \rightarrow (M_0\delta_0/\pi^2)(-\epsilon)^{-p}\tilde{\Omega}^\dagger\tilde{\Omega}, \quad (\text{A5})$$

$$Q^{-1} \rightarrow (\delta_0/2\pi\hbar)\Omega^\dagger\Omega, \quad \Omega = M_0^{1/2}\tilde{\Omega}. \quad (\text{A6})$$

The matrix Ω is a $d_T M \times d_T M$ submatrix of a $d_E M_0 \times d_E M_0$ unitary matrix, rescaled by a factor $\sqrt{M_0}$. In the relevant limit $M_0/M \rightarrow \infty$ this matrix has independent Gaussian elements,

$$P(\Omega) \propto \exp\left[-\frac{1}{2}\beta(d_E/d_T)\text{Tr}' \Omega^\dagger\Omega\right]$$

$$= \exp\left(-\frac{1}{2}\beta t_0 \text{Tr}' Q^{-1}\right), \quad (\text{A7})$$

with $t_0 = (2\pi\hbar/\delta_0)(d_E/d_T)$. The prime in the trace, and in the determinants appearing below, indicates that the d_T -fold degenerate eigenvalues are only counted once. The symmetry index β counts the number of independent degrees of freedom of the matrix elements of Ω , real in class D ($\beta = 1$) and quaternion in class C ($\beta = 4$). The positive-definite matrix Q^{-1} of the form (A6) is called a Wishart matrix in random-matrix theory [69].

Using Eq. (24), an infinitesimal deviation of S_0 from -1 can be expressed as

$$V\Omega(S_0 + 1)\Omega^\dagger V^\dagger = A, \quad (\text{A8})$$

with A a $d_T M \times d_T M$ anti-Hermitian matrix, $A = -A^\dagger$. The matrix A is a submatrix of iH , so its matrix elements are real in class D and quaternion in class C. The unitary matrix V has been inserted so that $P(A) = \text{constant}$ near $A = 0$. Since the transformation $\Omega \mapsto V\Omega$ has no effect on $P(\Omega)$ and leaves Q unaffected, we may in what follows omit V .

The joint distribution $P(S_0, Q^{-1})$ follows from $P(\Omega)P(A)$ upon multiplication by two Jacobian determinants,

$$\begin{aligned} P(S_0, Q^{-1}) &= P(\Omega)P(A) \left\| \frac{\partial \Omega}{\partial Q^{-1}} \right\| \times \left\| \frac{\partial A}{\partial S_0} \right\| \\ &\propto \exp\left(-\frac{1}{2}\beta t_0 \text{Tr}' Q^{-1}\right) \left\| \frac{\partial \Omega^\dagger \Omega}{\partial \Omega} \right\|^{-1} \left\| \frac{\partial \Omega^{-1} A \Omega^{\dagger-1}}{\partial A} \right\|^{-1}. \end{aligned} \quad (\text{A9})$$

The Jacobians can be evaluated using textbook methods [69,70],

$$\left\| \frac{\partial \Omega^\dagger \Omega}{\partial \Omega} \right\|^{-1} \propto (\text{Det}' \Omega^\dagger \Omega)^{-1+\beta/2}, \quad (\text{A10})$$

$$\left\| \frac{\partial \Omega^{-1} A \Omega^{\dagger-1}}{\partial A} \right\|^{-1} \propto (\text{Det}' \Omega^\dagger \Omega)^{\alpha+1+(M-1)\beta/2}. \quad (\text{A11})$$

Here $\alpha + 1$ equals the number of degrees of freedom of a diagonal element of A , while an off-diagonal element has β degrees of freedom (see Table III). So $\alpha + 1 = 0$, $\beta = 1$ for a real antisymmetric matrix A (class D), while $\alpha + 1 = 3$, $\beta = 4$ for a quaternion anti-Hermitian A (class C).

TABLE III. Upper table: Representation of the Hamiltonian H in the four Altland-Zirnbauer symmetry classes. All coefficients q_n , a, b are real. The Pauli matrices $\boldsymbol{\tau} = (\tau_x, \tau_y, \tau_z)$ act on the electron-hole degree of freedom, while the $\boldsymbol{\sigma}$'s act on the spin degree of freedom. The symmetry indices β and $\alpha + 1$ from Tables I and II count, respectively, the number of degrees of freedom of the off-diagonal and diagonal components of the Hermitian matrix H , in the Majorana basis for classes D, DIII and in the electron-hole basis for classes C, CI. For completeness and comparison, we show in the lower table the corresponding listing for the three Wigner-Dyson symmetry classes.

	C	D	CI	DIII
H_{nm} ($n \neq m$)	$iq_0 + \mathbf{q} \cdot \boldsymbol{\tau}$ $\beta = 4$	iq_0 $\beta = 1$	$a\tau_x + b\tau_z$ $\beta = 2$	$ia\sigma_x + ib\sigma_z$ $\beta = 2$
H_{nn}	$\mathbf{q} \cdot \boldsymbol{\tau}$ $\alpha + 1 = 3$	0 $\alpha + 1 = 0$	$a\tau_x + b\tau_z$ $\alpha + 1 = 2$	0 $\alpha + 1 = 0$
	A	AI	AII	
H_{nm} ($n \neq m$)	$a + ib$ $\beta = 2$	a $\beta = 1$	$q_0 + i\mathbf{q} \cdot \boldsymbol{\sigma}$ $\beta = 4$	
H_{nn}	a $\alpha + 1 = 1$	a $\alpha + 1 = 1$	q_0 $\alpha + 1 = 1$	

Collecting results, we arrive at the distribution

$$P(S_0, Q^{-1}) \propto \exp\left(-\frac{1}{2}\beta t_0 \text{Tr}' Q^{-1}\right) (\text{Det}' Q^{-1})^{\alpha+M\beta/2}. \quad (\text{A12})$$

The distribution (18) of the eigenvalues γ_n of Q^{-1} follows upon multiplication by one more Jacobian, from matrix elements to eigenvalues.

3. Preserved time-reversal symmetry, classes CI and DIII

The time-reversal operator acts in a different way in classes CI and DIII. In class CI the action is the transpose, so that $S = S^T$, $H = H^T$ are symmetric matrices. In class DIII these matrices are self-dual, $S = \sigma_y S^T \sigma_y \equiv S^D$, where the Pauli matrix σ_y acts on the spin degree of freedom. It is convenient to use a unified notation $\tilde{U}$ to denote the transpose U^T of a matrix in class CI and the dual U^D in class DIII. Unitary invariance of the circular ensemble then amounts to

$$P[S(E)] = P[\tilde{U} \cdot S(E) \cdot U], \quad (\text{A13})$$

for energy-independent unitary matrices U .

Time-reversal symmetry allows us to “take the square root” of the Fermi-level scattering matrix (Takagi factorization [71]),

$$S_0 = \tilde{S}_{1/2} S_{1/2}. \quad (\text{A14})$$

In class DIII the sign of the determinant of $S_{1/2}$ is a topological quantum number [72],

$$\text{Det } S_{1/2} = \text{Pf}(i\sigma_y S_0) = \pm 1, \quad (\text{A15})$$

equal to -1 when the quantum dot contains a Kramers pair of Majorana zero modes. The symmetrized time-delay matrix is defined in terms of this square root,

$$Q = -i\hbar \lim_{E \rightarrow 0} \tilde{S}_{1/2}^\dagger \frac{dS}{dE} S_{1/2}^\dagger. \quad (\text{A16})$$

The definition (3) of the matrix Q used in classes C and D, without time-reversal symmetry, gives the same eigenvalues as definition (A16), but would introduce a spurious correlation between S and Q . With the definition (A16) the unitary invariance (A13) allows us to equate $P(S_0, Q) = P(-1, Q)$, by taking $U = S_{1/2}^\dagger i\sigma_x$ in class CI and $U = S_{1/2}^\dagger \sigma_x$ in class DIII.

Comparing to the derivation of the previous subsection, what changes is that the matrix elements of Ω and A are equivalent to complex numbers $a + ib$, rather than being real or quaternion. Specifically, Ω has matrix elements of the form $a\sigma_0 + ib\sigma_y$ in both classes CI and DIII (to ensure that $\Omega^\dagger = \tilde{\Omega}$), while the matrix elements of A are of the form $ia\sigma_x + ib\sigma_z$ in class CI and of the form $a\sigma_x + b\sigma_z$ in class DIII (to ensure that $A^\dagger = -\tilde{A}$). The Jacobian (A10) still applies, now with $\beta = 2$, while the Jacobian (A11) evaluates to

$$\left\| \frac{\partial \Omega^{-1} A \Omega^{\dagger-1}}{\partial A} \right\|^{-1} \propto \begin{cases} (\text{Det}' \Omega^\dagger \Omega)^{M+1} & \text{in class CI,} \\ (\text{Det}' \Omega^\dagger \Omega)^{M-1} & \text{in class DIII.} \end{cases} \quad (\text{A17})$$

Collecting results, we arrive at

$$P(S_0, Q^{-1}) \propto \exp(-t_0 \text{Tr}' Q^{-1}) (\text{Det}' Q^{-1})^{M\pm 1}, \quad (\text{A18})$$

with exponent $M + 1$ in class CI and $M - 1$ in class DIII. As before, the primed trace and determinant count degenerate eigenvalues only once. The distribution (18) of the eigenvalues γ_n of Q^{-1} follows with $\beta = 2$ and $\alpha = \pm 1$.

APPENDIX B: DETAILS OF THE CALCULATION OF THE THERMOPOWER DISTRIBUTION

1. Invariant measure on the unitary, orthogonal, or symplectic groups

For later reference, we record explicit expressions for the invariant measure $dU = P(\{\alpha_n\}) \prod_n d\alpha_n$ (the Haar measure) in parametrizations $U(\{\alpha_n\})$ of the unitary group $SU(N)$, as well as the orthogonal or unitary symplectic subgroups $SO(N)$, $Sp(2N)$. (We will only need results for small N .)

The invariant measure is determined by the metric tensor

$$g_{mn} = -\text{Tr} U^\dagger (\partial U / \partial \alpha_m) U^\dagger (\partial U / \partial \alpha_n), \quad (\text{B1})$$

via $P(\{\alpha_n\}) \propto \sqrt{\det g}$. The function P represents the probability distribution of the α_n 's when the matrix U is drawn randomly and uniformly from the unitary group (circular unitary ensemble, CUE), or from the orthogonal and symplectic subgroups (circular real and quaternion ensembles, CRE and CQE).

For $SO(2)$ we have trivially

$$R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \Rightarrow P(\theta) = \text{constant}. \quad (\text{B2})$$

For $SU(2) = Sp(2)$ we can choose different parametrizations:

$$U = \exp[i\beta(\tau_z \cos \theta + \tau_x \sin \theta \cos \phi + \tau_y \sin \theta \sin \phi)] \\ \Rightarrow P(\beta, \theta, \phi) \propto \sin^2 \beta \sin \theta, \quad (\text{B3a})$$

$$U = e^{i\alpha\tau_z} \exp[i\beta(\tau_x \cos \phi + \tau_y \sin \phi)] \\ \Rightarrow P(\alpha, \beta, \phi) \propto \sin 2\beta, \quad (\text{B3b})$$

$$U = e^{i\alpha\tau_z} R(\theta) e^{i\alpha'\tau_z} \Rightarrow P(\alpha, \alpha', \theta) \propto \sin 2\theta. \quad (\text{B3c})$$

For the group of 3×3 orthogonal matrices we will use the Euler angle parametrization

$$O_\pm = \begin{pmatrix} R(\alpha) & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \pm 1 & 0 \\ 0 & R(\theta) \end{pmatrix} \begin{pmatrix} R(\alpha') & 0 \\ 0 & 1 \end{pmatrix} \\ \Rightarrow P(\alpha, \alpha', \theta) \propto \sin \theta. \quad (\text{B4})$$

The $\pm$ sign distinguishes the sign of the determinant $\text{Det } O_\pm = \pm 1$, with $SO(3)$ corresponding to O_+ .

Finally, for $Sp(4)$ we use the polar decomposition

$$U = \begin{pmatrix} U_1 & 0 \\ 0 & U_2 \end{pmatrix} \begin{pmatrix} \tau_0 \cos \theta & -\tau_0 \sin \theta \\ \tau_0 \sin \theta & \tau_0 \cos \theta \end{pmatrix} \begin{pmatrix} \tau_0 & 0 \\ 0 & U_3 \end{pmatrix} \\ \Rightarrow P(\theta) = \sin^3 2\theta. \quad (\text{B5})$$

The matrices U_p are independently and uniformly distributed in $SU(2)$; see Eq. (B3). There are only three independent U_p 's, with 3 free parameters each, because one of the four blocks can be absorbed in the three others, so we have set it to the unit τ_0 without loss of generality. [One can check that the total number $N(2N + 1) \mapsto 10$ of free parameters of $Sp(2N)$ agrees: $3 + 3 + 3$ from the U_p 's plus θ makes 10.]

2. Elimination of eigenvector components

The thermopower expressions (14) and (15) depend on the transmission eigenvalues T_n and delay times D_n , but in addition there is a dependence on eigenvectors. Many of the eigenvector degrees of freedom can be eliminated by using the invariance of the distribution of the time-delay matrix under the unitary transformation $Q \mapsto U^\dagger Q U$, following from Eq. (16).

a. Class C

In class C we proceed as follows. The 4×4 unitary symplectic scattering matrix S_0 has the polar decomposition (B5), which we write in the form

$$S_0 = \begin{pmatrix} U_1 & 0 \\ 0 & U_2 \end{pmatrix} \begin{pmatrix} \tau_0 \sqrt{1-T} & -\tau_0 \sqrt{T} \\ \tau_0 \sqrt{T} & \tau_0 \sqrt{1-T} \end{pmatrix} \begin{pmatrix} \tau_0 & 0 \\ 0 & U_3 \end{pmatrix}, \quad (\text{B6})$$

$$U_n = e^{i\alpha_n \tau_z} \exp[i\beta_n(\tau_x \cos \phi_n + \tau_y \sin \phi_n)]. \quad (\text{B7})$$

We ignore the spin degree of freedom, which plays no role in the calculation. The remaining twofold degeneracy of the transmission eigenvalue T comes from the electron-hole degree of freedom.

The time-delay matrix is Hermitian with quaternion elements,

$$Q = \begin{pmatrix} a\tau_0 & q \\ q^\dagger & b\tau_0 \end{pmatrix}, \quad q = q_0\tau_0 + iq_1\tau_x + iq_2\tau_y + iq_3\tau_z. \quad (\text{B8})$$

With some trial and error, we found the unitary symplectic transformation

$$Q \mapsto U^\dagger Q U, \quad U = (U_0)^2 \begin{pmatrix} \tau_0 & 0 \\ 0 & e^{-i\alpha_3 \tau_z} U_3^\dagger e^{i\alpha_3 \tau_z} \end{pmatrix}, \quad (\text{B9})$$

$$U_0 = \begin{pmatrix} U_1 & 0 \\ 0 & \tau_0 \end{pmatrix} \begin{pmatrix} \tau_0 \sqrt{1-T} & -\tau_0 \sqrt{T} \\ \tau_0 \sqrt{T} & \tau_0 \sqrt{1-T} \end{pmatrix}, \quad (\text{B10})$$

that eliminates most of the eigenvector components from the class-C thermopower expression (15). We are left with

$$S/S_0 = -\hbar^{-1} \frac{2q_3 \sqrt{T(1-T)}}{1 - (1-T) \cos 2\beta_1}. \quad (\text{B11})$$

The probability distribution of the eigenvector parameter β_1 follows from Eq. (B3b),

$$P(\beta_1) = \sin 2\beta_1, \quad 0 < \beta_1 < \pi/2. \quad (\text{B12})$$

b. Class D

The algebra is simpler in class D, where the matrix elements are real rather than quaternion. We use the Euler angle parametrization (B4) of the 3×3 orthogonal matrix S_0 with determinant $\text{Det } S_0 = \pm 1$. Substitution of the orthogonal transformation

$$Q \mapsto \begin{pmatrix} R(-\alpha') & 0 \\ 0 & 1 \end{pmatrix} Q \begin{pmatrix} R(\alpha') & 0 \\ 0 & 1 \end{pmatrix} \quad (\text{B13})$$

into the class-D thermopower expression (14) leads directly to

$$\frac{S}{S_0} = \frac{Q_{13}}{\hbar} \times \begin{cases} -\cotan(\theta/2) & \text{if } \text{Det } S_0 = +1, \\ \tan(\theta/2) & \text{if } \text{Det } S_0 = -1, \end{cases} \quad (\text{B14a})$$

$$P(\theta) = \frac{1}{2} \sin \theta, \quad 0 < \theta < \pi. \quad (\text{B14b})$$

The transmission eigenvalue is $T = \sin^2 \theta$. Since $P(\theta) = P(\pi - \theta)$ the probability distribution of the thermopower does not depend on the sign of $\text{Det } S_0$.

3. Marginal distribution of an element of the time-delay matrix

The two expressions (B11) and (B14a) for the thermopower contain a single off-diagonal element of the time-delay matrix Q . We can calculate its marginal distribution, using the eigenvalue distribution of Sec. III and the fact that the eigenvectors of Q are uniformly distributed with the invariant measure of the symplectic group (class C) or the orthogonal group (class D).

a. Class C

In class C the 4×4 time-delay matrix Q is diagonalized by a unitary symplectic matrix U ,

$$Q = U \begin{pmatrix} D_1 \tau_0 & 0 \\ 0 & D_2 \tau_0 \end{pmatrix} U^\dagger. \quad (\text{B15})$$

Each of the eigenvalues D_1 and D_2 of Q has a twofold degeneracy from the electron-hole degree of freedom. (As before, we can ignore the spin degree of freedom.) The matrix U has the polar decomposition (B5).

The quaternion Q_{12} is given in this parametrization by

$$Q_{12} = \frac{1}{2}(D_1 - D_2)(\sin 2\theta)U_1 U_2^\dagger, \quad (\text{B16})$$

and since q_3 from Eq. (B8) equals $-\frac{1}{2}i \text{Tr } \tau_z Q_{12}$, we have

$$q_3 = \frac{1}{4}(D_1 - D_2)(\sin 2\theta) \text{Tr } U_0. \quad (\text{B17})$$

The matrix $U_0 = -i\tau_z U_1 U_2^\dagger$ is uniformly distributed in $\text{SU}(2)$. Using the invariant measures (B3a) and (B5) we

arrive at

$$q_3 = \frac{1}{2}(D_1 - D_2) \cos \beta \sin 2\theta, \quad (\text{B18})$$

$$P(\beta, \theta) = (6/\pi) \sin^2 \beta \sin^3 2\theta, \quad 0 < \beta, \quad \theta < \pi/2.$$

The two angular variables β, θ can be combined into a single variable ξ :

$$q_3 = \frac{1}{2}(D_1 - D_2)\xi, \quad (\text{B19})$$

$$P(\xi) = \frac{3}{4}(1 - \xi^2), \quad -1 < \xi < 1.$$

The marginal distribution of q_3 then follows upon integration.

Collecting results, we have the following probability distributions for the variables appearing in the class-C thermopower:

$$S/S_0 = t_0 \hbar^{-1} \frac{(D_2 - D_1)\xi \sqrt{T(1-T)}}{1 - (1-T) \cos 2\beta}, \quad (\text{B20})$$

$$P(\beta) = \sin 2\beta, \quad 0 < \beta < \pi/2, \quad (\text{B21})$$

$$P(\xi) = \frac{3}{4}(1 - \xi^2), \quad -1 < \xi < 1, \quad (\text{B22})$$

$$P(T) = 6T(1-T), \quad 0 < T < 1, \quad (\text{B23})$$

$$P(D_1, D_2) = \frac{32}{42525}(D_1 - D_2)^4 (D_1 D_2)^{-12} \times \exp[-2/D_1 - 2/D_2], \quad D_1, D_2 > 0, \quad (\text{B24})$$

where for notational convenience we measure the delay times in units of t_0 .

b. Class D

The 3×3 time-delay matrix in class D is diagonalized by $Q = O_+ \text{diag}(D_1, D_2, D_3) O_+^\top$, with $O_+ \in \text{SO}(3)$ parameterized as in Eq. (B4). In terms of these parameters, the matrix element Q_{13} is given by

$$Q_{13} = X \cos \alpha + Y \sin \alpha, \quad (\text{B25})$$

$$X = \frac{1}{2}(D_1 - D_2) \sin \theta' \sin 2\alpha',$$

$$Y = \frac{1}{2}[(D_3 - D_2) \cos^2 \alpha' + (D_3 - D_1) \sin^2 \alpha'] \times \sin 2\theta',$$

$$P(\alpha, \alpha', \theta') = (8\pi^2)^{-1} \sin \theta',$$

$$0 < \alpha, \quad \alpha' < 2\pi, \quad 0 < \theta < \pi.$$

The thermopower distribution follows upon integration, using Eqs. (18), (B14), and (B25).

-
- [1] A. Altland and M. R. Zirnbauer, *Phys. Rev. B* **55**, 1142 (1997).
 - [2] *Handbook on Random Matrix Theory*, edited by G. Akemann, J. Baik, and P. Di Francesco (Oxford University Press, Oxford, 2011).
 - [3] G. Volovik, *JETP Lett.* **70**, 609 (1999).
 - [4] N. Read and D. Green, *Phys. Rev. B* **61**, 10267 (2000).
 - [5] M. Bocquet, D. Serban, and M. R. Zirnbauer, *Nucl. Phys. B* **578**, 628 (2000).
 - [6] D. A. Ivanov, *J. Math. Phys.* **43**, 126 (2002); [arXiv:cond-mat/0103089](https://arxiv.org/abs/cond-mat/0103089).
 - [7] R. Jackiw and C. Rebbi, *Phys. Rev. D* **13**, 3398 (1976).

- [8] D. Bagrets and A. Altland, *Phys. Rev. Lett.* **109**, 227005 (2012).
- [9] J. Liu, A. C. Potter, K. T. Law, and P. A. Lee, *Phys. Rev. Lett.* **109**, 267002 (2012).
- [10] M. A. Skvortsov, P. M. Ostrovsky, D. A. Ivanov, and Ya. V. Fominov, *Phys. Rev. B* **87**, 104502 (2013).
- [11] T. D. Stanescu and S. Tewari, *Phys. Rev. B* **87**, 140504(R) (2013).
- [12] J. D. Sau and S. Das Sarma, *Phys. Rev. B* **88**, 064506 (2013).
- [13] P. A. Ioselevich and M. V. Feigel'man, *New J. Phys.* **15**, 055011 (2013).

- [14] P. Neven, D. Bagrets, and A. Altland, *New J. Phys.* **15**, 055019 (2013).
- [15] D. A. Ivanov, P. M. Ostrovsky, and M. A. Skvortsov, [arXiv:1307.0372](#).
- [16] C.-Y. Hou, K. Shtengel, and G. Refael, *Phys. Rev. B* **88**, 075304 (2013).
- [17] P. W. Brouwer, K. M. Frahm, and C. W. J. Beenakker, *Phys. Rev. Lett.* **78**, 4737 (1997).
- [18] C. W. J. Beenakker, J. P. Dahlhaus, M. Wimmer, and A. R. Akhmerov, *Phys. Rev. B* **83**, 085413 (2011).
- [19] C. W. J. Beenakker, *Lect. Notes Phys.* **667**, 131 (2005).
- [20] T. Dirks, T. L. Hughes, S. Lal, B. Uchoa, Y.-F. Chen, C. Chialvo, P. M. Goldbart, and N. Mason, *Nat. Phys.* **7**, 386 (2011).
- [21] E. J. H. Lee, X. Jiang, R. Aguado, G. Katsaros, C. M. Lieber, and S. De Franceschi, *Phys. Rev. Lett.* **109**, 186802 (2012).
- [22] W. Chang, V. E. Manucharyan, T. S. Jespersen, J. Nygard, and C. M. Marcus, *Phys. Rev. Lett.* **110**, 217005 (2013).
- [23] J. P. Dahlhaus, B. Béri, and C. W. J. Beenakker, *Phys. Rev. B* **82**, 014536 (2010).
- [24] M. Cutler and N. F. Mott, *Phys. Rev.* **181**, 1336 (1969).
- [25] M. S. Kalenkov, A. D. Zaikin, and L. S. Kuzmin, *Phys. Rev. Lett.* **109**, 147004 (2012).
- [26] A. Ozaeta, P. Virtanen, F. S. Bergeret, and T. T. Heikkilä, *Phys. Rev. Lett.* **112**, 057001 (2014).
- [27] N. R. Claughton and C. J. Lambert, *Phys. Rev. B* **53**, 6605 (1996).
- [28] J. Eom, C.-J. Chien, and V. Chandrasekhar, *Phys. Rev. Lett.* **81**, 437 (1998).
- [29] R. Seviour and A. F. Volkov, *Phys. Rev. B* **62**, R6116(R) (2000).
- [30] D. A. Dikin, S. Jung, and V. Chandrasekhar, *Phys. Rev. B* **65**, 012511 (2001).
- [31] A. Parsons, I. A. Sosnin, and V. T. Petrashov, *Phys. Rev. B* **67**, 140502 (2003).
- [32] A. F. Volkov and V. V. Pavlovskii, *Phys. Rev. B* **72**, 014529 (2005).
- [33] P. Virtanen and T. T. Heikkilä, *Phys. Rev. Lett.* **92**, 177004 (2004); *Appl. Phys. A* **89**, 625 (2007).
- [34] G. Srivastava, I. Sosnin, and V. T. Petrashov, *Phys. Rev. B* **72**, 012514 (2005).
- [35] Ph. Jacquod and R. S. Whitney, *Europhys. Lett.* **91**, 67009 (2010).
- [36] P. Machon, M. Eschrig, and W. Belzig, *Phys. Rev. Lett.* **110**, 047002 (2013); [arXiv:1402.7373](#).
- [37] E. P. Wigner, *Phys. Rev.* **98**, 145 (1955).
- [38] F. T. Smith, *Phys. Rev.* **118**, 349 (1960).
- [39] Y. V. Fyodorov and D. V. Savin, in Ref. [2].
- [40] S. A. van Langen, P. G. Silvestrov, and C. W. J. Beenakker, *Superlatt. Microstruct.* **23**, 691 (1998).
- [41] S. F. Godijn, S. Möller, H. Buhmann, L. W. Molenkamp, and S. A. van Langen, *Phys. Rev. Lett.* **82**, 2927 (1999).
- [42] Y. Takane and H. Ebisawa, *J. Phys. Soc. Jpn.* **61**, 2858 (1992).
- [43] The transformation (8) from electron-hole basis to Majorana basis assumes that there is an even number of modes at each contact. This number is odd if the superconductor has an unpaired Majorana mode, in which case we have to work in the Majorana basis from the very beginning.
- [44] M. R. Zirnbauer, in Ref. [2].
- [45] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, *New J. Phys.* **12**, 065010 (2010).
- [46] M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [47] X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
- [48] We recall the definition of a quaternion, $q = q_0\tau_0 + iq_1\tau_x + iq_2\tau_y + iq_3\tau_z$, with real coefficients q_n . A symplectic matrix U is unitary, $UU^\dagger = 1$, and satisfies $U^* = \tau_y U \tau_y$. Since $q^* = \tau_y q \tau_y$, a symplectic matrix is a unitary matrix with quaternion elements (just like an orthogonal matrix is a unitary matrix with real elements).
- [49] E. P. Wigner, *Ann. Math.* **53**, 36 (1951); **55**, 7 (1952).
- [50] P. W. Brouwer, K. M. Frahm, and C. W. J. Beenakker, *Waves in Random Media* **9**, 91 (1999).
- [51] Ref. [17] uses a modified definition of the time-delay matrix, with a symmetrized energy derivative, to ensure the independence of S_0 and Q also in the presence of time-reversal symmetry. This modification is not needed for the class C and D ensembles considered in the main text, so we can stay with the usual unsymmetrized definition (3). The more general case is considered in Appendix A 3.
- [52] In differential geometry, $\alpha + 1$ and β from Table I appear as the root multiplicities m_ℓ and m_o of the symmetric space of transfer matrices; see P. W. Brouwer, A. Furusaki, C. Mudry, and S. Ryu, [arXiv:cond-mat/0511622](#). An alternative algebraic interpretation, in terms of the number of degrees of freedom of matrix elements of the Hamiltonian, is given in Table III.
- [53] The mean level spacing δ_0 includes the electron-hole degree of freedom, so the single-electron Hamiltonian has mean level spacing $2\delta_0$. Since δ_0 is the mean spacing of *distinct* levels, the mean spacing of all levels is δ_0/d_E .
- [54] To understand why the degeneracies d_E and d_T of energy and transmission eigenvalues may differ in the presence of particle-hole symmetry, we recall that the Kramers degeneracy applies to Hermitian operators that commute with an antiunitary operator squaring to -1 . The Hamiltonian H anticommutes with the particle-hole conjugation operator $\mathcal{C}$, so the Kramers theorem does not apply. In contrast, the transmission matrix product $tt^\dagger$ commutes with $\mathcal{C}$, so when $\mathcal{C}^2 = -1$ its eigenvalues have a Kramers degeneracy.
- [55] M. L. Mehta, *Random Matrices* (Elsevier, Amsterdam, 2004).
- [56] V. L. Lyuboshits, *Phys. Lett. B* **72**, 41 (1977).
- [57] The result (22) for the average density of states in the Altland-Zirnbauer ensembles follows upon integration of the probability distribution (18). This can be achieved with the help of the general integral formulas of F. Mezzadri and N. J. Simm, *J. Math. Phys.* **52**, 103511 (2011), but it is easier to start from the zero- α equation $(Mt_0)^{-1}(\sum_n D_n) = 1/M$ and note that a nonzero α amounts to the substitution $M \mapsto M + 2\alpha/\beta$ on the right-hand side.
- [58] T. Guhr, A. Müller-Groeling, and H. A. Weidenmüller, *Phys. Rep.* **299**, 189 (1998).
- [59] C. W. J. Beenakker, *Rev. Mod. Phys.* **69**, 731 (1997); [arXiv:0904.1432](#).
- [60] E. Akkermans, A. Auerbach, J. E. Avron, and B. Shapiro, *Phys. Rev. Lett.* **66**, 76 (1991).
- [61] P. W. Brouwer, S. A. van Langen, K. M. Frahm, M. Büttiker, and C. W. J. Beenakker, *Phys. Rev. Lett.* **79**, 913 (1997). In this study of charging effects on normal quantum dots the canonical and grand-canonical averages are simply related by $\langle \cdots \rangle_{\text{canon}} = t_0^{-1} \langle \cdots \times \text{Tr} Q \rangle_{\text{grand-canon}}$. To include charging effects in the superconducting quantum dot considered here, the quasiparticle density of states $\text{Tr} Q$ should be replaced by the charge density $\text{Tr} \tau_z Q$.

- [62] A. Mackenzie and Y. Maeno, [Rev. Mod. Phys. **75**, 657 \(2003\)](#).
- [63] S. Lederer, W. Huang, E. Taylor, S. Raghu, and C. Kallin, [arXiv:1404.4637](#).
- [64] A. Furusaki, M. Matsumoto, and M. Sigrist, [Phys. Rev. B **64**, 054514 \(2001\)](#).
- [65] M. Stone and R. Roy, [Phys. Rev. B **69**, 184511 \(2004\)](#).
- [66] I. Serban, B. Béri, A. R. Akhmerov, and C. W. J. Beenakker, [Phys. Rev. Lett. **104**, 147001 \(2010\)](#).
- [67] J. A. Sauls, [Phys. Rev. B **84**, 214509 \(2011\)](#).
- [68] P. W. Brouwer, [Phys. Rev. B **51**, 16878 \(1995\)](#).
- [69] P. J. Forrester, *Log-Gases and Random Matrices* (Princeton University Press, Princeton, NJ, 2010).
- [70] A. M. Mathai, *Jacobians of Matrix Transformations and Functions of Matrix Argument* (World Scientific Publishing, Singapore, 1997).
- [71] R. A. Horn and J. R. Johnson, *Matrix Analysis* (Cambridge University Press, Cambridge, 1985).
- [72] I. C. Fulga, F. Hassler, A. R. Akhmerov, and C. W. J. Beenakker, [Phys. Rev. B **83**, 155429 \(2011\)](#).