



Universiteit
Leiden
The Netherlands

Safety-efficacy balance of S-ketamine and S-norketamine in acute and chronic pain

Noppers, I.M.

Citation

Noppers, I. M. (2011, September 7). *Safety-efficacy balance of S-ketamine and S-norketamine in acute and chronic pain*. Retrieved from <https://hdl.handle.net/1887/17811>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/17811>

Note: To cite this publication please use the final published version (if applicable).

Curriculum Vitae

Ingeborg Marieke Noppers was born on the 29th of October 1981 in Emmen, the Netherlands. She obtained her VWO diploma at the Hondsrug College in Emmen in June 2000. Later that year she entered Medical School at the Rijksuniversiteit Groningen. She received her Medical Degree in September 2006 and began working as a resident in the Intensive Care department of the University Medical Center Groningen. Early 2008 she was appointed as a PhD student at the department of Anesthesiology in the Leiden University Medical Center under the supervision of Prof. Dr. A Dahan and Dr. EY Sarton and started the investigations described in this thesis. She started her residency in Anesthesiology in October 2010. As of April 2011 she started her medical specialist training at the department of Anesthesiology in the Leiden University Medical Center (Chairman: Prof. Dr. LPHJ Aarts).

List of publications

Droogh JM, Noppers IM, Tulleken JE, Meertens JH, Ligtenberg JJ, Monteban-Kooistra WE and Zijlstra JG. Comment on Mackenzie and Woodhouse: C-reactive protein concentrations during bacteraemia: a comparison between patients with and without liver dysfunction. *Intensive Care Med* 2007; 33(3):561.

Noppers IM, Nieuwenhuis JA, Bremmer R, Huisman G, Wijkstra PJ and Zijlstra JG. Respiratoire insufficiëntie bij patiënten met progressieve neuromusculaire ziekte soms goed te behandelen door ademhalingsondersteuning. *Ned Tijdschr Geneesk* 2007; 151(34):1861-1864.

Noppers IM and Zijlstra JG. Undulating blood pressure. *Neth J Med* 2008; 66(5):212-214.

Morariu AM, Noppers IM, van Dorp ELA, Sigtermans MJ, Olofsen E and Aarts LPHJ. Take-home messages from 2007/2008. *Nederlands tijdschrift voor anesthesiologie* 2008; 20(4):27-31.

Noppers IM and Bauer MCR. Neuropathische pijn, het werkt op je zenuwen! *Modern Medicine* 2009; 33(5a):16-19.

Sigtermans M, Noppers I, Sarton E, Bauer M, Mooren R, Olofsen E and Dahan A. An observational study on the effect of S(+)-ketamine on chronic pain versus experimental acute pain in Complex Regional Pain Syndrome type 1 patients. *Eur J Pain* 2010; 14(3):302-307.

Noppers IM, Bauer MCR and Dahan A. S(+)-ketamine als analgeticum voor acute en chronische pijn. *A&I* 2009; 1(2):34-38.

Dahan A, Olofsen E, Sigtermans M, Noppers I, Nieters M, Aarts L, Bauer M and Sarton E. Population pharmacokinetic-pharmacodynamic modeling of ketamine-induced pain relief of chronic pain. *Eur J Pain* 2011; 15(3):258-267.

Noppers I, Nieters M, Aarts L, Smith T, Sarton E and Dahan A. Ketamine for the treatment of chronic non-cancer pain. *Expert Opin Pharmacother* 2010; 11(14):2417-2429.

List of publications

Niesters M, Dahan A, Swartjes M, Noppers I, Fillingim RB, Aarts L and Sarton EY. Effect of ketamine on endogenous pain modulation in healthy volunteers. *Pain* 2011; 152(3):656-663.

Noppers I, Olofsen E, Niesters M, Aarts L, Mooren R, Dahan A, Kharasch E and Sarton E. Effect of rifampicin on S-ketamine and S-norketamine plasma concentrations in healthy volunteers after intravenous S-ketamine administration. *Anesthesiology* 2011; 114(6):1435-1445.

Noppers IM, Niesters M, Aarts LPHJ, Bauer MCR, Drewes AM, Dahan A and Sarton EY. Drug-induced liver injury following a repeated course of ketamine treatment for chronic pain in CRPS type 1 patients: A report of 3 cases. *Pain, in press.*

Noppers I, Niesters M, Swartjes M, Bauer M, Aarts L, Geleijnse G, Mooren R, Dahan A and Sarton E. Absence of long-term analgesic effect from a short-term S-ketamine infusion on fibromyalgia pain: A randomized, prospective, double blind, active placebo-controlled trial. *Eur J Pain, in press.*