



Universiteit
Leiden
The Netherlands

Analysis of gene expression in the outer cell layers of Arabidopsis roots during lateral root development

Veth-Tello, L.M.

Citation

Veth-Tello, L. M. (2005, March 2). *Analysis of gene expression in the outer cell layers of Arabidopsis roots during lateral root development*. Retrieved from <https://hdl.handle.net/1887/2315>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/2315>

Note: To cite this publication please use the final published version (if applicable).

References

- Abel S, Oeller PW, Theologis A (1994) Early auxin-induced genes encode short-lived nuclear proteins. *Proc Natl Acad Sci* 91: 326-330
- Bak S, Tax FE, Feldmann KA, Galbraith DW, Feyereisen R (2001) CYP83B1, a cytochrome P450 at the metabolic branch point in auxin and indole glucosinolate biosynthesis in *Arabidopsis*. *Plant Cell* 13: 101-111
- Barlier I, Kowalczyk M, Marchant A, Ljung K, Bhalerao R, Bennett M, Sandberg G, Bellini C (2000) The *SUR2* gene of *Arabidopsis thaliana* encodes the cytochrome P450 CYP83B1, a modulator of auxin homeostasis. *Proc Natl Acad Sci* 97: 14819-14824
- Beaudoin N, Rothstein SJ (1997) Developmental regulation of two tomato lipoxygenase promoters in transgenic tobacco and tomato. *Plant Mol Biol* 33: 835-846
- Beeckman T, Burssens S, Inzé D (2001) The *peri-cell-cycle* in *Arabidopsis*. *J Exp Bot* 52:403-411
- Beers EP, Jones AM, Dickerman AW (2004) The S8 serine, C1A cysteine and A1 aspartic protease families in *Arabidopsis*. *Phytochemistry* 65: 43-58
- Bennett MJ, Marchant A, Green HG, May ST, Ward SP, Millner PA, Walker AR, Schulz B, Feldmann KA (1996) *Arabidopsis AUX1* gene: a permease-like regulator of root gravitropism. *Science* 273: 948-950
- Berger D, Altmann T (2000) A subtilisin-like serine protease involved in the regulation of stomatal density and distribution in *Arabidopsis thaliana*. *Genes Dev* 14: 1119-1131
- Bergmann DC (2004) Integrating signals in stomatal development. *Curr Opin Plant Biol* 7: 26-32

- Bhalerao RP, Eklöf J, Ljung K, Marchant A, Bennett M, Sandberg G (2002) Shoot-derived auxin is essential for early lateral root emergence in *Arabidopsis* seedlings. *Plant J* 29: 325-332
- Boerjan W, Cervera MT, Delarue M, Beeckman T, Dewitte W, Bellini C, Caboche M, Van Onckelen H, Van Montagu M, Inzé D (1995) *superroot*, a recessive mutation in *Arabidopsis*, confers auxin overproduction. *Plant Cell* 7: 1405-1419
- Bogacheva AM (1999) Plant subtilisins. *Biochemistry (Mosc)* 64: 287-293
- Bonifer C (1999) Long-distance chromatin mechanisms controlling tissue-specific gene locus activation. *Gene* 238: 277-289
- Cahn MD, Zobel RW, Bouldin DR (1989) Relationship between root elongation rate and diameter and duration of growth of lateral roots of maize. *Plant Soil* 119: 271-279.
- Carpita NC, Gibeaut DM (1993) Structural models of primary cell walls in flowering plants: consistency of molecular structure with the physical properties of the walls during growth. *Plant J* 3: 1-30
- Casimiro I, Marchant A, Bhalerao RP, Beeckman T, Dhooge S, Swarup R, Graham N, Inzé D, Sandberg G, Casero PJ, Bennett M (2001) Auxin transport promotes *Arabidopsis* lateral root initiation. *Plant Cell* 13: 843-852
- Casimiro I, Beeckman T, Graham N, Bhalerao R, Zhang H, Casero P, Sandberg G, Bennett MJ (2003) Dissecting *Arabidopsis* lateral root development. *Trends Plant Sci* 8:165-171
- Celenza JL, Grisafi PL, Fink GR (1995) A pathway for lateral root formation in *Arabidopsis thaliana*. *Genes Dev* 9: 2131-2142
- Chang C, Shockey JA (1999) The ethylene-response pathway: signal perception to gene regulation. *Curr Opin Plant Biol* 2: 352-358

- Charlton WA (1996) Lateral root initiation. In Y Waisel, A Eshel, U Kafkafi, eds, *Plant Roots: The hidden half*, Ed 2. M Dekker Inc, New-York, pp 149-173
- Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16: 735-743
- Cosgrove DJ, Van Volkenburgh E, Cleland RE (1984) Stress relaxation of cell walls and the yield threshold for growth: demonstration and measurement by micro-pressure probe and psychrometer. *Planta* 162: 46-54
- Cosgrove DJ (1998) Cell wall loosening by expansins. *Plant Physiol* 118: 333-339
- Creelman RA, Mullet JE (1997) Biosynthesis and action of jasmonates in plants. *Annu Rev Plant Physiol Plant Mol Biol* 48: 355-381
- Creelman RA, Rao MV (September 30, 2002) The oxylipin pathway in *Arabidopsis*. In CR Somerville, EM Meyerowitz, eds, *The Arabidopsis Book*, American Society of Plant Biologists, Rockville, MD, doi/10.1199/tab0012, <http://www.aspb.org/publications/arabidopsis/>
- Delarue M, Prinsen E, Van Onckelen H, Caboche M, Bellini C (1998) *Sur2* mutations of *Arabidopsis thaliana* define a new locus involved in the control of auxin homeostasis. *Plant J* 14: 603-611
- Delledonne M, Xia Y, Dixon RA, Lamb C (1998) Nitric oxide functions as a signal in plant disease resistance. *Nature* 394: 585-588
- Denault JB, Leduc R (1996) Furin/PACE/SPC1: a convertase involved in exocytic and endocytic processing of precursor proteins. *FEBS Lett* 379: 113-116
- Deshaies RJ (1999) SCF and Cullin/Ring H2-based ubiquitin ligases. *Annu Rev Cell Dev Biol* 15: 435-467

- De Smet I, Signora L, Beeckman T, Inzé D, Foyer CH, Zhang H (2003) An abscisic acid-sensitive checkpoint in lateral root development of *Arabidopsis*. *Plant J* 33: 543-555
- Deutch CE, Winicov I (1995) Post-transcriptional regulation of a salt-inducible alfalfa gene encoding a putative chimeric proline-rich cell wall protein. *Plant Mol Biol* 27: 411-418
- De Veylder L, de Almeida Engler J, Burssens S, Manevski A, Lescure B, Van Montagu M, Engler G, Inzé D (1999) A new D-type cyclin of *Arabidopsis thaliana* expressed during lateral root primordia formation. *Planta* 208: 453-462
- Devoto A, Nieto-Rostro M, Xie D, Ellis C, Harmston R, Patrick E, Davis J, Sherratt L, Coleman M, Turner JG (2002) COI1 links jasmonate signalling and fertility to the SCF ubiquitin-ligase complex in *Arabidopsis*. *Plant J* 32: 457-466
- Devoto A, Turner JG (2003) Regulation of jasmonate-mediated plant responses in *Arabidopsis*. *Ann Bot* 92: 329-337
- DiDonato RJ, Arbuckle E, Buker S, Sheets J, Tobar J, Totong R, Grisafi P, Fink GR, Celenza JL (2004) *Arabidopsis ALF4* encodes a nuclear-localized protein required for lateral root formation. *Plant J* 37: 340-353
- Ditta G, Stanfield S, Corbin D, Helinski DR (1980) Broad host range DNA cloning system for gram-negative bacteria: construction of a gene bank of *Rhizobium meliloti*. *Proc Natl Acad Sci* 77: 7347-7351
- Dolan L, Janmaat K, Willemsen V, Linstead P, Poethig S, Roberts K, Scheres B (1993) Cellular organization of the *Arabidopsis thaliana* root. *Development* 119: 71-84.
- Dolan L (1997) The role of ethylene in the development of plant form. *J Exp Bot* 48: 201-210
- Drew MC, Saker LR, Ashley TW (1973) Nutrient supply and the growth of the seminal root system in barley. I. The effect of nitrate concentration on the growth of axes and laterals. *J Exp Bot* 24: 1189-1202

- Dubrovsky JG, Rost TL, Colón-Carmona A, Doerner P (2001) Early primordium morphogenesis during lateral root initiation in *Arabidopsis thaliana*. *Planta* 214: 30-36
- Ezaki B, Yamamoto Y, Matsumoto H (1995) Cloning and sequencing of the cDNAs induced by aluminium treatment and P_i starvation in cultured tobacco cells. *Physiol Plant* 93: 11-18
- Ezaki B, Tsugita S, Matsumoto H (1996) Expression of a moderately anionic peroxidase is induced by aluminum treatment in tobacco cells: possible involvement of peroxidase isozymes in aluminum ion stress. *Physiol Plant* 96: 21-28
- Farmer EE, Caldelari D, Pearce G, Walker-Simmons MK, Ryan CA (1994) Diethylthiocarbamic acid inhibits the octadecanoid signaling pathway for the wound induction of proteinase inhibitors in tomato leaves. *Plant Physiol* 106: 337-342
- Feng S, Ma L, Wang X, Xie D, Dinesh-Kumar SP, Wei N (2003) The COP9 signalosome interacts physically with SCF^{COI1} and modulates jasmonate response. *Plant Cell* 15: 1083-1094
- Ferreira PCG, Hemerly AS, de Almeida Engler J, Van Montagu M, Engler G, Inzé D (1994) Developmental expression of the *Arabidopsis* cyclin gene *cyc1At*. *Plant Cell* 6: 1763-1774
- Feys BJB, Benedetti CE, Penfold CN, Turner JG (1994) *Arabidopsis* mutants selected for resistance to the phytotoxin coronatine are male sterile, insensitive to methyl jasmonate, and resistant to a bacterial pathogen. *Plant Cell* 6: 751-759
- Forde BG, Lorenzo H (2001) The nutritional control of root development. *Plant and Soil* 232: 51-68
- Forde BG (2002) Local and long-range signaling pathways regulating plant responses to nitrate. *Annu Rev Plant Biol* 53: 203-224

- Fowler TJ, Bernhardt C, Tierney ML (1999) Characterization and expression of four proline-rich cell wall protein genes in *Arabidopsis* encoding two distinct subsets of multiple domain proteins. *Plant Physiol* 121: 1081-1091
- Fukaki H, Tameda S, Masuda H, Tasaka M (2002) Lateral root formation is blocked by a gain-of-function mutation in the *SOLITARY-ROOT/IAA14* gene of *Arabidopsis*. *Plant J* 29: 153-168
- Gil P, Dewey E, Friml J, Zhao Y, Snowden KC, Putterill J, Palme K, Estelle M, Chory J (2001) BIG: a calossin-like protein required for polar auxin transport in *Arabidopsis*. *Genes Dev* 15: 1985-1997
- Gleave AP (1992) A versatile binary vector system with a T-DNA organizational structure conducive to efficient integration of cloned DNA into the plant genome. *Plant Mol Biol* 20: 1203-1207
- Golldack D, Vera P, Dietz KJ (2003) Expression of subtilisin-like serine proteases in *Arabidopsis thaliana* is cell-specific and responds to jasmonic acid and heavy metals with developmental differences. *Physiol Plant* 118: 64-73
- Goodwin W, Pallas JA, Jenkins GI (1996) Transcripts of a gene encoding a putative cell wall-plasma membrane linker protein are specifically cold-induced in *Brassica napus*. *Plant Mol Biol* 31: 771-781
- Gray WM, del Pozo JC, Walker L, Hobbie L, Risseuw E, Banks T, Crosby WL, Yang M, Ma H, Estelle M (1999) Identification of an SCF ubiquitin-ligase complex required for auxin response in *Arabidopsis thaliana*. *Genes Dev* 13: 1678-1691
- von Groll U, Berger D, Altmann T (2002) The subtilisin-like serine protease SDD1 mediates cell-to-cell signaling during *Arabidopsis* stomatal development. *Plant Cell* 14: 1527-1539
- Guo Z, Ohta Y (1994) Effect of ethylene biosynthesis on the accumulation of 6-methoxymellein induced by elicitors in carrot cells. *J Plant Physiol* 144: 700-704

- Hagen G, Guilfoyle T (2002) Auxin-responsive gene expression: genes, promoters and regulatory factors. *Plant Mol Biol* 49: 373-385
- Hellmann H, Estelle M (2002) Plant development: Regulation by protein degradation. *Science* 297: 793-797
- Himanen K, Boucheron E, Vanneste S, de Almeida Engler J, Inzé D, Beeckman T (2002) Auxin-mediated cell cycle activation during early lateral root initiation. *Plant Cell* 14: 2339-2351
- Holk A, Klumpp L, Scherer GFE (2002) A cell wall protein down-regulated by auxin suppresses cell expansion in *Daucus carota* (L.). *Plant Mol Biol* 50: 295-305
- Hood EE, Gelvin SB, Melchers LS, Hoekema A (1993) New *Agrobacterium* helper plasmids for gene transfer to plants. *Transgenic Res* 2: 208-218
- Irving HR, Dyson G, McConchie R, Parish RW, Gehring CA (1999) Effects of exogenously applied jasmonates on growth and intracellular pH in maize coleoptile segments. *J Plant Growth Regul* 18: 93-100
- Janzik I, Macheroux P, Amrhein N, Schaller A (2000) LeSBT1, a subtilase from tomato plants: Overexpression in insect cells, purification, and characterization. *J. Biol. Chem.* 275: 5193-5199
- Jefferson RA, Kavanagh TA, Bevan MW (1987) GUS fusions: β -glucuronidase as a sensitive and versatile fusion marker in higher plants. *EMBO J* 6: 3901-3907
- Kayes JM, Clark SE (1998) CLAVATA2, a regulator of meristem and organ development in Arabidopsis. *Development* 125: 3843-3851
- Kim J, Harter K, Theologis A (1997) Protein-protein interactions among the Aux/IAA proteins. *Proc Natl Acad Sci* 94: 11786-11791

- King JJ, Stimart DR, Fisher RH, Bleecker AB (1995) A mutation altering auxin homeostasis and plant morphology in *Arabidopsis*. *Plant Cell* 7: 2023-2037
- Kisu Y, Harada Y, Goto M, Esaka M (1997) Cloning of the pumpkin ascorbate oxidase gene and analysis of a *cis*-acting region involved in induction by auxin. *Plant Cell Physiol* 38: 631-637
- Knee EM, Hangarter RP (1996) Differential IAA dose response relations of the *axr1* and *axr2* mutants of *Arabidopsis*. *Physiol Plant* 98: 320-324
- Konishi M, Sugiyama M (2003) Genetic analysis of adventitious root formation with a novel series of temperature-sensitive mutants of *Arabidopsis thaliana*. *Development* 130: 5637-5647
- Krysan PJ, Young JC, Sussman MR (1999) T-DNA as an insertional mutagen in *Arabidopsis*. *Plant Cell* 11: 2283-2290
- Lamb C, Dixon RA (1997) The oxidative burst in plant disease resistance. *Annu Rev Plant Physiol Plant Mol Biol* 48: 251-275
- Laskowski MJ, Williams ME, Nusbaum HC, Sussex IM (1995) Formation of lateral root meristems is a two-stage process. *Development* 121: 3303-3310
- Lenhard M, Laux T (2003) Stem cell homeostasis in the *Arabidopsis* shoot meristem is regulated by intercellular movement of CLAVATA3 and its sequestration by CLAVATA1. *Development* 130: 3163-3173.
- Levine A, Tenhaken R, Dixon R, Lamb C (1994) H₂O₂ from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell* 79: 583-593
- Leyser HMO, Lincoln CA, Timpte C, Lammer D, Turner J, Estelle M (1993) *Arabidopsis* auxin-resistance gene *AXR1* encodes a protein related to ubiquitin-activating enzyme E1. *Nature* 364: 161-164

- Lin X, et al., (1999) Sequence analysis of chromosome 2 of the plant *Arabidopsis thaliana*. *Nature* 402: 761-768
- Lincoln C, Britton JH, Estelle M (1990) Growth and development of the *axr1* mutants of *Arabidopsis*. *Plant Cell* 2: 1071-1080
- Linkohr B, Williamson LC, Fitter AH, Leyser HMO (2002) Nitrate and phosphate availability and distribution have different effects on root system architecture of *Arabidopsis*. *Plant J* 29: 751-760
- López-Bucio J, Hernández-Abreu E, Sánchez-Calderón L, Nieto-Jacobo MF, Simpson J, Herrera-Estrella L (2002) Phosphate availability alters architecture and causes changes in hormone sensitivity in the *Arabidopsis* root system. *Plant Physiol* 129: 244-256
- Lyapina S, Cope G, Shevchenko A, Serino G, Tsuge T, Zhou C, Wolf DA, Wei N, Shevchenko A, Deshaies RJ (2001) Promotion of NEDD8-CUL1 conjugate cleavage by COP9 signalosome. *Science* 292: 1382-1385
- Malamy JE, Benfey PN (1997) Organization and cell differentiation in lateral roots of *Arabidopsis thaliana*. *Development* 124: 33-44
- Malamy JE, Ryan KS (2001) Environmental regulation of lateral root initiation in *Arabidopsis*. *Plant Physiol* 127: 899-909
- Marchant A, Bhalerao R, Casimiro I, Eklöf J, Casero PJ, Bennett M, Sandberg G (2002) AUX1 promotes lateral root formation by facilitating indole-3-acetic acid distribution between sink and source tissues in the *Arabidopsis* seedling. *Plant Cell* 14: 589-597
- Mayer K, et al (1999) Sequence analysis of chromosome 4 of the plant *Arabidopsis thaliana*. *Nature* 402: 769-777
- Menke FLH, Champion A, Kijne JW, Memelink J (1999a) A novel jasmonate- and elicitor-responsive element in the periwinkle secondary metabolite biosynthetic gene *Str*

interacts with a jasmonate- and elicitor-inducible AP2-domain transcription factor, ORCA2. *EMBO J* 18: 4455-4463

Menke FLH, Parchmann S, Mueller MJ, Kijne JW, Memelink J (1999b) Involvement of the octadecanoid pathway and protein phosphorylation in fungal elicitor-induced expression of terpenoid indole alkaloid biosynthetic genes in *Catharanthus roseus*. *Plant Physiol* 119: 1289-1296

Mortlock DP, Guenther C, Kingsley DM (2003) A general approach for identifying distant regulatory elements applied to the *Gdf6* gene. *Genome Research* 13: 2069-2081

Neuteboom LW, Ng JMY, Kuyper M, Clijdesdale OR, Hooykaas PJJ, van der Zaal BJ (1999a) Isolation and characterization of cDNA clones corresponding with mRNAs that accumulate during auxin-induced lateral root formation. *Plant Mol Biol* 39: 273-287

Neuteboom LW, Veth-Tello LM, Clijdesdale OR, Hooykaas PJJ, van der Zaal BJ (1999b) A novel subtilisin-like protease gene from *Arabidopsis thaliana* is expressed at sites of lateral root emergence. *DNA Research* 6: 13-19

Neuteboom LW (2000) Gene expression during lateral root development. PhD thesis. Leiden University. The Netherlands

Ozawa S, Yasutani I, Fukuda H, Komamine A, Sugiyama M (1998) Organogenic responses in tissue culture of *srd* mutants of *Arabidopsis thaliana*. *Development* 125: 135-142

Pages L (1995) Growth patterns of the lateral roots of young oak (*Quercus robur*) tree seedlings: relationship with apical diameter. *New Phytol* 130: 503-509.

Pagnussat GC, Simontacchi M, Puntarulo S, Lamattina L (2002) Nitric oxide is required for root organogenesis. *Plant Physiol* 129: 954-956

Parsons R, Sunley R (2001) Nitrogen nutrition and the role of root-shoot nitrogen signalling particularly in symbiotic systems. *J Exp Bot* 52: 435-443

- Pasquali G, Ouwerkerk PBF, Memelink J (1994) Versatile transformation vectors to assay the promoter activity of DNA elements in plants. *Gene* 149: 373-374
- Peña-Cortés H, Albrecht T, Prat S, Weiler EW, Willmitzer L (1993) Aspirin prevents wound-induced gene expression in tomato leaves by blocking jasmonic acid biosynthesis. *Planta* 191: 123-128
- del Pozo JC, Timpte C, Tan S, Callis J, Estelle M (1998) The ubiquitin-related protein RUB1 and auxin response in *Arabidopsis*. *Science* 280: 1760-1763
- del Pozo JC, Estelle M (1999) The *Arabidopsis* cullin AtCUL1 is modified by the ubiquitin-related protein RUB1. *Proc Natl Acad Sci* 96: 15342-15347
- del Pozo JC, Dharmasiri S, Hellmann H, Walker L, Gray WM, Estelle M (2002) AXR1-ECR1-dependent conjugation of RUB1 to the *Arabidopsis* cullin AtCUL1 is required for auxin response. *Plant Cell* 14: 421-433
- Preuss D, Copenhaver G, Keit K. September 21, 2000. Plant chromosome compositions and methods. Patent: WO 0055325-A 272; The University of Chicago (US)
- Rahman A, Amakawa T, Goto N, Tsurumi S (2001) Auxin is a positive regulator for ethylene-mediated response in the growth of *Arabidopsis* roots. *Plant Cell Physiol* 42: 301-307
- Reed RC, Brady SR, Muday GK (1998) Inhibition of auxin movement from the shoot into the root inhibits lateral root development in *Arabidopsis*. *Plant Physiol* 118: 1369-1378
- Rhee SY, Beavis W, Berardini TZ, Chen G, Dixon D, Doyle A, Garcia-Hernandez M, Huala E, Lander G, Montoya M, Miller N, Mueller LA, Mundodi S, Reiser L, Tacklind J, Weems DC, Wu Y, Xu I, Yoo D, Yoon J, Zhang P (2003) The *Arabidopsis* Information Resource (TAIR): a model organism database providing a centralized, curated gateway to *Arabidopsis* biology, research materials and community. *Nucleic Acids Res*

31:224

- Ribeiro A, Akkermans ADL, van Kammen A, Bisseling T, Pawlowski K (1995) A nodule-specific gene encoding a subtilisin-like protease is expressed in early stages of actinorhizal nodule development. *Plant Cell* 7: 785-794
- Richards KD, Gardner RC (1995) pEARLI 1: An Arabidopsis member of a conserved gene family (accession No. L43080) (PGR 95-099). *Plant Physiol* 109: 1497
- Richards KD, Schott EJ, Sharma YK, Davis KR, Gardner RC (1998) Aluminum induces oxidative stress genes in *Arabidopsis thaliana*. *Plant Physiol* 116: 409-418
- Rouster J, Leah R, Mundy J, Cameron-Mills V (1997) Identification of a methyl jasmonate-responsive region in the promoter of a lipoxygenase 1 gene expressed in barley grain. *Plant J* 11: 513-523
- Ruegger M, Dewey E, Hobbie L, Brown D, Bernasconi P, Turner J, Muday G, Estelle M (1997) Reduced naphthylphthalamic acid binding in the *tir3* mutant of Arabidopsis is associated with a reduction in polar auxin transport and diverse morphological defects. *Plant Cell* 9: 745-757
- Ruegger M, Dewey E, Gray WM, Hobbie L, Turner J, Estelle M (1998) The TIR1 protein of *Arabidopsis* functions in auxin response and is related to human SKP2 and yeast Grr1p. *Genes Dev* 12: 198-207
- Sharma VK, Ramirez J, Fletcher JC (2003) The Arabidopsis *CLV3*-like (*CLE*) genes are expressed in diverse tissues and encode secreted proteins. *Plant Mol Biol* 51: 415-425
- Scholz A, Truss M, Beato M (1999) Hormone-dependent recruitment of NF-Y to the uteroglobin gene enhancer associated with chromatin remodeling in rabbit endometrial epithelium. *J Biol Chem* 274: 4017-4026

- Schwechheimer C, Serino G, Callis J, Crosby WM, Lyapina S, Deshaies RJ, Gray WM, Estelle M, Deng X-W (2001) Interactions of the COP9 signalosome with the E3 ubiquitin-ligase SCF^{TIR1} in mediating auxin response. *Science* 292: 1379-1382
- Schwechheimer C, Serino G, Deng X-W (2002) Multiple ubiquitin ligase-mediated processes required COP9 signalosome and AXR1 function. *Plant Cell* 14: 2553-2563
- Seidah NG, Chretien M (1999) Proprotein and prohormone convertases: a family of subtilases generating diverse bioactive polypeptides. *Brain Res* 848: 45-62
- Serino G, Deng X-W (2003) The COP9 signalosome: regulating plant development through the control of proteolysis. *Ann Rev Plant Biol* 54: 165-182
- Serna L, Fenoll C (2002) Reinforcing the idea of signalling in the stomatal pathway. *Trends Genet* 18: 597-600
- Shopfer P (2001) Hydroxyl radical-induced cell-wall loosening *in vitro* and *in vivo*: implications for the control of elongation growth. *Plant J* 28: 679-688
- Showalter AM, (1993) Structure and function of plant cell wall proteins. *Plant Cell* 5: 9-23
- Signora L, Smet ID, Foyer CH, Zhang H (2001) ABA plays a central role in mediating the regulatory effects of nitrate on root branching in *Arabidopsis*. *Plant J* 28: 655-662
- van Slogteren GMS, Hoge JHC, Hooykaas PJJ, Schilperoort RA (1983) Clonal analysis of heterogenous crown gall tumor tissue induced by wild type and shooter mutant strains of *Agrobacterium tumefaciens* - expression of T-DNA genes. *Plant Mol Biol* 2: 321-333
- Struhl K (1998) Histone acetylation and transcriptional regulatory mechanism. *Genes Dev.* 12: 599-606
- Suttle JC (1988) Effect of ethylene treatment on polar IAA transport, net IAA uptake and specific binding of *N*-1-naphthylphthalamic acid in tissues and microsomes isolated from etiolated pea epicotyls. *Plant Physiol* 88: 795-799

- Swarup R, Friml J, Marchant A, Ljung K, Sandberg G, Palme K, Bennett M (2001) Localization of the auxin permease AUX1 suggests two functionally distinct hormone transport pathways operate in the *Arabidopsis* root apex. *Genes Dev* 15: 2648-2653
- Tanaka H, Onouchi H, Kondo M, Hara-Nishimura I, Nishimura M, Machida C, Machida Y (2001) A subtilisin-like serine protease is required for epidermal surface formation in *Arabidopsis* embryos and juvenile plants. *Development* 128: 4681-4689
- Tang Z, Sadka A, Morishige DT, Mullet JE (2001) Homeodomain leucine zipper proteins bind to the phosphate response domain of the soybean *VspB* tripartite promoter. *Plant Physiol* 125: 797-809
- The Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796-815
- Timpte C, Lincoln C, Pickett FB, Turner J, Estelle M (1995) The *AXR1* and *AUX1* genes of *Arabidopsis* function in separate auxin-response pathways. *Plant J* 8: 561-569
- Turner JG, Ellis C, Devoto A (2002) The jasmonate signal pathway. *Plant Cell* S153-S164
- Ulmasov T, Murfett J, Hagen G, Guilfoyle T (1997) Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. *Plant Cell* 9: 1963-1971
- Vera P, Lamb C, Doerner PW (1994) Cell-cycle regulation of hydroxyproline-rich glycoprotein *HRGPnt3* gene expression during the initiation of lateral root meristems. *Plant J* 6: 717-727
- Vergunst AC, de Waal EC, Hooykaas PJJ (1998) Root transformation by *Agrobacterium tumefaciens*. In J Martinez Zapaler, J Salinas, eds, *Methods in Molecular Biology*, Vol 82: *Arabidopsis* Protocols, Humana Press Inc, Totowa NJ, pp 227-244

- Ward SP, Estelle M (2001) Auxin signaling involves regulated protein degradation by the ubiquitin-proteasome pathway. *J Plant Growth Regul* 20: 265-273
- Weijers D, Franke-van Dijk M, Vencken RJ, Quint A, Hooykaas P, Offringa R (2001) An *Arabidopsis* Minute-like phenotype caused by a semi-dominant mutation in a *RIBOSOMAL PROTEIN S5* gene. *Development* 128: 4289-4299
- Xiang C, Miao Z-H, Lam E (1996) Coordinated activation of *as-1*-type elements and a tobacco glutathione *S*-transferase gene by auxins, salicylic acid, methyl-jasmonate and hydrogen peroxide. *Plant Mol Biol* 32: 415-426
- Xie DX, Feys BF, James S, Nieto-Rostro M, Turner JG (1998) *COI1*: an *Arabidopsis* gene required for jasmonate-regulated defense and fertility. *Science* 280: 1091-1094
- Xie Q, Frugis G, Colgan D, Chua N-H (2000) *Arabidopsis* NAC1 transduces auxin signal downstream of TIR1 to promote lateral root development. *Genes Dev* 14: 3024-3036
- Xie Q, Guo H-S, Dallman G, Fang S, Weissman AM, Chua N-H (2002) SINAT5 promotes ubiquitin-related degradation of NAC1 to attenuate auxin signals. *Nature* 419: 167-170
- Xu L, Liu F, Lechner E, Genschik P, Crosby WL, Ma H, Peng W, Huang D, Xie D (2002) The SCF^{COI1} ubiquitin-ligase complexes are required for jasmonate response in *Arabidopsis*. *Plant Cell* 14: 1919-1935
- Yang SF, Hoffman NE (1984) Ethylene biosynthesis and its regulation in higher plants. *Ann Rev Plant Physiol* 35: 155-189
- Yang M, Sack FD (1995) The *too many mouths* and *four lips* mutations affect stomatal production in *Arabidopsis*. *Plant Cell* 7: 2227-2239
- Yazaki K, Matsuoka H, Shimomura K, Bechthold A, Sato F (2001) A novel dark-inducible protein, LeDI-2, and its involvement in root-specific secondary metabolism in *Lithospermum erythrorhizon*. *Plant Physiol* 125: 1831-1841

- Zenser N, Ellsmore A, Leasure C, Callis J (2001) Auxin modulates the degradation rate of Aux/IAA proteins. *Proc Natl Acad Sci* 98: 11795-11800
- Zenser N, Dreher KA, Edwards SR, Callis J (2003) Acceleration of Aux/IAA proteolysis is specific for auxin and independent of *AXR1*. *Plant J* 35: 285-294
- Zhang H, Forde BG (1998) An *Arabidopsis* MADS box gene that controls nutrient-induced changes in root architecture. *Science* 279: 407-409
- Zhang H, Jennings A, Barlow P, Forde BG (1999) Dual pathways for regulation of root branching by nitrate. *Proc Natl Acad Sci* 96: 6529-6534
- Zhang H, Forde BG (2000) Regulation of *Arabidopsis* root development by nitrate availability. *J Exp Bot* 51: 51-59
- Zhao C, Johnson BJ, Kositsup B, Beers EP (2000) Exploiting secondary growth in *Arabidopsis*. Construction of xylem and bark cDNA libraries and cloning of three xylem endopeptidases. *Plant Physiol* 123: 1185-1196
- Zizamalova E, Napier RM (2003) Points of regulation for auxin action. *Plant Cell Rep* 21: 625-634