



Universiteit
Leiden
The Netherlands

Predictors of clinical outcome in total hip and knee replacement : a methodological appraisal of implants and patient factors

Keurentjes, J.C.

Citation

Keurentjes, J. C. (2014, September 30). *Predictors of clinical outcome in total hip and knee replacement : a methodological appraisal of implants and patient factors*. Retrieved from <https://hdl.handle.net/1887/28958>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/28958>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/28958> holds various files of this Leiden University dissertation

Author: Keurentjes, Johan Christiaan

Title: Predictors of clinical outcome in total hip and knee replacement : a methodological appraisal of implants and patient factors

Issue Date: 2014-09-30

Bibliography

- 1 Ethgen O, Bruyère O, Richy F, Dardennes C, Reginster JY. Health-related quality of life in total hip and total knee arthroplasty. A qualitative and systematic review of the literature. *J Bone Joint Surg Am.* 2004;86-A(5):963–74.
- 2 Havelin LI, Fenstad AM, Salomonsson R, Mehnert F, Furnes O, Overgaard Sr, Pedersen AB, Herberts P, Kärrholm J, Garellick G. The Nordic Arthroplasty Register Association: a unique collaboration between 3 national hip arthroplasty registries with 280,201 THRs. *Acta Orthop.* 2009;80(4):393–401.
- 3 Robertsson O, Bizjajeva S, Fenstad AM, Furnes O, Lidgren L, Mehnert F, Odgaard A, Pedersen AB, Havelin LI. Knee arthroplasty in Denmark, Norway and Sweden. *Acta Orthop.* 2010;81(1):82–9.
- 4 Chang RW, Pellisier JM, Hazen GB. A cost-effectiveness analysis of total hip arthroplasty for osteoarthritis of the hip. *JAMA.* 1996;275(11):858–65.
- 5 O’Shea K, Bale E, Murray P. Cost analysis of primary total hip replacement. *Ir Med J.* 2002;95(6):177–180.
- 6 Räsänen P, Paavolainen P, Sintonen H, Koivisto AM, Blom M, Ryynänen OP, Roine RP. Effectiveness of hip or knee replacement surgery in terms of quality-adjusted life years and costs. *Acta Orthop.* 2007;78(1):108–115.
- 7 Hamilton DF, Clement ND, Burnett R, Patton JT, Moran M, Howie CR, Simpson AHRW, Gaston P. Do modern total knee replacements offer better value for money? A health economic analysis. *Int Orthop.* 2013;37(11):2147–52.
- 8 LROI. Annual reports. Insight into quality of orthopaedic care in the Netherlands.
URL <http://www.lroi.nl/en/annual-reports>
- 9 Otten R, van Roermund PM, Picavet HSJ. [Trends in the number of knee and hip arthroplasties: considerably more knee and hip prostheses due to osteoarthritis in 2030]. *Ned Tijdschr Geneeskdl.* 2010;154:A1534.
- 10 Advice by the Board of the NOV on the use of primary hip prostheses.
URL <http://www.orthopeden.org/>

- 11 Beswick AD, Wylde V, Gooberman-Hill R, Blom A, Dieppe P. What proportion of patients report long-term pain after total hip or knee replacement for osteoarthritis? A systematic review of prospective studies in unselected patients. *BMJ open*. 2012;2(1):e000435.
- 12 Gandhi R, Davey JR, Mahomed NN. Predicting patient dissatisfaction following joint replacement surgery. *J Rheumatol*. 2008;35(12):2415–8.
- 13 Noble PC, Conditt MA, Cook KF, Mathis KB. The John Insall Award: Patient expectations affect satisfaction with total knee arthroplasty. *Clin Orthop*. 2006;452:35–43.
- 14 Brokelman RBG, van Loon CJM, Rijnberg WJ. Patient versus surgeon satisfaction after total hip arthroplasty. *J Bone Joint Surg Br*. 2003;85-B(4):495–498.
- 15 Nilsdotter AK, Toksvig-Larsen S, Roos EM. Knee arthroplasty: are patients' expectations fulfilled? A prospective study of pain and function in 102 patients with 5-year follow-up. *Acta Orthop*. 2009;80(1):55–61.
- 16 Lingard EA, Sledge CB, Learmonth ID. Patient expectations regarding total knee arthroplasty: differences among the United States, United Kingdom, and Australia. *J Bone Joint Surg Am*. 2006;88(6):1201–1207.
- 17 Garellick G, Malchau H, Herberts P. Survival of hip replacements. A comparison of a randomized trial and a registry. *Clin Orthop Relat Res*. 2000;(375):157–167.
- 18 Baker PN, van der Meulen JH, Lewsey J, Gregg PJ. The role of pain and function in determining patient satisfaction after total knee replacement. Data from the National Joint Registry for England and Wales. *J Bone Joint Surg Br*. 2007;89(7):893–900.
- 19 Mancuso CA, Salvati EA, Johanson NA, Peterson MG, Charlson ME. Patients' expectations and satisfaction with total hip arthroplasty. *J Arthroplasty*. 1997;12(4):387–96.
- 20 Vissers MM, de Groot IB, Reijman M, Bussmann JB, Stam HJ, Verhaar JA. Functional capacity and actual daily activity do not contribute to patient satisfaction after total knee arthroplasty. *BMC Musculoskelet Disord*. 2010;11:121.
- 21 Keurentjes JC, Blane D, Bartley M, Keurentjes JJB, Fiocco M, Nelissen RGR. Socio-Economic Position has no effect on Improvement in Health-Related Quality of Life and Patient Satisfaction in Total Hip and Knee Replacement: A Cohort Study. *PLoS ONE*. 2013;8(e56785):e56785.
- 22 Kaplan E, Meier P. Nonparametric Estimation from Incomplete Observations. *J Am Stat Assoc*. 1958;53(282):457–481.
- 23 Van Gelder M, Pijpe A. E - epidemiology: a comprehensive update. *OA Epidemiology*. 2013;1(2):1.

- 24 Saleh KJ, Radosevich DM, Kassim RA, Moussa M, Dykes D, Bottolfson H, Gioe TJ, Robinson H. Comparison of commonly used orthopaedic outcome measures using palm-top computers and paper surveys. *J Orthop Res.* 2002;20:1146–1151.
- 25 So-Osman C, Nelissen R, Te Slaa R, Coene L, Brand R, Brand A. A randomized comparison of transfusion triggers in elective orthopaedic surgery using leucocyte-depleted red blood cells. *Vox Sang.* 2010;98(1):56–64.
- 26 So-Osman C, Nelissen R, Brand R, Faber F, Slaa RT, Stiggelbout A, Brand A. The impact of a restrictive transfusion trigger on post-operative complication rate and well-being following elective orthopaedic surgery: a post-hoc analysis of a randomised study. *Blood Transfus.* 2013;11(2):289–95.
- 27 Keurentjes JC, Fiocco M, So-Osman C, Onstenk R, Koopman-Van Gemert AWMM, Pöll RG, Kroon HM, Vliet Vlieland TPM, Nelissen RG. Patients with Severe Radiographic Osteoarthritis Have a Better Prognosis in Physical Functioning after Hip and Knee Replacement: A Cohort-Study. *PLoS ONE.* 2013;8(4):e59500.
- 28 Keurentjes JC, Fiocco M, Nelissen RG. Willingness to undergo surgery again validated clinically important differences in health-related quality of life after total hip replacement or total knee replacement surgery. *J Clin Epidemiol.* 2014;67(1):114–20.
- 29 Murray DW, Carr AJ, Bulstrode CJ. Which primary total hip replacement? *J Bone Joint Surg Br.* 1996;78(3):510.
- 30 Bozic KJ, Durbakula S, Berry DJ, Naessens JM, Rappaport K, Cisternas M, Saleh KJ, Rubash HE. Differences in patient and procedure characteristics and hospital resource use in primary and revision total joint arthroplasty: a multicenter study. *J Arthroplasty.* 2005;20(7 Suppl 3):17–25.
- 31 Bozic KJ, Katz P, Cisternas M, Ono L, Ries MD, Showstack J. Hospital resource utilization for primary and revision total hip arthroplasty. *J Bone Joint Surg Am.* 2005;87(3):570–576.
- 32 Ong KL, Mowat FS, Chan N, Lau E, Halpern MT, Kurtz SM. Economic burden of revision hip and knee arthroplasty in Medicare enrollees. *Clin Orthop.* 2006;446(446):22–28.
- 33 Vanhegan IS, Malik AK, Jayakumar P, UI Islam S, Haddad FS. A financial analysis of revision hip arthroplasty: the economic burden in relation to the national tariff. *J Bone Joint Surg Br.* 2012;94(5):619–23.
- 34 Schreurs BW, Keurentjes JC, Gardeniers JWM, Verdonschot N, Slooff TJJH, Veth RPH. Acetabular revision with impacted morsellised cancellous bone grafting and a cemented acetabular component: a 20- to 25-year follow-up. *J Bone Joint Surg Br.* 2009;91(9):1148–53.

- 35 National Institute for Health and Care Excellence: Review of TA2 hip disease replacement prostheses and TA44 hip disease metal on metal resurfacing.
URL <http://publications.nice.org.uk/>
- 36 Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535–b2535.
- 37 Pijls BG, Dekkers OM, Middeldorp S, Valstar ER, van der Heide HJL, Van der Linden-Van der Zwaag HMJ, Nelissen RGHH. AQUILA: assessment of quality in lower limb arthroplasty. An expert Delphi consensus for total knee and total hip arthroplasty. *BMC Musculoskelet Disord*. 2011;12:173.
- 38 Thiessen Philbrook H, Barrowman N, Garg AX. Imputing variance estimates do not alter the conclusions of a meta-analysis with continuous outcomes: a case study of changes in renal function after living kidney donation. *J Clin Epidemiol*. 2007;60(3):228–40.
- 39 Furukawa TA, Barbui C, Cipriani A, Brambilla P, Watanabe N. Imputing missing standard deviations in meta-analyses can provide accurate results. *J Clin Epidemiol*. 2006;59(1):7–10.
- 40 Fiocco M, Putter H, van Houwelingen JC. Meta-analysis of pairs of survival curves under heterogeneity: a Poisson correlated gamma-frailty approach. *Stat Med*. 2009; 28(30):3782–97.
- 41 Putter H, Fiocco M, Stijnen T. Meta-analysis of diagnostic test accuracy studies with multiple thresholds using survival methods. *Biom J*. 2010;52(1):95–110.
- 42 Pijls BG, Valstar ER, Nouta KA, Plevier JW, Fiocco M, Middeldorp S, Nelissen RG. Early migration of tibial components is associated with late revision: a systematic review and meta-analysis of 21,000 knee arthroplasties. *Acta Orthop*. 2012;83(6):614–24.
- 43 *R: A Language and Environment for Statistical Computing*.
URL <http://www.r-project.org/>
- 44 Svenska Höftprotesregistret.
URL <http://www.shpr.se/en/Publications/DocumentsReports.aspx>
- 45 Eskelinen A, Remes V, Helenius I, Pulkkinen P, Nevalainen J, Paavolainen P. Uncemented total hip arthroplasty for primary osteoarthritis in young patients: a mid-to long-term follow-up study from the Finnish Arthroplasty Register. *Acta Orthop*. 2006;77(1):57–70.
- 46 Rogers A, Kulkarni R, Downes EM. The ABG hydroxyapatite-coated hip prosthesis: one hundred consecutive operations with average 6-year follow-up. *J Arthroplasty*. 2003; 18(5):619–25.
- 47 Hamilton WG, Calendine CL, Beykirch SE, Hopper RH, Engh CA. Acetabular fixation options: first-generation modular cup curtain calls and caveats. *J Arthroplasty*. 2007; 22(4 Suppl 1):75–81.

- 48 Philippe M, Ameil M. Survival analysis at 10 years of a cohort of 297 atlas total hip prostheses. *Eur J Orthop Surg Traumatol.* 2007;17(6):573–578.
- 49 Ritter MA. The cemented acetabular component of a total hip replacement. All polyethylene versus metal backing. *Clin Orthop.* 1995;(311):69–75.
- 50 Amstutz HC, Le Duff MJ, Campbell PA, Gruen TA, Wisk LE. Clinical and radiographic results of metal-on-metal hip resurfacing with a minimum ten-year follow-up. *J Bone Joint Surg Am.* 2010;92(16):2663–71.
- 51 Veitch SW, Whitehouse SL, Howell JR, Hubble MJW, Gie GA, Timperley AJ. The concentric all-polyethylene Exeter acetabular component in primary total hip replacement. *J Bone Joint Surg Br.* 2010;92(10):1351–5.
- 52 Vassan UT, Sharma S, Chowdary KP, Bhamra MS. Uncemented metal-on-metal acetabular component: follow-up of 112 hips for a minimum of 5 years. *Acta Orthop.* 2007;78(4):470–8.
- 53 Keisu K, Lindgren JU. Outcome of total hip arthroplasties with cement. Results after implantation of the Harris Design 2 prosthesis, 1983–1985. *J Arthroplasty.* 1996; 11(7):789–96.
- 54 Petersen MB, Poulsen IH, Thomsen J, Solgaard S. The hemispherical Harris-Galante acetabular cup, inserted without cement. The results of an eight to eleven-year follow-up of one hundred and sixty-eight hips. *J Bone Joint Surg Am.* 1999;81(2):219–24.
- 55 Metzner G, Strassl R, Neumann DRP, Hitzl W, Ramsauer T, Dorn U. [Long-term results of the cementless Hofer-Imhof threaded titanium cup]. *Z Orthop Unfall.* 1997;147(2):166–74.
- 56 Shetty AA, Slack R, Tindall A, James KD, Rand C. Results of a hydroxyapatite-coated (Furlong) total hip replacement: a 13- to 15-year follow-up. *J Bone Joint Surg Br.* 2005; 87(8):1050–4.
- 57 Partio E, von Bonsdorff H, Wirta J, Avikainen V. Survival of the Lubinus hip prosthesis. An eight- to 12-year follow-up evaluation of 444 cases. *Clin Orthop.* 1994;(303):140–6.
- 58 Bourne RB, Rorabeck CH, Patterson JJ, Guerin J. Tapered titanium cementless total hip replacements: a 10- to 13-year followup study. *Clin Orthop.* 2001;(393):112–20.
- 59 Berli BJ, Schäfer D, Morscher EW. Ten-year survival of the MS-30 matt-surfaced cemented stem. *J Bone Joint Surg Br.* 2005;87(7):928–33.
- 60 Gwynne-Jones DP, Garneti N, Wainwright C, Matheson JA, King R. The Morscher Press Fit acetabular component: a nine- to 13-year review. *J Bone Joint Surg Br.* 2009; 91(7):859–64.

- 61 Farizon F, de Lavison R, Azoulai JJ, Bousquet G. Results with a cementless alumina-coated cup with dual mobility. A twelve-year follow-up study. *Int Orthop.* 1998;22(4):219–24.
- 62 Emms NW, Stockley I, Hamer AJ, Wilkinson JM. Long-term outcome of a cementless, hemispherical, press-fit acetabular component: survivorship analysis and dose-response relationship to linear polyethylene wear. *J Bone Joint Surg Br.* 2010;92(6):856–61.
- 63 Diks M, van den Broek C, Anderson P, van Limbeek J, Spruit M. The uncemented, titanium-coated rm cup: Survival and analyses of failures. *Hip Int;*(Vol. 15 Issue 2 (April - June 2005)):71–77.
- 64 Levy BA, Berry DJ, Pagnano MW. Long-term survivorship of cemented all-polyethylene acetabular components in patients > 75 years of age. *J Arthroplasty.* 2000;15(4):461–7.
- 65 Keisu KS, Orozco F, Sharkey PF, Hozack WJ, Rothman RH, McGuigan FX. Primary cementless total hip arthroplasty in octogenarians. Two to eleven-year follow-up. *J Bone Joint Surg Am.* 2001;83-A(3):359–63.
- 66 Suckel A, Geiger F, Kinzl L, Wulker N, Garbrecht M. Long-term results for the uncemented Zweymüller/Alloclassic hip endoprosthesis. A 15-year minimum follow-up of 320 hip operations. *J Arthroplasty.* 2009;24(6):846–53.
- 67 Dohle J, Becker W, Braun M. [Radiological analysis of osseointegration after implantation of the Zweymüller-Alloclassic total hip system]. *Z Orthop Unfall;*139(6):517–24.
- 68 Ochs U, Eingartner C, Volkmann R, Ochs BG, Huber C, Weller S, Weise K. Prospective long-term follow-up of the cementless bicontact hip stem with plasmapore coating. *Z Orthop Unfall;*145 Suppl:S3–8.
- 69 Bodén H, Salemyr M, Sköldenberg O, Ahl T, Adolphson P. Total hip arthroplasty with an uncemented hydroxyapatite-coated tapered titanium stem: results at a minimum of 10 years' follow-up in 104 hips. *J Orthop Sci.* 2006;11(2):175–9.
- 70 Meding JB, Keating EM, Ritter MA, Faris PM, Berend ME. Minimum ten-year follow-up of a straight-stemmed, plasma-sprayed, titanium-alloy, uncemented femoral component in primary total hip arthroplasty. *J Bone Joint Surg Am.* 2004;86-A(1):92–7.
- 71 Rasquinha VJ, Ranawat CS. Durability of the cemented femoral stem in patients 60 to 80 years old. *Clin Orthop Relat Res.* 2004;419:115–123.
- 72 Chatelet JC, Setiey L. [Long term bone behavior in total primary hip arthroplasty with a fully hydroxyapatite-coated femoral stem: a continuous series of 120 cases with twelve years follow-up]. *Orthop Traumatol Surg Res.* 2004;90(7):628–35.
- 73 Young L, Duckett S, Dunn A. The use of the cemented Exeter Universal femoral stem in a District General Hospital: a minimum ten-year follow-up. *J Bone Joint Surg Br.* 2009; 91(2):170–5.

- 74 Mannan K, Freeman MAR, Scott G. The Freeman femoral component with hydroxyapatite coating and retention of the neck: an update with a minimum follow-up of 17 years. *J Bone Joint Surg Br.* 2010;92(4):480–5.
- 75 Datir SP, Kurta IC, Wynn-Jones CH. Ten-year survivorship of rough-surfaced femoral stem with geometry similar to Charnley femoral stem. *J Arthroplasty.* 2006;21(3):392–7.
- 76 Jewett BA, Collis DK. Radiographic failure patterns of polished cemented stems. *Clin Orthop.* 2006;453:132–6.
- 77 Grant P, Nordsletten L. Total hip arthroplasty with the Lord prosthesis. A long-term follow-up study. *J Bone Joint Surg Am.* 2004;86-A(12):2636–41.
- 78 Savilahti S, Myllyneva I, Pajamäki KJ, Lindholm TS. Survival of Lubinus straight (IP) and curved (SP) total hip prostheses in 543 patients after 4-13 years. *Arch Orthop Trauma Surg.* 1997;116(1-2):10–3.
- 79 Emerson RH, Head WC, Emerson CB, Rosenfeldt W, Higgins LL. A comparison of cemented and cementless titanium femoral components used for primary total hip arthroplasty: a radiographic and survivorship study. *J Arthroplasty.* 2002;17(5):584–91.
- 80 Lombardi AV, Berend KR, Mallory TH, Skeels MD, Adams JB. Survivorship of 2000 tapered titanium porous plasma-sprayed femoral components. *Clin Orthop.* 2009; 467(1):146–54.
- 81 Capello WN, D'Antonio JA, Jaffe WL, Geesink RG, Manley MT, Feinberg JR. Hydroxyapatite-coated femoral components: 15-year minimum followup. *Clin Orthop.* 2006;453:75–80.
- 82 Mohajer MA, Hofmann F, Graf R. Retrospective and consecutive analysis of the long-term outcomes of the SBG stem: a 15-year follow-up study. *Arch Orthop Trauma Surg.* 2010; 130(2):185–90.
- 83 Santori FS, Santori N. Mid-term results of a custom-made short proximal loading femoral component. *J Bone Joint Surg Br.* 2010;92(9):1231–7.
- 84 Parvizi J, Keisu KS, Hozack WJ, Sharkey PF, Rothman RH. Primary total hip arthroplasty with an uncemented femoral component: a long-term study of the Taperloc stem. *J Arthroplasty.* 2004;19(2):151–6.
- 85 Johnsson R, Thorngren KG, Persson BM. Revision of total hip replacement for primary osteoarthritis. *J Bone Joint Surg Br.* 1988;70(1):56–62.
- 86 Espehaug B, Furnes O, Havelin LI, Engesaeter LB, Vollset SE. The type of cement and failure of total hip replacements. *J Bone Joint Surg Br.* 2002;84(6):832–8.

- 87 Espehaug B, Furnes O, Engesaeter LB, Havelin LI. 18 years of results with cemented primary hip prostheses in the Norwegian Arthroplasty Register: concerns about some newer implants. *Acta Orthop.* 2009;80(4):402–412.
- 88 Hop JD, Callaghan JJ, Olejniczak JP, Pedersen DR, Brown TD, Johnston RC. The Frank Stinchfield Award. Contribution of cable debris generation to accelerated polyethylene wear. *Clin Orthop.* 1997;(344):20–32.
- 89 Skutek M, Bourne RB, Rorabeck CH, Burns A, Kearns S, Krishna G. The twenty to twenty-five-year outcomes of the Harris design-2 matte-finished cemented total hip replacement. A concise follow-up of a previous report. *J Bone Joint Surg Am.* 2007; 89(4):814–8.
- 90 Neumann DRP, Thaler C, Hitzl W, Huber M, Hofstädter T, Dorn U. Long-term results of a contemporary metal-on-metal total hip arthroplasty: a 10-year follow-up study. *J Arthroplasty.* 2010;25(5):700–8.
- 91 Macheras G, Kateros K, Kostakos A, Koutsostathis S, Danomaras D, Papagelopoulos PJ. Eight- to ten-year clinical and radiographic outcome of a porous tantalum monoblock acetabular component. *J Arthroplasty.* 2009;24(5):705–9.
- 92 de Jong PT, van der Vis HM, de Man FHR, Marti RK. Weber rotation total hip replacement: a prospective 5- to 20-year followup study. *Clin Orthop.* 2004;(419):107–14.
- 93 Zwartele R, Peters A, Brouwers J, Olsthoorn P, Brand R, Doets C. Long-term results of cementless primary total hip arthroplasty with a threaded cup and a tapered, rectangular titanium stem in rheumatoid arthritis and osteoarthritis. *Int Orthop.* 2008;32(5):581–7.
- 94 Delaunay C, Kapandji AI. Survival analysis of cementless grit-blasted titanium total hip arthroplasties. *J Bone Joint Surg Br.* 2001;83(3):408–13.
- 95 Kobayashi S, Saito N, Horiuchi H, Iorio R, Takaoka K. Poor bone quality or hip structure as risk factors affecting survival of total-hip arthroplasty. *Lancet.* 2000;355(9214):1499–504.
- 96 von Schewelov T, Sanzén L, Besjakov J, Carlsson A. The Elite-Plus stem migrates more than the flanged Charnley stem. *Acta Orthop.* 2010;81(3):280–5.
- 97 Mäkelä K, Eskelinen A, Pulkkinen P, Paavolainen P, Remes V. Cemented total hip replacement for primary osteoarthritis in patients aged 55 years or older: results of the 12 most common cemented implants followed for 25 years in the Finnish Arthroplasty Register. *J Bone Joint Surg Br.* 2008;90(12):1562–9.
- 98 Hook S, Moulder E, Yates PJ, Burston BJ, Whitley E, Bannister GC. The Exeter Universal stem: a minimum ten-year review from an independent centre. *J Bone Joint Surg Br.* 2006;88(12):1584–90.

- 99 Skinner JA, Todo S, Taylor M, Wang JS, Pinskerova V, Scott G. Should the cement mantle around the femoral component be thick or thin? *J Bone Joint Surg Br.* 2003; 85(1):45–51.
- 100 Journeaux SF, Morgan DA, Donnelly WJ. The medium-term results of a cemented Freeman femoral neck-retaining prosthesis. *J Bone Joint Surg Br.* 2000;82(2):188–91.
- 101 Skinner JA, Kroon PO, Todo S, Scott G. A femoral component with proximal HA coating. An analysis of survival and fixation at up to ten years. *J Bone Joint Surg Br.* 2003; 85(3):366–70.
- 102 Räber DA, Czaja S, Morscher EW. Fifteen-year results of the Müller CoCrNiMo straight stem. *Arch Orthop Trauma Surg.* 2001;121(1-2):38–42.
- 103 Riede U, Lüem M, Ilchmann T, Eucker M, Ochsner PE. The M.E Müller straight stem prosthesis: 15 year follow-up. Survivorship and clinical results. *Arch Orthop Trauma Surg.* 2007;127(7):587–92.
- 104 Baumann B, Hendrich C, Barthel T, Bockholt M, Walther M, Eulert J, Rader CP. 9- to 11-year results of cemented titanium mueller straight stem in total hip arthroplasty. *Orthopedics.* 2007;30(7):551–7.
- 105 Kovac S, Trebse R, Milosev I, Pavlovic V, Pisot V. Long-term survival of a cemented titanium-aluminium-vanadium alloy straight-stem femoral component. *J Bone Joint Surg Br.* 2006;88(12):1567–73.
- 106 Jaffe WL, Hawkins CA. Normalized and proportionalized cemented femoral stem survivorship at 15 years. *J Arthroplasty.* 1999;14(6):708–13.
- 107 Engesaeter LB, Lie SA, Espehaug B, Furnes O, Vollset SE, Havelin LI. Antibiotic prophylaxis in total hip arthroplasty: effects of antibiotic prophylaxis systemically and in bone cement on the revision rate of 22,170 primary hip replacements followed 0-14 years in the Norwegian Arthroplasty Register. *Acta Orthop Scand.* 2003;74(6):644–51.
- 108 Thanner J, Freij-Larsson C, Kärrholm J, Malchau H, Wesslén B. Evaluation of Boneloc. Chemical and mechanical properties, and a randomized clinical study of 30 total hip arthroplasties. *Acta Orthop Scand.* 1995;66(3):207–14.
- 109 Muirhead-Allwood SK. Lessons of a hip failure. *BMJ.* 1998;316(7132):644.
- 110 Faro LM, Huiskes R. Quality assurance of joint replacement. Legal regulation and medical judgement. *Acta Orthop Scand Suppl.* 1992;250:1–33.
- 111 Cohen D. Out of joint: the story of the ASR. *BMJ.* 2011;342:d2905.
- 112 Franklin J, Robertsson O, Gestsson J, Lohmander LS, Ingvarsson T. Revision and complication rates in 654 Exeter total hip replacements, with a maximum follow-up of 20 years. *BMC Musculoskelet Disord.* 2003;4:6.

- 113 Sundfeldt M, Carlsson LV, Johansson CB, Thomsen P, Gretzer C. Aseptic loosening, not only a question of wear: a review of different theories. *Acta Orthop.* 2006;77(2):177–197.
- 114 Valstar ER, Nelissen RG, Reiber JH, Rozing PM. The use of Roentgen stereophotogrammetry to study micromotion of orthopaedic implants. *ISPRS J Photogramm Remote Sens.* 2002;56(5-6):376–389.
- 115 Nelissen RG, Brand R, Rozing PM. Survivorship analysis in total condylar knee arthroplasty. A statistical review. *J Bone Joint Surg Am.* 1992;74(3):383–389.
- 116 Keurentjes JC, Fiocco M, Schreurs BW, Pijls BG, Nouta KA, Nelissen RGHH. Revision surgery is overestimated in hip replacement. *Bone Joint Res.* 2012;1(10):258–262.
- 117 Labek G, Sekyra K, Pawelka W, Janda W, Stöckl B. Outcome and reproducibility of data concerning the Oxford unicompartmental knee arthroplasty: a structured literature review including arthroplasty registry data. *Acta Orthop.* 2011;82(2):131–5.
- 118 Wilmshurst P. The regulation of medical devices. *BMJ.* 2011;342:d2822.
- 119 Nelissen RGHH, Pijls BG, Kärrholm J, Malchau H, Nieuwenhuijse MJ, Valstar ER. RSA and registries: the quest for phased introduction of new implants. *J Bone Joint Surg Am.* 2011;93 Suppl 3:62–5.
- 120 Sedrakyan A. Hip resurfacing: a complex challenge for device regulation. *Lancet.* 2012;380(9855):1720–2.
- 121 McCulloch P, Altman DG, Campbell WB, Flum DR, Glasziou P, Marshall JC, Nicholl J, Aronson JK, Barkun JS, Blazeby JM, Boutron IC, Clavien PA, Cook JA, Ergina PL, Feldman LS, Maddern GJ, Reeves BC, Seiler CM, Strasberg SM, Meakins JL, Ashby D, Black N, Bunker J, Burton M, Campbell M, Chalkidou K, Chalmers I, de Leval M, Deeks J, Grant A, Gray M, Greenhalgh R, Jenicek M, Kehoe S, Lilford R, Littlejohns P, Loke Y, Madhock R, McPherson K, Meakins J, Rothwell P, Summerskill B, Taggart D, Tekkis P, Thompson M, Treasure T, Trohler U, Vandenbroucke J. No surgical innovation without evaluation: the IDEAL recommendations. *Lancet.* 2009;374(9695):1105–12.
- 122 McCulloch P, Barkun J, Sedrakyan A. Implantable device regulation in Europe. *Lancet.* 2012;380(9843):729.
- 123 Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am.* 2007;89(4):780–785.
- 124 Krieger N. A glossary for social epidemiology. *J Epidemiol Community Health.* 2001;55(10):693–700.
- 125 Lostao L, Regidor E, Gimeno D, Netuveli G, Blane D. Socioeconomic patterns in health services use in Great Britain and Spain before and after the health system reforms of the 1990s. *Health Place.* 2011;17(3):830–5.

- 126 Quintana JM, Escobar A, Aguirre U, Lafuente I, Arenaza JC. Predictors of health-related quality-of-life change after total hip arthroplasty. *Clin Orthop.* 2009;467(11):2886–94.
- 127 Escobar A, Quintana JM, Bilbao A, Azkárate J, Güenaga JI, Arenaza JC, Gutierrez LF. Effect of patient characteristics on reported outcomes after total knee replacement. *Rheumatology.* 2007;46(1):112–9.
- 128 Gill TM, Feinstein AR. A critical appraisal of the quality of quality-of-life measurements. *JAMA*;272(8):619–26.
- 129 Muldoon MF, Barger SD, Flory JD, Manuck SB. What are quality of life measurements measuring? *BMJ.* 1998;316(7130):542–5.
- 130 Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care.* 1992;30(6):473–83.
- 131 Stewart AL, Hays RD, Ware JE. The MOS short-form general health survey. Reliability and validity in a patient population. *Med Care.* 1988;26(7):724–35.
- 132 Jenkinson C, Coulter A, Wright L. Short form 36 (SF36) health survey questionnaire: normative data for adults of working age. *BMJ.* 1993;306(6890):1437–40.
- 133 Aaronson NK, Muller M, Cohen PD, Essink-Bot ML, Fekkes M, Sanderman R, Sprangers MA, te Velde A, Verrips E. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol.* 1998;51(11):1055–68.
- 134 Ostendorf M, van Stel HF, Buskens E, Schrijvers AJP, Marting LN, Verbout AJ, Dhert WJA. Patient-reported outcome in total hip replacement. A comparison of five instruments of health status. *J Bone Joint Surg Br.* 2004;86(6):801–808.
- 135 Grosse Frie K, van der Meulen J, Black N. Single item on patients' satisfaction with condition provided additional insight into impact of surgery. *J Clin Epidemiol.* 2012; 65(6):619–26.
- 136 Charnley J. The long-term results of low-friction arthroplasty of the hip performed as a primary intervention. *J Bone Joint Surg Br.* 1972;54(1):61–76.
- 137 Dunbar MJ, Robertsson O, Ryd L. What's all that noise? The effect of co-morbidity on health outcome questionnaire results after knee arthroplasty. *Acta Orthop Scand.* 2004; 75(2):119–26.
- 138 Dempster AP, Laird NM, Rubin DB. Maximum Likelihood Estimation from Incomplete Data Via the EM Algorithm. *J R Stat Soc Series B Stat Methodol.* 1977;39(1):1–38.
- 139 Honaker J, King G, Blackwell M. AMELIA II : A Program for Missing Data. *J Stat Softw.* 2010;45(7):1–53.

- 140 King G, Honaker J, Joseph A, Scheve K. Analyzing Incomplete Political Science Data: An Alternative Algorithm for Multiple Imputation. *Am Polit Sci Rev.* 2001;95(1):49–69.
- 141 Keurentjes JC, Van Tol FR, Fiocco M, Schoones JW, Nelissen RG. Minimal clinically important differences in health-related quality of life after total hip or knee replacement. *Bone Joint Res.* 2012;1(5):71–77.
- 142 Rubin DB. *Multiple Imputation for Nonresponse in Surveys*, vol. 519. Wiley. 1987.
- 143 Malchau H, Garellick G, Eisler T, Kärrholm J, Herberts P. Presidential guest address: the Swedish Hip Registry: increasing the sensitivity by patient outcome data. *Clin Orthop.* 2005;441:19–29.
- 144 Hartge P. Participation in population studies. *Epidemiology.* 2006;17(3):252–4.
- 145 Galea S, Tracy M. Participation rates in epidemiologic studies. *Ann Epidemiol.* 2007; 17(9):643–53.
- 146 Jones J. The effects of non-response on statistical inference. *J Health Soc Policy.* 1996; 8(1):49–62.
- 147 Bruyère O, Ethgen O, Neuprez A, Zégels B, Gillet P, Huskin JP, Reginster JY. Health-related quality of life after total knee or hip replacement for osteoarthritis: a 7-year prospective study. *Arch Orthop Trauma Surg.* 2012;132(11):1583–7.
- 148 Ng CY, Ballantyne JA, Brenkel IJ. Quality of life and functional outcome after primary total hip replacement. A five-year follow-up. *J Bone Joint Surg Br.* 2007;89(7):868–73.
- 149 Vissers MM, Bussmann JB, Verhaar JAN, Arends LR, Furlan AD, Reijman M. Recovery of physical functioning after total hip arthroplasty: systematic review and meta-analysis of the literature. *Phys Ther.* 2011;91(5):615–29.
- 150 Allen Butler R, Rosenzweig S, Myers L, Barrack RL. The Frank Stinchfield Award: the impact of socioeconomic factors on outcome after THA: a prospective, randomized study. *Clin Orthop.* 2011;469(2):339–47.
- 151 Davis ET, Lingard EA, Schemitsch EH, Waddell JP. Effects of socioeconomic status on patients' outcome after total knee arthroplasty. *Int J Qual Health Care.* 2008;20(1):40–6.
- 152 Mujica-Mota RE, Tarricone R, Ciani O, Bridges JF, Drummond M. Determinants of demand for total hip and knee arthroplasty: a systematic literature review. *BMC Health Serv Res.* 2012;12(1):225.
- 153 Agabiti N, Picciotto S, Cesaroni G, Bisanti L, Forastiere F, Onorati R, Pacelli B, Pandolfi P, Russo A, Spadea T, Perucci Ca. The influence of socioeconomic status on utilization and outcomes of elective total hip replacement: a multicity population-based longitudinal study. *Int J Qual Health Care.* 2007;19(1):37–44.

- 154 Brennan SL, Stanford T, Wluka AE, Henry MJ, Page RS, Graves SE, Kotowicz Ma, Nicholson GC, Pasco Ja. Cross-sectional analysis of association between socioeconomic status and utilization of primary total hip joint replacements 2006-7: Australian Orthopaedic Association National Joint Replacement Registry. *BMC Musculoskelet Disord.* 2012;13(1):63.
- 155 Dixon T, Shaw M, Ebrahim S, Dieppe P. Trends in hip and knee joint replacement: socioeconomic inequalities and projections of need. *Ann Rheum Dis.* 2004;63(7):825–30.
- 156 Hawkins K, Escoto KH, Ozminkowski RJ, Bhattacharai GR, Migliori RJ, Yeh CS. Disparities in major joint replacement surgery among adults with Medicare supplement insurance. *Popul Health Manag.* 2011;14(5):231–8.
- 157 Judge A, Welton NJ, Sandhu J, Ben-Shlomo Y. Geographical variation in the provision of elective primary hip and knee replacement: the role of socio-demographic, hospital and distance variables. *J Public Health.* 2009;31(3):413–22.
- 158 Judge A, Welton NJ, Sandhu J, Ben-Shlomo Y. Equity in access to total joint replacement of the hip and knee in England: cross sectional study. *BMJ.* 2010;341(aug11 1):c4092–c4092.
- 159 Majeed A, Eliaho J, Bardsley M, Morgan D, Bindman AB. Variation in coronary artery bypass grafting, angioplasty, cataract surgery, and hip replacement rates among primary care groups in London: association with population and practice characteristics. *J Public Health Med.* 2002;24(1):21–6.
- 160 Peterson MG, Hollenberg JP, Szatrowski TP, Johanson NA, Mancuso CA, Charlson ME. Geographic variations in the rates of elective total hip and knee arthroplasties among Medicare beneficiaries in the United States. *J Bone Joint Surg Am.* 1992;74(10):1530–9.
- 161 Steel N, Melzer D, Gardener E, McWilliams B. Need for and receipt of hip and knee replacement—a national population survey. *Rheumatology.* 2006;45(11):1437–41.
- 162 Yong PFK, Milner PC, Payne JN, Lewis Pa, Jennison C. Inequalities in access to knee joint replacements for people in need. *Ann Rheum Dis.* 2004;63(11):1483–9.
- 163 Wolterbeek N, Garling EH, Mertens BJ, Nelissen RGHH, Valstar ER. Kinematics and early migration in single-radius mobile- and fixed-bearing total knee prostheses. *Clin Biomech.* 2012;27(4):398–402.
- 164 Roos EM. Effectiveness and practice variation of rehabilitation after joint replacement. *Curr Opin Rheumatol.* 2003;15(2):160–2.
- 165 Bini SA, Sidney S, Sorel M. Slowing demand for total joint arthroplasty in a population of 3.2 million. *J Arthroplasty.* 2011;26(6 Suppl):124–8.

- 166 Scott CEH, Bugler KE, Clement ND, Macdonald D, Howie CR, Biant LC. Patient expectations of arthroplasty of the hip and knee. *J Bone Joint Surg Br.* 2012;94(7):974–81.
- 167 Jaeschke R, Singer J, Guyatt GH. Measurement of health status. Ascertaining the minimal clinically important difference. *Control Clin Trials.* 1989;10(4):407–15.
- 168 King MT. A point of minimal important difference (MID): a critique of terminology and methods. *Expert Rev Pharmacoecon Outcomes Res.* 2011;11(2):171–84.
- 169 Quintana JM, Escobar A, Bilbao A, Aróstegui I, Lafuente I, Vidaurreta I. Responsiveness and clinically important differences for the WOMAC and SF-36 after hip joint replacement. *Osteoarthritis Cartilage.* 2005;13(12):1076–83.
- 170 Escobar A, Quintana JM, Bilbao A, Aróstegui I, Lafuente I, Vidaurreta I. Responsiveness and clinically important differences for the WOMAC and SF-36 after total knee replacement. *Osteoarthritis Cartilage.* 2007;15(3):273–80.
- 171 Nilsdotter AK, Aurell Y, Siösteen AK, Lohmander LS, Roos HP. Radiographic stage of osteoarthritis or sex of the patient does not predict one year outcome after total hip arthroplasty. *Ann Rheum Dis.* 2001;60(3):228–32.
- 172 Meding JB, Anderson AR, Faris PM, Keating EM, Ritter MA. Is the preoperative radiograph useful in predicting the outcome of a total hip replacement? *Clin Orthop.* 2000;(376):156–60.
- 173 Judge A, Javaid MK, Arden NK, Cushnaghan J, Reading I, Croft P, Dieppe PA, Cooper C. Clinical tool to identify patients who are most likely to achieve long-term improvement in physical function after total hip arthroplasty. *Arthritis Care Res.* 2012;64(6):881–9.
- 174 Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthrosis. *Ann Rheum Dis.* 1957;16(4):494–502.
- 175 Hobbs T. Twelve year report of The New Zealand Joint Registry.
URL <http://www.cdhb.govt.nz/njr/reports/A2D65CA3.pdf>
- 176 Gossec L, Paternotte S, Bingham CO, Clegg DO, Coste P, Conaghan PG, Davis AM, Giacovelli G, Gunther KP, Hawker G, Hochberg MC, Jordan JM, Katz JN, Kloppenburg M, Lanzarotti A, Lim K, Lohmander LS, Mahomed NN, Maillefert JF, Manno RL, March LM, Mazzuca SA, Pavelka K, Punzi L, Roos EM, Rovati LC, Shi H, Singh JA, Suarez-Almazor ME, Tajana-Messi E, Dougados M. OARSI/OMERACT initiative to define states of severity and indication for joint replacement in hip and knee osteoarthritis. An OMERACT 10 Special Interest Group. *J Rheumatol.* 2011;38(8):1765–9.
- 177 de Beer J, Petruccelli D, Adili A, Piccirillo L, Wismer D, Winemaker M. Patient perspective survey of total hip vs total knee arthroplasty surgery. *J Arthroplasty.* 2012;27(6):865–9.e1–5.

- 178 Dobbs HS. Survivorship of total hip replacements. *J Bone Joint Surg Br.* 1980;62-B(2):168–73.
- 179 Dorey F, Amstutz HC. Survivorship analysis in the evaluation of joint replacement. *J Arthroplasty.* 1986;1(1):63–9.
- 180 Putter H, Fiocco M, Geskus RB. Tutorial in biostatistics: competing risks and multi-state models. *Stat Med.* 2007;26(11):2389–430.
- 181 Yan Y, Moore R, Hoover D. Competing risk adjustment reduces overestimation of opportunistic infection rates in AIDS. *J Clin Epidemiol.* 2000;53(8):817–822.
- 182 Southern DA, Faris PD, Brant R, Galbraith PD, Norris CM, Knudtson ML, Ghali WA. Kaplan-Meier methods yielded misleading results in competing risk scenarios. *J Clin Epidemiol.* 2006;59(10):1110–4.
- 183 Fiocco M, Putter H, van de Velde CJH, van Houwelingen JC. Reduced rank proportional hazards model for competing risks: An application to a breast cancer trial. *J Stat Plan Inference.* 2006;136(5):1655–1668.
- 184 Fiocco M, Putter H, Van Houwelingen JC. Reduced rank proportional hazards model for competing risks. *Biostatistics.* 2005;6(3):465–78.
- 185 Bernoulli D. Essai d'une nouvelle analyse de la mortalite causee par la petite verole et des avantages de l'inoculation pour la prevenir. *Mém Math Phys Acad Roy Sci.* 1760;pp. 1–45.
- 186 Blower S, Bernoulli D. An attempt at a new analysis of the mortality caused by smallpox and of the advantages of inoculation to prevent it. 1766. *Rev Med Virol.* 2004;14(5):275–88.
- 187 de Wreede LC, Fiocco M, Putter H. mstate: An R Package for the Analysis of Competing Risks and Multi-State Models. *J Stat Softw.* 2011;38(7):1–30.
- 188 de Wreede LC, Fiocco M, Putter H. The mstate package for estimation and prediction in non- and semi-parametric multi-state and competing risks models. *Comput Methods Programs Biomed.* 2010;99(3):261–74.
- 189 Keurentjes JC, Kuipers RM, Wever DJ, Schreurs BW. High incidence of squeaking in THAs with alumina ceramic-on-ceramic bearings. *Clin Orthop Relat Res.* 2008; 466:1438–1443.
- 190 Harris WH, Sledge CB. Total hip and total knee replacement. *N Engl J Med.* 1990; 323(11):725–31.
- 191 Singh J, Sloan JA, Johanson NA. Challenges with health-related quality of life assessment in arthroplasty patients: problems and solutions. *J Am Acad Orthop Surg.* 2010;18(2):72–82.

- 192 Guyatt GH, Osoba D, Wu AW, Wyrwich KW, Norman GR. Methods to explain the clinical significance of health status measures. *Mayo Clin Proc.* 2002;77(4):371–83.
- 193 Revicki D, Hays RD, Cella D, Sloan J. Recommended methods for determining responsiveness and minimally important differences for patient-reported outcomes. *J Clin Epidemiol.* 2008;61(2):102–9.
- 194 Angst F, Aeschlimann A, Stucki G. Smallest detectable and minimal clinically important differences of rehabilitation intervention with their implications for required sample sizes using WOMAC and SF-36 quality of life measurement instruments in patients with osteoarthritis of the lower ex. *Arthritis Rheum.* 2001;45(4):384–391.
- 195 Wyrwich KW, Wolinsky FD. Identifying meaningful intra-individual change standards for health-related quality of life measures. *J Eval Clin Pract.* 2000;6(1):39–49.
- 196 Cohen J. *Statistical power analysis for the behavioral sciences*, vol. 2. Erlbaum. 1988.
- 197 Ware J. *SF-36 health survey : manual and interpretation guide*. Boston: The Health Institute New England Medical Center. 1993.
- 198 Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. URL http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
- 199 Shi HY, Mau LW, Chang JK, Wang JW, Chiu HC. Responsiveness of the Harris Hip Score and the SF-36: five years after total hip arthroplasty. *Qual Life Res.* 2009;18(8):1053–60.
- 200 Zhang W, Moskowitz RW, Nuki G, Abramson S, Altman RD, Arden N, Bierma-Zeinstra S, Brandt KD, Croft P, Doherty M, Dougados M, Hochberg M, Hunter DJ, Kwok K, Lohmander LS, Tugwell P. OARSI recommendations for the management of hip and knee osteoarthritis, part I: critical appraisal of existing treatment guidelines and systematic review of current research evidence. *Osteoarthritis Cartilage.* 2007;15(9):981–1000.
- 201 Zhang W, Moskowitz RW, Nuki G, Abramson S, Altman RD, Arden N, Bierma-Zeinstra S, Brandt KD, Croft P, Doherty M, Dougados M, Hochberg M, Hunter DJ, Kwok K, Lohmander LS, Tugwell P. OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartilage.* 2008;16(2):137–62.
- 202 Zhang W, Nuki G, Moskowitz RW, Abramson S, Altman RD, Arden NK, Bierma-Zeinstra S, Brandt KD, Croft P, Doherty M, Dougados M, Hochberg M, Hunter DJ, Kwok K, Lohmander LS, Tugwell P. OARSI recommendations for the management of hip and knee osteoarthritis: part III: Changes in evidence following systematic cumulative update of research published through January 2009. *Osteoarthritis Cartilage.* 2010;18(4):476–99.

- 203 Chesworth BM, Mahomed NN, Bourne RB, Davis AM. Willingness to go through surgery again validated the WOMAC clinically important difference from THR/TKR surgery. *J Clin Epidemiol.* 2008;61(9):907–18.
- 204 Walther BA, Moore JL. The concepts of bias, precision and accuracy, and their use in testing the performance of species richness estimators, with a literature review of estimator performance. *Ecography.* 2005;28(6):815–829.
- 205 Arden NK, Kiran A, Judge A, Biant LC, Javaid MK, Murray DW, Carr AJ, Cooper C, Field RE. What is a good patient reported outcome after total hip replacement? *Osteoarthritis Cartilage.* 2011;19(2):155–62.
- 206 Tubach F, Wells GA, Ravaud P, Dougados M. Minimal clinically important difference, low disease activity state, and patient acceptable symptom state: methodological issues. *J Rheumatol.* 2005;32(10):2025–9.
- 207 Tubach F, Ravaud P, Baron G, Falissard B, Logeart I, Bellamy N, Bombardier C, Felson D, Hochberg M, van der Heijde D, Dougados M. Evaluation of clinically relevant states in patient reported outcomes in knee and hip osteoarthritis: the patient acceptable symptom state. *Ann Rheum Dis.* 2005;64(1):34–7.
- 208 Dawson J, Fitzpatrick R, Carr A, Murray D. Questionnaire on the perceptions of patients about total hip replacement. *J Bone Joint Surg Br.* 1996;78:185–190.
- 209 Dawson J, Fitzpatrick R, Murray D, Carr A. Questionnaire on the perceptions of patients about total knee replacement. *J Bone Joint Surg Br.* 1998;80:63–69.
- 210 Judge A, Arden NK, Kiran A, Price A, Javaid MK, Beard D, Murray D, Field RE. Interpretation of patient-reported outcomes for hip and knee replacement surgery: identification of thresholds associated with satisfaction with surgery. *J Bone Joint Surg Br.* 2012;94(3):412–8.
- 211 Gosens T, Hoefnagels NHM, de Vet RCW, Dhert WJA, van Langelaan EJ, Bulstra SK, Geesink RGT. The "Oxford Heup Score": the translation and validation of a questionnaire into Dutch to evaluate the results of total hip arthroplasty. *Acta Orthop.* 2005;76(2):204–11.
- 212 Haverkamp D, Breugem SJM, Sierevelt IN, Blankevoort L, van Dijk CN. Translation and validation of the Dutch version of the Oxford 12-item knee questionnaire for knee arthroplasty. *Acta Orthop.* 2005;76(3):347–52.
- 213 Farrar JT, Young JP, LaMoreaux L, Werth JL, Poole RM. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain.* 2001;94(2):149–58.

- 214 Judge A, Arden NK, Price A, Glyn-Jones S, Beard D, Carr AJ, Dawson J, Fitzpatrick R, Field RE. Assessing patients for joint replacement: can pre-operative Oxford hip and knee scores be used to predict patient satisfaction following joint replacement surgery and to guide patient selection? *J Bone Joint Surg Br.* 2011;93(12):1660–4.
- 215 Kvien TK, Heiberg T, Hagen KB. Minimal clinically important improvement/difference (MCII/MCID) and patient acceptable symptom state (PASS): what do these concepts mean? *Ann Rheum Dis.* 2007;66 Suppl 3(suppl_3):iii40–1.
- 216 Clement ND, Macdonald D, Simpson AHRW. The minimal clinically important difference in the Oxford knee score and Short Form 12 score after total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2013;22(8):1933–9.
- 217 Paulsen A, Roos EM, Pedersen AB, Overgaard S. Minimal clinically important improvement (MCII) and patient-acceptable symptom state (PASS) in total hip arthroplasty (THA) patients 1 year postoperatively. *Acta Orthop.* 2014;85(1):39–48.
- 218 Browne JP, van der Meulen JH, Lewsey JD, Lamping DL, Black N. Mathematical coupling may account for the association between baseline severity and minimally important difference values. *J Clin Epidemiol.* 2010;63(8):865–74.
- 219 Dougados M. It's good to feel better but it's better to feel good. *J Rheumatol.* 2005; 32(1):1–2.
- 220 Fitzpatrick R, Davey C, Buxton MJ, Jones DR. Evaluating patient-based outcome measures for use in clinical trials. *Health Technol Assess.* 1998;2(14):i–iv, 1–74.
- 221 Rolfson O, Kärrholm J, Dahlberg LE, Garellick G. Patient-reported outcomes in the Swedish Hip Arthroplasty Register: results of a nationwide prospective observational study. *J Bone Joint Surg Br.* 2011;93(7):867–75.
- 222 Kvien TK, Mowinckel P, Heiberg T, Dammann KL, Dale O, Aanerud GJ, Alme TN, Uhlig T. Performance of health status measures with a pen based personal digital assistant. *Ann Rheum Dis.* 2005;64(10):1480–4.
- 223 Bellamy N, Wilson C, Hendrikz J, Whitehouse SL, Patel B, Dennison S, Davis T. Osteoarthritis Index delivered by mobile phone (m-WOMAC) is valid, reliable, and responsive. *J Clin Epidemiol.* 2011;64(2):182–90.
- 224 Manfreda K, Bosnjak M, Berzelak J, Haas I, Vehovar V. Web surveys versus other survey modes: A meta-analysis comparing response rates. *Int J Market Res.* 2008; 50(1):79–104.
- 225 Rolfson O, Salomonsson R, Dahlberg LE, Garellick G. Internet-based follow-up questionnaire for measuring patient-reported outcome after total hip replacement surgery-reliability and response rate. *Value Health*;14(2):316–21.

- 226 Ryan JM, Corry JR, Attewell R, Smithson MJ. A comparison of an electronic version of the SF-36 General Health Questionnaire to the standard paper version. *Qual Life Res.* 2002;11(1):19–26.
- 227 Evangelou E, Kerkhof HJ, Styrkarsdottir U, Ntzani EE, Bos SD, Esko T, Evans DS, Metrustry S, Panoutsopoulou K, Ramos YFM, Thorleifsson G, Tsilidis KK, Arden N, Aslam N, Bellamy N, Birrell F, Blanco FJ, Carr A, Chapman K, Day-Williams AG, Deloukas P, Doherty M, Engström G, Helgadottir HT, Hofman A, Ingvarsson T, Jonsson H, Keis A, Keurentjes JC, Kloppenburg M, Lind PA, McCaskie A, Martin NG, Milani L, Montgomery GW, Nelissen RGHH, Nevitt MC, Nilsson PM, Ollier WE, Parimi N, Rai A, Ralston SH, Reed MR, Riancho JA, Rivadeneira F, Rodriguez-Fontenla C, Southam L, Thorsteinsdottir U, Tsezou A, Wallis GA, Wilkinson JM, Gonzalez A, Lane NE, Lohmander LS, Loughlin J, Metspalu A, Uitterlinden AG, Jonsdottir I, Stefansson K, Slagboom PE, Zeggini E, Meulenbelt I, Ioannidis JP, Spector TD, van Meurs JBJ, Valdes AM. A meta-analysis of genome-wide association studies identifies novel variants associated with osteoarthritis of the hip. *Ann Rheum Dis.* 2013;0:1–7.
- 228 Nagelkerke NJD. A note on a general definition of the coefficient of determination. *Biometrika.* 1991;78(3):691–692.
- 229 Steyerberg EW, Eijkemans MJ, Harrell FE, Habbema JD. Prognostic modelling with logistic regression analysis: a comparison of selection and estimation methods in small data sets. *Stat Med.* 2000;19(8):1059–79.
- 230 Steyerberg EW, Bleeker SE, Moll HA, Grobbee DE, Moons KGM. Internal and external validation of predictive models: a simulation study of bias and precision in small samples. *J Clin Epidemiol.* 2003;56(5):441–7.
- 231 Steyerberg E. *Clinical Prediction Models: A Practical Approach to Development, Validation and Updating.* Springer. 2009.
- 232 Deutskens E, de Ruyter K, Wetzels M, Oosterveld P. Response Rate and Response Quality of Internet-Based Surveys: An Experimental Study. *Mark Lett.* 2004;15(1):21–36.
- 233 Brandtzaeg PB, Heim J, Karahasanović A. Understanding the new digital divide: A typology of Internet users in Europe. *Int J Hum Comput Stud.* 2011;69(3):123–138.
- 234 Lee SJ, Kavanaugh A, Lenert L. Electronic and computer-generated patient questionnaires in standard care. *Best Pract Res Clin Rheumatol.* 2007;21(4):637–47.
- 235 Nieuwenhuijse MJ, Valstar ER, Kaptein BL, Nelissen RGHH. Good diagnostic performance of early migration as a predictor of late aseptic loosening of acetabular cups: results from ten years of follow-up with Roentgen stereophotogrammetric analysis (RSA). *J Bone Joint Surg Am.* 2012;94(10):874–80.

- 236 Mancuso CA, Graziano S, Briskie LM, Peterson MGE, Pellicci PM, Salvati EA, Sculco TP. Randomized trials to modify patients' preoperative expectations of hip and knee arthroplasties. *Clin Orthop.* 2008;466(2):424–31.
- 237 Ranstam J, Kärrholm J, Pulkkinen P, Mäkelä K, Espehaug B, Pedersen AB, Mehnert F, Furnes O. Statistical analysis of arthroplasty data. II. Guidelines. *Acta Orthop.* 2011; 82(3):258–67.
- 238 Werner F, Werner K, Oberzaucher J. *Tablets for Seniors: An Evaluation of a Current Model (iPad)*. Advanced Technologies and Societal Change. Berlin, Heidelberg: Springer Berlin Heidelberg. 2012.