



Universiteit
Leiden
The Netherlands

Allogeneic cellular immunotherapy for chronic B-cell leukemia

Hoogendoorn, M.

Citation

Hoogendoorn, M. (2007, February 15). *Allogeneic cellular immunotherapy for chronic B-cell leukemia*. Department of Hematology, Faculty of Medicine, Leiden University Medical Center (LUMC), Leiden University. Retrieved from <https://hdl.handle.net/1887/11408>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/11408>

Note: To cite this publication please use the final published version (if applicable).

Allogeneic cellular immunotherapy for chronic B-cell leukemia

Allogeneic cellular immunotherapy for chronic B-cell leukemia

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van de Rector Magnificus prof.mr.dr. P.F. van der Heijden,
volgens besluit van het College voor Promoties
te verdedigen op donderdag 15 februari 2007
klokke 16.15 uur

door

Mels Hoogendoorn

geboren te Zoetermeer
in 1965

Promotiecommissie

Promotoren: Prof. Dr. J.H.F. Falkenburg
Prof. Dr. R. Willemze

Referent: Prof. Dr. M.H.J. van Oers, Universiteit van Amsterdam

Overigen leden: Prof. Dr. J.C. Kluin-Nelemans, Rijksuniversiteit Groningen
Prof. Dr. R.M. Egeler
Prof. Dr. W.E. Fibbe

Cover: Klaas Koster
Lay-out: Bart Nijmeijer
Printed by: GildePrint

Financial support for the printing of this thesis was provided by:
Amgen BV, Bristol-Myers Squibb, Bronovo Research Fonds, Novartis BV, Pfizer BV, Roche BV,
Schering BV, Ortho Biotec

© M.Hoogendoorn, Suawoude, The Netherlands, 2007

Contents

| | |
|--|-----|
| Chapter 1 | 9 |
| Introduction | |
| Chapter 2 | 31 |
| Generation of B-cell chronic lymphocytic leukemia (CLL)-reactive T-cell lines and clones from HLA class I-matched donors using modified CLL cells as stimulators: implications for adoptive immunotherapy. <i>Leukemia. 2004;18:1278-1287</i> | |
| Chapter 3 | 53 |
| Minor Histocompatibility Antigen (mHag)-specific And B-cell Chronic Lymphocytic Leukemia (CLL)-reactive T Cells Can Be Derived From Matched Sibling Donors Using Transformed CLL Cells As Stimulator Cells. <i>Submitted</i> | |
| Chapter 4 | 71 |
| Primary allogeneic T-cell responses against mantle cell lymphoma antigen presenting cells for adoptive immunotherapy after stem cell transplantation. <i>Clin Cancer Res. 2005;11:5310-5318</i> | |
| Chapter 5 | 93 |
| Characterization of graft-versus-leukemia responses in patients treated for advanced chronic lymphocytic leukemia with donor lymphocyte infusions after <i>in vitro</i> T-cell depleted allogeneic stem cell transplantation following reduced-intensity conditioning. <i>Submitted</i> | |
| Chapter 6 | 113 |
| Summary and General Discussion | |
| Chapter 7 | 127 |
| Nederlandse Samenvatting | |
| Nawoord | 139 |
| Curriculum Vitae | 141 |

