



Universiteit
Leiden
The Netherlands

Prognosis in monoclonal proteinaemia

Schaar, C.G.

Citation

Schaar, C. G. (2006, November 9). *Prognosis in monoclonal proteinaemia*. Retrieved from <https://hdl.handle.net/1887/4983>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4983>

Note: To cite this publication please use the final published version (if applicable).

4 Monoclonal proteinaemia and solid tumours

Schaar CG¹, Snijder S², Oostindiër MJ², Rosendaal FR^{1;3},
Willemze R¹, Kluin-Nelemans JC⁴



European Journal of Cancer 2004; 40: 1539-1544

-
1. Department of Hematology, Leiden University Medical Centre Leiden
 2. Comprehensive Cancer Centre West, Leiden
 3. Department of Clinical Epidemiology, Leiden University Medical Centre
 4. Department of Hematology, Groningen University Hospital, Groningen
All in The Netherlands

Abstract

A higher prevalence of solid tumours in patients with M(onoclonal) proteinaemia without a co-existing haematological malignancy has been reported. We investigated this association by linking a population-based registry of patients with newly diagnosed M-proteinaemia (n=1464) with the Regional Cancer Registry. Patients were followed for a median of 7.4 years for those alive. In total 167 (11%) patients with 173 solid tumours were compared to 861 patients with 'M-proteinaemia only' (without a haematological malignancy). The M-protein isotype or level or clinical parameters did not differ between the groups. M-protein isotype was not associated with a specific tumour type. Standardized morbidity ratios (SMR) for nearly all solid tumours were elevated in the year of M-protein discovery, but the excess risk disappeared during follow-up suggesting selection through diagnostic investigations rather than a causal role. In this large series of patients with both newly diagnosed M-proteinaemia and a solid tumour no relationship could be established.

Introduction

For many decades, it has been assumed that in patients with monoclonal (M)-proteinaemia and without any evidence of a co-existing haematological malignancy, the prevalence of solid tumours is increased suggestive of a causal relationship¹⁻⁸. Further analysis of this relationship could determine more exactly the incidence of this phenomenon in patients with a solid tumour and vice versa, thereby establishing the relevance of screening in cases of M-proteinaemia. To this end we linked a population-based M-protein database^{9;10} to the regional cancer registry and compared solid tumour M-protein patients with other groups in order to try to establish a time relationship between both diagnoses.

Patients and methods

Patient population

From 1991 until 1993, a population-based registry on M-proteinaemia was set up in the region of the Comprehensive Cancer Centre West (CCCW), a geographical area with 1.6 million inhabitants. Clinical chemists, internists, haematologists, pathologists and other physicians reported all patients with newly diagnosed M-proteinaemia or multiple myeloma in the CCCW-area. Information on patient characteristics, laboratory tests results, and results of bone marrow examination and skeletal x-rays were documented. The M-protein-related diagnosis, comorbidity and therapy were recorded. Follow-up was done annually. At follow-up, clinical data, any evolution into a haematological malignancy, appearance of any solid tumour, M-protein levels and other relevant laboratory tests were collected from the patients' hospital charts or from the general physician. In total, 1464 patients have been registered. The set-up and contents of this registry have been described previously^{9;10}.

Diagnostic criteria for monoclonal proteinaemia and solid tumours

M-proteins were either detected by agarose or by cellulose acetate electrophoresis depending on the method used in the various hospitals involved. For inclusion in the registry, each M-protein had to be confirmed by immunotyping (immunofixation). The presence of a solid tumour preferably had to be confirmed by histology, otherwise a clinical diagnosis had to be based on at least radiological evidence of a tumour. Cancer sites were grouped according to the World Health Organization's International Classification of Diseases for Oncology (ICD-O)¹¹.

Linkage to the Regional Cancer Registry

For verification and to ensure the completeness of our data on solid tumours, the database was linked to the regional database of the Netherlands Regional Cancer Registry¹². In this cancer registry, all patients with newly diagnosed malignancies living in the CCCW region reported by the pathology laboratories are entered. The date of the cytological or histological confirmation constitutes the date of diagnosis. In addition, all hospitals employ a separate registry of the discharge diagnoses. For the present study, patient data were linked if name, sex and date of birth were identical in both databases to exclude the probability of false-positive or false negative linkages.

Solid tumour prevalence and incidence analysis

Follow-up started at registration (between 1991 and 1993) and is still ongoing. For the solid cancer linkage study, complete coverage with the Regional Cancer Registry was guaranteed until January 1st 1998. Endpoints were the development of a (haematological) malignancy or death, and patients still alive were censored for all other events on January 1st 2002. First the prevalence of a solid tumour at first diagnosis was calculated. Patients were diagnosed with an M-protein-related solid tumour if the tumour was diagnosed within the time frame of two years, one preceding or following the discovery of the M-protein. Thus, all malignancies that could be associated with the M-protein, but were not present anymore due to treatment were included, as well as any asymptomatic multiple myeloma (MM), other haematological malignancy or solid tumours that developed later on. In case of a simultaneous haematological malignancy or solid tumour during this period, the M-protein was considered to be associated with the former and not with the solid tumour.

Secondly, Standardized morbidity ratios (SMR) for the most prevalent tumours were determined for the period between registration (1991-1993) until January 1, 1998. Patients with newly diagnosed M-proteinaemia were at risk until the diagnosis of a solid tumour, MM, other haematological malignancy was made or until they died. Multiplication of person-years under observation by the age-, sex-, and period-specific incidence rates yielded the number of solid tumours expected in the M-protein cohort if they experienced the same risk as was prevalent in the region of the CCCW. With this method, standardized incidence rates between patient and reference group were compared (indirect standardization) and expressed as the ratio of the incidence rates (SMR), which may be viewed as a relative risk. Confidence limits for the SMR were based on a Poisson distribution for the observed number of deaths¹³.

Statistical methods

Statistical methods to compare the 'M-proteinaemia Only' versus 'Solid Tumour group' included Mann Whitney's test and in the case of case-control design, chi-square test when appropriate. Analyses were performed using SPSS version 10. Data were entered in the database using SPSS Data Entry version 2 (both SPSS Inc. Chicago, Il., USA).

Results

Prevalence of solid tumours at first diagnosis of M-proteinaemia

The database consisted of 1464 patients with an initial diagnosis of M-proteinaemia. The frequency of newly discovered cases was 31/100,000 inhabitants and 189/100,000 for people above 70 years of age⁹. In 271 patients, a diagnosis of MM was made, in 164 another haematological malignancy was diagnosed, but in the large majority no explanation was found (provisional⁹ or definite monoclonal gammopathy of unknown significance, MGUS) (Table 1).

In total, 173 solid tumours without any evidence of MM or other haematological malignancy were diagnosed in 167 (11%) patients. The types of tumour found are depicted in Figure 1. Nearly all tumours (n=167; 96%) were (adeno)carcinomas, the other six malignancies consisted of melanomas (n=4), sarcoma (n=1) and leiomyosarcoma (n=1).

Table 1. Clinical characteristics in all diagnostic groups with M-proteinaemia.

Monoclonal proteinaemia only		Solid tumour	P	Multiple myeloma ^a	Other hematological malignancies ^b
Number	861 (59)	167 (11)		271 (19)	164 (11)
Sex					
M (%)	423 (49)	103 (62)	0.004	138 (51)	96 (58)
F (%)	438 (51)	64 (38)		133 (49)	69 (42)
Median age (range, years)					
M	73 (17-103)	75 (37-95)	0.05	71 (28-93)	72 (21-94)
F	72 (20-103)	75 (37-95)	0.03		
	75 (17-98)	76 (47-92)	0.91		
M-protein type & level (g/l)					
IgG (%)	618 (72)	29 (76)	0.45	155 (57)	75 (46)
Median (range)	10 (1-30)	10.5 (<1-85)		32.5 (6-117)	12.3 (<1-34)
IgA (%)	80 (9)	10 (6)	0.71	75 (28)	5 (3)
Median (range)	8.4 (4-31)	10 (2-47)		28.1 (5-81)	15.6 (2-25)
IgM (%)	159 (19)	30 (18)	0.19	4 (2)	81 (49)
Median (range)	7 (1-21)	10 (<1-30)		22.1 (2-57)	13.8 (2-110)

In the patient with prostate carcinoma, two haematological malignancies were observed: chronic lymphocytic leukaemia (CLL) and Hodgkin's lymphoma. M, male; F, female; MGUS, monoclonal gammopathy of unknown significance.

^{a)} In five patients, a solid tumour was present (carcinoma of lung, breast, thyroid, ovary, and unknown primary site).

^{b)} In eight patients, a solid tumour was present (melanoma, carcinoma of unknown primary site, bladder, prostate, colon, larynx, skin, and liver).

Men (n=103)**Head & Neck: 4 (4%)**

Meninges
Hypopharynx
Nasopharynx
Larynx

Respiratory tract: 36 (35%)

Lung: 35 (34%)
Pleura

Gastrointestinal tract: 25 (24%)

Oesophagus
Stomach: 5 (5%)
Liver
Biliary system: 2 (2%)
Pancreas: 4 (4%)
Colon & rectosigmoid: 12 (12%)

Urogenital tract: 24 (23%)

Kidney: 3 (3%)
Urether & bladder: 8 (8%)
Prostate: 13 (13%)

Skin: 3 (3%)**Melanoma: 4 (4%)****Unknown primary: 8 (8%)****Women (n=64)****Head & Neck: 2 (3%)**

Gingiva
Thyroid

Respiratory tract: 5 (7%)

Lung: 5 (7%)

Breast: 19 (28%)**Gastrointestinal tract: 21 (31%)**

Oesophagus
Stomach: 3 (4%)
Biliary system: 2 (3%)
Pancreas: 6 (9%)
Colon & rectosigmoid: 10 (15%)

Urogenital tract: 12 (17%)

Kidney: 2 (3%)
Ovary: 5 (7%)
Urether & bladder: 2 (3%)
Uterus
Cervix
Vulva


Skin: 2 (3%)**Unknown primary: 7 (10%)**

Figure 1. Distribution of solid tumours and M-proteinaemia. Two tumours were diagnosed in six patients: bilateral breast (2), breast + pancreas, breast carcinoma + colon leiomyosarcoma, squamous skin + melanoma, pancreas + gingiva. No percentage between brackets means only one tumour found. Modified from Goffinet and colleagues¹⁹.

Solid tumour and M-protein isotype

To study whether specific tumour types were related to specific M-protein isotypes, we selected patients with one of the six most frequently occurring solid tumours, lung (n=40), colon (n=21), breast (n=15), prostate (n=13), bladder (n=10) and pancreas (n=9). In all, the IgG isotype predominated (60-85%), followed by IgM (7-30%) and IgA (0-10%). No clear preferential M-protein types were seen within specific tumour groups.

'Solid Tumour Group' versus 'M-proteinaemia Only Group/MGUS'

To study whether patients with M-proteinaemia and a solid tumour (i.e. 'Solid Tumour Group') had specific characteristics in demographics or M-protein isotype and levels, we compared this group with the patients without any malignancy (Table 1). In the tumour group, male gender predominated. In addition, small, but significant, differences in age were seen. The distribution of M-protein isotypes and levels was identical with the majority of patients expressing a median of 10 g/l IgG monoclonal proteinaemia.

Response of the M-protein on cancer treatment

When M-proteins are causally related to a solid tumour, one would expect to find an increase of the M-protein levels during tumour progression and a decrease after tumour disappearance. In the Solid Tumour group, 64 out of 167 patients died within one year after the detection of the M-protein leaving 103 patients with follow-up data. In 25 patients (median follow-up 37 months, range 3-95 months) the M-protein was measured at least once after the first detection and therapy (if any) of the solid tumour. Since in four patients a haematological malignancy developed (see 'Follow-up since entry in the M-protein database'), a relationship (rising or lowering of the M-protein in correspondence with the progression or decrease of the tumour) could be studied in only 21 patients. In this small group of patients, no convincing relationship was seen between the behaviour of the solid tumour and the M-protein levels (data not shown).

Follow-up since entry in the M-protein database

During follow-up (last analysis 1-1-2002) a new solid tumour was detected in 23 patients in addition to those already diagnosed in the 167 patients. All of these tumours occurred in the period 1992-1998 (see Table 2). In the years thereafter, no additional tumours were found. Out of the 167 patients in whom a solid tumour was diagnosed simultaneously with the M-protein, three patients developed MM (14, 56 and 65 months after the detection of the M-protein) and one patient developed a non-Hodgkin's lymphoma (NHL) (28 months after the detection of the M-protein). For comparison, in the Monoclonal proteinaemia only /MGUS group, 28 developed MM, and another 17 developed haematological malignancies, consisting of NHL (n=12), myelodysplastic syndrome or acute myeloid leukaemia (n=3) or myeloproliferative disease (n=2).

Table 2. Standardized morbidity ratios (SMR) for the most prevalent solid tumors in the year of discovery of the M-protein and the years after.

Solid tumour (all (adeno)carcinoma)	Year of M-protein discovery (1991,1992, 1993)				Follow-up (1992-1997)			
	Observed number	Expected number	SMR	95% conf. interval	Observed number	Expected number	SMR	95% conf. interval
Men								
Lung	19	0.9	21.1	12.5-31.9	5	4.1	1.2	0.4-2.6
Colon & rectosigmoid	4	0.4	10	2.5-22.5	1	1.7	0.6	0-2.4
Prostate	5	0.9	5.6	1.7-11.6	3	4.5	0.7	0.2-1.7
Unknown primary	5	0.2	25	7.6-52.4	2	0.8	2.5	0.2-7.3
Pancreas	1	0.1	10	0-40	0	0.25	0	0-4
Stomach	2	0.3	6.7	0.6-19.4	0	0.8	0	0-1.3
Bladder	3	0.2	15	2.7-37.3	1	1.0	1	0-4.0
Women								
Lung	3	0.1	30	5.4-74.6	0	0.6	0	0-1.7
Colon & rectosigmoid	4	0.4	10	2.5-22.5	5	1.6	3.1	1.0-6.5
Breast	3	0.6	5	0.9-12.4	3	2.9	1.0	0.2-2.6
Unknown primary	5	0.1	50	15.2-104.7	2	0.6	3.3	0.3-9.7
Pancreas	4	0.1	40	10-90	2	0.4	5.0	0.4-14.6
Stomach	2	0.1	20	1.7-58.3	0	0.7	0	0-1.4
Bladder	0	0.1	0	0-40	1	0.29	3.5	0-13.8

Standardized Morbidity Ratio

Cumulative follow-up of all 1464 patients during this selected period (1991-1998) was 3060 person-years with a median follow-up of 1.3 years (range 0-7 years) and a median follow-up of 7.4 years (range 10 months-7 years) for those still alive. Cumulative follow-up (measuring the time interval between the date of diagnosis of the M-proteinaemia and the date of diagnosis of the solid tumour) for the Solid Tumour group was 24 person-years (median less than 1 day, range 0-2.8 years). In conclusion, most solid tumours were diagnosed simultaneously with the detection of the M-protein (median follow-up less than one day, see above). In the first year after the detection of the M-protein, SMR for nearly all solid tumours showed an increased risk (range 0-50). However, all declined sharply or normalised during subsequent follow-up years (Table 2).

Discussion

In this population-based registry on patients with newly diagnosed M-proteinaemia we describe the largest series collected thus far of patients with both a solid tumour and a M-protein but without any evidence of a co-existing haematological malignancy. Since 1928, investigators reported an increased prevalence of solid tumours in patients with M-proteinaemia suggesting a paraneoplastic phenomenon. For comparison with our cohort, we selected only studies with more than 100 patients, with a description of the related malignancy including concise histopathology and information on the determination of the M-protein, and were left with 8¹⁻⁸. Identical to our series, nearly all solid tumours described were (adeno)carcinomas. M-proteins were mostly of the IgG isotype and levels (if investigated) were generally lower than 30 g/l.

The co-existing tumours in this M-protein database were manifest at the diagnosis of the M-protein in the large majority of patients. During follow-up, only a small additional number of solid tumours were detected. Kyle and colleagues *et al* observed the development of a second tumour in 15 of 241 MGUS-patients during a 20-35 year follow-up¹⁴ and Pasqualetti and colleagues reported 31 out of 263 similar patients who died due to a solid tumour during a median follow-up of 11.5 years¹⁵. In contrast, in the only prospective study investigating the incidence of haematological and solid malignancies in patients with M-proteinaemia, Gregersen and colleagues did not observe an increased risk of solid tumours in 1229 patients during follow-up (mean 4.8 years, range 0-15.7 years)¹⁶. Similarly to our findings, the risk for developing a solid tumour was increased in the first year of follow-up though this risk diminished thereafter¹⁶. Approaching the probable relationship between M-proteinaemia and cancer the other way around yielded no association either: in two cross-sectional stud-

ies the prevalence of M-proteinaemia in patients with non-haematological tumours was not increased when compared to the prevalence in the general population^{17;18}.

In conclusion, we did not observe differences in clinical characteristics between patients with 'M-proteinaemia only/MGUS' and patients with 'Solid tumour and M-proteinaemia'. There was no relation between specific solid tumours and M-protein isotype nor did the serum level of the M-protein change after anti-tumour therapy (although the number of patients was small in this analysis). Although risks for nearly all solid tumours found were initially elevated in patients with newly diagnosed M-proteinaemia, these decreased in the year after suggesting a diagnostic selection of patients rather than a causal role.

Acknowledgements

We are indebted to mr. K.G. van der Ham, Department of Clinical Pathology for his assistance in the creation of Figure 1.

Reference list

1. Creyssel R, Fine JM, Morel P. Etude biochimique de quelques formes atypiques de dysproteïnémies. *Rev Hematol* 1959;14:238-249.
2. Osserman EF, Takatsuki K. Plasma cell myeloma: gamma globulin synthesis and structure. *Medicine* 1963;42:357-385.
3. Hällén J. Discrete gammaglobulin (M-) components in serum. Clinical study of 150 subjects without myelomatosis. *Acta Med Scand* 1966;462:1-127.
4. Danon F, Clauvel JP, Seligmann M. Les 'paraprotéines' de type IgG et IgA en dehors de la maladie de Kahler. *Rev Fr Etud Clin Biol* 1967;12:681-701.
5. Pick AI, Yeshurun D, Schreiber S, Fleminger R, Kessler H. Monoclonal gammopathy in patients with epithelial tumors. *Harefuah* 1974;86:5-13.
6. Colls BM, Lorier MA. Immunocytoma, Cancer, and other associations of monoclonal gammopathy: A review of 224 cases. *N Z Med J* 1975;82:221-226.
7. Ameis A, Ko HS, Pruzanski W. M-components - a review of 1242 cases. *Can Med Ass J* 1976;114:889-92, 895.
8. Bonomo L, Dammacco F, Miglietta A. Paraproteinemia and neoplasia. *Ric Clin Lab* 1985;15:99-104.
9. Ong F, Hermans J, Noordijk EM et al. Developing a population-based registry for patients with paraproteinemias or multiple myeloma. *J Clin Epidemiol* 1997;50:909-915.
10. Ong F, Hermans J, Noordijk EM et al. A population-based registry on paraproteinaemia in The Netherlands. Comprehensive Cancer Centre West, Leiden, The Netherlands. *Br J Haematol* 1997;99:914-920.
11. World Health Organisation (WHO). International Classification of Diseases for Oncology. Geneva; 1976.

12. Visser, O., Coebergh, J. W. W., Schouten, L. J., and Dijck, J. A. A. M. Incidence of cancer in the Netherlands 1994. Visser, O., Coebergh, J. W. W., Schouten, L. J., and Dijck, J. A. A. M. 6. 1997. Utrecht, Vereniging van Integrale Kankercentra.
13. Vandenbroucke JP. A shortcut method for calculating the 95 per cent confidence interval of the standardized mortality ratio. *Am J Epidemiol* 1982;115:303-304.
14. Kyle RA. 'Benign' monoclonal gammopathy – after 20 to 35 years of follow-up. *Mayo Clin Proc* 1993;68:26-36.
15. Pasqualetti P, Festuccia V, Collacciani A, Casale R. The natural history of monoclonal gammopathy of undetermined significance. A 5- to 20-year follow-up of 263 cases. *Acta Haematol* 1997;97:174-179.
16. Gregersen H, Mellemkjaer L, Salling IJ et al. Cancer risk in patients with monoclonal gammopathy of undetermined significance. *Am J Hematol* 2000;63:1-6.
17. Migliore PJ, Alexanian R. Monoclonal gammopathy in human neoplasia. *Cancer* 1968;21:1127-1131.
18. Talerman A, Haije WG. The frequency of M-components in sera of patients with solid malignant neoplasms. *Br J Cancer* 1973;27:276-282.
19. Goffinet DR, Warnke R, Dunnick NR et al. Clinical and surgical (laparotomy) evaluation of patients with non- Hodgkin's lymphomas. *Cancer Treat Rep* 1977;61:981-992.