

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/21704> holds various files of this Leiden University dissertation

**Author:** Gupta, Vikas

**Title:** Multimodality cardiac image analysis for the assessment of coronary artery disease

**Issue Date:** 2013-09-11

# Bibliography

- [1] MATLAB (version 7.5). *The MathWorks Inc.,Natick, Massachusetts*, 2007.
- [2] MATLAB (version 7.13). *The MathWorks Inc.,Natick, Massachusetts*, 2011. URL <http://www.mathworks.com>.
- [3] N. Ablitt, J. Gao, P. Gatehouse, and G.Z. Yang. Motion decoupling and registration for 3d magnetic resonance myocardial perfusion imaging. In *Proceedings of the International Conference on Computational Science-Part III*, pages 285–294, 2002.
- [4] G. Adluru, E.V.R. DiBella, and M.C. Schabel. Model-based registration for dynamic cardiac perfusion mri. *Journal of Magnetic Resonance Imaging*, 24(5):1062–1070, 2006.
- [5] G. Adluru, E.V.R. DiBella, and R.T. Whitaker. Automatic segmentation of cardiac short axis slices in perfusion. In *Proceedings of International Symposium on Biomedical Imaging (ISBI)*, pages 133–136. IEEE, 2006.
- [6] N. Al-Saadi, E. Nagel, M. Gross, A. Bornstedt, B. Schnackenburg, C. Klein, W. Klimek, H. Oswald, and E. Fleck. Noninvasive detection of myocardial ischemia from perfusion reserve based on cardiovascular magnetic resonance. *Circulation*, 101(12):1379, 2000.
- [7] N. Al-Saadi, E. Nagel, M. Gross, B. Schnackenburg, and C. and Fleck E. Paetsch, I. and Klein. Improvement of myocardial perfusion reserve early after coronary intervention: assessment with cardiac magnetic resonance imaging. *Journal of the American College of Cardiology*, 36(5):1557–64, 2000.
- [8] N. Al-Saadi, M. Gross, A. Bornstedt, B. Schnackenburg, C. Klein, E. Fleck, and E. Nagel. Comparison of various parameters for determining an index of myocardial perfusion reserve in detecting coronary stenosis with cardiovascular magnetic resonance tomography. *Zeitschrift für Kardiologie*, 90(11):824, 2001.
- [9] M.S. Albert, W. Haung, J.H. Lee, C.S. Patlak, and C.S. Springer Jr. Susceptibility changes following bolus injections. *Magnetic Resonance in Medicine*, 29(5):700–708, 1993.
- [10] A.E. Arai. Magnetic resonance first-pass myocardial perfusion imaging. *Topics in Magnetic Resonance Imaging*, 11(6):383, 2000.
- [11] American Heart Association. Heart disease and stroke statistics - 2010 update: A report from the american heart association. *Circulation*, 2010.
- [12] D. Atkinson, D.L.G. Hill, P.N.R. Stoye, P.E. Summers, and S.F. Keevil. Automatic correction of motion artifacts in magnetic resonance images using an entropy focus criterion. *IEEE Transactions on Medical Imaging*, 16(6):903–910, 1997.
- [13] D.J. Atkinson, D. Burstein, and R.R. Edelman. First-pass cardiac perfusion: evaluation with ultrafast MR imaging. *Radiology*, 174(3):757, 1990.
- [14] A.K. Attili, A. Schuster, E. Nagel, J.H.C. Reiber, and R.J. van der Geest. Quantification in cardiac MRI: advances in image acquisition and processing. *The International Journal of Cardiovascular Imaging*, 26:27–40, 2010.

- [15] L. Axel, J. Costantini, and J. Listerud. Intensity correction in surface-coil MR imaging. *American Journal of Roentgenology*, 148(2):418–420, 1987.
- [16] R.J. Bache and J.S. Schwartz. Effect of perfusion pressure distal to a coronary stenosis on transmural myocardial blood flow. *Circulation*, 65(5):928, 1982.
- [17] N. Baka, J. Milles, E. A. Hendriks, A. Suinesiaputra, M.J. Jerosh-Herold, J. H. C. Reiber, and B. P. F. Lelieveldt. Segmentation of myocardial perfusion MR sequences with multi-band active appearance models driven by spatial and temporal features. In *Proceedings of SPIE*, volume 6914, pages 691415–1–10, 2008.
- [18] C. Baldy, P. Douek, P. Croisille, I.E. Magnin, D. Revel, and M. Amiel. Automated myocardial edge detection from breath-hold cine-MR images: evaluation of left ventricular volumes and mass. *Magnetic Resonance Imaging*, 12(4):589–598, 1994.
- [19] R. Bansal and G. Funka-Lea. Integrated image registration for cardiac MR perfusion data. In *MICCAI*, pages 659–666, 2002.
- [20] J. Bardanova, I. Provaznik, and M Novakova. Hidden markov model in wavelet analysis of myocardial ischemia in rabbit. In *Computers in Cardiology*, pages 419–421, 2000.
- [21] W. Bauer. Myocardial perfusion and capillary blood volume during left ventricular remodelling. *Magnetic Resonance Materials in Biology, Physics, and Medicine*, 6(2-3):148–149, 1998.
- [22] J.J. Bax, R.S. Beanlands, F.J. Klocke, J. Knuuti, A.A. Lammertsma, H.R. Schaeffers, M.A. Schelbert, G.K. Von Schulthess, L.J Shaw, G.Z. Yang, et al. Diagnostic and clinical perspectives of fusion imaging in cardiology: is the total greater than the sum of its parts? *Heart*, 93(1):16–22, 2007.
- [23] F. Behloul, J.P. Roux, and R. Unterreiner. Automatic boundary detection of the left ventricle and measurement of local myocardial perfusion in MRI. In *Proceedings of Computers in Cardiology*, volume 24, pages 145–148, 1997.
- [24] U. Behrens, J. Teubnerb, C.J.G. Evertsz, M. Walzb, H. Jiirgens, and H.O. Peitgena. Computer assisted dynamic evaluation of contrast-enhanced breast-MRI. In *Proceedings of the International Symposium on Computer and Communication Systems for Image Guided Diagnosis and Therapy, Paris, June 1996*, pages 362–367, 1996.
- [25] B. Belaroussi, J. Milles, S. Carme, Y.M. Zhu, and H. Benoit-Cattin. Intensity non-uniformity correction in MRI: Existing methods and their validation. *Medical Image Analysis*, 10(2): 234–246, 2006.
- [26] P. Beliveau, R.M. Setser, F. Cherieta, R.D. White, and T. O'Donnell. Computation of coronary perfusion territories from CT angiography. In *Proc. of Computers in Cardiology*, volume 34, pages 753–756, 2007.
- [27] G.A. Beller. Recent advances and future trends in multimodality cardiac imaging. *Heart, Lung and Circulation*, 19(3):193–209, 2010.
- [28] M. Bellin and A. J. Van Der Molen. Extracellular gadolinium-based contrast media: An overview. *European Journal of Radiology*, 66(2):160 – 167, 2008.
- [29] H. Benali, I. Buvat, F. Frouin, J.P. Bazin, and R.D. Paola. A statistical model for the determination of the optimal metric in factor analysis of medical image sequences (FAMIS). *Physics in Medicine and Biology*, 38:1065, 1993.

- [30] L. M. Bidaut and J. P. Vallée. Automated registration of dynamic MR images for the quantification of myocardial perfusion. *Journal of Magnetic Resonance Imaging*, 13(4):648–55, 2001.
- [31] D.E. Bild, D.A. Bluemke, G.L. Burke, R. Detrano, A.V. Diez Roux, A.R. Folsom, P. Greenland, et al. Multi-ethnic study of atherosclerosis: objectives and design. *American journal of epidemiology*, 156(9):871, 2002.
- [32] S. Bock, C. Kühnel, T. Boskamp, and H.O. Peitgen. Robust vessel segmentation. In *Proceedings of SPIE*, volume 6915, page 691539, 2008.
- [33] JG Bosch, SC Mitchell, BPF Lelieveldt, F. Nijland, O. Kamp, M. Sonka, and JHC Reiber. Automatic segmentation of echocardiographic sequences by active appearance motion models. *IEEE Transactions on Medical Imaging*, 21(11):1374–1383, 2002.
- [34] T. Boskamp, D. Rinck, F. Link, B. Kümmerlen, G. Stamm, and P. Mildenerger. New vessel analysis tool for morphometric quantification and visualization of vessels in CT and MR imaging data sets. *Radiographics*, 24(1):287–297, 2004.
- [35] A. O. Boudraa, F. Behloul, M. Janier, E. Canet, J. Champier, J. P. Roux, and D. Revel. Temporal covariance analysis of first-pass contrast-enhanced myocardial magnetic resonance images. *Computers in Biology and Medicine*, 31(2):133–42, 2001.
- [36] L. Bracoud, F. Vincent, C. Pachai, E. Canet, P. Croisille, D. Revel, E. Canet, P. Croisille, and D. Revel. Automatic registration of MR first-pass myocardial perfusion images. In *Proceedings of Functional Imaging and Modeling of the Heart (FIMH)*, pages 215–223, 2003.
- [37] M. Breeuwer, M. Quist, L. Spreeuwers, I. Paetsch, N. Al-Saadi, and E. Nagel. Towards automatic quantitative analysis of cardiac mr perfusion images. In *International Congress Series*, volume 1230, pages 967–973, 2001.
- [38] M. Breeuwer, L. Spreeuwers, and M. Quist. Automatic quantitative analysis of cardiac MR perfusion images. In *Proceedings of SPIE*, volume 4322, pages 733–742, 2001.
- [39] J. J. Brown and C. B. Higgins. Myocardial paramagnetic contrast agents for MR imaging. *American Journal of Roentgenology*, 151(5):865–871, 1988.
- [40] D. Burstein, E. Taratuta, and W.J. Manning. Factors in myocardial perfusion imaging with ultrafast MRI and Gd-DTPA administration. *Magnetic Resonance in Medicine*, 20(2):299–305, 1991.
- [41] S. Busch, T.R.C. Johnson, K. Nikolaou, F. Von Ziegler, A. Knez, M.F. Reiser, and C.R. Becker. Visual and automatic grading of coronary artery stenoses with 64-slice CT angiography in reference to invasive angiography. *European Radiology*, 17(6):1445–1451, 2007.
- [42] I. Buvat, H. Benali, and R. Paola. Statistical distribution of factors and factor images in factor analysis of medical image sequences. *Physics in Medicine and Biology*, 43:1695–1711, 1998.
- [43] V.D. Calhoun, T. Adali, L.K. Hansen, J. Larsen, and J.J. Pekar. ICA of functional MRI data: an overview. In *Proceedings of the International Workshop on Independent Component Analysis and Blind Signal Separation*, pages 281–288, 2003.
- [44] E. Canet, D. Revel, L. Sebbag, M. de Lorgeril, C. Baldy, C. Delabre, and M. Amiel. Noninvasive assessment of no-reflow phenomenon in a canine model of reperfused infarction by contrast-enhanced magnetic resonance imaging. *American Heart Journal*, 130(5):949–956, 1995.

- [45] T.J. Carroll, V.M. Haughton, H.A. Rowley, and D. Cordes. Confounding effect of large vessels on MR perfusion images analyzed with independent component analysis. *American Journal of Neuroradiology*, 23(6):1007, 2002.
- [46] J.E. Cates, A.E. Lefohn, and R.T. Whitaker. GIST: An interactive, GPU-based level set segmentation tool for 3D medical images. *Medical Image Analysis*, 8(3):217–231, 2004.
- [47] M.D. Cerqueira, N.J. Weissman, V. Dilsizian, A.K. Jacobs, S. Kaul, W.K. Laskey, D.J. Pennell, J.A. Rumberger, T. Ryan, and M.S. Verani. Standardized myocardial segmentation and nomenclature for tomographic imaging of the heart. *Circulation*, 105(4):539–542, 2002.
- [48] C. Chefdh tel, G. Hermosillo, and O. Faugeras. Flows of diffeomorphisms for multimodal image registration. In *Proceedings of International Symposium on Biomedical Imaging (ISBI)*, pages 753–756, 2002.
- [49] A.S.H. Cheng, T.J. Pegg, T.D. Karamitsos, N. Searle, M. Jerosch-Herold, R.P. Choudhury, A.P. Banning, S. Neubauer, M.D. Robson, and J.B. Selvanayagam. Cardiovascular magnetic resonance perfusion imaging at 3-Tesla for the detection of coronary artery disease : A comparison with 1.5-Tesla. *Journal of the American College of Cardiology*, 49(25):2440–2449, 2007.
- [50] T.F. Christian, D.W. Rettmann, A.H. Aletras, S.L. Liao, J.L. Taylor, R.S. Balaban, and A.E. Arai. Absolute myocardial perfusion in canines measured by using dual-bolus first-pass MR imaging. *Radiology*, 232(3):677–684, 2004.
- [51] T.F. Christian, A.H. Aletras, and A.E. Arai. Estimation of absolute myocardial blood flow during first-pass MR perfusion imaging using a dual-bolus injection technique : Comparison to single-bolus injection method. *Journal of Magnetic Resonance Imaging*, 27(6):1271–1277, 2008.
- [52] J.P. Christiansen, T.D. Karamitsos, S.G. Myerson, J.M. Francis, and S. Neubauer. Stress perfusion imaging using cardiovascular magnetic resonance: A review. *Heart, Lung and Circulation*, 19(12):697–705, 2010.
- [53] N.C.F. Codella, J.W. Weinsaft, M.D. Cham, M. Janik, M.R. Prince, and Y. Wang. Left Ventricle: Automated segmentation by using myocardial effusion threshold reduction and intravoxel computation at MR imaging. *Radiology*, 248:1004–1012, 2008.
- [54] N.C.F. Codella, M.D. Cham, R. Wong, C. Chu, J.K. Min, M.R. Prince, Y. Wang, and J.W. Weinsaft. Rapid and accurate left ventricular chamber quantification using a novel CMR segmentation algorithm: a clinical validation study. *Journal of Magnetic Resonance Imaging*, 31(4):845–853, 2010.
- [55] P Comon. Independent component analysis, A new concept? *Signal Processing*, 36(3): 287–314, 1994.
- [56] A. Comte, A. Lalande, S. Aho, P.M. Walker, and F. Brunotte. Realignment of myocardial first-pass MR perfusion images using an automatic detection of the heart-lung interface. *Magnetic Resonance Imaging*, 22(7):1001–1009, 2004.
- [57] T.F. Cootes, G.J. Edwards, and C.J. Taylor. Active appearance models. In *Proceedings of the 5th European Conference on Computer Vision-Volume II*, pages 484–498, 1998.
- [58] J H Cullen, M a Horsfield, C R Reek, G R Cherryman, D B Barnett, and N J Samani. A myocardial perfusion reserve index in humans using first-pass contrast-enhanced magnetic resonance imaging. *Journal of the American College of Cardiology*, 33(5):1386–94, 1999.

- [59] R.H. Davies, C.J. Twining, T.F. Cootes, J.C. Waterton, and C.J. Taylor. A minimum description length approach to statistical shape modeling. *IEEE Transactions on Medical Imaging*, 21(5): 525–537, 2002.
- [60] M. Debruyne, M. Hubert, and J.A.K. Suykens. Model selection in kernel based regression using the influence function. *Journal of Machine Learning Research (JMLR)*, 9:2377–2400, 2008.
- [61] T. Delzescaux, F. Frouin, A. De Cesare, S. Philipp-Foliguuet, R. Zeboudj, M. Janier, A. Todd-Pokropek, and A. Herment. Adaptive and self-evaluating registration method for myocardial perfusion assessment. *Magnetic Resonance Materials in Biology, Physics, and Medicine*, 13(1): 28–39, 2001.
- [62] Thierry Delzescaux, Frédérique Frouin, Alain de Cesare, Sylvie Philipp-Foliguuet, Andrew Todd-Pokropek, Alain Herment, and Marc Janier. Using an adaptive semiautomated self-evaluated registration technique to analyze MRI data for myocardial perfusion assessment. *Journal of Magnetic Resonance Imaging*, 18(6):681–690, 2003.
- [63] E. Di Bella and A. Sitek. Time curve analysis techniques for dynamic contrast MRI studies. In *Information Processing in Medical Imaging*, pages 211–217, 2001.
- [64] E. V. R. Di Bella, D. L. Parker, and A. J. Sinusas. On the dark rim artifact in dynamic contrast-enhanced MRI myocardial perfusion studies. *Magnetic Resonance in Medicine*, 54(5): 1295–1299, 2005.
- [65] M.F. Di Carli, S. Dorbala, J. Meserve, G. El Fakhri, A. Sitek, and S.C. Moore. Clinical myocardial perfusion PET/CT. *Journal of Nuclear Medicine*, 48(5):783, 2007.
- [66] R. Di Paola, J.P/ Bazin, F. Aubry, A. Aurengo, F. Cavailloles, J.Y. Herry, and E. Kahn. Handling of dynamic sequences in nuclear medicine. *IEEE Transactions on Nuclear Science*, 29(4): 1310–1321, 1982.
- [67] A. Discher, N. Rougon, and F. Preteux. An unsupervised approach for measuring myocardial perfusion in MR image sequences. In *Proceedings of SPIE*, volume 5916, pages 126–137, 2005.
- [68] C. Dornier, M.K. Ivancevic, P. Thevenaz, and J.P. Vallée. Improvement in the quantification of myocardial perfusion using an automatic spline-based registration algorithm. *Journal of Magnetic Resonance Imaging*, 18(2):160–168, 2003.
- [69] A. Dromigny-Badin, Y.M. Zhu, I. Magnin, and D. Revel. Fusion of cine magnetic resonance and contrast-enhanced first-pass magnetic resonance data in patients with coronary artery disease : A feasibility study. *Investigative radiology*, 33(1):12, 1998.
- [70] G.J. Edwards, C.J. Taylor, and T.F. Cootes. Interpreting face images using active appearance models. In *3rd International Conference on Automatic Face and Gesture Recognition*, pages 300–305, 1998.
- [71] A.C. Eichenberger, E. Schuiki, V.D. Köhli, F.W. Amann, G.C. McKinnon, and G.K. von Schulthess. Ischemic heart disease: Assessment with gadolinium-enhanced ultrafast MR imaging and dipyridamole stress. *Journal of Magnetic Resonance Imaging*, 4(3):425–431, 1994.
- [72] F.H. Epstein, V. Fuster, L. Badimon, J.J. Badimon, and J.H. Chesebro. The pathogenesis of coronary artery disease and the acute coronary syndromes. *New England Journal of Medicine*, 326(4):242–250, 1992.

- [73] T.L. Faber, C.A. Santana, E.V. Garcia, J. Candell-Riera, R.D. Folks, J.W. Peifer, A. Hopper, S. Aguade, J. Angel, and J.L. Klein. Three-dimensional fusion of coronary arteries with myocardial perfusion distributions: Clinical validation. *J Nucl Med*, 45(5):745–753, 2004.
- [74] E.P. Ficaro, B.C. Lee, J.N. Kritzman, and J.R. Corbett. Corridor4dm: the michigan method for quantitative nuclear cardiology. *Journal of nuclear cardiology*, 14(4):455–465, 2007.
- [75] F. Fidler, C.M. Wacker, C. Dueren, M. Weigel, P.M. Jakob, W.R. Bauer, and A. Haase. Myocardial perfusion measurements by spin-labeling under different vasodynamic states. *Journal of Cardiovascular Magnetic Resonance*, 6(2):509–516, 2004.
- [76] D.S. Fieno, S.M. Shea, Y. Li, K.R. Harris, J.P. Finn, and D. Li. Myocardial perfusion imaging based on the blood oxygen level-dependent effect using T2-prepared steady-state free-precession magnetic resonance imaging. *Circulation*, 110(10):1284, 2004.
- [77] D.J. Fleet and A.D. Jepson. Computation of component image velocity from local phase information. *International Journal of Computer Vision*, 5(1):77–104, 1990.
- [78] R.D. Folks. Interpretation and reporting of myocardial perfusion spect: A summary for technologists. *Journal of nuclear medicine technology*, 30(4):153–163, 2002.
- [79] H. Frank and S. Globits. Magnetic resonance imaging evaluation of myocardial and pericardial disease. *Journal of Magnetic Resonance Imaging*, 10(5):617–626, 1999.
- [80] F. Frouin, J.P. Bazin, M. Di Paola, O. Jolivet, and R. Di Paola. FAMIS: a software package for functional feature extraction from biomedical multidimensional images. *Computerized Medical Imaging and Graphics*, 16(2):81–91, 1992.
- [81] O. Gaemperli, T. Schepis, V. Kalff, M. Namdar, I. Valenta, L. Stefani, L. Desbiolles, S. Leschka, L. Husmann, H. Alkadhi, et al. Validation of a new cardiac image fusion software for three-dimensional integration of myocardial perfusion spect and stand-alone 64-slice ct angiography. *European journal of nuclear medicine and molecular imaging*, 34(7):1097–1106, 2007.
- [82] O. Gaemperli, T. Schepis, I. Valenta, L. Husmann, H. Scheffel, V. Duerst, F.R. Eberli, T.F. Luscher, H. Alkadhi, and P.A. Kaufmann. Cardiac image fusion from stand-alone spect and ct: clinical experience. *Journal of Nuclear Medicine*, 48(5):696–703, 2007.
- [83] O. Gaemperli, L. Husmann, T. Schepis, P. Koepfli, I. Valenta, W. Jenni, H. Alkadhi, T.F. Lüscher, and P.A. Kaufmann. Coronary ct angiography and myocardial perfusion imaging to detect flow-limiting stenoses: a potential gatekeeper for coronary revascularization? *European heart journal*, 30(23):2921–2929, 2009.
- [84] C. M Gallippi, C. M. Kramer, Y. Hu, D. A. Vido, N. Reichek, and W. J. Rogers. Fully automated registration and warping of contrast-enhanced first-pass perfusion images. *Journal of Cardiovascular Magnetic Resonance*, 4(4):459–469, 2002.
- [85] Caterina M Gallippi and Gregg E Trahey. Automatic Image Registration for MR and Ultrasound Cardiac Images. *Ultrasound*, pages 148–154, 2001.
- [86] J. Gao, N. Ablitt, A. Elkington, and G. Z. Yang. In *Proceedings of the International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, pages 612–619, 2002.

- [87] S. Geman and D. Geman. Stochastic relaxation, Gibbs distributions and the Bayesian restoration of images. *Journal of Applied Statistics*, 20(5):25–62, 1993.
- [88] B. Gerber, S. Raman, K. Nayak, F. Epstein, P. Ferreira, L. Axel, and D. Kraitchman. Myocardial first-pass perfusion cardiovascular magnetic resonance : history, theory, and current state of the art. *Journal of Cardiovascular Magnetic Resonance*, 10(1):18, 2008.
- [89] R.T. Go, T.H. Marwick, W.J. MacIntyre, G.B. Saha, D.R. Neumann, D.A. Underwood, and C.C. Simpfordorfer. A prospective comparison of rubidium-82 PET and thallium-201 SPECT myocardial perfusion imaging utilizing a single dipyridamole stress in the diagnosis of coronary artery disease. *Journal of Nuclear Medicine*, 31(12):1899, 1990.
- [90] J.C. Gower. Generalized procrustes analysis. *Psychometrika*, 40(1):33–51, 1975.
- [91] M. Guizar-Sicairos, S.T. Thurman, and J.R. Fienup. Efficient sub-pixel image registration algorithms. *Optics letters*, 33(2):156–158, 2008. URL <http://www.mathworks.com/matlabcentral/fileexchange/18401-efficient-subpixel-image-registration-by-cross-correlation>.
- [92] S.N. Gupta, M. Solaiyappan, G.M. Beache, A.E. Arai, and T.K.F. Foo. Fast method for correcting image misregistration due to organ motion in time-series MRI data. *Magnetic resonance in medicine*, 49(3):506–514, 2003.
- [93] V. Gupta, E.A. Hendriks, J. Milles, R.J. Van Der Geest, M. Jerosch-Herold, J.H.C. Reiber, and B.P.F. Lelieveldt. Fully automatic registration and segmentation of first-pass myocardial perfusion MR image sequences. *Academic radiology*, 17(11):1375–1385, 2010.
- [94] V. Gupta, H.A. Kirişli, E.A. Hendriks, R.J. van der Geest, M. van de Giessen, W. Niessen, J.H.C. Reiber, and B.P.F. Lelieveldt. Cardiac MR perfusion image processing techniques: A survey. *Medical Image Analysis*, 16(4):767–785, 2012.
- [95] V. Gupta, M. Van de Giessen, H.A. Kirişli, S. W. Kirschbaum, W. J. Niessen, and B.P.F. Lelieveldt. Robust motion correction in the frequency domain of cardiac MR stress perfusion sequences. pages 667–674, 2012.
- [96] R. Hachamovitch, D.S. Berman, L.J. Shaw, H. Kiat, I. Cohen, J.A. Cabico, J. Friedman, and G.A. Diamond. Incremental prognostic value of myocardial perfusion single photon emission computed tomography for the prediction of cardiac death: differential stratification for risk of cardiac death and myocardial infarction. *Circulation*, 97(6):535–543, 1998.
- [97] G.K. Hansson. Inflammation, atherosclerosis, and coronary artery disease. *New England Journal of Medicine*, 352(16):1685–1695, 2005.
- [98] E. Heiberg, J. Sjögren, M. Ugander, M. Carlsson, H. Engblom, and H. Arheden. Design and validation of segment-freely available software for cardiovascular image analysis. *BMC medical imaging*, 10(1):1, 2010.
- [99] A. Hennemuth, T. Boskamp, D. Fritz, C. Kühnel, S. Bock, D. Rinck, M. Scheuering, and H.O. Peitgen. One-click coronary tree segmentation in CT angiographic images. In *Proceedings of International Congress Series*, volume 1281, pages 317–321, 2005.
- [100] A. Hennemuth, S. Behrens, C. Kuehnel, S. Oeltze, O. Konrad, and H.O. Peitgen. Novel methods for parameter-based analysis of myocardial tissue in MR images. In *Proceedings of SPIE*, volume 6511, page 54, 2007.



- [101] A. Hennemuth, A. Seeger, O. Friman, S. Miller, B. Klumpp, S. Oeltze, and H.O. Peitgen. A comprehensive approach to the analysis of contrast enhanced cardiac MR images. *IEEE Transactions on Medical Imaging*, 27(11):1592–1610, 2008.
- [102] A. Hennemuth, A. Seeger, O. Friman, S. Miller, and HO Peitgen. Automatic detection and quantification of non-viable myocardium in late enhancement images. In *Proceedings of International Society for Magnetic Resonance in Medicine*, 2008.
- [103] B. Hesse, K. Tägil, A. Cuocolo, C. Anagnostopoulos, M. Bardies, J. Bax, F. Bengel, E. Busemann Sokole, G. Davies, M. Dondi, et al. EANM/ESC procedural guidelines for myocardial perfusion imaging in nuclear cardiology. *European Journal of Nuclear Medicine and Molecular Imaging*, 32(7):855–897, 2005.
- [104] K.H. Hiller, M. Bock, CM Wacker, LR Schad, C. Waller, A. Haase, G. Van Kaick, G. Ertl, and WR Bauer. MR-perfusion measurements: Basic methodology and current status. *Magnetic Resonance Materials in Physics, Biology and Medicine*, 6(2):98–99, 1998.
- [105] AE Holland, JW Goldfarb, and RR Edelman. Diaphragmatic and cardiac motion during suspended breathing: preliminary experience and implications for breath-hold MR imaging. *Radiology*, 209(2):483, 1998.
- [106] A. Hyvärinen and E. Oja. Independent component analysis: algorithms and applications. *Neural Networks*, 13(4-5):411–430, 2000.
- [107] K Iwata and K Ogasawara. Comparison of the cost-effectiveness of stress myocardial perfusion mri and spect in patients with suspected coronary artery disease. *Radiological Physics and Technology*, pages 1–7, 2012.
- [108] E. Jackson, N. Bellenger, M. Seddon, S. Harden, and C. Peebles. Ischaemic and non-ischaemic cardiomyopathies—cardiac MRI appearances with delayed enhancement. *Clinical Radiology*, 62(5):395–403, 2007.
- [109] M.F. Janier, A.N. Mazzadi, M. Lionnet, F. Frouin, X. Andre-Fouet, L. Cinotti, D. Revel, and P. Croisille. Factor analysis of medical image sequences improves evaluation of first-pass mr imaging acquisitions for myocardial perfusion. *Academic Radiology*, 9(1):26–39, 2002.
- [110] M. Jerosch-Herold. Quantification of myocardial perfusion by cardiovascular magnetic resonance. *Journal of Cardiovascular Magnetic Resonance*, 12(1):1–16, 2010.
- [111] M. Jerosch-Herold, C. Swingen, and R.T. Seethamraju. Myocardial blood flow quantification with MRI by model-independent deconvolution. *Medical Physics*, 29:886, 2002.
- [112] M. Jerosch-Herold, R.T. Seethamraju, C.M. Swingen, N.M. Wilke, and A.E. Stillman. Analysis of myocardial perfusion MRI. *Journal of Magnetic Resonance Imaging*, 19(6):758–770, 2004.
- [113] R Jogiya, A Schuster, A Zaman, Y Samaroo, E Nagel, S Kozerke, and S Plein. Feasibility of three-dimensional (3d) balanced steady-state-free-precession (bssfp) myocardial perfusion mri at 3 tesla using local rf shimming with dual-source rf transmission. *Journal of Cardiovascular Magnetic Resonance*, 15(Suppl 1):P23, 2013.
- [114] T. A. Kaandorp, H. .J.. Lamb, J. .J. Bax, E. E. van der Wall, and A. de Roos. Magnetic resonance imaging of coronary arteries, the ischemic cascade, and myocardial infarction. *American Heart Journal*, 149(2):200–208, 2005.

- [115] P.A. Kaufmann. Cardiac hybrid imaging: state-of-the-art. *Annals of nuclear medicine*, 23(4): 325–331, 2009.
- [116] Kazunori Kawakami, Kenya Murase, Masayuki Kumashiro, Atsushi Kawakami, Nobuyoshi Tanki, Youichi Yamazaki, Takashi Tabuchi, Masako Nagayama, and Yuji Watanabe. Automatic motion correction for quantification of myocardial perfusion with dynamic magnetic resonance imaging. *Magnetic Resonance in Medical Sciences*, 3(3):105–117, 2004.
- [117] J.T. Keijzer, A.C. van Rossum, M.J. van Eenige, A.J.P. Karreman, M. Hofman, J. Valk, and C.A. Visser. Semiquantitation of regional myocardial blood flow in normal human subjects by first-pass magnetic resonance imaging. *American Heart Journal*, 130(4):893–901, 1995.
- [118] P. Kellman and A.E. Arai. Imaging sequences for first pass perfusion - A review. *Journal of Cardiovascular Magnetic Resonance*, 9(3):525–537, 2007.
- [119] H. A. Kirişli, V. Gupta, S. Kirschbaum, L. Neefjes, R. J. van Geuns, N. R. A. Mollet, B. P. F. Lelieveldt, J. H. C. Reiber, T. van Walsum, and W.J. Niessen. Integrated visualization and analysis of coronary arteries and myocardial perfusion. *International Journal of Computer Assisted Radiology and Surgery (CARS)*, 5(1):S323, 2010.
- [120] H.A. Kirişli, M. Schaap, S. Klein, S.L. Papadopoulou, M. Bonardi, C.H. Chen, A.C. Weustink, N.R.A. Mollet, E. P. A. Vonken, R.J. van der Geest, T. van Walsum, and W.J. Niessen. Evaluation of a multi-atlas based method for segmentation of cardiac CTA data: a large-scale, multi-center and multi-vendor study. *Medical Physics*, 37(12):6279–6292, 2010.
- [121] H.A. Kirişli, V. Gupta, S.W. Kirschbaum, L. Neefjes, R.J. van Geuns, N. Mollet, B.P.F. Lelieveldt, J.H.C. Reiber, T. van Walsum, and W.J. Niessen. A patient-specific visualization tool for comprehensive analysis of coronary CTA and perfusion MRI data. In *Proc. of SPIE Medical Imaging*, 2011.
- [122] H.A. Kirişli, V. Gupta, SW Kirschbaum, A. Rossi, CT Metz, M. Schaap, RJ van Geuns, N. Mollet, B.P.F. Lelieveldt, J.H.C. Reiber, et al. Comprehensive visualization of multimodal cardiac imaging data for assessment of coronary artery disease: first clinical results of the SMARTVis tool. *International journal of computer assisted radiology and surgery*, 7(4):557–571, 2012.
- [123] S.W. Kirschbaum and R.J. van Geuns. Cardiac magnetic resonance imaging to detect and evaluate ischemic heart disease. *Hell J Cardiol*, 50:119–126, 2009.
- [124] S.W. Kirschbaum, T. Baks, M. van den Ent, G. Sianos, G.P. Krestin, P.W. Serruys, P.J. de Feyter, and R.J.M. van Geuns. Evaluation of left ventricular function three years after percutaneous recanalization of chronic total coronary occlusions. *The American journal of cardiology*, 101(2):179–185, 2008.
- [125] S.W. Kirschbaum, T. Springeling, A. Rossi, E. Duckers, J.L. Gutierrez-Chico, E. Regar, P.J. de Feyter, and R.J. van Geuns. Comparison of adenosine magnetic resonance perfusion imaging with invasive coronary flow reserve and fractional flow reserve in patients with suspected coronary artery disease. *Int J Cardiol*, 147:184–186, 2011.
- [126] D.E. Kivelitz, K.G. Bis, N.M. Wilke, M. Jerosch-Herold, J.E. Juni, and B.K. Hamm. Quantitative MR first pass perfusion imaging demonstrates coronary artery patency post interventions. *Radiology*, 205:254, 1997.
- [127] M.A. Klein, B.D. Collier, R.S. Hellman, and V.S. Bamrah. Detection of chronic coronary artery disease: value of pharmacologically stressed, dynamically enhanced turbo-fast low-angle shot MR images. *American Journal of Roentgenology*, 161(2):257, 1993.

- [128] F.J. Klocke and D. Li. Testing coronary flow reserve without a provocative stress: A "bold" approach. *Journal of the American College of Cardiology*, 41(5):841, 2003.
- [129] H. Knutsson and M. Andersson. Morphons: Segmentation using elastic canvas and paint on priors. In *Proceedings of International Conference on Image Processing*, volume 2, 2005.
- [130] M. Kocher, JL Daire, P. Thevenaz, T. Schindler, PF Keller, D. Didier, and JP Vallee. Myocardial perfusion assessment by use of system identification method in a one-compartment model. In *Proceedings of 29th Annual International Conference of the IEEE- Engineering in Medicine and Biology Society*, pages 4492–4495, 2007.
- [131] H. Köstler, C. Ritter, M. Reiss-Zimmermann, M. Beer, D. Hahn, and J. Sandstede. Correction for partial volume errors in MR heart perfusion imaging. *Magnetic Resonance in Medicine*, 51(4):848–852, 2004.
- [132] C. Kühnel, A. Hennemuth, S. Oeltze, T. Boskamp, and H.O. Peitgen. Enhanced cardio vascular image analysis by combined representation of results from dynamic MRI and anatomic CTA. In *Proc. of SPIE Medical Imaging*, volume 6918, 2008.
- [133] C. Kühnel, A. Hennemuth, H.O. Peitgen, and A.H. Mahnken. New analysis tools for the comprehensive assessment of the coronary arteries and myocardial viability in CT data sets. In *Proc. of Computers in Cardiology*, volume 35, pages 733–736, 2008.
- [134] Henrik BW Larsson, Thomas Fritz-Hansen, Egill Rostrup, Lars Søndergaard, Poul Ring, and Ole Henriksen. Myocardial perfusion modeling using mri. *Magnetic resonance in medicine*, 35(5):716–726, 2005.
- [135] D. C. Lee and N. P. Johnson. Quantification of absolute myocardial blood flow by magnetic resonance perfusion imaging. *JACC: Cardiovascular Imaging*, 2(6):761, 2009.
- [136] T.W. Lee. *Independent component analysis: theory and applications*. 1998.
- [137] G.N. Levine, E.R. Bates, J.C. Blankenship, S.R. Bailey, J.A. Bittl, B. Cercek, C.E. Chambers, S.G. Ellis, R.A. Guyton, S.M. Hollenberg, et al. 2011 accf/aha/scai guideline for percutaneous coronary intervention: a report of the american college of cardiology foundation/american heart association task force on practice guidelines and the society for cardiovascular angiography and interventions. *Journal of the American College of Cardiology*, 58(24):e44–e122, 2011.
- [138] C. Li and Y. Sun. Nonrigid registration of myocardial perfusion MRI using pseudo ground truth. In *Proceedings of the International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, volume 12, pages 165–72, 2009.
- [139] S. Li. Markov random field models in computer vision. In *European Conference on Computer Vision*, pages 361–370, 1994.
- [140] P. Libby. Inflammation in atherosclerosis. *Nature*, 420(6917):868–874, 2002.
- [141] L. Lorenzo, R. S. MacLeod, R. T. Whitaker, G. Adluru, and E. V. R. DiBella. Level sets and shape models for segmentation of cardiac perfusion MRI. In *Proceedings of SPIE*, volume 6144, pages 1320–1331, 2006.
- [142] T. Mäkelä, P. Clarysse, O. Sipilä, N . Pauna, Q. C. Pham, and I. E Katila, T .and Magnin. A review of cardiac image registration methods. *IEEE Transactions on Medical Imaging*, 21(9): 1011–21, 2002.

- [143] W.J. Manning and D.J. Pennell. *Cardiovascular magnetic resonance*, volume 18. 2002.
- [144] W.J. Manning, D.J. Atkinson, W. Grossman, S. Paulin, and R.R. Edelman. First-pass nuclear magnetic resonance imaging studies using gadolinium-DTPA in patients with coronary artery disease. *Journal of the American College of Cardiology*, 18(4):959–965, 1991.
- [145] M.L. Marcus, R.F. Wilson, and C.W. White. Methods of measurement of myocardial blood flow in patients: a critical review. *Circulation*, 76(2):245–53, 1987.
- [146] A.L. Martel, A.R. Moody, S.J. Allder, G.S. Delay, and P.S. Morgan. Extracting parametric images from dynamic contrast-enhanced MRI studies of the brain using factor analysis. *Medical Image Analysis*, 5(1):29, 2001.
- [147] K. McLeish, D.L.G. Hill, D. Atkinson, J.M. Blackall, and R. Razavi. A study of the motion and deformation of the heart due to respiration. *IEEE Transactions on Medical Imaging*, 21(9):1142–1150, 2002.
- [148] MEDIS. QMass MR. 2005.
- [149] M. Mellor and M. Brady. Non-rigid multimodal image registration using local phase. In *Proceedings of the International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, pages 789–796, 2004.
- [150] M. Mellor and M. Brady. Phase mutual information as a similarity measure for registration. *Medical Image Analysis*, 9(4):330–343, 2005.
- [151] C.T. Metz, M. Schaap, A.C. Weustink, N.R.A. Mollet, T. van Walsum, and W.J. Niessen. Coronary centerline extraction from CT coronary angiography images using a minimum cost path approach. *Medical Physics*, 36(12):5568–5579, 2009.
- [152] MICCAI Challenges. Grand challenges in Medical Image Analysis. 2007-2010.
- [153] D.D. Miller, G. Holmvang, J.B. Gill, D. Dragotakes, H.L. Kantor, R.D. Okada, and T.J. Brady. MRI detection of myocardial perfusion changes by gadolinium-DTPA infusion during dipyridamole hyperemia. *Magnetic Resonance in Medicine*, 10(2):246–255, 1989.
- [154] J. Milles, R. J. van der Geest, M. Jerosch-Herold, J. H. C. Reiber, and B. P. F. Lelieveldt. Fully automated motion correction in first-pass myocardial perfusion MR image sequences. *IEEE Transactions on Medical Imaging*, 27(11):1611–21, 2008.
- [155] M Minoves, A Garcia, J Pavia, and R Herranz. Evaluation of myocardial perfusion defects by means of "bull's eye" images. *Clinical Cardiology*, 22:16–22.
- [156] S.C. Mitchell, B.P.F. Lelieveldt, R.J. van der Geest, H.G. Bosch, JHC Reiver, and M. Sonka. Multistage hybrid active appearance model matching: segmentation of left and right ventricles in cardiac MR images. *IEEE Transactions on medical imaging*, 20(5):415–423, 2001.
- [157] S.C. Mitchell, J.G. Bosch, B.P.F. Lelieveldt, R.J. van der Geest, J.H.C. Reiber, and M. Sonka. 3-D active appearance models: segmentation of cardiac MR and ultrasound images. *IEEE Transactions on Medical Imaging*, 21(9):1167–1178, 2002.
- [158] G. Mowatt, E. Cummins, N. Waugh, S. Walker, J.A. Cook, X. Jia, G.S. Hillis, and C.M. Fraser. Systematic review of the clinical effectiveness and cost-effectiveness of 64-slice or higher computed tomography angiography as an alternative to invasive coronary angiography in the investigation of coronary artery disease. *Health Technology Assessment*, 12:17, 2008.

- [159] O.M. Mühling, Y. Wang, P. Panse, M. Jerosch-Herold, M.M. Cayton, L.S. Wann, M.M. Mirhoseini, and N.M. Wilke. Transmyocardial laser revascularization preserves regional myocardial perfusion: an MRI first pass perfusion study. *Cardiovascular Research*, 57(1):63, 2003.
- [160] W. Müller, T. Nocke, and H. Schumann. Enhancing the visualization process with principal component analysis to support the exploration of trends. In *Proceedings of the 2006 Asia-Pacific Symposium on Information Visualisation-Volume 60*, pages 121–130, 2006.
- [161] E. Nagel, R. Underwood, D. Pennell, UP Sechtem, S. Neubauers, S. Wickline, OM Hess, M. Schwaiger, and E. Fleck. New developments in non-invasive cardiac imaging: critical assessment of the clinical role of cardiac magnetic resonance imaging. *European Heart Journal*, 19(9):1286–1293, 1998.
- [162] E. Nagel, C. Klein, I. Paetsch, S. Hettwer, B. Schnackenburg, K. Wegscheider, and E. Fleck. Magnetic resonance perfusion measurements for the noninvasive detection of coronary artery disease. *Circulation*, 108(4):432–7, 2003.
- [163] T. Nakaura, D. Utsunomiya, S. Shiraishi, S. Tomiguchi, T. Honda, H. Ogawa, K. Awai, and Y. Yamashita. Three-dimensional cardiac image fusion using new ct angiography and spect methods. *American Journal of Roentgenology*, 185(6):1554–1557, 2005.
- [164] O. Nestares and D.J. Heeger. Robust multiresolution alignment of MRI brain volumes. *Magnetic Resonance in Medicine*, 43(5):705–715, 2000.
- [165] R.W. Nesto and G.J. Kowalchuk. The ischemic cascade: temporal sequence of hemodynamic, electrocardiographic and symptomatic expressions of ischemia. *The American Journal of Cardiology*, 59(7):C23–C30, 1987.
- [166] T Netsch and A Van Muiswinkel. Automated registration and outlier identification for the improvement of cardiac MR perfusion quantification. In *19th annual meeting of the ESMRMB, Cannes, France*, page 105, 2002.
- [167] S. Oeltze and B. Preim. Visualization of vasculature with convolution surfaces: method, validation and evaluation. *IEEE Transactions on Medical Imaging*, 24(4):540–548, 2005.
- [168] S Oeltze, F Grothues, A Hennemuth, A Kuß, and B Preim. Integrated visualization of morphologic and perfusion data for the analysis of coronary artery disease. In *Proceedings of Eurographics/IEEE-VGTC Symposium on Visualization*, pages 131–138, 2006.
- [169] S. Oeltze, H. Doleisch, H. Hauser, P. Muigg, and B. Preim. Interactive visual analysis of perfusion data. *IEEE Transactions on Visualization and Computer Graphics*, 13(6):1392–1399, 2007.
- [170] S. Oeltze, A. Hennemuth, S. Glaßer, C. Kühnel, and B. Preim. Glyph-based visualization of myocardial perfusion data and enhancement with contractility and viability information. In *Proceedings of Eurographics Workshop on Visual Computing for Biomedicine (2008)*, 2008.
- [171] H. Ólafsdóttir, M. B. Stegmann, B.K. Ersbøll, and H. B. W. Larsson. A comparison of FFD-based nonrigid registration and AAMs applied to myocardial perfusion MRI. In *Proceedings of SPIE*, volume 6144, pages 388–396, 2006.
- [172] S. Osher and J.A. Sethian. Fronts propagating with curvature-dependent speed: algorithms based on hamilton-jacobi formulations. *Journal of Computational Physics*, 79(1):12–49, 1988.

- [173] M. Oudkerk, A.E. Stillman, S.S. Halliburton, W.A. Kalender, S. Möhlenkamp, C.H. McCollough, R. Vliegenthart, L.J. Shaw, W. Stanford, A.J. Taylor, et al. Coronary artery calcium screening: current status and recommendations from the european society of cardiac radiology and north american society for cardiovascular imaging. *The International Journal of Cardiovascular Imaging (formerly Cardiac Imaging)*, 24(6):645–671, 2008.
- [174] N. Paragios. A level set approach for shape-driven segmentation and tracking of the left ventricle. *IEEE Transactions on Medical Imaging*, 22(6):773–776, 2003.
- [175] H. Penzkoffer, BJ Wintersperger, A. Knez, J. Weber, and M. Reiser. Assessment of myocardial perfusion using multisection first-pass MRI and color-coded parameter maps: A comparison to 99mTc Sesta MIBI SPECT and systolic myocardial wall. *Magnetic Resonance Imaging*, 17(2):161–170, 1999.
- [176] O. Pereztol-Valdés, J. Candell-Riera, C. Santana-Boado, J. Angel, S. Aguadé-Bruix, J. Castell-Conesa, E.V. Garcia, and J. Soler-Soler. Correspondence between left ventricular 17 myocardial segments and coronary arteries. *Eur Heart J*, 26(24):637–2643, 2005.
- [177] O. Pereztol-Valdés, J. Candell-Riera, C. Santana-Boado, J. Angel, S. Aguadé-Bruix, J. Castell-Conesa, E.V. Garcia, and J. Soler-Soler. Correspondence between left ventricular 17 myocardial segments and coronary arteries. *European heart journal*, 26(24):2637–2643, 2005.
- [178] NH Pijls, GJ Uijen, A. Hoevelaken, T. Arts, WR Aengevaeren, HS Bos, JH Fast, KL Van Leeuwen, and T. Van der Werf. Mean transit time for the assessment of myocardial perfusion by videodensitometry. *Circulation*, 81(4):1331–1340, 1990.
- [179] S. Plein, J. Greenwood, and J.P. Ridgway. *Cardiovascular MR Manual*. 2010.
- [180] C. Pluempitiwiriyaewj and S. Sotthivirat. Active contours with automatic initialization for myocardial perfusion analysis. In *27th Annual International Conference of the Engineering in Medicine and Biology Society*, pages 3332–3335, 2005.
- [181] C. Pluempitiwiriyaewj, J.M.F. Moura, Y.J.L. Wu, and C. Ho. STACS: New active contour scheme for cardiac MR image segmentation. *IEEE Transactions on Medical Imaging*, 24(5): 593–603, 2005.
- [182] B. Preim, S. Oeltze, M. Mlejnek, E. Gröller, A. Hennemuth, and S. Behrens. Survey of the visual exploration and analysis of perfusion data. *IEEE Transactions on Visualization and Computer Graphics*, 15(2):205–220, 2009.
- [183] K.P. Pruessmann, M. Weiger, M.B. Scheidegger, and P. Boesiger. SENSE: Sensitivity encoding for fast MRI. *Magnetic Resonance in Medicine*, 42(5):952–962, 1999.
- [184] G.L. Raff, A. Abidov, S. Achenbach, D.S. Berman, L.M. Boxt, M.J. Budoff, V. Cheng, T. DeFrance, J.C. Hellinger, R.P. Karlsberg, et al. Scct guidelines for the interpretation and reporting of coronary computed tomographic angiography. *Journal of cardiovascular computed tomography*, 3(2):122–136, 2009.
- [185] A. Ranhosky and J. Kempthorne-Rawson. The safety of intravenous dipyridamole thallium myocardial perfusion imaging. Intravenous dipyridamole thallium imaging study group. *Circulation*, 81(4):1205–1209, 1990.

- [186] JH Reiber, P.W. Serruys, C.J. Kooijman, W. Wijns, C.J. Slager, JJ Gerbrands, JC Schuurbijs, A. Den Boer, and PG Hugenholz. Assessment of short-, medium-, and long-term variations in arterial dimensions from computer-assisted quantitation of coronary cineangiograms. *Circulation*, 71(2):280–288, 1985.
- [187] V.L. Roger, A.S. Go, D.M. Lloyd-Jones, E.J. Benjamin, J.D. Berry, W.B. Borden, D.M. Bravata, S. Dai, E.S. Ford, C.S. Fox, et al. Heart disease and stroke statistics—2012 update a report from the american heart association. *Circulation*, 125(1):e2–e220, 2012.
- [188] N. Rougon, A. Discher, and F. Preteux. Region-driven statistical nonrigid registration: application to model-based segmentation and tracking of the heart in perfusion MRI. In *Proc. SPIE*, volume 5916, pages 148–159, 2005.
- [189] N. Rougon, A. Discher, and F. Preteux. Region-driven statistical nonrigid registration: application to model-based segmentation and tracking of the heart in perfusion MRI. In *Proceedings of SPIE*, volume 5916, pages 148–159, 2005.
- [190] N. Rougon, C. Petitjean, F. Prêteux, P. Cluzel, and P. Grenier. A non-rigid registration approach for quantifying myocardial contraction in tagged MRI using generalized information measures. *Medical Image Analysis*, 9(4):353–375, 2005.
- [191] D. Rueckert, L.I. Sonoda, C. Hayes, D.L.G. Hill, M.O. Leach, and D.J. Hawkes. Nonrigid registration using free-form deformations: application to breast MR images. *IEEE Transactions on Medical Imaging*, 18(8):712–721, 1999.
- [192] V.M. Runge, W. Schoerner, H.P. Niendorf, M. Laniado, D. Koehler, C. Claussen, R. Felix, and A.E. James Jr. Initial clinical evaluation of gadolinium DTPA for contrast-enhanced magnetic resonance imaging. *Magnetic Resonance Imaging*, 3(1):27–35, 1985.
- [193] N.K. Sabharwal and A. Lahiri. Role of myocardial perfusion imaging for risk stratification in suspected or known coronary artery disease. *Heart*, 89(11):1291, 2003.
- [194] D. M. Sait, E. Yılmaz, S. Doğan, B. Akdeniz, N. Barış, U. Eomete, and L. Iyilikçi. Evaluation of myocardial ischemia in coronary artery disease with cardiac mr perfusion method: comparison with the results of catheter or ct angiography. *Medicinski glasnik: official publication of the Medical Association of Zenica-Doboj Canton, Bosnia and Herzegovina*, 10(1): 63, 2013.
- [195] H. Sakuma, N. Suzawa, Y. Ichikawa, K. Makino, T. Hirano, K. Kitagawa, and K. Takeda. Diagnostic accuracy of stress first-pass contrast-enhanced myocardial perfusion MRI compared with stress myocardial perfusion scintigraphy. *American Journal of Roentgenology*, 185(1):95, 2005.
- [196] A.D. Sam, M.D. Morasch, J. Collins, G. Song, R. Chen, and F.S. Pereles. Safety of gadolinium contrast angiography in patients with chronic renal insufficiency. *Journal of Vascular Surgery*, 38(2):313–318, 2003.
- [197] C.A. Santana, E.V. Garcia, T.L. Faber, G.K.R. Sirineni, F.P. Esteves, R. Sanyal, R. Halkar, M. Ornelas, L. Verdes, S. Lerakis, et al. Diagnostic performance of fusion of myocardial perfusion imaging (mpi) and computed tomography coronary angiography. *Journal of nuclear cardiology*, 16(2):201–211, 2009.
- [198] M. Santarelli. Automated cardiac MR image segmentation: theory and measurement evaluation. *Medical Engineering & Physics*, 25(2):149–159, 2003.

- [199] A. Sato, T. Nozato, H. Hikita, S. Miyazaki, Y. Takahashi, T. Kuwahara, A. Takahashi, M. Hiroe, and K. Aonuma. Incremental value of combining 64-slice computed tomography angiography with stress nuclear myocardial perfusion imaging to improve noninvasive detection of coronary artery disease. *Journal of nuclear cardiology*, 17(1):19–26, 2010.
- [200] M. Schaap, L. Neeffjes, C. Metz, A. van der Giessen, A. Weustink, N. Mollet, J. Wentzel, T. van Walsum, and W. Niessen. Coronary lumen segmentation using graph cuts and robust kernel regression. In *Information Processing in Medical Imaging*, pages 528–539, 2009.
- [201] M. Schaap, T. van Walsum, L. Neeffjes, C.T. Metz, E. Capuano, M. de Bruijne, and W.J. Niessen. Robust shape regression for supervised vessel segmentation and its application to coronary segmentation in CTA. *IEEE Transactions on Medical Imaging*, in press, 2011.
- [202] S. Schaefer, R.A. Lange, D.P. Gutekunst, R.W. Parkey, J.T. Willerson, and R.M. Peshock. Contrast-enhanced magnetic resonance imaging of hypoperfused myocardium. *Investigative Radiology*, 26(6):551, 1991.
- [203] A. Schenk, G. Prause, and H.O. Peitgen. Efficient semiautomatic segmentation of 3d objects in medical images. In *Proceedings of the International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, pages 71–131, 2000.
- [204] T.H. Schindler, N. Magosaki, M. Jeserich, U. Oser, T. Krause, R. Fischer, E. Moser, E. Nitzsche, M. Zehender, H. Just, et al. Fusion imaging: combined visualization of 3D reconstructed coronary artery tree and 3D myocardial scintigraphic image in coronary artery disease. *The International Journal of Cardiac Imaging*, 15(5):357–368, 1999.
- [205] A Schöllhuber. Fully Automatic Segmentation of the Myocardium in Cardiac Perfusion MRI. In *Proceedings of Central European Seminar on Computer Graphics*, 2008.
- [206] A.J. Scholte, C.J. Roos, and J.M. van Werkhoven. Function and anatomy: SPECT-MPI and MSCT coronary angiography. *EuroIntervention*, 6(G):94–100, 2010.
- [207] W.G. Schreiber, M. Schmitt, P. Kalden, O.K. Mohrs, K.F. Kreitner, and M. Thelen. Dynamic contrast-enhanced myocardial perfusion imaging using saturation-prepared TrueFISP. *Journal of Magnetic Resonance Imaging*, 16(6):641–652, 2002.
- [208] S. Schroeder, A.F. Kopp, A. Baumbach, C. Meisner, A. Kuettner, C. Georg, B. Ohnesorge, C. Herdeg, C.D. Claussen, and K.R. Karsch. Noninvasive detection and evaluation of atherosclerotic coronary plaques with multislice computed tomography1. *Journal of the American College of Cardiology*, 37(5):1430–1435, 2001.
- [209] J.D. Schuijff, W. Wijns, J.W. Jukema, D.E. Atsma, A. de Roos, H.J. Lamb, M.P.M. Stokkel, P. Dibbets-Schneider, I. Decramer, P. De Bondt, et al. Relationship between noninvasive coronary angiography with multi-slice computed tomography and myocardial perfusion imaging. *Journal of the American College of Cardiology*, 48(12):2508–2514, 2006.
- [210] C.L. Schultz, RJ Alfydi, A.D. Nelson, S.Y. Kopywoda, and M.E. Clampitt. The effect of motion on two-dimensional fourier transformation magnetic resonance images. *Radiology*, 152(1): 117, 1984.
- [211] M. Schwaiger. Myocardial perfusion imaging with PET. *Journal of Nuclear Medicine*, 35(4): 693, 1994.



- [212] J. Schwitter, D. Nanz, S. Kneifel, K. Bertschinger, M. Buchi, PR Knusel, B. Marincek, TF Luscher, and GK Von Schulthess. Assessment of myocardial perfusion in coronary artery disease by magnetic resonance: a comparison with positron emission tomography and coronary angiography. *Circulation*, 103(18):2230, 2001.
- [213] J. Schwitter, C. M. Wacker, A. C. Van Rossum, M. Lombardi, N. Al-Saadi, H. Ahlstrom, T. Dill, H. B. W. Larsson, S. D. Flamm, Moritz M., et al. MR-IMPACT: comparison of perfusion-cardiac magnetic resonance with single-photon emission computed tomography for the detection of coronary artery disease in a multicentre, multivendor, randomized trial. *European heart journal*, 29(4):480–489, 2008.
- [214] J. Schwitter, C. M. Wacker, N. Wilke, N. Al-Saadi, E. Sauer, K. Huettle, S. O. Schönberg, A. Luchner, O. Strohm, H. Ahlstrom, et al. Mr-impact ii: Magnetic resonance imaging for myocardial perfusion assessment in coronary artery disease trial: perfusion-cardiac magnetic resonance vs. single-photon emission computed tomography for the detection of coronary artery disease: a comparative multicentre, multivendor trial. *European Heart Journal*, 2012.
- [215] A. D. Scott, J. Keegan, and D. N. Firmin. Motion in cardiovascular MR imaging. *Radiology*, 250(2):331, 2009.
- [216] I. M. Scott, T. Cootes, and C. Taylor. Improving appearance model matching using local image structure. In *Proceedings of Information Processing in Medical Imaging*, pages 258–269, 2003.
- [217] D. Selle, B. Preim, A. Schenk, and H.O. Peitgen. Analysis of vasculature for liver surgical planning. *IEEE Transactions on Medical Imaging*, 21(11):1344–1357, 2003.
- [218] R.M. Setser, T.P. O'Donnell, N.G. Smedira, J.F. Sabik, S.S. Halliburton, A.E. Stillman, and R.D. White. Coregistered MR imaging myocardial viability maps and multi-detector row CT coronary angiography displays for surgical revascularization planning: Initial experience. *Radiology*, 237(2):465, 2005.
- [219] R. Shahzad, T. van Walsum, H.A. Kirisli, H. Tang, C.T. Metz, M. Schaap, L. J. van Vliet, and W.J. Niessen. Automatic detection, quantification and lumen segmentation of the coronary arteries using a two point centerline extraction scheme. In *3D Cardiovascular Imaging: A MICCAI segmentation challenge: MICCAI 2012 workshop proceedings*, 2012.
- [220] R. Shahzad, T. van Walsum, M. Schaap, A. Rossi, S. Klein, A.C. Weustink, P. de Feyter, L. J. van Vliet, and W.J. Niessen. Vessel specific coronary artery calcium scoring: An automatic system. *Academic Radiology*, 20(1):1–9, 2013.
- [221] P.J. Slomka, V.Y. Cheng, D. Dey, J. Woo, A. Ramesh, S. Van Krieking, Y. Suzuki, Y. Elad, R. Karlsberg, D.S. Berman, et al. Quantitative analysis of myocardial perfusion spect anatomically guided by coregistered 64-slice coronary ct angiography. *Journal of Nuclear Medicine*, 50(10):1621–1630, 2009.
- [222] D.K. Sodickson and W.J. Manning. Simultaneous acquisition of spatial harmonics (SMASH): fast imaging with radiofrequency coil arrays. *Magnetic Resonance in Medicine*, 38(4):591–603, 1997.
- [223] P. Soille. *Morphological image analysis: Principles and applications*. 1999.
- [224] S. Sourbron. Technical aspects of MR perfusion. *European Journal of Radiology*, 76(3): 304–313, 2010.

- [225] L. J. Spreeuwens, M. Breeuwer, W. J. Niessen, and M.A. Viergever. Automatic detection of myocardial boundaries in MR cardio perfusion images. *Lecture notes in computer science*, 2208:1228–1231, 2001.
- [226] M B Stegmann, H Ólafsdóttir, and H B W Larsson. Unsupervised motion-compensation of multi-slice cardiac perfusion MRI. *Medical Image Analysis*, 9(4):394–410, 2005.
- [227] M.B. Stegmann and R. Larsen. Multi-band modelling of appearance. *Image and Vision Computing*, 21(1):61–68, 2003.
- [228] M.B. Stegmann and D. Pedersen. Bi-temporal 3D active appearance models with applications to unsupervised ejection fraction estimation. *Proceedings of SPIE Medical Image Processing*, 5747:336–350, 2005.
- [229] P. Stolzmann, H. Alkadhi, H. Scheffel, A. Hennemuth, C. Kuehnel, S. Baumueller, S. Kozerke, V. Falk, B. Marincek, and O.F. Donati. Image fusion of coronary CT angiography and cardiac perfusion MRI: a pilot study. *European radiology*, 20(5):1174–1179, 2010.
- [230] P. Storey, Q. Chen, W. Li, R.R. Edelman, and P.V. Prasad. Band artifacts due to bulk motion. *Magnetic Resonance in Medicine*, 48(6):1028–1036, 2002.
- [231] Y. Sun, M.P. Jolly, and J.M.F. Moura. Contrast-invariant registration of cardiac and renal MR perfusion images. *Lecture Notes in Computer Science*, pages 903–910, 2004.
- [232] L. Tautz, A. Hennemuth, M. Andersson, A. Seeger, H. Knutsson, and O. Friman. Phase-based non-rigid registration of myocardial perfusion MR image sequences. In *Proceedings of International Symposium on Biomedical Imaging (ISBI)*, pages 516–519, 2010.
- [233] M. Termeer, J.O. Bescós, M. Breeuwer, A. Vilanova, F. Gerritsen, and M.E. Gröller. CoViCAD: comprehensive visualization of coronary artery disease. *IEEE Transactions on Visualization and Computer Graphics*, 13(6):1632–1639, 2007.
- [234] M. Termeer, J. Oliván Bescós, M. Breeuwer, A. Vilanova, F. Gerritsen, M. E. Gröller, and E. Nagel. Visualization of myocardial perfusion derived from coronary anatomy. *IEEE Transactions on Visualization and Computer Graphics*, (6):1595–602, 2008.
- [235] J. Theiler and G. Gisler. A contiguity-enhanced k-means clustering algorithm for unsupervised multispectral image segmentation. In *Proceedings of SPIE*, volume 3159, pages 108–118, 1997.
- [236] P. Thevenaz, U.E. Ruttimann, and M. Unser. A pyramid approach to subpixel registration based on intensity. *IEEE Transactions on Image Processing*, 7(1):27–41, 2002.
- [237] J. P. Thirion. Image matching as a diffusion process: an analogy with Maxwell’s demons. *Medical Image Analysis*, 2(3):243–260, 1998.
- [238] H.S. Thomsen. Gadolinium-based contrast media may be nephrotoxic even at approved doses. *European Radiology*, 14(9):1654–1656, 2004.
- [239] K. Thygesen, J.S. Alpert, H.D. White, A.S. Jaffe, F.S. Apple, M. Galvani, H.A. Katus, L.K. Newby, J. Ravkilde, B. Chaitman, et al. Universal definition of myocardial infarction. *European heart journal*, 28(20):2525–2538, 2007.

- [240] C.Y. Tong, F.S. Prato, G. Wisenberg, T.Y. Lee, E. Carroll, D. Sandler, and J. Wills. Techniques for the measurement of the local myocardial extraction efficiency for inert diffusible contrast agents such as gadopentate dimeglumine. *Magnetic Resonance in Medicine*, 30(3):332–336, 1993.
- [241] C.Y. Tong, F.S. Prato, G. Wisenberg, T.Y. Lee, E. Carroll, D. Sandler, J. Wills, and D. Drost. Measurement of the extraction efficiency and distribution volume for Gd-DTPA in normal and diseased canine myocardium. *Magnetic Resonance in Medicine*, 30(3):337–346, 1993.
- [242] N.V. Tsekos, Y. Zhang, H. Merkle, N. Wilke, M. Jerosch-Herold, A. Stillman, and K. Ugurbil. Fast anatomical imaging of the heart and assessment of myocardial perfusion with arrhythmia insensitive magnetization preparation. *Magnetic Resonance in Medicine*, 34(4):530–536, 1995.
- [243] B.M.W. Tsui, E.C. Frey, K.J. LaCroix, D.S. Lalush, W.H. McCartney, M.A. King, and G.T. Gullberg. Quantitative myocardial perfusion spect. *Journal of Nuclear Cardiology*, 5(5):507–522, 1998.
- [244] I. Vajda. *Theory of Statistical Evidence and Information*. 1989.
- [245] J.P.M. Vallée, H.D. Sostman, J.R. Macfall, T.R. Degrado, J. Zhang, L. Sebbag, F.R. Cobb, T. Wheeler, L.W. Hedlund, T.G. Turkington, et al. Quantification of myocardial perhsion by MRI after coronary occlusion. *Magnetic Resonance in Medicine*, 40(2):287–297, 1998.
- [246] R.J. van der Geest and J.H.C. Reiber. Quantification in cardiac MRI. *Journal of Magnetic resonance imaging*, 10(5):602–608, 1999.
- [247] J.M. van Werkhoven, J.D Schuijf, O. Gaemperli, J.W. Jukema, E. Boersma, W. Wijns, P. Stolzmann, H. Alkadhi, I. Valenta, M.P. Stokkel, L.J. Kroft, A. de Roos, G. Pundziute, A. Scholte, E.E. van der Wall, P.A. Kaufmann, and J.J. Bax. Computed tomography and gated single-photon emission computed tomography in patients with suspected coronary artery disease. *J Am Coll Cardiol*, 53:623–632, 2009.
- [248] S. Veerer, M.J. Dunn, and G.Z. Yang. Multiresolution image registration for two-dimensional gel electrophoresis. *Proteomics*, 1(7):856–870, 2001.
- [249] M. Vöhringer, H. Mahrholdt, A. Yilmaz, and U. Sechtem. Significance of late gadolinium enhancement in cardiovascular magnetic resonance imaging (CMR). *Herz*, 32(2):129–137, 2007.
- [250] A. Wagner, H. Mahrholdt, T.A. Holly, M.D. Elliott, M. Regenfus, M. Parker, F.J. Klocke, R.O. Bonow, R.J. Kim, and R.M. Judd. Contrast-enhanced MRI and routine single photon emission computed tomography (SPECT) perfusion imaging for detection of subendocardial myocardial infarcts: an imaging study. *The Lancet*, 361(9355):374–379, 2003.
- [251] H. Wang, L. Dong, J. O'Daniel, R. Mohan, A.S. Garden, K. Kian Ang, D.A. Kuban, M. Bonnen, J.Y. Chang, and R. Cheung. Validation of an accelerated 'demons' algorithm for deformable image registration in radiation therapy. *Physics in Medicine and Biology*, 50:2887–2905, 2005.
- [252] Y. Wang, S.J. Riederer, and R.L. Ehman. Respiratory motion of the heart: kinematics and the implications for the spatial resolution in coronary imaging. *Magnetic Resonance in Medicine*, 33(5):713–719, 1995.

- [253] H.J. Weinmann, R.C. Brasch, W.R. Press, and G.E. Wesbey. Characteristics of gadolinium-DTPA complex: a potential NMR contrast agent. *American Journal of Roentgenology*, 142(3): 619, 1984.
- [254] A.M. Weissler, W.S. Harris, and C.D. Schoenfeld. Systolic time intervals in heart failure in man. *Circulation*, 37(2):149–159, 1968.
- [255] G.E. Wesbey, C.B. Higgins, M.T. McNamara, B.L. Engelstad, M.J. Lipton, R. Sievers, R.L. Ehman, J. Lovin, and R.C. Brasch. Effect of gadolinium-DTPA on the magnetic relaxation times of normal and infarcted myocardium. *Radiology*, 153(1):165, 1984.
- [256] A. Weustink and P. de Feyter. The role of multi-slice computed tomography in stable angina management - a current perspective. *Neth Heart Journal*, in press:–, 2011.
- [257] A C Weustink and P J De Feyter. The role of multi-slice computed tomography in stable angina management - a current perspective. *Netherlands Heart Journal*, 19:336–343, 2011.
- [258] N. Wilke and M. Jerosch-Herold. Assessing myocardial perfusion in coronary artery disease with magnetic resonance first-pass imaging. *Cardiology Clinics*, 16(2):227–246, 1998.
- [259] R.F. Wilson, K. Wyche, BV Christensen, S. Zimmer, and D. D. Laxson. Effects of adenosine on human coronary arterial circulation. *Circulation*, 82(5):1595–1606, 1990.
- [260] H. Wold. Soft modeling by latent variables: the nonlinear iterative partial least squares approach. *Perspectives in Probability and Statistics, papers in honour of MS Bartlett*, pages 520–540, 1975.
- [261] S.D. Wolff and R.S. Balaban. Magnetization transfer contrast (MTC) and tissue water proton relaxation in vivo. *Magnetic Resonance in Medicine*, 10(1):135–144, 1989.
- [262] S.D. Wolff, J. Eng, and R.S. Balaban. Magnetization transfer contrast: method for improving contrast in gradient-recalled-echo images. *Radiology*, 179(1):133, 1991.
- [263] G. Wollny, M. J. Ledesma-Carbayo, P. Kellman, and A. Santos. Exploiting quasiperiodicity in motion correction of free-breathing myocardial perfusion MRI. *IEEE Transactions on Medical Imaging*, 29(8):1516–27, 2010.
- [264] G. Wollny, P. Kellman, A. Santos, and M.J. Ledesma-Carbayo. Automatic motion compensation of free breathing acquired myocardial perfusion data by using independent component analysis. *Medical Image Analysis*, 16, 2012.
- [265] K Wong, E Wu, M Ng, Y Wu, H Tse, C Lau, G Lo, and E Yang. Image registration in myocardial perfusion MRI. In *Proceedings of IEEE - Engineering in Medicine and Biology Society*, volume 1, pages 453–4, 2005.
- [266] C. Xu and J.L. Prince. Snakes, shapes, and gradient vector flow. *IEEE Transactions on Image Processing*, 7(3):359–369, 2002.
- [267] H. Xue, J. Guehring, L. Srinivasan, S. Zuehlsdorff, K. Saddi, C. Chefdh tel, J. V. Hajnal, and D. Rueckert. Evaluation of rigid and non-rigid motion compensation of cardiac perfusion MRI. In *Proceedings of the International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, volume 11, pages 35–43, 2008.

- [268] H. Xue, S. Zuehlsdorff, P. Kellman, A. Arai, S. Nielles-Vallespin, C. Chefdhóotel, C. H. Lorenz, and J. Guehring. Unsupervised inline analysis of cardiac perfusion MRI. In *Proceedings of the International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, volume 12, pages 741–9, 2009.
- [269] G Z Yang, P Burger, J Panting, P D Gatehouse, D Rueckert, D J Pennell, and D N Firmin. Motion and deformation tracking for short-axis echo-planar myocardial perfusion imaging. *Medical Image Analysis*, 2(3):285–302, 1998.
- [270] R. K. Yin. *Case Study Research: Design and Methods*. SAGE Publications, 4th edition, 2009.
- [271] P. Zanzonico, L.N. Rothenberg, and H.W. Strauss. Radiation exposure of computed tomography and direct intracoronary angiography: risk has its reward. *Journal of the American College of Cardiology*, 47(9):1846, 2006.
- [272] A. Zavaljevski, S.K. Holland, A.P. Dhawan, R.S. Dunn, and W.S. Ball. Multilevel computed hemodynamic parameter maps from dynamic perfusion MRI. *IEEE Transactions on Instrumentation and Measurement*, 48(3):711–720, 1999.
- [273] K. Zierler. Indicator dilution methods for measuring blood flow, volume, and other properties of biological systems: a brief history and memoir. *Annals of Biomedical Engineering*, 28(8): 836–848, 2000.
- [274] K.L. Zierler. Theoretical basis of indicator-dilution methods for measuring flow and volume. *Circulation Research*, 10(3):393, 1962.
- [275] K.L. Zierler. Equations for measuring blood flow by external monitoring of radioisotopes. *Circulation Research*, 16(4):309–321, 1965.
- [276] B. Zitova and J. Flusser. Image registration methods: a survey. *Image and vision computing*, 21(11):977–1000, 2003.