On the pathology of preeclampsia : genetic variants, complement dysregulation and angiogenesis

Buurma, A.A.J.

BIBLIOGRAPHY

Buurma A*, Penning ME*, Prins F, Schutte JM, Bruijn JA, Wilhelmus S, Rajakumar A, Bloemenkamp KW, Karumachi SA, Baelde HJ. Preeclampsia is associated with the presence of transcriptionally active placental fragments in the maternal lung. *Hypertension*, in press.

Buurma A, Turner RJ, Driessen JH, Mooyaart AL, Schoones JW, Bruijn JA, Bloemenkamp KW, Dekkers OM, Baelde HJ. Genetic variants in pre-eclampsia: a meta-analysis. *Hum Reprod Update* 2013; 19(3):289-303

Buurma A, Cohen D, Veraar K, Schonkeren D, Claas FH, Bruijn JA, Bloemenkamp KW, Baelde HJ. Preeclampsia is characterized by placental complement dysregulation. *Hypertension* 2012; 60(5):1332-1337.

Cohen D, **Buurma A**, Goemaere NN, Girardi G, le Cessie S, Scherjon S, Bloemenkamp KW, de Heer E, Bruijn JA, Bajema IM. Classical complement activation as a footprint for murine and human antiphospholipid antibody-induced fetal loss. *J Pathol* 2011; 225(4):502-511.

Buurma A, Cohen D. Ovariectomie of ovaria in situ laten bij hysterectomie? *Ned Tijdschr Geneeskd* 2011; 155:A3673

*Both authors contributed equally.