

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/29935> holds various files of this Leiden University dissertation

Author: Jacobs, Chris G.C.

Title: Surviving embryogenesis : the extraembryonic serosa protects the insect egg against desiccation and infection

Issue Date: 2014-12-04

Chapter 5

The extraembryonic serosa is a frontier epithelium providing the insect egg with a full-range innate immune response

Chris G.C. Jacobs, Herman P. Spaink and Maurijn van der Zee

eLife - accepted for publication

Drosophila larvae and adults possess a potent innate immune response, but the response of *Drosophila* eggs is poor. In contrast to *Drosophila*, eggs of the beetle *Tribolium* are protected by a serosa, an extraembryonic epithelium that is present in all insects except higher flies. Here, we test a possible immune function of this frontier epithelium using *Tc-zen1* RNAi-mediated deletion. First, we show that bacteria propagate twice as fast in serosa-less eggs. Then, we compare the complete transcriptomes of wild-type, control RNAi and *Tc-zen1* RNAi eggs before and after sterile or septic injury. Infection induces genes involved in Toll and IMD-signaling, melanisation, production of reactive oxygen species and antimicrobial peptides in wild-type eggs, but not in serosa-less eggs. Finally, we demonstrate constitutive and induced immune gene expression in the serosal epithelium using *in situ* hybridization. We conclude that the serosa provides insect eggs with a full-range innate immune response.

Keywords: RNA sequencing; *Tribolium castaneum*; innate immunity; serosa; NF-kappaB; Toll

Introduction

To combat infection, insects rely on humoral and local immune responses. The humoral immune response is characterized by the massive secretion of antimicrobial peptides into the hemolymph and is mainly exerted by the fat body. Epithelia and hemocytes play the main role in local immune defenses that comprise melanisation, local AMP production, phagocytosis and encapsulation (Davis & Engstrom, 2012; Ferrandon, 2013; Ganesan et al., 2011; Lemaitre & Hoffmann, 2007; Ligoxygakis, 2013; Wang et al., 2014). The mechanisms regulating these innate immune responses have largely been uncovered with the aid of genetic and molecular studies in the fruit fly *Drosophila melanogaster*. When microbes invade the fly, their released peptidoglycans are sensed by peptidoglycan recognition proteins (PGRPs) and Gram-negative binding proteins (GNBPs) leading to the activation of the main immune signaling pathways. The mesodiaminopimelic acid-type (DAP-type) peptidoglycans of Gram-negative bacteria activate the IMD pathway, whereas the Lys-type peptidoglycans of Gram-positive bacteria activate the Toll pathway. The activation of the Toll-pathway is mediated by a proteolytic cascade of serine proteases leading to the cleavage of the cytokine Spätzle, the ligand of the transmembrane receptor Toll. Activation



of the immune signaling pathways leads to nuclear localization of the NF-kappaB factors Dorsal, Dif or Relish that induce antimicrobial peptides (AMPs). Other upregulated genes are prophenoloxidases (proPOs which mediate melanisation) and dual oxidase (DUOX which produces reactive oxygen species).

Drosophila has been extremely helpful uncovering those mechanisms, but research in other insects, such as the mealworm beetle *Tenebrio molitor* has also generated insightful results (see Park et al., 2010). The biochemical details of pathway activation, for instance, have mainly been unraveled using this beetle (see Park et al., 2010 for review). With the availability of tools such as RNAseq and RNAi, more insect species are being established as model organism for innate immunity research (Altincicek & Vilcinskas, 2007; Gerardo et al., 2010; Johnston et al., 2013; Johnston & Rolff, 2013; Waterhouse et al., 2007; Zhu et al., 2013). In particular the red flour beetle (*Tribolium castaneum*), has received much attention in innate immune studies (Altincicek et al., 2013; Altincicek et al., 2008; Behrens et al., 2014; Contreras et al., 2013; Milutinović et al., 2013; Roth et al., 2010; Zhong et al., 2013; Zou et al., 2007). Comparative genome analysis has revealed that components of intracellular immune signaling pathways (Toll, IMD and JAK/STAT) in *Drosophila* are 1:1 conserved in *Tribolium* (Zou et al., 2007). The RNAi knockdown technology has shown that the IMD and Toll pathway are largely functionally conserved (Shrestha & Kim, 2010; Yokoi et al., 2012a; Yokoi et al., 2012b). Their activity does, however, not strictly depend on either Gram-negative or Gram-positive bacteria (Yokoi et al., 2012a; Yokoi et al., 2012b), but this distinction is also not completely black and white in *Drosophila* (Leone et al., 2008; Leulier et al., 2003). Nevertheless, species-specific family expansion and sequence divergence in the PGRP and AMP families indicate species-specific differences, possibly required for effective recognition and elimination of evolving pathogens (Altincicek et al., 2008; Christophides et al., 2002; Park et al., 2010; Zou et al., 2007).

Not only larvae and adults, but also insect eggs are constantly threatened by pathogens (see Blum & Hilker, 2008; Kellner, 2008 for review). *Serratia* bacteria, for instance, have been found inside eggs of corn earworms and corn borers (Bell, 1969; Lynch et al., 1976) and can infect eggs in the laboratory (Sikowski et al., 2001). We have also shown that *Serratia* infection leads to reduced egg survival in the burying beetle *Nicrophorus vespilloides* (Jacobs et al., 2014). Maternal investments have been proposed to counter microbial infections. Female medflies, for example, cover their eggs with antimicrobial secretions (Marchini et al., 1997) and in the absence of maternal care, eggs of earwigs die of fungal infection (Boos et al., 2014). Two studies focusing on transgenerational immune priming, however, have shown that the antimicrobial activity of eggs is of internal origin (Sadd & Schmid-Hempel, 2007; Zanchi et al., 2012). This is often implicitly interpreted as maternal loading of antimicrobials into the egg (Moreau et al., 2012), but maternal transfer of bacteria to the eggs also leaves zygotic investment as possibility (Freitag et al., 2014; Trauer & Hilker, 2013). Overall, it is ecologically relevant to gain a better understanding of the immune system in insect eggs.

The zygotic response in *Drosophila* eggs, however, seems poor. It is not until late stage 15, (one of the latest stages in development when ectoderm and trachea have differentiated) that eggs show up to 25-fold upregulation of antimicrobial peptides (Tan et al., 2014). This is incomparable to the upregulation in adult flies that is at least an order of magnitude larger. Except for Cecropin (Tingvall et al., 2001), stage 11 embryos do not show



any induction of antimicrobial peptides, and cannot contain an infection of non-pathogenic bacteria, leading to reduced survival (Tan et al., 2014). In strong contrast, we have shown that eggs of *Tribolium* which were not even half way during development could upregulate several AMPs to levels comparable to the adult (Jacobs & van der Zee, 2013). This upregulation depends on the serosa, an extraembryonic epithelium that envelopes yolk and embryo (Jacobs & van der Zee, 2013). This membrane is present in all insects, but was lost in a small group of higher Diptera (the Schizophora) to which *Drosophila* belongs (Rafiqi et al., 2008; Schmidt-Ott, 2000). Although two maternal extracellular coverings, the chorion and the vitelline membrane envelop the insect egg, the serosa is the first cellular epithelium surrounding the egg at the interface between the microbe rich external milieu on the one side and the yolk and embryo at the other side. Thus, the serosa could function as an immune competent barrier epithelium. This has been suggested before, as the NF-kappaB factor Dorsal is highly expressed in the presumptive serosa (Chen et al., 2000). The absence of the serosa might account for the poor immune response in *Drosophila* eggs.

To gain deeper insights into the role of the serosa, we chose *Tribolium castaneum*, a beetle that possesses a serosa, like all non-Schizophoran insects. In this beetle, we can prevent development of the serosa by parental RNA interference with *Tc-zerknüllt1* (*Tc-zen1*). This technique generates *Tribolium* eggs with an amnion at the dorsal side, but without a serosa (van der Zee et al., 2005). At the relative humidity of the air of the laboratory, normal larvae hatch from these eggs (Jacobs et al., 2013). As *Tc-zen1* is only expressed in the early serosa (van der Zee et al., 2005) and is not expressed anymore by the time the experiments are performed (see discussion), we expect only to find effects that are a consequence of the absence of the serosa. We investigated growth of bacteria in serosa-less and wild-type eggs, sequenced the whole transcriptome of naive and immune-challenged eggs with and without a serosal epithelium and confirmed constitutive and induced gene expression in the serosa by *in situ* hybridization. We conclude that the serosa is a frontier epithelium that provides immune competence to the insect egg.

Results

Bacteria propagate twice as fast in serosa-less eggs

To examine the influence of the serosa on bacterial growth in infected eggs, and to standardize our infection method, we counted colony forming units (cfu's) directly after infection (t=0) and 6 hours later (t=6) (Figure 5-1). We pricked 24-40h old eggs (i.e. up to half-way during development) with a tungsten needle dipped in a concentrated mix of *Escherichia coli* and *Micrococcus luteus* cultures (see Materials and Methods). To determine cfu's, we shortly treated eggs with 0.5% hypochlorite to sterilize the outside. Untreated eggs did hardly contain bacteria that grow on LB agar plates (on average 3 cfu's were found). Sterile injury did not increase this number (Figure 5-1, lower lines). In contrast, septic injury introduced on average 53 bacteria into wild-type eggs and 49 into serosa-less eggs. These numbers increased on average to 747 cfu's in wild-type eggs and to 7260 cfu's in serosa-less eggs. When we use the formula $N_{(t)} = N_{(0)} * e^{kt}$, the specific bacterial growth rate k in wild-type eggs is 0.44 h⁻¹, whereas $k = 0.83$ h⁻¹ in serosa-less eggs. This means that bacteria grow twice as fast in serosa-less eggs and suggests that the serosa exerts an immune function.






Figure 5-1: Counts of colony forming units (cfu's) after sterile and septic injury. Green lines represent bacterial growth in wild-type eggs. Red lines represent bacterial growth in *Tc-zen1* RNAi (serosa-less) eggs. Sterile injury did not introduce bacteria (lower lines: average of 2 cfu's found at t=0, and an average of 5 cfu's found at t=6). Septic injury introduced on average 53 bacteria into wild-type eggs and 49 into serosa-less eggs. These numbers increased to 747 +/- 106 cfu's in wild-type eggs (green upper line) and to 7260 +/- 1698 cfu's in serosa-less eggs (red upper line) at t=6. This means that bacteria propagate twice as fast in serosa-less eggs ($p < 0.01$, as determined by a Pearson's chi-square test). Suspensions of 10 eggs were used per LB agar plate (see Materials and Methods), and 10 plates were analyzed per treatment and time point, giving rise to the error bars presented in the graph (standard error).


RNAseq reveals a full-range immune response in *Tribolium* eggs

To characterize this immune function, we sequenced the whole transcriptome of wild-type eggs, *Tc-zen1* RNAi (serosa-less) eggs and control RNAi eggs without injury, after sterile injury and after septic injury (Figure 5-2). The control RNAi consists of a injection of a 500bp dsRNA derived from a vector sequence without target in the *Tribolium castaneum* genome. For these 9 different treatments, three biological replicates were carried out (independent RNAi, independent injury) giving a total of 27 samples (Figure 5-2). Illumina next generation sequencing resulted in over 970 million cDNA reads with over 49 billion bp sequence information. Approximately 72% of the reads could be mapped to *Tribolium* gene models built on the 3.0 genome assembly (Richards et al., 2008) (Supplementary Table 5-1). We found expression of 14,903 of the total of 16,541 predicted genes, of which 13,464 genes were expressed in wild-type, control and *Tc-zen1* eggs and 1440 genes were expressed in a subset of these treatments.

Table 5-1: Number of differentially expressed immune genes in *Tribolium castaneum* eggs (Green = induction ; Red = repression)

	Wild-type Sterile injury		Wild-type Septic injury		Control Sterile injury		Control Septic injury		<i>Tc-zen1</i> Sterile injury		<i>Tc-zen1</i> Septic injury	
Microbial recognition	4	1	7	2	6	0	8	3	1	0	0	0
Extracellular signal transduction and modulation	27	6	32	10	33	4	34	10	4	5	2	0
Intracellular transduction pathways (Toll/IMD/JNK/JAK-STAT)	2	1	3	2	2	2	6	3	3	2	2	1
Execution / Stress	12	0	20	2	16	4	24	7	5	2	3	1
Total	45	8	62	16	57	10	72	23	13	9	7	2
	up	down	up	down	up	down	up	down	up	down	up	down





b

	Naive eggs	Sterile Injury	Septic Injury
Wild-type	3x	3x	3x
Control RNAi	3x	3x	3x
<i>Tc-zen1</i> RNAi	3x	3x	3x

Figure 5-2: Experimental setup a) We collected eggs from wild-type, control RNAi and *Tc-zen1* RNAi beetles overnight. These eggs were incubated for 24h at 30°C to ensure development of the serosa. Eggs are then maximally 40h old, while total developmental time is close to 85h at 30°C. Eggs were pricked with a sterile needle (sterile injury), pricked with a mix of *E. coli* and *M. luteus* (septic injury), or remained untreated (naive). They were incubated for another 6 hours at 30°C before total RNA was extracted for RNAseq. To analyze the immune response, the transcriptomes of sterilely injured eggs and of septicly injured eggs were compared to naive eggs. This was done for wild-type, control and *Tc-zen1* RNAi eggs. b) We collected 3 biological samples for each combination of eggtype (wild-type, Control RNAi or *Tc-zen1* RNAi) and treatment (naive, sterile injury or septic injury) giving a total of 27 biological samples.

First, we identified the immune-responsive genes by determining differential expression of genes between naive eggs on the one hand and sterilely injured eggs or septicly injured eggs on the other hand. We only considered genes with at least a 2-fold change in expression and an adjusted p-value smaller than 0,01. This gave a total of 415 differentially expressed genes in sterilely injured eggs compared to the naive eggs, and a total of 538 differentially regulated genes in septicly injured eggs compared to naive eggs. This shows that *Tribolium* eggs possess an extensive transcriptional response upon infection.

To obtain a global impression of the kind of genes differentially regulated upon infection in wild-type and control RNAi eggs, we assigned gene ontology terms (GO-terms) to all *Tribolium* genes. As no GO-term annotation is available for *Tribolium*, we blasted *Tribolium* genes against *Drosophila* and used the *Drosophila* GO-terms of the best hit. Using the Wallenius approximation (Young et al., 2010), we found several highly overrepresented GO-term categories with a p-value below 0.001 in both wild-type eggs (Figure 5-3a) and control RNAi eggs (Figure 5-3b). The overrepresented categories are mostly immune related. This indicates that our approach does not depend on artefacts generated by pricking eggs (e.g. delayed development), but mainly identifies genes involved in the innate immune response.

To obtain a more detailed analysis of the immune response in wild-type and control eggs, we focused on 368 genes that have been annotated as immune genes (Altincicek et al., 2013; Zou et al., 2007) (Supplementary files 4-9, available online). Of these genes, 78 were differentially regulated in wild-type eggs upon septic injury (Table 5-1 and Supplementary Table 5-3 and file 5), while 95 immune genes were differentially regulated in control RNAi eggs (Table 5-1 and Supplementary Table 5-3 and file 7). This



indicates that RNAi itself leads to an increased number of differentially regulated genes upon bacterial challenge, but, more importantly, shows that *Tribolium* eggs possess an elaborate immune response. In the following sections, we take a closer look at the exact genes involved in this extensive immune response.

Recognition of microbes and extracellular signal transduction

Of the 7 predicted peptidoglycan recognition proteins (PGRPs) in *Tribolium* we found significant induction of *PGRP-LA*, *LC*, *SA* and *SB* (Supplementary Table 5-2). Of these PGRPs, *PGRP-SA* and *SB* were induced over 200-fold (Figure 4a, Supplementary Table 5-2). Thus, it could be that these PGRPs rather function as effectors digesting Gram-positive bacteria, as shown for human *PGRP-S* (Dziarski et al., 2003). At least *PGRP-SB* shows all the amino acid residues characteristic for catalytic PGRPs (Kim et al., 2003). No induction was found for *PGRP-LE* and *LD*. These findings strongly resemble the response of *Tribolium* adults, in which the same PGRPs responded to infection (Altincicek et al., 2013). Of the Gram-negative binding proteins (GNBP), we found induction of *GNBP2* and *GNBP3* (Supplementary Table 5-2). In *Tribolium* adults and in *Drosophila*, however, only *GNBP3* is immune-inducible (Altincicek et al., 2013; Lemaitre & Hoffmann, 2007), whereas *GNBP1* and *GNBP3* are immune-inducible in *Tenebrio* (Johnston et al., 2013).




Figure 5-3: Types of genes that are differentially regulated a) Significantly over-represented GO-terms in the genes induced in wild-type eggs after septic injury ($p < 0.001$). b) Significantly over-represented GO-terms in the genes induced in control RNAi eggs after septic injury ($p < 0.001$). These categories indicate that the detected differential regulation does not result from artefacts induced by treatments (such as death or delayed development) and show that *Tribolium* eggs display an elaborate immune response.

Thioester-containing proteins (TEPs) have also been suggested to function as pattern recognition proteins, possibly targeting microbes for phagocytosis (Stroschein-Stevenson et al., 2005; Wang & Wang, 2013). We did not find induction of thioester-containing proteins (TEPs) but rather repression, for instance of *TEP-D* (Supplementary Table 5-2). This is surprising, since TEPs are upregulated in *Tribolium* larvae and adults (Altincicek et al., 2013; Behrens et al., 2014); and *Drosophila* (Stroschein-Stevenson et al., 2005; Wang & Wang, 2013). Similar to *Drosophila*, however, we did find induction of a putative TEP/complement-binding receptor-like protein (*LpR2*). We also found induction of *C-type lectin 6* and repression of *C-type lectin 1* and *13*. These lectins are thought to be involved in microbial recognition but no induction or repression has been found in *Drosophila* or *Tribolium* adults (Altincicek et al., 2013; De Gregorio et al., 2001).



The serine proteases and serpins have significantly expanded in number in *Tribolium* (Zou et al., 2007), similar to *Anopheles* (Christophides et al., 2002). Interestingly, most of them seem to be functional in the immune response as we found induction of 36 serine proteases and serpins, and repression of another 10 upon infection (Supplementary Table 5-2). This number is much higher than previously reported for adults (Altincicek et al., 2013). Of the Spaetzle ligands, we found induction of *spz1* and *spz2* and repression of *spz4* and 5 (Supplementary Table 5-2). In larvae and adults, however, different Spaetzles were induced or repressed, indicating specific use at different stages of the life cycle (Altincicek et al., 2013; Behrens et al., 2014).

In total, 51 of the 78 immune genes that are differentially regulated in wild-type eggs are involved in bacterial recognition and extracellular signal transduction, showing the prominence of these extracellular processes in the modulation of the immune response of the *Tribolium* egg.

Transmembrane and intracellular signal transduction

We found induction of several intracellular signaling components of the Toll, IMD and JNK pathways (Figure 5-4a, Supplementary Table 5-2) upon immune-challenge of *Tribolium* eggs. This suggests that these pathways are largely functionally conserved between *Drosophila* and *Tribolium*, although we could hardly detect expression of *dredd*, the endoprotease that cleaves Relish for nuclear translocation. Similar to larvae and adults (Altincicek et al., 2013; Behrens et al., 2014), JAK-STAT pathway components were not differentially regulated. Interestingly, we found significant upregulation of the *toll3* receptor upon infection. This was also found in larvae and adults (Altincicek et al., 2013; Behrens et al., 2014) and suggests that it is not *toll1*, but *toll3* that plays a major role in the innate immune response of *Tribolium*.

Execution mechanisms

As expected, we found the highest induction amongst the antimicrobial peptides. We detected generally more than 500-fold upregulation of defensins, attacins, coleopterins, cecropins and thaumatin (Figure 5-4a, Supplementary Table 5-2). This means that *Tribolium* eggs can induce AMPs to comparable levels as larvae and adults (Altincicek et al., 2013; Behrens et al., 2014). We also found upregulation of *prophenoloxidase1* (*proPO1*), a gene involved in melanisation, and of *heme peroxidase 11*, a dual oxidase (*DUOX*) ortholog involved in the production of reactive oxygen species (Supplementary Table 5-2). This shows that *Tribolium* eggs are indeed able to respond with the full complement of immune defense mechanisms.

Currently, 19 AMPs are recognized in *Tribolium*, based on homology with known AMPs. However, due to the presence of species-specific AMPs and extreme sequence diversity of these molecules, homology based searches have probably missed several AMPs (Yang et al., 2011; Zou et al., 2007). AMPs are generally small (less than 30 kDa), cationic, hydrophobic and possibly have high glycine and/or proline content (Bulet & Stöcklin, 2005; Philippe Bulet et al., 2004). Based on the antimicrobial peptide database (Wang et al., 2009), many proteins encoded in the *Tribolium* genome fulfil those criteria and are identified as candidate antimicrobial peptides. Using our RNA sequencing data, however, we could select those candidate proteins that exhibit at least a two-fold induction upon infection. Based on these criteria, we found 20 potential new AMPs






Figure 5-4: Immune-responsive genes in wild-type, control and *Tc-zen1* RNAi eggs. a) Schematic representation of the immune signaling pathways in *Tribolium* as described in (Zou et al., 2007). Significantly induced genes after septic injury in wild-type or control RNAi eggs are indicated in green; significantly repressed genes after septic injury in wild-type or control RNAi eggs are indicated in red. Genes not differentially expressed are black. The size of the gene names represents the fold change (small = 1.5-10 fold, medium = 10-500 fold, large = 500+ fold expression). b) Venn-diagram showing the number of differentially expressed genes in septicallly injured eggs as compared to naive eggs (FDR<0.01). In total, 538 genes are differentially expressed upon infection, of which 394 in wild-type eggs, 435 in control RNAi eggs and only 57 in *Tc-zen1* RNAi eggs. This means that *Tribolium* eggs display an extensive transcriptional response upon infection and that this response is largely abolished in eggs without a serosa.



Table 5-2: Antimicrobial properties of known and potential new antimicrobial peptides in *Tribolium castaneum*. In the table are known antimicrobial peptides and those proteins that show at least a two-fold induction upon infection, are smaller than 200 amino acids and are not negatively charged. TC009336 was included because of the high glycine content.

Gene ID	Molecular weight (kDa)	Peptide length (AA)	Hydrophobic ratio	Net charge	Glycine content	Proline content	Foldchange Wild-type	Foldchange Control
Cecropin1 / TC000499	3.67	31	58%	+5	6%	0%	Inf	Inf
Cecropin3 / TC000500	9.80	90	43%	+2	6%	13%	Inf	49x
attacin2 / TC007738	15.80	145	37%	+7	12%	4%	3098x	2190x
Coleoptericin1 / TC005093	15.99	141	30%	-1	9%	7%	2392x	18067x
Defensin2 / TC010517	8.73	79	50%	+6	6%	1%	1183x	Inf
Defensin3 / TC012469	9.42	83	50%	+7	3%	1%	908x	Inf
attacin1 / TC007737	17.49	165	28%	+9	18%	3%	869x	3696x
TC007858	20.14	182	35%	0	11%	3%	484x	54x
Defensin1 / TC006250	14.91	132	46%	+11	4%	3%	187x	1551x
TC011036	12.89	109	39%	+13	2%	6%	138x	11x
Coleoptericin2 / TC005096	15.96	141	30%	-1	9%	7%	91x	227x
TC015479	13.00	120	42%	+6	5%	1%	80x	26x
TC007763	16.87	158	37%	+4	6%	17%	47x	67x
TC004646	15.04	135	34%	+2	7%	7%	40x	29x
TC008806	15.83	142	33%	+2	10%	2%	31x	37x
TC009336	13.50	137	30%	-4	39%	2%	15x	7x
TC014565	20.73	176	38%	+17	2%	2%	14x	9x
TC001030	14.62	137	29%	+9	10%	12%	9x	16x
TC001784	13.54	150	27%	+7	43%	2%	8x	6x
TC005478	13.70	122	45%	+10	4%	1%	6x	7x
TC015612	20.32	182	36%	+7	6%	6%	6x	2x
TC007901	7.25	64	25%	+5	7%	10%	6x	5x
TC015304	19.30	180	38%	+2	6%	9%	5x	no hit
TC011733	11.89	106	46%	+3	2%	0%	5x	5x
TC003374	12.22	124	61%	+3	1%	9%	2x	9x
TC008557	17.82	172	31%	+2	18%	0%	3x	5x
TC015754	15.69	140	34%	+5	4%	7%	2x	2x
TC000435	11.84	105	37%	+5	5%	0%	2x	2x
TC009096	12.84	111	16%	+16	9%	6%	2x	2x

(Table 5-2, we included the properties of several known AMPs as a reference). Although the antimicrobial properties of these peptides still have to be experimentally verified, this shows the strength of unbiased approaches to find novel immune genes.






Figure 5-5: *In situ* hybridization showing expression of AMP genes in the serosa upon septic injury. a-f *thaumatin1* *in situ* hybridization. a) Superficial view. *thaumatin1* is expressed around the site of injury (asterix). Brown melanisation is observed around the site of injury. a') Magnification of the expression area shown in a. Asterix marks the site of injury. b) DAPI counterstaining of the same egg as in a. The large polyploid serosal nuclei can be distinguished from the oversaturated DAPI signal from the germ band. Head lobes to the left. b') magnification of b. c) Overlay of the *in situ* hybridization shown in a and the DAPI staining shown in b. The *thaumatin1* expression associates with the large polyploid serosal nuclei and is not found in the embryo proper. c') magnification of the expression area shown in c. d) Focal plane through the egg. *Thaumatin1* is expressed in a thin outer layer at the surface of the egg. d') Magnification of the expression area shown in d. e) DAPI staining of the same egg shown in d. The embryo is brightly visible. Head to the left. e') Magnification of e. f) Overlay of the *in situ* hybridization shown in d and the DAPI staining shown in e. f') Magnification of the expression area. g) *attacin1* *in situ* hybridization. Brown melanisation is visible around the site of injury (arrowhead). g') Magnification of the anterior region of the egg shown in g. h) DAPI staining of the same egg shown in g. The germ band is brightly stained (head to the left) and the separate large serosal nuclei are visible. h') Magnification of the anterior of the egg shown in h. i) Overlay of the *in situ* hybridization shown in g and the DAPI staining shown in h. *Attacin1* is expressed in the large serosal cells covering the germ band and is not expressed in the dense cells of the germ band. i') Magnification of the anterior of the egg shown in i. The *attacin1* staining associates with the large serosal nuclei.



The immune response is dependent on the extraembryonic serosa

To investigate the role of the serosa in the immune response, we compared the transcriptional response of wild-type and control eggs to the response of serosa-less eggs. Of all 538 genes differentially regulated upon bacterial challenge, 481 genes are only responsive in eggs with a serosa. The vast majority, 276 genes, are differentially regulated in both wild-type and control eggs, but not in serosa-less eggs (Figure 5-4b). In the serosa-less *Tc-zen1* RNAi eggs merely 57 genes are differentially regulated upon microbial challenge, despite our finding that RNAi rather increases the number of immune responsive genes. Of all 368 *Tribolium* genes that are annotated as immune genes (Altincicek et al., 2013; Zou et al., 2007), only 9 were differentially regulated upon infection in serosa-less eggs (Table 5-1 and Supplementary Table 5-2). Except for *serpin24*, all of the other 8 genes were also differentially regulated in response to sterile injury, indicating that they do not respond to infection but to wounding. Notably, none of the AMPs is induced upon infection in serosa-less eggs, neither *proPO1*, nor the DUOX ortholog *Hpx11* (Supplementary Table 5-2). Thus, the serosa is essential for the early immune response of the *Tribolium* egg.




Figure 5-6: Constitutive expression of immune genes in the serosa. a-c *toll3* *in situ* hybridization. a) *toll3* is expressed in the flat and thin serosal cells (partly detached from the egg), but also in the germ rudiment (head lobes to the right). a') Magnification of the area indicated with an arrow in a. b) DAPI staining of the same egg shown in a. The bright staining of the germ band can be distinguished from the large nuclei of the serosa. c) Overlay of the *in situ* hybridization shown in a and the DAPI staining shown in b. c') Magnification of c. *Toll3* is expressed in cells of the serosa. d-f *scavenger receptor B5* *in situ* hybridization. d) *scavenger receptor B5* shows expression in every serosal cell at the surface. d') Magnification of d. e) DAPI staining of the same egg shown in d. The germband is brightly stained (head to the left) and the staining of the serosal nuclei is clearly visible when not overwhelmed by staining of the dense nuclei of the germ band. e') Magnification of e. The serosal nuclei are visible. Bright staining of the germband to the right. f) Overlay of the *in situ* hybridization shown in d and the DAPI staining shown in e. *Scavenger receptor B5* expression follows the serosal nuclei, and is not detected in the germ band. f') Magnification of f. *Scavenger receptor B5* mRNA is detected around the large polyploid serosal nuclei, and not around the dense nuclei of the germ rudiment.



These data corroborate our previous qPCR study showing that AMP and PGRP upregulation upon infection is abolished in serosa-less eggs (Jacobs & van der Zee, 2013). To see if we could also independently confirm serosa-dependent induction of some of our newly identified candidates, we performed qPCR on the transmembrane recognition protein of the IMD pathway *PGRP-LC*, the serine proteases *cS-P8*, *SPH-H57*, *SPH-H70*, the serine protease inhibitors *serpin24* and *serpin26*, the Toll receptor *toll3* and the novel potential AMPs *TC004646*, *TC007763*, *TC007857*, *TC008806* and *TC015479* (Supplementary Figure 5-1). The fold-changes detected by qPCR after sterile and septic injury of wild-type eggs match the values found in the RNAseq data. The largest deviation was found for the potential AMP *TC007858* that is upregulated 156 times upon septic injury in our qPCRs, but 484 times according to the RNAseq data (Supplementary Figure 5-1j). Most importantly, all qPCRs convincingly showed absence of induction in *Tc-zen1* RNAi eggs, thus providing independent support for our conclusion that the serosa is required for the immune response in *Tribolium* eggs.

To investigate if it is the serosal epithelium itself that expresses the identified immune genes and to exclude indirect effects, we performed *in situ* hybridization on two AMPs (*thaumatin1* and *attacin1*) of which mRNA length permitted *in situ* detection. In naive eggs, we could not detect *thaumatin1* or *attacin1* expression. In contrast, expression was obvious in challenged eggs (Figure 5-5). In these eggs, brown melanisation was found at the site of injury (asterix in Figure 5-5a and a' and arrowhead in Figure 5-5g) and the individual nuclei of the serosa can be distinguished from the oversaturated DAPI signal marking the germband (Figure 5-5b, e, h) (Handel et al., 2000). The *thaumatin1* expression clearly associates with the large polyploid serosal nuclei and not with the dense cells of the germ band (overlay in Figure 5-5c and c'). A deeper focal plane of a different egg demonstrates exclusive expression in the overlying serosa on the outer surface (Figure 5-5d, d'), and not in the underlying embryo proper (Figure 5-5e, f). Also *attacin1* expression consistently associated with the large polyploid serosal nuclei (Figure 5-5g-i').

Thus, it is the serosal epithelium itself that expresses these AMPs upon infection. Although we cannot exclude an indirect role of the serosa in the expression of the other identified immune genes, we propose that the serosa itself expresses these genes and thus regulates the described immune response involving melanisation, the generation of reactive oxygen species and the massive production of AMPs.

The serosa constitutively expresses some immune genes

To discover immune genes that are constitutively expressed in the serosa, we compared the transcriptomes of naive *Tc-zen1* eggs to naive wild-type eggs. We found 44 immune genes that have serosa-dependent expression (Table 5-3). Of these genes, more than 75% is involved in the recognition of microbes and extracellular signal transduction (Table 5-3) such as *PGRP-LA*, many serine proteases and *Spz4* and *Spz5*. In contrast, most of the genes of the intracellular signal transduction were present in *Tc-zen1* eggs at similar levels as in wild-type eggs. Notably, the transmembrane receptor *toll3* exhibits higher expression in unchallenged eggs with a serosa than in eggs without a serosa. These data indicate that the serosa is an immune competent epithelium that expresses many genes involved in bacterial recognition and transduction of this recognition to receptor activation.

To confirm constitutive expression of these identified genes, we performed *in*



situ hybridization on naive eggs. We chose the receptor *toll3* that shows 2 times higher expression in eggs with a serosa, and the *scavenger receptor B5* that shows 30 times higher expression in eggs with a serosa (Table 5-3). We found ubiquitous expression of *toll3* in the egg (Figure 5-6a). Although *toll3* was clearly expressed in the serosa (partly detached from the egg Figure 5-6a'), we detected also expression in the embryo. As *in situ* hybridization is not a quantitative technique, and because the serosal cells are flat and thin, it is possible that we could not detect the two-fold higher expression in the serosa. For *scavenger receptor B5* that has a 30-fold higher expression in eggs with a serosa, we did find clear expression in the serosal epithelium (Figure 5-6d), whereas the underlying germ band was not stained (Figure 5-6f and f'). We propose that all genes listed in Table 5-3 are constitutively expressed in the serosa and thus make the serosa an immune-competent frontier epithelium.

Taken together, we have shown that eggs of the beetle *Tribolium castaneum* display an extensive transcriptional immune response. This response is entirely dependent on the serosa, an extraembryonic epithelium that envelops yolk and embryo. This immune competent frontier epithelium constitutively expresses some immune genes and can induce massive amounts of AMPs.

Discussion

We have provided the first characterization of the complete transcriptional immune response in an insect egg. We identified 538 immune responsive genes in the *Tribolium* egg, of which 481 are only found in eggs with a serosal epithelium. The number of immune-responsive genes found in the *Tribolium* egg is comparable to the number found in larvae (Behrens et al., 2014) and higher than what was found in adults (Altincicek et al., 2013), but this might be due to differences in sequence coverage. We cannot exclude that some expression differences we found might be due to somewhat delayed development after pricking the eggs. However, the GO-categories of the differentially regulated genes (shown in Figure 5-3) are mainly immune-related, suggesting that an effect of delayed development is negligible.

The induction of several genes from both the Toll- and IMD-pathway indicates that both pathways are utilized in the immune response of the *Tribolium* egg. It is striking that Toll signaling seems to be involved in innate immunity in the egg, because Toll signaling at this stage has only been associated with developmental functions until now (Leulier & Lemaitre, 2008; Nunes da Fonseca et al., 2008). In *Drosophila*, Toll1 has been described as the essential immune-related Toll receptor (Leulier & Lemaitre, 2008). Other Tolls are not essential for the immune response, except for an antiviral function of Toll7 (Nakamoto et al., 2012). Interestingly, *toll3* is significantly upregulated upon infection of the egg, and not *toll1*. *Toll3* is also upregulated in infected adults and larvae (Altincicek et al., 2013; Behrens et al., 2014), suggesting a novel role for *toll3* in *Tribolium* innate immunity. It should be noted that *toll1-4* in *Tribolium* are all closely related to *Drosophila toll1*, and more distantly to *Drosophila toll3* (Nunes da Fonseca et al., 2008; Zou et al., 2007). Thus, subfunctionalization into developmental and immune-related functions might have occurred among the *Tribolium toll1-4* paralogs.



Table 5-3: Differentially regulated immune genes in naive wild-type eggs compared to naive *Tc-zen1* eggs.
SP = Serine protease ; SPH = Non catalytic serine protease ; cSP = Clip-domain serine protease

Gene ID	Description	Fold change	FDR adjusted p-value	Gene ID	Description	Fold change	FDR adjusted p-value
Extracellular signal transduction and modulation							
TC000247	cSPH-H2	2.70	<0.01	TC005754	serpin22	5.26	<0.01
TC000248	cSPH-H3	4.11	<0.01	TC006255	serpin24	0.69	0.03
TC000249	cSPH-H4	5.16	<0.01	TC011718	serpin27	1.62	<0.01
TC000740	SPH-H17	9.28	<0.01	TC006726	Spz4	3.00	<0.01
TC000829	SPH-H18	8.26	<0.01	TC013304	Spz5	122.56	<0.01
TC007026	cSPH-H78	29.79	<0.01	Microbial recognition			
TC012390	SPH-H129	1.60	<0.01	TC002789	PGRP-LA	3.95	0.02
TC000495	cSP-P8	6.57	<0.01	TC014664	TEP-B	2.90	0.02
TC000497	cSP-P10	4.50	<0.01	TC005976	PSH	3.43	<0.01
TC000547	SP-P13	2.41	<0.01	TC006978	C-type lectin1	14.52	<0.01
TC000635	SP-P16	2.54	<0.01	TC013911	C-type lectin 13	18.21	<0.01
TC004160	cSP-P44	9.79	<0.01	Toll-signalling pathway			
TC004624	cSP-P52	0.52	<0.01	TC004438	Toll3	2.28	<0.01
TC004635	cSP-P53	51.56	<0.01	IMD-signalling pathway			
TC005230	cSP-P61	250.00	<0.01	TC014708	NFAT	2.01	<0.01
TC006033	SP-P68	1.54	<0.01	Execution mechanisms			
TC009090	cSP-P91	2.80	<0.01	TC005375	hexamerin2	0.38	<0.01
TC009092	cSP-P93	3.00	<0.01	TC005493	Heme peroxidase 1	3.84	<0.01
TC009093	cSP-P94	27.76	<0.01	TC015234	Heme peroxidase 2	6.30	<0.01
TC013277	cSP-P136	3.04	<0.01	TC010356	Scavenger receptor-B13	0.60	0.03
TC013415	SP-P141	11.85	<0.01	TC015854	Scavenger receptor-B2	1.91	<0.01
TC000760	serpin1	5.29	<0.01	TC014946	Scavenger receptor-B5	29.29	<0.01
TC005750	serpin18	1.92	<0.01	TC000948	Scavenger receptor-B6	163.90	<0.01
TC005752	serpin20	2.30	<0.01	TC014954	Scavenger receptor-B9	1.96	<0.01

In *Tribolium*, only 19 AMPs have been identified (Zou et al., 2007). This is in strong contrast to another beetle species, *Harmonia axyridis*, in which more than 50 putative AMPs have been recognized (Vilcinskas et al., 2013). We were able to identify 20 new potential AMPs based on the antimicrobial properties of known AMPs (Bulet et al., 1999; Bulet & Stöcklin, 2005). Additional AMPs might still be discovered, as we have not investigated peptides longer than 200 amino acids. Some of these long peptides, for instance the Thaumatinins, are known to have antimicrobial properties (Altincicek et al., 2013; Altincicek et al., 2008). We might also have missed AMPs because some might be specifically expressed at other stages, for instance in larvae or adults. Although activity assays against bacteria and fungi are needed to verify antimicrobial properties, the discovery of 20 new potential AMPs shows the power of our experimental strategy for getting an unbiased understanding of insect immunity.

The most important conclusion of our study is that the immune response in *Tribolium* eggs depends on the extraembryonic serosa. To delete the serosa, we used parental *Tc-zen1* RNAi (van der Zee et al., 2005). Formally, it is possible that the lack of the immune response we reported is not caused by absence of the serosa, but by a more direct effect of *Tc-zen1* RNAi, for instance if the transcription factor Zen would directly regulate immune genes in the embryo. This is highly unlikely, as *Tc-zen1* is only expressed in the early serosa (van der Zee et al., 2005) and is not expressed anymore by the time



we performed infection. Indeed, we found only three RNAseq reads that map to *Tc-zen1*, confirming that *Tc-zen1* is practically not expressed at the time we performed experiments. Thus, we are confident to conclude that the lack of the full-range immune response after *Tc-zen1* RNAi is exclusively due to the absence of the serosa.

Eggs with a serosa express crucial bacterial recognition genes, such as *PGRP-LA*, and many extracellular signaling components, such as serine proteases, at higher levels than serosa-less eggs, indicating constitutive expression in the serosa. It could be that these components activate receptors elsewhere in the egg, for instance the *toll3* receptor that is more ubiquitously expressed. However, our *in situ* hybridizations unambiguously demonstrate that it is the serosal epithelium itself that expresses AMPs upon infection, indicating that it is the serosal epithelium itself that harbors the functional immune response reducing bacterial propagation in infected eggs (Figure 5-1).

Overall, bacterial infection of *Tribolium* eggs induces genes involved in melanisation, the acute-phase oxidative response and AMP production, and differentially regulates many other immune genes. This response is completely abolished in eggs without a serosa, the extraembryonic epithelium that envelopes yolk and embryo at the interface with the microbe-rich external milieu. Barrier epithelia like the midgut have recently been highlighted as key players in the local immune defenses in insects (Davis & Engstrom, 2012; Ferrandon, 2013). We conclude that the serosa is a frontier epithelium that provides the insect egg with a full-range immune response.

Interestingly, the separation of the serosal cells from the germ rudiment is the first morphological distinction that can be made in the blastoderm of the developing egg (Handel et al., 2000). The serosal cells will have enveloped the complete embryo before the ectoderm starts to differentiate. These serosal cells can provide the insect egg with an innate immune response long before the embryonic ectoderm or trachea are immune responsive. In addition, the polyploid nuclei allow the serosal cells to quickly synthesize large amounts of proteins providing protection for the vulnerable developing embryo. Thus, the serosa is well suited to provide early immune protection to the egg. *Drosophila* eggs do not develop a serosa, as this extraembryonic membrane was lost in the Schizophoran flies (Rafiqi et al., 2008; Schmidt-Ott, 2000). A trade-off with developmental speed might have driven the loss of the serosa in these flies living on ephemeral food sources (Jacobs et al., 2014). We suggest that the absence of the serosa in the Schizophora accounts for the poor immune response of *Drosophila* eggs. Since all other insects possess a serosa, we propose that early immune competence is a general property of insect eggs.

Conclusions

Tribolium castaneum eggs can mount a full-range innate immune response involving antimicrobial peptides, melanisation and the production of reactive oxygen species. This response depends entirely on the extraembryonic serosa, an immune competent frontier epithelium that is absent in *Drosophila*.

Materials and methods

Beetles and *Tc-zen1* RNAi

The *Tribolium* stock used for this study was the *T. castaneum* wild-type strain San Bernardino. Stock keeping and *Tc-zen1* RNAi was performed as described in (van der Zee



et al., 2005). The control dsRNA was synthesized from a 500bp vector sequence cloned from the pCRII vector (Invitrogen) using the primers 5'-TGCCGGATCAAGAGCTACCAA-3' and 5'-TGTGAGCAAAAGGCCAGCAA-3' and has no targets in the *Tribolium castaneum* genome (see also Jacobs et al., 2013; Jacobs & van der Zee, 2013).

Infection

Infection experiments were performed as described in (Jacobs & van der Zee, 2013). 24-40h old eggs (total developmental time is close to 85h) were pricked with a sterile tungsten needle or with a tungsten needle dipped in a concentrated mix of *E. coli* and *M. luteus* cultures (bacteria provided by D. Ferrandon, Strasbourg) or were not pricked at all. To allow comparison to the extensive body of work in *Drosophila*, we have used the same strains of *E. coli* and *M. luteus* as are traditionally used in *Drosophila* (Ferrandon et al., 2007). 6h later, eggs were used for RNA isolation or *in situ* hybridization.

Cfu counts

Cfu's were determined directly after infection (t=0) or 6h after infection (t=6). Eggs were shortly washed for 15 seconds in a 0.5% hypochlorite solution to sterilize the outside, and rinsed with water. 10 eggs were pooled and homogenized in 100 µl water with a sterile pestle. For t=0, 25 µl of this suspension was directly plated on LB agar plates; for t=6 these 100 microliters were either diluted 50 times in 50 µl water (for wild-type eggs), or 500 times in 50 µl water (for *Tc-zen1* eggs). Of these dilutions 25 µl was plated on LB agar plates. Colonies were counted after an overnight incubation at 37 °C, and average numbers of cfu's were calculated per egg. For each combination of time and treatment, the cfu's were measured 10 times. Statistical significance was determined by performing a Pearson's chi-square test. Bacterial load of wild-type eggs increased to on average 32,975 cfu's after 24h, but at this time point comparisons to *Tc-zen1* RNAi eggs were unreliable as bacteria might have reached a maximum. At t=6, bacteria were still in their exponential growth phase and the formula $N_{(t)} = N_{(0)} * e^{kt}$ could be used to calculate the specific growth rate.

Sample collection for transcriptional analysis

For RNAseq and qPCR, total RNA of approximately 300 eggs was extracted using TRIzol extraction (Invitrogen) after which the RNA was purified and DNA digested on column with the RNeasy kit (Qiagen). We collected 3 biological samples for each of the 9 treatments, giving a total of 27 biological samples (Figure 5-2). cDNA library synthesis and sequencing was performed by the ZF-screens (Leiden, the Netherlands) sequencing company on an Illumina HiSeq2500 sequencer.

Data analysis and bioinformatics

Sequencing reads were mapped with CLC genomics workbench 6 using the first 51 bp with highest sequencing quality and score values over 20, allowing 2 mismatches to the reference sequence of the *Tribolium* genome 3.0 which was obtained from Ensemble (Flicek et al., 2013). The mismatch cost was set to 2, the insertion cost to 3, the deletion cost to 3, the length fraction to 0.5 and the similarity fraction was set at 0.8. To calculate statistical differences of the expression levels of genes between treatments we utilized the DESeq package (Anders & Huber, 2010) in Bioconductor (Gentleman et al., 2004) in R (R Development Core Team, 2009). The P values were adjusted for multiple testing with the



Benjamini-Hochberg procedure, which determines false discovery rate (FDR). We trimmed the data to only contain genes that are induced more than 2 fold or repressed more than 2 fold. To minimize false discovery rate we set the cut-off value for significant genes to an FDR of <0.01. DESeq was used to normalize the count data, calculate mean values, fold changes, size factors, variance and P values (raw and adjusted) of a test for differential gene expression based on generalized linear models using negative binomial distribution errors.

Sequence annotation

Sequence homology searches of predicted reference gene sequences and subsequent functional annotation by gene ontology terms (GO) and InterPro terms (InterProScan, EBI) were determined using the BLAST2GO software suite v2.6.6 (Conesa et al., 2005). First, homology searches were performed through BLASTX against sequences of the *Drosophila* protein database with a cut-off value of 1.0E-10. Subsequently, GO classification annotations were created after which InterPro searches on the InterProEBI web server were performed remotely by utilizing BLAST2GO.

qPCR

RNA was collected as described under “Sample collection for transcriptional analysis”. The quality of RNA preparation was confirmed spectro-photometrically and on gel. One microgram of total RNA was used for cDNA synthesis. First strand cDNA was made using the Cloned AMV First Strand Synthesis kit (Invitrogen). Each qRT-PCR mixture (25 µl) contained 2.5 ng of cDNA, and the real-time detection and analyses were done based on SYBR green dye chemistry using the qPCR kit for SYBR Green I (Eurogentec) and a CFX96 thermocycler (Bio-rad). Thermal cycling conditions used were 50 °C for 2 min, 95 °C for 10 min, then 50 cycles of 95 °C for 15 s, 60 °C for 30 s, 72 °C for 30 s; this was followed by dissociation analysis of a ramp from 65 to 95 °C with a read every 0.5 °C. Relative quantification for each mRNA was done using the Livak-method (Livak & Schmittgen, 2001). The values obtained for each mRNA were normalized by *RPL13a* mRNA amount for *Tribolium* (primers as in Lord et al., 2010). Total RNA for each treatment was isolated two times (biological replication) and each sample was measured by qRT-PCR twice (technical replication). The primers used for qPCR are in Supplementary Table 5-3.

In situ hybridizations

In situ hybridizations involving alkaline phosphatase-based visualization of DIG-labelled probes were essentially performed as described in (Tautz & Pfeifle, 1989), but without the proteinaseK step. Eggs were fixed for 20 minutes in a 1:1 mix of heptane and 3,7% formaldehyde in PBST. As the serosa tightly associates with the vitelline membrane, we used *Tc-CHS1* RNAi eggs (Jacobs et al., 2013), making it possible to manually dissect eggs containing the serosa from the vitelline membrane. The following primers were used to amplify 500bp fragments of *thaumatin1*, *attacin1*, *toll3* and *scavenger receptor B5*. Thaumatin1 FW 5'-CTAAGCGAAGGGGGTTTCGT-3' RV 5'-TTTGTGGTCATCGTAGGCGT-3' Attacin1 FW 5'-ATCGTCCAAGACCAGCAAGG-3' RV 5'-GAAGCGGTGGCTAACTGGA-3' Toll3 FW 5'-AACTGGGAGGTTTTGCACAC-3' RV 5'-AACTCCATTTCCCCCAAAC-3' SR-B5 FW 5'-AGCCAGGGAGTTCATGTTTCG-3' RV 5'-TGATTTGGTAACGGACGGCA-3' PCR fragments were cloned into the TOPO II vector (Invitrogen) according to the



manufacturer's protocol. From these plasmids, templates for probe synthesis were amplified using M13 primers. DIG-labelled probes were synthesized using the MEGAscript kit (Ambion) according to the manufacturer's protocol, but with Roche RNA-labelling mix (Roche).

Data access

The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (Barrett et al., 2013) and are accessible through GEO Series accession number GSE54018 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE54018>).

Acknowledgments

We thank Nora Braak, Romée de Blois and O.L. van de Peppel for help with the *in situ* hybridizations. Heiko Vogel from the Max Planck Institute for Chemical Ecology, Jena, Germany for critically reading the manuscript. MvdZ was funded by NWO VENI grant 863.09.014.

References

- Altincicek, B., Elashry, A., Guz, N., Grundler, F. M., Vilcinskas, A., & Dehne, H. W. (2013). Next generation sequencing based transcriptome analysis of septic-injury responsive genes in the beetle *Tribolium castaneum*. *PLoS ONE*, 8(1), e52004. doi: 10.1371/journal.pone.0052004
- Altincicek, B., Knorr, E., & Vilcinskas, A. (2008). Beetle immunity: Identification of immune-inducible genes from the model insect *Tribolium castaneum*. *Developmental and Comparative Immunology*, 32(5), 585-595. doi: 10.1016/j.dci.2007.09.005
- Altincicek, B., & Vilcinskas, A. (2007). Analysis of the immune-inducible transcriptome from microbial stress resistant, rat-tailed maggots of the drone fly *Eristalis tenax*. *BMC Genomics*, 8, 326. doi: 10.1186/1471-2164-8-326
- Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biology*, 11(10), R106.
- Barrett, T., Wilhite, S. E., Ledoux, P., Evangelista, C., Kim, I. F., Tomashevsky, M., . . . Soboleva, A. (2013). NCBI GEO: archive for functional genomics data sets—update. *Nucleic Acids Research*, 41(D1), D991-D995. doi: 10.1093/nar/gks1193
- Behrens, S., PeuSZ, R., Milutinovi, B., Eggert, H., Esser, D., Rosenstiel, P., . . . Kurtz, J. (2014). Infection routes matter in population-specific responses of the red flour beetle to the entomopathogen *Bacillus thuringiensis*. *BMC Genomics*, 15(1), 445.
- Bell, J. V. (1969). *Serratia marcescens* found in eggs of *Heliothis zea*: Tests against *Trichoplusia ni*. *Journal of Invertebrate Pathology*, 13(1), 151-152.
- Blum, M. S., & Hilker, M. (2008). Chemical Protection of Insect Eggs. In M. Hilker & T. Meinert (Eds.), *Chemoecology of Insect Eggs and Egg Deposition* (pp. 61-90): Blackwell Publishing Ltd.
- Boos, S., Meunier, J., Pichon, S., & Kölliker, M. (2014). Maternal care provides antifungal protection to eggs in the European earwig. *Behavioral Ecology*, 25(4), 754-761. doi: 10.1093/beheco/aru046
- Bulet, P., Hetru, C., Dimarcq, J. L., & Hoffmann, D. (1999). Antimicrobial peptides in insects; structure and function. *Developmental and Comparative Immunology*, 23(4-5), 329-344.
- Bulet, P., & Stöcklin, R. (2005). Insect Antimicrobial Peptides: Structures, Properties and Gene Regulation. *Protein and Peptide Letters*, 12, 3-11.
- Bulet, P., Stöcklin, R., & Menin, L. (2004). Anti-microbial peptides: from invertebrates to vertebrates. *Immunological Reviews*, 198(1), 169-184. doi: 10.1111/j.0105-2896.2004.0124.x
- Chen, G., Handel, K., & Roth, S. (2000). The maternal NF-kappa B/Dorsal gradient of *Tribolium castaneum*: dynamics of early dorsoventral patterning in a short-germ beetle. *Development*, 127(23), 5145-5156.
- Christophides, G. K., Zdobnov, E., Barillas-Mury, C., Birney, E., Blandin, S., Blass, C., . . . Kafatos, F. C. (2002). Immunity-Related Genes and Gene Families in *Anopheles gambiae*. *Science*, 298(5591), 159-165. doi: 10.1126/science.1077136



- Conesa, A., Götz, S., García-Gómez, J. M., Terol, J., Talón, M., & Robles, M.** (2005). Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics*, 21(18), 3674-3676. doi: 10.1093/bioinformatics/bti610
- Contreras, E., Rausell, C., & Real, M. D.** (2013). Proteome Response of *Tribolium castaneum* Larvae to *Bacillus thuringiensis* Toxin Producing Strains. *PLoS ONE*, 8(1), e55330. doi: 10.1371/journal.pone.0055330
- Davis, M. M., & Engstrom, Y.** (2012). Immune response in the barrier epithelia: lessons from the fruit fly *Drosophila melanogaster*. *J Innate Immun*, 4(3), 273-283. doi: 10.1159/000332947
- De Gregorio, E., Spellman, P. T., Rubin, G. M., & Lemaitre, B.** (2001). Genome-wide analysis of the *Drosophila* immune response by using oligonucleotide microarrays. *Proceedings of the National Academy of Sciences*, 98(22), 12590-12595. doi: 10.1073/pnas.221458698
- Dziarski, R., Platt, K. A., Gelius, E., Steiner, H., & Gupta, D.** (2003). Defect in neutrophil killing and increased susceptibility to infection with nonpathogenic gram-positive bacteria in peptidoglycan recognition protein-S (PGRP-S)-deficient mice. *Blood*, 102(2), 689-697. doi: 10.1182/blood-2002-12-3853
- Ferrandon, D.** (2013). The complementary facets of epithelial host defenses in the genetic model organism *Drosophila melanogaster*: from resistance to resilience. *Current Opinion in Immunology*, 25(1), 59-70. doi: <http://dx.doi.org/10.1016/j.coi.2012.11.008>
- Ferrandon, D., Imler, J. L., Hetru, C., & Hoffmann, J. A.** (2007). The *Drosophila* systemic immune response: sensing and signalling during bacterial and fungal infections. *Nature Reviews Immunology*, 7(11), 862-874. doi: 10.1038/nri2194
- Flicke, P., Ahmed, I., Amode, M. R., Barrell, D., Beal, K., Brent, S., . . . Searle, S. M. J.** (2013). Ensembl 2013. *Nucleic Acids Research*, 41(D1), D48-D55. doi: 10.1093/nar/gks1236
- Freitag, D., Schmidtberg, H., Dickel, F., Lochnit, G., Vogel, H., & Vilcinskas, A.** (2014). The maternal transfer of bacteria can mediate trans-generational immune priming in insects. *Virulence*, 5(4), 547-554.
- Ganesan, S., Aggarwal, K., Paquette, N., & Silverman, N.** (2011). NF-kappaB/Rel proteins and the humoral immune responses of *Drosophila melanogaster*. *Curr Top Microbiol Immunol*, 349, 25-60. doi: 10.1007/82_2010_107
- Gentleman, R., Carey, V., Bates, D., Bolstad, B., Dettling, M., Dudoit, S., . . . Zhang, J.** (2004). Bioconductor: open software development for computational biology and bioinformatics. *Genome Biology*, 5(10), R80.
- Gerardo, N. M., Altincicek, B., Anselme, C., Atamian, H., Barribeau, S. M., de Vos, M., . . . Vilcinskas, A.** (2010). Immunity and other defenses in pea aphids, *Acyrthosiphon pisum*. *Genome Biology*, 11(2), R21. doi: 10.1186/gb-2010-11-2-r21
- Handel, K., Grunfelder, C. G., Roth, S., & Sander, K.** (2000). *Tribolium* embryogenesis: a SEM study of cell shapes and movements from blastoderm to serosal closure. *Development Genes and Evolution*, 210(4), 167-179.
- Jacobs, C. G. C., Rezende, G. L., Lamers, G. E. M., & van der Zee, M.** (2013). The extraembryonic serosa protects the insect egg against desiccation. *Proceedings of the Royal Society B: Biological Sciences*, 280(1764). doi: 10.1098/rspb.2013.1082
- Jacobs, C. G. C., & van der Zee, M.** (2013). Immune competence in insect eggs depends on the extraembryonic serosa. *Developmental & Comparative Immunology*, 41(2), 263-269. doi: <http://dx.doi.org/10.1016/j.dci.2013.05.017>
- Jacobs, C. G. C., Wang, Y., Vogel, H., Vilcinskas, A., Van der Zee, M., & Rozen, D. E.** (2014). Egg survival is reduced by grave-soil microbes in the carrion beetle, *Nicrophorus vespilloides*. *BMC Evolutionary Biology*, in press.
- Johnston, P. R., Makarova, O., & Rolff, J.** (2013). Inducible Defenses Stay Up Late: Temporal Patterns of Immune Gene Expression in *Tenebrio molitor*. G3 (Bethesda). doi: 10.1534/g3.113.008516
- Johnston, P. R., & Rolff, J.** (2013). Immune- and wound-dependent differential gene expression in an ancient insect. *Developmental & Comparative Immunology*, 40(3-4), 320-324. doi: 10.1016/j.dci.2013.01.012
- Kellner, R. L. L.** (2008). The Role of Microorganisms for Eggs and Progeny. In M. Hilker & T. Meinert (Eds.), *Chemoecology of Insect Eggs and Egg Deposition* (pp. 149-167): Blackwell Publishing Ltd.
- Kim, M.-S., Byun, M., & Oh, B.-H.** (2003). Crystal structure of peptidoglycan recognition protein LB from *Drosophila melanogaster*. *Nature Immunology*, 4(8), 787-793.
- Lemaitre, B., & Hoffmann, J.** (2007). The Host Defense of *Drosophila melanogaster*. *Annual Review of Immunology*, 25(1), 697-743. doi: 10.1146/annurev.immunol.25.022106.141615
- Leone, P., Bischoff, V., Kellenberger, C., Hetru, C., Royet, J., & Roussel, A.** (2008). Crystal structure of *Drosophila* PGRP-SD suggests binding to DAP-type but not lysine-type peptidoglycan. *Mol Immunol*, 45(9), 2521-2530. doi: 10.1016/j.molimm.2008.01.015
- Leulier, F., & Lemaitre, B.** (2008). Toll-like receptors- taking an evolutionary approach. *Nature Reviews Genetics*, 9(3), 165-178.
- Leulier, F., Parquet, C., Pili-Floury, S., Ryu, J. H., Caroff, M., Lee, W. J., . . . Lemaitre, B.** (2003). The *Drosophila*




- immune system detects bacteria through specific peptidoglycan recognition. *Nature Immunology*, 4(5), 478-484. doi: 10.1038/ni922
- Ligoxygakis, P.** (2013). Genetics of immune recognition and response in *Drosophila* host defense. *Adv Genet*, 83, 71-97. doi: 10.1016/b978-0-12-407675-4.00002-x
- Livak, K. J., & Schmittgen, T. D.** (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, 25(4), 402-408. doi: 10.1006/meth.2001.1262
- Lord, J. C., Hartzer, K., Toutges, M., & Oppert, B.** (2010). Evaluation of quantitative PCR reference genes for gene expression studies in *Tribolium castaneum* after fungal challenge. *Journal of Microbiological Methods*, 80(2), 219-221. doi: <http://dx.doi.org/10.1016/j.mimet.2009.12.007>
- Lynch, R. E., Lewis, L. C., & Brindley, T. A.** (1976). Bacteria associated with eggs and first-instar larvae of the European corn borer: Identification and frequency of occurrence. *Journal of Invertebrate Pathology*, 27(2), 229-237. doi: [http://dx.doi.org/10.1016/0022-2011\(76\)90150-6](http://dx.doi.org/10.1016/0022-2011(76)90150-6)
- Marchini, D., Marri, L., Rosetto, M., Manetti, A. G. O., & Dallai, R.** (1997). Presence of Antibacterial Peptides on the Laid Egg Chorion of the Medfly *Ceratitis capitata*. *Biochemical and Biophysical Research Communications*, 240(3), 657-663. doi: <http://dx.doi.org/10.1006/bbrc.1997.7694>
- Milutinović, B., Stolpe, C., Peuß, R., Armitage, S. A. O., & Kurtz, J.** (2013). The Red Flour Beetle as a Model for Bacterial Oral Infections. *PLoS ONE*, 8(5), e64638. doi: 10.1371/journal.pone.0064638
- Moreau, J., Martinaud, G., Troussard, J.-P., Zanchi, C., & Moret, Y.** (2012). Trans-generational immune priming is constrained by the maternal immune response in an insect. *Oikos*, 121(11), 1828-1832. doi: 10.1111/j.1600-0706.2011.19933.x
- Nakamoto, M., Moy, R. H., Xu, J., Bambina, S., Yasunaga, A., Shelly, S. S., . . . Cherry, S.** (2012). Virus recognition by Toll-7 activates antiviral autophagy in *Drosophila*. *Immunity*, 36(4), 658-667. doi: 10.1016/j.immuni.2012.03.003
- Nunes da Fonseca, R., von Levetzow, C., Kalscheuer, P., Basal, A., van der Zee, M., & Roth, S.** (2008). Self-Regulatory Circuits in Dorsoventral Axis Formation of the Short-Germ Beetle *Tribolium castaneum*. *Developmental cell*, 14(4), 605-615.
- Park, J. W., Kim, C. H., Rui, J., Park, K. H., Ryu, K. H., Chai, J. H., . . . Lee, B. L.** (2010). Beetle Immunity. In K. Söderhall (Ed.), *Invertebrate Immunity* (Vol. 708, pp. 163-179): Landes Bioscience and Springer Science+Business Media.
- R Development Core Team.** (2009). R: A language and environment for statistical computing (Version 2.9.0). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.R-project.org>
- Rafiqi, A. M., Lemke, S., Ferguson, S., Stauber, M., & Schmidt-Ott, U.** (2008). Evolutionary origin of the amnioserosa in cyclorrhaphan flies correlates with spatial and temporal expression changes of zen. *Proceedings of the National Academy of Sciences of the United States of America*, 105(1), 234-239. doi: 10.1073/pnas.0709145105
- Richards, S., Gibbs, R. A., Weinstock, G. M., Brown, S. J., Denell, R., Beeman, R. W., . . . Zdobnov, E. M.** (2008). The genome of the model beetle and pest *Tribolium castaneum*. *Nature*, 452(7190), 949-955.
- Roth, O., Joop, G., Eggert, H., Hilbert, J., Daniel, J., Schmid-Hempel, P., & Kurtz, J.** (2010). Paternally derived immune priming for offspring in the red flour beetle, *Tribolium castaneum*. *Journal of Animal Ecology*, 79(2), 403-413. doi: 10.1111/j.1365-2656.2009.01617.x
- Sadd, B. M., & Schmid-Hempel, P.** (2007). Facultative but persistent transgenerational immunity via the mother's eggs in bumblebees. *Current Biology*, 17(24), R1046-R1047. doi: 10.1016/j.cub.2007.11.007
- Schmidt-Ott, U.** (2000). The amnioserosa is an apomorphic character of cyclorrhaphan flies. *Development Genes and Evolution*, 210(7), 373-376.
- Shrestha, S., & Kim, Y.** (2010). Activation of immune-associated phospholipase A2 is functionally linked to Toll/Imd signal pathways in the red flour beetle, *Tribolium castaneum*. *Developmental and Comparative Immunology*, 34(5), 530-537. doi: 10.1016/j.dci.2009.12.013
- Sikowski, P. P., Lawrence, A. M., & Inglis, G. D.** (2001). Effects of *Serratia marcescens* on Rearing of the Tobacco Budworm (Lepidoptera: Noctuidae). *American entomologist*, 47(1), 51-60.
- Stroschein-Stevenson, S. L., Foley, E., O'Farrell, P. H., & Johnson, A. D.** (2005). Identification of *Drosophila* Gene Products Required for Phagocytosis of *Candida albicans*. *PLoS Biology*, 4(1), e4. doi: 10.1371/journal.pbio.0040004
- Tan, Kiri L., Vlisidou, I., & Wood, W.** (2014). Ecdysone Mediates the Development of Immunity in the *Drosophila* Embryo. *Current Biology*, 24(10), 1145-1152. doi: <http://dx.doi.org/10.1016/j.cub.2014.03.062>
- Tautz, D., & Pfeifle, C.** (1989). A non-radioactive insitu hybridization method for the localization of specific RNAs in *Drosophila* embryos reveals translational control of the segmentation gene hunchback. *Chromosoma*, 98(2), 81-85. doi: 10.1007/bf00291041



- Tingvall, T. O., Roos, E., & Engstrom, Y.** (2001). The GATA factor Serpent is required for the onset of the humoral immune response in *Drosophila* embryos. *Proceedings of the National Academy of Sciences of the United States of America*, 98(7), 3884-3888. doi: 10.1073/pnas.061230198
- Trauer, U., & Hilker, M.** (2013). Parental Legacy in Insects: Variation of Transgenerational Immune Priming during Offspring Development. *PLoS ONE*, 8(5), e63392. doi: 10.1371/journal.pone.0063392
- van der Zee, M., Berns, N., & Roth, S.** (2005). Distinct functions of the *Tribolium* zerknüllt genes in serosa specification and dorsal closure. *Current Biology*, 15(7), 624-636.
- Vilcinskis, A., Mukherjee, K., & Vogel, H.** (2013). Expansion of the antimicrobial peptide repertoire in the invasive ladybird *Harmonia axyridis*. *Proceedings of the Royal Society B: Biological Sciences*, 280(1750), 20122113. doi: 10.1098/rspb.2012.2113
- Wang, G., Li, X., & Wang, Z.** (2009). APD2: the updated antimicrobial peptide database and its application in peptide design. *Nucleic Acids Research*, 37(suppl 1), D933-D937. doi: 10.1093/nar/gkn823
- Wang, L., Kounatidis, I., & Ligoxygakis, P.** (2014). *Drosophila* as a model to study the role of blood cells in inflammation, innate immunity and cancer. *Front Cell Infect Microbiol*, 3, 113. doi: 10.3389/fcimb.2013.00113
- Wang, X.-W., & Wang, J.-X.** (2013). Pattern recognition receptors acting in innate immune system of shrimp against pathogen infections. *Fish & Shellfish Immunology*, 34(4), 981-989. doi: http://dx.doi.org/10.1016/j.fsi.2012.08.008
- Waterhouse, R. M., Kriventseva, E. V., Meister, S., Xi, Z., Alvarez, K. S., Bartholomay, L. C., . . . Christophides, G. K.** (2007). Evolutionary dynamics of immune-related genes and pathways in disease-vector mosquitoes. *Science*, 316(5832), 1738-1743. doi: 10.1126/science.1139862
- Yang, W., Cheng, T., Ye, M., Deng, X., Yi, H., Huang, Y., . . . Xia, Q.** (2011). Functional Divergence among Silkworm Antimicrobial Peptide Paralogs by the Activities of Recombinant Proteins and the Induced Expression Profiles. *PLoS ONE*, 6(3), e18109. doi: 10.1371/journal.pone.0018109
- Yokoi, K., Koyama, H., Ito, W., Minakuchi, C., Tanaka, T., & Miura, K.** (2012a). Involvement of NF- κ B transcription factors in antimicrobial peptide gene induction in the red flour beetle, *Tribolium castaneum*. *Developmental and Comparative Immunology*, 38(2), 342-351. doi: 10.1016/j.dci.2012.06.008
- Yokoi, K., Koyama, H., Minakuchi, C., Tanaka, T., & Miura, K.** (2012b). Antimicrobial peptide gene induction, involvement of Toll and IMD pathways and defense against bacteria in the red flour beetle, *Tribolium castaneum*. *Results in Immunology*, 2(0), 72-82. doi: 10.1016/j.rinim.2012.03.002
- Young, M., Wakefield, M., Smyth, G., & Oshlack, A.** (2010). Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biology*, 11(2), R14.
- Zanchi, C., Troussard, J. P., Moreau, J., & Moret, Y.** (2012). Relationship between maternal transfer of immunity and mother fecundity in an insect. *Proceedings of the Royal Society B: Biological Sciences*, 279(1741), 3223-3230. doi: 10.1098/rspb.2012.0493
- Zhong, D., Wang, M. H., Pai, A., & Yan, G.** (2013). Transcription profiling of immune genes during parasite infection in susceptible and resistant strains of the flour beetles (*Tribolium castaneum*). *Exp Parasitol*. doi: 10.1016/j.exppara.2013.01.014
- Zhu, J.-Y., Yang, P., Zhang, Z., Wu, G.-X., & Yang, B.** (2013). Transcriptomic Immune Response of *Tenebrio molitor* Pupae to Parasitization by *Scleroderma guani*. *PLoS ONE*, 8(1), e54411. doi: 10.1371/journal.pone.0054411
- Zou, Z., Evans, J. D., Lu, Z. Q., Zhao, P. C., Williams, M., Sumathipala, N., . . . Jiang, H. B.** (2007). Comparative genomic analysis of the *Tribolium* immune system. *Genome Biology*, 8(8). doi: 10.1186/gb-2007-8-8-r177



Supplementary information



Supplementary Figure 5-1: RT-qPCR verification of immune gene expression. The expression levels of several immune genes was verified by RT-qPCR. Expression shown relative to the expression in naive eggs, the mean fold change of the biological replicates (based on 2 technical replicates) is plotted and error bars show the standard error. Black bars represent expression after sterile injury, white bars represent expression after septic injury. Expression levels measured by RT-qPCR show very similar results as the expression levels measured by RNAseq (see Table S1). A) *PGRP-LC*, B) *SPH-H57*, C) *SPH-H70*, D) *cSP-P8*, E) *serpin 24*, F) *serpin 26*, G) *toll3*, H) *TC004646*, I) *TC007763*, J) *TC007858*, K) *TC008806*, L) *TC015479*. See Methods for experimental details.



Supplementary Table 5-1: Significant differently expressed immune genes in *Tribolium castaneum* eggs. (part 1/3)

Green = induction ; Red = repression ; Black = not significant

SP = Serine protease ; SPH = Non catalytic serine protease ; cSP = Clip-domain serine protease

Gene ID	Description	Wildtype eggs				Control eggs				Tc-zen1 eggs			
		Sterile injury		Septic injury		Sterile injury		Septic injury		Sterile injury		Septic injury	
		Fold change	FDR adjusted p-value	Fold change	FDR adjusted p-value	Fold change	FDR adjusted p-value	Fold change	FDR adjusted p-value	Fold change	FDR adjusted p-value	Fold change	FDR adjusted p-value
Microbial recognition													
TC0002789	PGRP-LA	1.63	0.31	3.12	<0.01	2.53	<0.01	3.33	<0.01	1.12	1.00	1.67	1.00
TC0002790	PGRP-LC	1.28	1.00	1.65	0.09	1.88	<0.01	2.43	<0.01	1.40	0.57	1.29	1.00
TC010611	PGRP-SA	4.53	0.96	241.06	<0.01	3.40	0.49	217.03	<0.01	1.11	1.00	9.38	0.89
TC013620	PGRP-SB	119.59	<0.01	937.86	<0.01	Inf	<0.01	Inf	<0.01	1.92	0.99	34.74	0.31
TC011529	βGRP2	2.65	<0.01	4.94	<0.01	2.53	<0.01	5.03	<0.01	1.13	1.00	1.29	1.00
TC003991	βGRP3	4.38	<0.01	4.90	<0.01	4.22	<0.01	5.47	<0.01	2.12	1.00	1.02	1.00
TC000808	TEP-D	1.29	1.00	1.18	1.00	0.64	0.19	0.57	0.05	0.71	0.80	0.70	1.00
TC0001981	LpR2	1.29	0.44	1.47	<0.01	1.26	0.06	1.40	<0.01	1.40	0.04	1.41	0.21
TC006978	C-type lectin1	0.83	1.00	0.40	<0.01	0.70	0.08	0.41	<0.01	0.84	1.00	0.72	1.00
TC003708	C-type lectin6	2.16	<0.01	2.38	<0.01	2.86	<0.01	2.99	<0.01	1.20	0.38	1.16	0.56
TC013911	C-type lectin13	0.22	<0.01	0.21	<0.01	0.51	0.07	0.38	<0.01	1.52	1.00	1.64	1.00
Extracellular signal transduction and modulation													
TC000248	cSPH-H3	0.75	0.54	0.48	<0.01	0.81	0.41	0.61	<0.01	0.78	0.65	0.84	1.00
TC000249	cSPH-H4	9.50	<0.01	16.54	<0.01	7.38	<0.01	12.31	<0.01	1.13	1.00	2.00	1.00
TC000252	cSPH-H6	1.20	0.81	1.23	0.16	1.11	0.83	1.16	0.41	1.37	0.02	1.31	0.31
TC000740	SPH-H17	0.58	<0.01	0.36	<0.01	0.60	0.04	0.52	<0.01	0.60	0.68	0.86	1.00
TC000829	SPH-H18	2.87	<0.01	3.92	<0.01	2.38	<0.01	3.02	<0.01	0.62	1.00	0.90	1.00
TC002150	cSPH-H34	1.09	1.00	2.40	0.60	4.84	0.13	5.91	0.04	2.91	0.33	3.01	0.80
TC004622	cSPH-H51	1.02	1.00	1.06	1.00	0.66	0.03	0.60	<0.01	0.84	0.61	0.98	1.00
TC004900	SPH-H57	6.42	0.02	22.34	<0.01	10.36	<0.01	14.76	<0.01	4.06	0.02	3.68	0.23
TC005908	SPH-H64	20.64	<0.01	16.37	<0.01	6.54	<0.01	7.57	<0.01	3.21	1.00	1.44	1.00
TC006246	SPH-H70	252.58	<0.01	495.16	<0.01	21.54	<0.01	59.55	<0.01	3.48	1.00	5.22	1.00
TC006247	SPH-H71	Inf	1.00	Inf	0.13	Inf	0.14	Inf	<0.01	0.00	1.00	0.00	1.00
TC007026	cSPH-H78	1.16	1.00	0.81	1.00	1.59	<0.01	1.13	0.78	1.05	1.00	0.95	1.00
TC011067	cSPH-H125	1.09	1.00	1.84	0.14	2.48	<0.01	2.26	0.02	2.00	0.04	1.94	0.23
TC000495	cSP-P8	7.51	<0.01	26.66	<0.01	10.81	<0.01	40.80	<0.01	1.43	1.00	1.48	1.00
TC000496	SP-P9	36.96	<0.01	110.13	<0.01	7.17	<0.01	13.49	<0.01	0.88	1.00	1.03	1.00
TC000497	cSP-P10	4.39	<0.01	7.19	<0.01	8.02	<0.01	12.89	<0.01	1.21	1.00	1.46	1.00
TC000547	SP-P13	0.67	<0.01	0.51	<0.01	0.68	<0.01	0.51	<0.01	0.72	0.02	0.75	0.23
TC002659	SP-P36	5.44	<0.01	5.06	<0.01	4.73	<0.01	4.15	<0.01	2.16	0.93	4.42	0.89
TC004160	cSP-P44	1.43	<0.01	1.97	<0.01	1.78	<0.01	2.24	<0.01	0.99	1.00	1.04	1.00
TC004523	SP-P46	4.69	<0.01	7.19	<0.01	5.38	<0.01	7.50	<0.01	1.26	0.92	1.51	0.47
TC004535	SP-P50	3.26	<0.01	4.40	<0.01	3.69	<0.01	4.90	<0.01	0.79	0.97	0.69	0.94
TC005976	cSP-PSH	2.08	<0.01	2.96	<0.01	2.54	<0.01	3.08	<0.01	1.60	0.41	1.64	0.80
TC006033	SP-P68	1.57	<0.01	1.74	<0.01	1.37	<0.01	1.48	<0.01	1.09	1.00	1.02	1.00
TC006034	SP-P69	1.38	<0.01	2.19	<0.01	1.57	<0.01	2.57	<0.01	1.52	<0.01	1.53	0.02
TC008653	cSP-P83	20.28	<0.01	17.73	<0.01	40.65	<0.01	33.53	<0.01	2.15	1.00	1.41	1.00
TC008657	cSP-P84	12.81	0.18	15.19	0.02	5.11	0.24	3.06	0.79	3.13	1.00	3.97	1.00
TC009089	cSP-P90	1.84	0.47	2.24	<0.01	1.37	0.74	2.14	<0.01	1.03	1.00	0.95	1.00
TC009090	cSP-P91	2.04	<0.01	4.23	<0.01	2.12	<0.01	5.62	<0.01	1.29	0.95	1.39	1.00
TC009092	cSP-P93	3.31	<0.01	7.32	<0.01	2.92	<0.01	7.04	<0.01	1.19	0.99	1.06	1.00
TC009093	cSP-P94	0.82	1.00	0.49	<0.01	1.16	1.00	0.72	0.11	0.72	1.00	0.68	1.00
TC009094	cSP-P95	1.50	0.08	1.36	0.11	2.35	<0.01	2.00	<0.01	1.55	0.39	1.55	0.87
TC011078	cSP-P126	7.69	<0.01	8.28	<0.01	4.11	<0.01	5.81	<0.01	0.92	1.00	1.37	1.00
TC013277	cSP-P136	4.38	<0.01	5.84	<0.01	6.27	<0.01	7.46	<0.01	1.37	0.61	1.59	0.37
TC013326	cSP-P140	1.34	0.07	1.50	<0.01	1.79	<0.01	1.82	<0.01	0.93	0.93	1.05	1.00
TC015110	SP-P153/SR-A4	1.15	1.00	1.20	0.81	1.68	0.04	1.38	0.32	0.67	0.53	0.74	1.00
TC015295	SP-P156	1.66	1.00	12.17	<0.01	2.45	<0.01	12.49	<0.01	1.66	0.66	1.54	0.85
TC015297	SP-P157	0.75	0.98	0.62	0.03	0.82	0.54	0.66	<0.01	0.48	<0.01	0.64	0.07
TC002085	serpin2	0.63	0.29	0.75	0.50	0.64	0.19	0.61	0.01	0.82	0.77	0.86	1.00
TC005750	serpin18	0.57	<0.01	0.51	<0.01	0.56	<0.01	0.56	<0.01	0.45	<0.01	0.71	0.21
TC005751	serpin19	0.76	0.97	0.97	1.00	0.73	0.61	0.94	1.00	0.32	0.04	0.43	0.33
TC005752	serpin20	0.53	<0.01	0.62	0.03	0.74	0.36	0.74	0.49	0.57	0.11	0.69	0.86
TC005753	serpin21	0.67	0.82	0.44	<0.01	0.57	0.13	0.50	0.01	0.40	<0.01	0.41	0.14
TC005754	serpin22	3.40	<0.01	4.09	<0.01	3.64	<0.01	4.32	<0.01	0.87	1.00	0.87	1.00
TC006255	serpin24	7.70	<0.01	13.68	<0.01	8.81	<0.01	13.71	<0.01	1.39	0.06	1.60	<0.01



Supplementary Table 5-1: Significant differently expressed immune genes in *Tribolium castaneum* eggs. (part 2/3)

Green = induction ; Red = repression ; Black = not significant

SP = Serine protease ; SPH = Non catalytic serine protease ; cSP = Clip-domain serine protease

TC007869	serpin26	6.38	<0.01	14.76	<0.01	7.56	<0.01	17.97	<0.01	1.09	1.00	1.33	1.00
TC011718	serpin27	3.00	<0.01	5.38	<0.01	3.15	<0.01	5.23	<0.01	1.03	1.00	0.90	1.00
TC013310	serpin28	2.93	<0.01	4.44	<0.01	2.82	<0.01	3.80	<0.01	1.04	1.00	1.08	1.00
TC014237	serpin30	8.42	<0.01	7.51	<0.01	5.64	<0.01	5.47	<0.01	1.04	1.00	1.22	1.00
TC000520	Spz1	1.41	0.42	1.76	<0.01	1.43	0.17	1.45	0.08	1.08	1.00	1.09	1.00
TC001054	Spz2	10.86	0.01	58.65	<0.01	12.16	<0.01	42.30	<0.01	2.64	1.00	2.32	1.00
TC006726	Spz4	0.56	0.02	0.42	<0.01	0.76	0.30	0.51	<0.01	0.71	0.60	0.81	1.00
TC013304	Spz5	0.42	<0.01	0.23	<0.01	0.61	0.14	0.35	<0.01	0.56	0.69	0.55	0.91
Toll-signalling pathway													
TC004438	Toll3	1.58	<0.01	1.96	<0.01	2.01	<0.01	2.49	<0.01	0.91	0.97	0.95	1.00
TC000625	Toll9	1.34	1.00	1.19	1.00	0.56	0.12	0.50	0.03	0.67	0.43	0.67	0.77
TC008202	ML1	0.53	0.46	0.43	0.02	0.68	<0.01	0.59	<0.01	0.64	0.12	0.90	1.00
TC002003	Cactus	1.18	1.00	1.52	0.06	1.27	0.07	1.28	0.03	1.08	1.00	1.09	1.00
IMD-signalling pathway													
TC010851	IMD	2.02	<0.01	3.31	<0.01	1.78	<0.01	3.39	<0.01	0.92	1.00	0.94	1.00
TC014026	casps4	0.77	0.70	0.74	0.21	0.90	0.98	0.88	0.79	0.71	0.02	0.76	0.13
TC011191	REL1	1.04	1.00	2.08	<0.01	1.06	1.00	2.39	<0.01	1.29	0.03	1.38	0.04
TC014708	NFAT	0.39	<0.01	0.31	<0.01	0.39	<0.01	0.30	<0.01	0.32	<0.01	0.42	<0.01
JNK-signalling pathway													
TC010766	Puckered	1.28	1.00	1.77	0.06	1.22	0.63	1.46	<0.01	1.54	0.04	1.46	0.40
TC011870	Kay	1.27	1.00	1.93	0.08	1.23	0.28	1.52	<0.01	1.43	<0.01	1.49	0.05
Execution mechanisms													
TC007737	Attacin1	16.69	0.10	869.41	<0.01	56.07	<0.01	3696.08	<0.01	4.02	0.70	44.43	0.18
TC007738	Attacin2	12.25	0.60	3098.01	<0.01	6.62	0.21	2190.29	<0.01	0.47	1.00	5.28	1.00
TC006250	Defensin1	1.29	1.00	187.05	0.07	5.93	1.00	1551.17	<0.01	0.51	1.00	6.30	1.00
TC010517	Defensin2	22.67	0.78	1183.49	<0.01	Inf	0.07	Inf	<0.01	0.60	1.00	2.56	1.00
TC012469	Defensin3	16.53	0.24	907.69	<0.01	Inf	0.10	Inf	<0.01	1.84	1.00	8.26	1.00
TC000517	Thaumatin1	56.16	<0.01	89.30	<0.01	71.76	<0.01	103.03	<0.01	2.17	0.28	1.58	1.00
TC000499	Cecropin1	no hit	no hit	Inf	0.65	no hit	no hit	Inf	0.01	no hit	no hit	no hit	no hit
TC000500	Cecropin3	Inf	<0.01	Inf	<0.01	21.26	<0.01	48.59	<0.01	no hit	no hit	no hit	no hit
TC005093	Coleoptericin1	7.61	0.60	2392.39	<0.01	24.39	0.02	18067.24	<0.01	0.46	1.00	12.60	1.00
TC005096	Coleoptericin2	10.09	1.00	91.36	0.38	32.45	0.18	227.32	<0.01	1.04	1.00	7.30	0.91
TC000325	ProPO1	1.58	0.34	1.80	0.03	1.38	0.36	1.52	0.03	0.93	0.95	0.96	1.00
TC014907	ProPO2	0.84	1.00	0.72	0.59	0.74	<0.01	0.73	<0.01	0.91	0.63	0.88	0.71
TC005377	hexamerin4	13.49	<0.01	13.73	<0.01	16.05	<0.01	10.85	<0.01	22.95	<0.01	23.87	<0.01
TC011090	catalase2	0.96	1.00	0.88	0.99	0.54	<0.01	0.59	0.01	0.73	0.65	0.68	0.96
TC010362	Glutathione oxidase 1	1.47	0.16	1.79	<0.01	1.44	0.41	1.64	0.02	1.50	0.17	1.38	0.78
TC005493	Heme peroxidase 1	1.03	1.00	1.31	0.30	1.33	0.89	1.89	0.02	1.21	1.00	1.19	1.00
TC004551	Heme peroxidase 5	2.01	<0.01	2.79	<0.01	2.03	<0.01	2.74	<0.01	2.66	<0.01	2.32	<0.01
TC000175	Heme peroxidase 7	2.14	<0.01	2.25	<0.01	2.17	0.02	2.08	0.01	1.06	1.00	1.28	1.00
TC001556	Heme peroxidase 9	0.96	1.00	1.02	1.00	0.62	<0.01	0.61	<0.01	0.62	<0.01	0.62	<0.01
TC004592	Heme peroxidase 11	7.36	<0.01	12.07	<0.01	5.57	<0.01	8.83	<0.01	1.65	0.51	1.41	1.00
TC011676	Superoxide dismutase 2	1.31	0.03	1.20	0.21	1.24	0.24	1.09	0.80	0.92	1.00	0.92	1.00
TC011675	Superoxide dismutase 4	1.20	1.00	1.57	0.67	2.84	0.05	2.59	0.08	2.21	0.96	1.65	1.00
TC010356	Savenger receptor B13	3.10	<0.01	4.80	<0.01	3.33	<0.01	4.68	<0.01	6.06	<0.01	4.95	<0.01
TC012758	Savenger receptor B16	0.85	1.00	0.68	0.11	0.84	0.68	0.71	0.03	0.85	0.51	1.02	1.00
TC015854	Savenger receptor B2	1.32	0.16	1.60	<0.01	1.50	<0.01	1.86	<0.01	0.72	0.10	0.68	0.32
TC008210	Savenger receptor B3	9.30	<0.01	17.12	<0.01	24.52	<0.01	38.10	<0.01	0.66	1.00	1.13	1.00
TC014946	Savenger receptor B5	1.97	<0.01	2.77	<0.01	2.91	<0.01	3.51	<0.01	1.38	1.00	1.50	0.95
TC000948	Savenger receptor B6	1.15	1.00	0.51	0.21	0.94	0.98	0.46	<0.01	1.13	1.00	0.34	1.00
TC014954	Savenger receptor B9	0.72	0.17	0.67	0.02	0.80	0.12	0.72	<0.01	1.08	1.00	1.13	0.86
TC015640	Savenger receptor C	1.11	1.00	1.39	0.04	1.27	0.79	1.369	0.37	1.63	<0.01	1.50	0.15
TC011427	Nimrod A	0.70	1.00	0.45	0.03	0.48	<0.01	0.32	<0.01	0.49	0.02	0.55	0.34



Supplementary Table 5-1: Significant differently expressed immune genes in *Tribolium castaneum* eggs. (part 3/3)
 Green = induction ; Red = repression ; Black = not significant
 SP = Serine protease ; SPH = Non catalytic serine protease ; cSP = Clip-domain serine protease

Stress-related immune-responsive genes											
TC015563	apoD	7.50	<0.01	15.03	<0.01	6.39	<0.01	12.51	<0.01	1.11	1.00
TC010172	Hsp68	6.06	<0.01	9.53	<0.01	4.86	<0.01	7.53	<0.01	2.25	0.23
TC005338	Hsp27	29.05	0.37	66.54	0.07	28.20	<0.01	67.72	<0.01	7.53	<0.01
Total number of immune genes		368		368		368		368		368	
Total upregulated		45		62		57		72		13	
Total downregulated		8		16		10		23		9	

Supplementary Table 5-3: Primers used for qRT-PCR.

Gene	Forward primer	Reverse primer
<i>Tribolium castaneum</i>		
<i>PGRP-LC</i>	5'-GAAGGAACGCTCGCTAACCC-3'	5'-CGACAGCACCATCACACTCA-3'
<i>SPH-H57</i>	5'-ACCCATCCACAATATGCGGG-3'	5'-CGCGGGATTATTTGGTCTCC-3'
<i>SPH-H70</i>	5'-GACAATTTCCGTGGCAGGTG-3'	5'-ACACCCACAGAATCCATTTTCAT-3'
<i>cSP-P8</i>	5'-CCGAATGTGGAGTCCAGGAAG-3'	5'-AATGAGAGTTCGCCACAGG-3'
<i>Serpin 24</i>	5'-TGCCACTGCTGTCAATTTTCC-3'	5'-AAACCTCGGGCGAAACAAC-3'
<i>Serpin 26</i>	5'-GGGTGTCTGAACGGACCAAA-3'	5'-AAGGGACGTCTTTTCGTGCT-3'
<i>Toll3</i>	5'-AACCAGATTACGGGCAACTACA-3'	5'-TGCAGCTCTTGTAAGCCTATGA-3'
<i>TC004646</i>	5'-GCTGATCCTCGCACTGTGTA-3'	5'-GCGAAAACGACGAAGAATTTTCATT-3'
<i>TC007763</i>	5'-AGTCTTTTGATGCGTAGCACTC-3'	5'-GTGCTGGTAGACGGGAAC-3'
<i>TC007858</i>	5'-GCACATGGCGAGCCAGATTA-3'	5'-GTCTCTCCACCCACAATGG-3'
<i>TC008806</i>	5'-ACGCTAGTGACTGTGTGGTC-3'	5'-CCAAAACTTTCCCGTTGCCT-3'
<i>TC015479</i>	5'-GACCCTCATCCTGTGCTGTC-3'	5'-GTTGTTGCACTCCGCTTCT-3'



