



Universiteit
Leiden
The Netherlands

Exploitation of host chemokine signalling by pathogenic mycobacteria

Torraca, V.

Citation

Torraca, V. (2016, November 17). *Exploitation of host chemokine signalling by pathogenic mycobacteria*. Retrieved from <https://hdl.handle.net/1887/44243>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/44243>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/44243> holds various files of this Leiden University dissertation.

Author: Torraca, V.

Title: Exploitation of host chemokine signalling by pathogenic mycobacteria

Issue Date: 2016-11-17

Chapter 1


Introduction and thesis outline

Chapter 1

Introduction and thesis outline

Bacterial infections remain a major cause of morbidity and mortality for mankind. In particular, the WHO Global Tuberculosis report (2015) estimated that one third of the world population is infected with *Mycobacterium tuberculosis* and that 9.6 million people have contracted tuberculosis (TB) in 2014 alone. An emergent problem is the prominence of multidrug and extensively drug-resistant strains. The evolution of antibiotic resistance is inevitable and requires continuous development of alternative therapeutic strategies. Therefore, a deeper understanding of the host-pathogen interface is needed. If the molecular mechanisms underlying host-pathogen interactions will be better understood, it may become possible to treat intracellular infections by targeting host cell functions that support bacterial growth, rather than targeting the pathogen itself. An important advantage of this approach is that it is more difficult for the pathogen to develop resistance to a host-targeted therapy. In this thesis, we investigated the host-pathogen dynamics during mycobacterial infection and we focused on the mechanisms by which mycobacteria can hijack chemokine signalling. Chemokines are small secreted signalling proteins that play multiple functions in a vast range of diseases. Chemokine receptors are highly expressed by immune cells and chemokine ligands are largely induced by inflammatory processes, including mycobacterial infections. However, since chemokine ligands and receptors are largely redundant in function and expression patterns, it has been difficult to address the relevance of different chemokine signalling axes for the onset and the progression of inflammatory diseases. Research on chemokine functions in TB has been particularly complicated in the most widely studied murine model. In this thesis, we alternatively applied the zebrafish (*Danio rerio*)-*Mycobacterium marinum* model to dissect how chemokines orchestrate the response of immune cells to mycobacterial infection. Several chemokine axes are conserved from fish to mammals and the similarities between human-*M. tuberculosis* and zebrafish-*M. marinum* pathology are remarkable. In this introduction, we briefly describe these aspects, which, together with the suitability of the zebrafish for microscopic analysis and genetic manipulations, have made us choose this model to investigate chemokine function in mycobacterial disease.

GRAPHICAL ABSTRACT



The human chemokine network as currently known consists of 25 receptors and 45 ligands. Ligand-receptor connecting lines indicate binding specificity. The AKCR family comprises atypical receptors that lack the motif for signalling through G-proteins and can antagonise the activities of other chemokine receptors (connected with orange lines) by binding one or more of their ligands. The chemokine receptor families include CXCR (green), XCR (cyan), CCR (blue), CX3CR (violet), ACKR (orange). Chemokine ligand families include CXCL (ELR+ chemokines in shades of green, homeostatic chemokines in shades of red, platelet chemokines in shades of orange, dual function chemokines in shades of pink), XCL (cyan), CCL (shades of blue) and CX3CL (violet).

The zebrafish model system for biomedical research

Zebrafish initially emerged as a powerful vertebrate model organism to study embryonic development and morphogenesis^{1,2,3,4,5} and this model species is now used to study the mechanisms of a wide variety of human diseases⁵. Due to their optical accessibility, the early embryonic and larval stages are ideally suited for non-invasive intravital imaging and longitudinal *in vivo* studies¹. The short generation time (3-4 months) together with the external fertilisation and the possibility to daily obtain a large number of fertilised eggs from a zebrafish adult colony has permitted the efficient application of genetic tools, thereby providing the zebrafish field with transgenic lines that fluorescently label distinct tissues, organs, cell types, but also the cellular ultrastructural components, such as the nucleus, autophagosomes and the cytoskeleton⁶.

The genome of zebrafish has been sequenced and forward genetics approaches have been successfully applied to mutate a vast array of genes^{3,4}. Additionally, reverse genetic approaches, including morpholino knockdown and mRNA injection-based overexpression, are widely used in zebrafish embryos and have permitted rapid screening of gene functions. The zebrafish model also keeps up with state-of-the-art technologies, such as novel genome editing techniques (e.g. TALENS and CRISPR/Cas9)^{7,8,9} and the application of cell/tissue-specific RNA-sequencing analysis¹⁰, which are contributing to exponentially increase the availability of tools and data for this model system.

Due to their continuous exchanges with the surrounding water, embryos and larvae are easily penetrated by a variety of drugs and vital dyes and are therefore chemically and pharmacologically tractable. With the increase of tools and successes in using zebrafish, we have gained a grounded knowledge on how to exploit this model to study diverse biological processes. Therefore, not surprisingly, the zebrafish field has promptly expanded from the initial niche in developmental research to other scientific branches, spanning from eco/toxicology to neuronal physiology, behavioural studies and immunology. The potential of zebrafish research and its translational value to develop new treatments for human diseases is also underscored by new therapeutics that emerged from initial studies in zebrafish. In particular, a milestone for the zebrafish translational research was achieved when the first drug developed through the use of zebrafish has passed phase 1 trial and advanced to phase 2 clinical studies^{11,12}.

Despite the many advantages of using the zebrafish as a model for human diseases, there are also several limitations that should be taken into account when extrapolating conclusions from experiments in this animal model. The first important aspect to consider is the evolutionary divergence. There are about 445 millions of years of evolutionary divergence between the zebrafish and the human species, which is approximately 5-fold higher than the rodents-human divergence (93 millions of years). While key aspects of the immune signalling architecture are maintained between human and zebrafish, some significant differences exist between fish and mammalian immunology. An example is the fact that the Toll-like receptor Tlr4, differently than in mammals, is not involved in bacterial lipopolysaccharide sensing in zebrafish and many other fish species, which explains also the high tolerance of fish to this compound¹³. Another difficulty when working with zebrafish, resides in the fact that the teleost genome has undergone genome duplication, which means that a variety of genes are present in two copies in these species. Some gene families have also experienced species-specific expansions in zebrafish *per se*

(e.g. the chemokine ligand families¹⁴). Adding to this, the different gene copies have sometimes undergone functional specialisation and have diverged to assume slightly different roles¹⁵.

Oftentimes regarded as an advantage, the small size of the zebrafish can be, in some circumstances, a limiting factor. In fact, the reduced dimension of zebrafish makes it difficult (in juveniles and adults) or nearly impossible (in embryos and larvae) to collect body fluids or organs, and therefore to precisely quantify bioactive molecules and chemicals in the blood or specific tissues. This has been a relevant limit, especially for the application of zebrafish larvae for high-throughput screening of drugs via bath exposure, including anti-tubercular therapeutics^{16,17}. In fact, not all the drugs can infiltrate through the skin and the impossibility of blood and tissue sampling made it impossible to distinguish inactive compounds from non-permeable false negatives. However, protocols for sampling the yolk sac have been described and it has been shown that dosage of compounds from these specimens can be used as an indication for the compound uptake, enabling therefore, to discriminate ineffective compounds from those that could not cross the skin barrier¹⁸.

The zebrafish community additionally suffers from the poor availability of immunological reagents, and from the inefficiency of site-directed genome manipulation strategies, which effectively apply to the murine model. However, the lack of antibodies is well compensated by the expanding availability of transgenic reporter lines and the possibility to apply whole mount RNA *in situ* hybridisation. Additionally, the recent advances in genome editing technology (e.g. CRISPR/Cas9 editing) has enabled precise gene disruptions and generation of conditional knockout, at par of those available in mice^{19,20}.

Finally, another limit of the zebrafish model consists in the fact that, differently from rodents, strict breeding and husbandry protocols are not widely applied in the zebrafish community and true inbred zebrafish lines are not adopted in the majority of studies. However, generation of clonal zebrafish lines is possible, either by conventional repeated sibling incrosses or even directly via gynogenesis²¹. Therefore, it is expected that the zebrafish community will increasingly adopt these lines and more standardised culturing protocols for future research.

Taken together, despite some drawbacks, the zebrafish model holds great promise for translational biomedical research as it concentrates in a single vertebrate system several useful features, including remarkable homology to mammals in biological processes, optical transparency, availability of genetic tools, chemical tractability, adaptability to high-throughput research methods and finally, suitability to address research questions at multiple levels, from the molecular level to the whole organism level.

Tuberculosis and mycobacterial infection in the zebrafish-*M. marinum* surrogate model

Mycobacterium tuberculosis (*Mtb*) is a respiratory pathogen responsible for tuberculosis (TB). The main pathologic feature of mycobacterial infection is the formation of granulomas, which are inflammatory collections of immune cells that contain the infection and that are often chronically maintained²². TB is a life threatening and global disease, with high prevalence in all human populations²³. Although many organs and tissues can be

infected with *M. tuberculosis*, the disease most often affects the lungs, with 75-80% of TB cases where the infection is exclusively pulmonary²⁴. Approximately 1.5 million people die of tuberculosis each year, and there are over nine million new cases annually²³. Although drug treatments are available, the therapeutic regimen is long and requires a combination of several pharmaceuticals. In addition, drug-resistant strains have evolved that do not respond to the currently available treatments. One third of the world population is estimated to be infected with *Mtb*. Only about 10% of infected individuals will manifest active TB disease while, in the vast majority of cases, the pathology is contained although not eliminated (latent TB). In case of latency, people are generally not contagious, and may never develop active TB. However, the infection can reactivate even many years later, especially in immune-suppressed/compromised patients, for example people subjected to chemotherapies or carrying HIV infection.

The granuloma, the histopathological hallmark of TB disease, is composed mostly of infected and non-infected immune cells that have migrated to the infectious focus. The signals responsible for granuloma formation and maintenance are only partly elucidated and it is clear that both host and pathogen signals integrate together to initiate the cellular and molecular programme that leads to granuloma formation²². As further elaborated on in **Chapter 2**, *M. marinum* (*Mm*) is a closely related species to *Mtb* that is a natural pathogen of zebrafish and other ectothermic animals. *Mm* causes necrotic granulomatous lesion formation in host tissues and these lesions are histologically very similar to those generated by *Mtb* in TB patients^{25,26,27}. Additionally, the *Mm*-zebrafish surrogate model has been proven to recapitulate key aspects of TB, including bacterial dormancy/reactivation, the coordination of host and pathogen components in the formation of granulomas, the dynamicity of these structures and the key role of the inflammatory status in controlling the infection^{28,29,30,31}. The striking conservation of molecular mechanisms, cellular dynamics and histopathological features between human and fish tuberculosis, together with the unique accessibility of the early stages of granuloma formation in zebrafish larvae, made the zebrafish-*Mm* host-pathogen pair a valuable platform to unravel the core pathogenic processes of mycobacterial infections. However, it must be taken into account that experimental fish embryo infections are generally performed using non-natural infection routes. Most commonly, the bacteria are delivered directly into the blood and generate a systemic infection that anatomically poorly represents the lung disease that is evoked in *M. tuberculosis*-infected patients. While use of a fish model makes it obviously impossible to model the organ-specific effects of TB pathology, a variety of alternative routes can be used to mimic and study specific aspects of the disease. For example, the importance of the epithelium for the initial steps of macrophage recruitment and granuloma formation has been addressed by delivering the bacteria into the hindbrain cavity³², which is delimited by an epithelial barrier that resembles the alveolar lumen. Similarly, the relevance of the vasculature response could be addressed by delivering the pathogens in the poorly vascularised trunk tissue, where progression of the infection substantially depends on the induction of local angiogenesis³³.

Human infections with *Mm* are rare and generally limited to granulomatous nodules in the tissues of the skin, owing to the optimal growth of this pathogen at lower temperatures than that of the human body. Hence, *Mm* does not represent a serious risk for health, when compared to *Mtb*. The limited safety concern and the possibility to handle *Mm* under biosafety level 2 instead of level 3 conditions greatly expands the research opportunities that can be explored using the zebrafish-*Mm* surrogate model.

Chemokines and chemokine receptors, the portrait of an “extended” family

Chemokines are chordate-specific signalling molecules. Their primary function is to control cell movements by activating specific heterotrimeric G-protein coupled receptors (GPCRs), a class of 7 transmembrane (7TM) receptors (**Figure 1, Table 1**). Chemokines and chemokine receptors play critical roles in a wide context of biological processes, both in physiological and pathological conditions, which include development³⁴, ontogenesis and homing of immune cells³⁵, angiogenesis³⁶, organogenesis³⁷, autoimmune reactions³⁸, infectious diseases³⁹ and cancer^{40,41}. These activities are not exclusively related to chemokine function in chemotaxis and cell migration, since multiple signal transduction pathways can be activated by GPCRs, including cell proliferation and differentiation⁴².

GPCRs are the largest and most diverse group of membrane receptors in eukaryotes⁴³ and their pleiotropic functions, despite similar protein and signalling architecture, have greatly influenced modern medical research. In fact, GPCRs mainly signal via well-described pathways, which has facilitated drug discovery and testing. Additionally, they are involved in a variety of biological processes and therefore therapeutic targeting of GPCRs is relevant for many diseases. It has been recently estimated that between 1/3 and 1/2 of all the drugs available on the market target GPCRs^{44,45}.

At the intracellular side of the plasma membrane, inactive GPCR/chemokine receptors constitutively interact with a three subunit G-protein complex, also known as heterotrimeric G-proteins. G-proteins are able to bind and exchange guanosine triphosphate (GTP) with guanosine diphosphate (GDP). Specifically, it is the $G\alpha$ subunit of the heterotrimer that exchanges GDP/GTP in response to conformational changes of the receptor⁴⁶. Receptor conformational switches are evoked by binding of cognate chemokine ligands to their receptors. In the inactive form, $G\alpha$ is bound to GDP. Activation of the receptor evokes physical replacement of GDP with GTP. This affects the conformation of the $G\alpha$ subunit, which in turn dissociates from both the receptor and the $G\beta\gamma$ heterodimer. Both $G\alpha$ -GTP and $G\beta\gamma$ possess lipid anchors and therefore diffuse laterally on the plasma membrane, where they interact with their downstream signalling partners⁴⁶. This phenomenon is transient and, as soon as $G\alpha$ -GTP decays to $G\alpha$ -GDP, it re-associates to the GPCR receptor and to $G\beta\gamma$. Despite this highly conserved signalling mechanism, vertebrate genomes encode multiple variants of G-proteins, which therefore can transduce diverse downstream signalling⁴⁷. Some chemokine receptors have also shown selectivity for coupling with G-protein complexes containing specific G isoforms⁴⁷.

The cascade initiated by GPCR/chemokines receptors commonly includes the activation of phosphoinositide 3-kinase (PI3K), which leads to changes in intracellular phosphatidylinositol-3,4,5-triphosphate, and mediates the recruitment of pleckstrin homology (PH) domain-containing proteins to the cell membrane⁴⁸. Several components of the Rho family of small GTPases, as well as their GTP/GDP exchange factors (GEFs) and kinases, also contain PH domains. Activation of the small GTPases (e.g. RAC1, CDC42, RHOA) is critical to link chemoattractant receptors to cytoskeletal changes and the formation/maintenance of cell polarity⁴⁹.




Figure 1. Downstream pathways activated by chemokine signalling. Most of the downstream signals evoked by chemokine ligand/receptor binding depend on receptor dissociation from heterotrimeric G-proteins ($G\alpha\beta\gamma$) and on the GDP-GTP (guanosine diphosphate and triphosphate) exchange in the $G\alpha$ subunit. Different families of G α -proteins exist (Gs, Gi, and Gq), which can activate RHOA (RAS homologue A), PLC or PLA2 (phospholipase C or A2) and PTK (protein tyrosine kinase). Gi (shown in figure) can additionally inhibit AC (adenylyl cyclase), while Gs can stimulate this enzyme and production of cAMP (cyclic adenosine monophosphate). PLA2 controls AA (arachidonic acid) release and affects the synthesis of pro/anti-inflammatory derivatives. PTK activates the MAPK (mitogen-activated protein kinases), while RHOA evokes actin polymerisation and sustains cell migration. $G\beta\gamma$ modulates GRB2 (growth factor receptor-bound protein 2), which via activation of RAS proteins, leads to downstream effects synergistic with the MAPK. $G\beta\gamma$ also activates PI3K (phosphoinositide 3-kinase) and production of PIP3 (phosphatidylinositol-3,4,5-triphosphate). PIP3 is key to maintain cell asymmetry by mediating cortical recruitment and activation of PH (pleckstrin homology) domain-containing proteins, including small GTPases and their regulators (e.g. TIAM1/RAC1, GEFs, ROCK, CDC42). PIP3 also activates AKT and PLC. AKT controls a phosphorylation cascade while PLC leads to IP3 (inositol-1, 4, 5-triphosphate) and DAG (diacylglycerol) production. IP3 mobilises Ca^{2+} from the ER (endoplasmic reticulum) and evokes profound changes in the cell, as Ca^{2+} controls cytoskeleton dynamics and modulates a variety of targets, such as PLA2 and PKC (protein kinase C). Calmodulin and MLCK (myosin light chain kinase) control contractility of actin fibres in a Ca^{2+} -dependent manner. Additionally, microtubule elongation depends on Ca^{2+} gradients. As a consequence, lysosomes and secretion granules that are trafficked on microtubule tracks via the motor protein kinesin accumulate towards the cell periphery. Ca^{2+} also controls the activity of RABs and synaptotagmins, which are involved in the formation of the SNARE (Soluble NSF attachment protein receptor) complexes, evoking membrane fusion and exocytosis. G-protein signalling terminates with the modulation of gene expression via an array of transcription factors, including NF κ B (nuclear factor kappa B) and others (e.g. c-FOS/c-JUN, ELK1). The effects are pleiotropic and span from proliferation/apoptosis to immune defence and metabolism. Chemokines can also induce G-protein independent signalling, for example via direct activation of JAK2/STAT5 (Janus kinase 2/signal transducer and activator of transcription 5) or via β -arrestin. The latter is also involved in clathrin-dependent internalisation of chemokine ligand/receptor complex, which ultimately leads to ligand degradation and recycling of the receptor to the plasma membrane.

Chemokine sensing also activates phospholipase C (PLC)⁴⁷, which hydrolyses phosphatidylinositol-(4,5)-biphosphate to produce inositol-1,4,5-triphosphate (IP3) and diacylglycerol (DAG). IP3 induces the mobilisation of intracellular calcium stores, which in turn has effects on calmodulin and myosin light-chain kinase (MLCK). Increased intracellular calcium is a hallmark of chemoattractant stimulation as this mediates, for example, cytoskeleton redistribution and relocation of focal adhesions⁵⁰. Ca^{2+} also controls microtubule stability and their directional elongation in the direction of the cell movement. Secretion granules and lysosomes, which are trafficked in the cells via kinesin (a microtubule-associated motor protein), follow microtubule tracks and tend to accumulate at the cortex of the leading edge. Here the coordinated activity of Ca^{2+} , RABs and synaptotagmins – the main components of the SNARE (Soluble NSF attachment protein receptor) complex – will mediate fusion and exocytosis⁵¹. This process will, in turn, deliver phospholipids to the plasma membrane to sustain emission of protrusions and will determine secretion of enzymes and signalling mediators into the extracellular space, to facilitate locomotion and to transmit signals to neighbouring cells.

Another target of G-proteins is adenylyl cyclase (AC), which catalyses synthesis of cyclic adenosine monophosphate (cAMP). Some versions of $G\alpha$ can activate this signalling while others (called inhibitory $G\alpha$ or G_i) can instead negatively control this enzyme. Most chemokine receptors seem to associate with G_i complexes and therefore tend to antagonise this signalling pathway⁵². The activated G-proteins can also have a variety of other effects. These include activation of RAS/mitogen-activated protein kinase (MAPK) pathways^{53,54} and modulation of phospholipase A2 (PLA2)⁵⁵, with downstream effects as diverse as apoptosis and cell proliferation⁵⁴, immune responses^{56,57}, arachidonic acid (AA) signalling⁵⁵ and metabolic control⁵⁸ (**Figure 1**).

Chemokine receptors were originally thought to act entirely through G-protein-mediated processes. However, we now know that some chemokine receptors can also initiate alternative signalling pathways, that do not require G-protein-coupling. For example, some chemokines can directly activate the JAK-STAT (Janus kinase-signal transducer and activator of transcription) signalling⁵⁹ or β -arrestin signalling^{60,61}, with the latter being associated with clathrin-mediated ligand/receptor complex internalisation and desensitisation. There is also a class of these receptors that are completely incapable of transmitting any G-signalling, known as atypical chemokine receptors which are further discussed below. Notably, their ligand-mediated internalisation/recycling remains intact (or even enhanced), since this requires only direct β -arrestin coupling. This aberrant signalling mechanism is responsible for the role of these receptors as ligand scavengers, as the internalisation of the ligand/receptor complex mediates degradation of the cargo and recycling of the empty receptor to the plasma membrane, where new cycles of scavenging can take place^{62,63}.

Chemokine ligands are classified into the CXC, CC, XC (or C) and CX3C subfamilies, according to the arrangement of four conserved cysteine residues, which are important for maintenance of their tridimensional structure^{64,65}. In mammals as in teleost fish, the CXC and the CC families are the two largest subfamilies, whereas the CX3C and C families only contain few members (**Table 1**). In zebrafish, another fish-specific chemokine subfamily was also identified, namely CX chemokines. These CX chemokines lack one of the two conserved cysteine residues in the N-terminus but retain the third and the

Chapter 1

fourth. This distinguishes them from the XC subfamily chemokines, which only retain the second and fourth of the signature cysteine residues¹⁴.

CCR	CCL	Function	Specificity	Significance of the axis in TB
CCR1	CCL3-3L1-5-6*-7-8-9*-13-14-15-16-23	Inflammatory	M(Φ), DC, B, T, mast cells, basophils, eosinophils	-
CCR2	CCL2-7-8-11-12*-13-16	Inflammatory	M(Φ), DC, T, NK	Controversial. Recruitment of M(Φ) and DCs via CCR2 permitted better containment of high <i>Mtb</i> doses when injected intravenously. However, CCR2 KO mice were indistinguishable from wt when exposed to lower doses, most likely due to compensations.
CCR3	CCL3L1-5-7-8-11-13-14-15-24-26-28	Inflammatory	T, eosinophils	-
<u>CCR4</u>	CCL2-4-3-5-17-22	Dual	M(Φ), DC, T, NK, NKT, basophils	Controversial. Differential recruitment/priming of T cells and iNKT evoked increased susceptibility of CCR4 KO mice to <i>M. bovis</i> -BCG. However, no impairment was found in CCR4 KO animals when exposed to low <i>Mtb</i> doses.
CCR5	CCL3-3L1-4-5-8-11-13-14-16	Inflammatory	M(Φ), microglia, DC, eosinophils,	Controversial. Pathology of CCR5 KO mice was indistinguishable from that of wt, most likely due to compensation by CCR1. However, studies performed with CCL5 KO animals suggest a beneficial effect of this axis early after exposure with low infection dose.
<u>CCR6</u>	CCL20-21	Dual	DC, B, T, NKT	Impaired recruitment of iNKT and T cells in CCR6 KO mice increased susceptibility to <i>M. bovis</i> -BCG. Follow-up experiments with <i>Mtb</i> still necessary.
<u>CCR7</u>	CCL19-21	Homoeostatic	DC, B, T	Beneficial. Reverse migration of DCs to the draining lymph nodes is delayed in CCR7 KO animals, which delays activation of T cell-mediated response against <i>Mtb</i> .
CCR8	CCL1-4-16-18	Dual	DC, B, T, NK	-
<u>CCR9</u>	CCL25	Homoeostatic	T	-
<u>CCR10</u>	CCL27-28	Homoeostatic	B, T, eosinophils	-

Table 1. Mammalian chemokine axes, and their roles in tuberculosis as emerging from murine knockout (KO) studies (continued on the next page). B: B-lymphocytes; T: T-lymphocytes; M(Φ): monocyte/macrophages; N Φ : neutrophils; DC: dendritic cells, NK: natural killers; NKT: natural killer T cells. Underlined names indicate the existence of zebrafish orthologues, based on synteny and sequence alignment. * CCL6-9-12 and CXCL15 are absent in humans but present in rodents.

CXCR	CXCL	Function	Specificity	Significance of the axis in TB
<u>CXCR1</u>	CXCL6-7-8	Inflammatory	NΦ,	-
<u>CXCR2</u>	CXCL1-2-3-5-6-7-8	Inflammatory	NΦ, NK	Detrimental. <i>CXCR2</i> KO animals presented reduced lung pathology, due to contained NΦ infiltration, which limited inflammation.
<u>CXCR3</u>	CXCL4-4L1-9-10-11-13	Dual	M(Φ), microglia, T, NK	Detrimental. <i>CXCR3</i> KO mice displayed attenuated lung pathology, attributed to differential T cell priming.
<u>CXCR4</u>	CXCL12-14	Homoeostatic	M(Φ), B, T	KO not available, since embryonic lethal.
<u>CXCR5</u>	CXCL13	Homoeostatic	B, T	Beneficial. Aberrant recruitment of lymphocytes limited <i>Mtb</i> control in <i>CXCR5</i> KO mice.
CXCR6	CXCL16	Dual	B, NKT	-
-	CXCL15*	Inflammatory	NΦ	-
<u>CXCR8</u> (GPR35)	CXCL17	Homoeostatic	M(Φ), DC	-

CX3CR/XCR	CX3CL/XCL	Function	Specificity	Significance of the axis in TB
CX3CR1	CX3CL1, CCL26	Inflammatory	M(Φ), microglia, DC, T	Dispensable. Following low-dose aerosol challenge with <i>Mtb</i> , <i>CX3CR1</i> KO mice were indistinguishable from wt in terms of survival and infection burden.
<u>XCR1</u>	XCL1-2	Dual	T	-

ACKR	ACKRL	Function	Specificity	Significance of the axis in TB
DARC (ACKR1)	CCL1-2-5-7-8-11-13-14-16-17-18, CXCL1-2-3-5-6-7-8-9-10-11-13	Atypical	ligand scavenger	-
<u>D6</u> (ACKR2)	CCL2-3-3L1-4-4L1-5-7-8-11-12-13-14-17-22-23-24-26	Atypical	ligand scavenger	Beneficial. <i>D6</i> KO mice are highly susceptible to low doses of <i>Mtb</i> . The phenotype is linked to an increased number of M(Φ)s, DCs and T cells infiltrating the infected tissues.
<u>CXCR7</u> (ACKR3)	CXCL11-12	Atypical	ligand scavenger/ gradient control	KO not available, since lethal <i>in utero</i> .
<u>CCRL1</u> (ACKR4)	CCL19-21-25, CXCL13	Atypical	ligand scavenger	-
<u>CCRL2</u> (ACKR5)	CCL2-19	Atypical	ligand scavenger	-
<u>PITPNM3</u> (ACKR6)	CCL18	Atypical	ligand scavenger	-

Table 1. Mammalian chemokine axes and their implication in tuberculosis as emerging from murine knockout (KO) studies (continued from the previous page).

In mammals, CXC chemokines can be further divided into ELR and non-ELR chemokines, based on the presence of the ELR (Glu-Leu-Arg) motif upstream of their CXC motif⁶⁴. In these species, the presence of the ELR motif distinguishes neutrophil-competent chemokines (ELR-positive) from non-neutrophil-competent chemokines (ELR-negative), which attract other myeloid and lymphoid cells (monocyte/macrophages, lymphocytes, dendritic cells and natural killer cells)^{66,67} (**Table 1**). Mammalian ELR+ chemokines are CXCL1-2-3-5-6-7-8-15. These act mostly through receptors CXCR1-2. ELR- chemokines include CXCL4-9-10-11-12-13-14-16-17 and interact with receptors CXCR3-4-5-6-8. In fish, the ELR motif is not conserved and is not essential for the attraction of neutrophils by fish neutrophil-chemotactic chemokines⁶⁸.

The chemokine receptors are also classified in subgroups, namely CXCR, CCR, XCR or CX3CR, depending on the ligands that they can bind (**Table 1**)^{64,65}. A CXR family (which should bind the zebrafish-specific CX chemokines) has not been identified and the interacting partners of CXL ligands remain unknown¹⁴. In addition to classical chemokine receptors, a new group of receptors (or more precisely, interceptors) has been introduced to include non-classical chemokine receptors. This group phylogenetically belongs to the same protein family, but differs functionally, due to the inability to mediate classical GPCR signal transduction by coupling with G-proteins. This heterogeneous group of “decoy” receptors mostly act as broad spectrum ligand scavengers and are designated as atypical chemokine receptors or ACKR⁶⁹. Most atypical receptors lack or display an altered E/DRY (Glu/Asp-Arg-Tyr) motif, which is key to couple 7TM receptors with G-proteins and mediate classical signalling (**Figure 1**).

Chemokines can be functionally classified as homeostatic (or constitutive) or inflammatory according to the regulation and pattern of their expression⁶⁵. The homeostatic chemokines are generally expressed by a variety of tissues in physiological situations, while the expression of inflammatory chemokines requires inflammation/exposure to pathogens. However, this classification is not strict and many chemokines (e.g. CXCL9-10-11) exert a dual function, by controlling both physiological and pathological responses (**Table 1**).

The number of chemokines identified and characterised has expanded rapidly in the past decades, both for human and animal models⁶⁵. In humans, 25 chemokine receptors (including ACKRs, **Table 1**) and 45 chemokine ligands have been identified, while in zebrafish at least 36 putative chemokine receptors and 75 putative chemokine ligands have been mapped⁶⁵. Studies have revealed high levels of redundancy in the function and expression patterns of both chemokines and chemokine receptors⁴¹. Several receptors bind multiple chemokines and several chemokines can also bind to more than one receptor (**Table 1**). During infection/inflammation and development, a variety of chemokine ligands can be expressed at a specific site and multiple chemokine receptors can be expressed by the chemoattracted cells. Taken together, chemokine ligands and receptors can be highly synergistic, complementary or, in some cases, even antagonistic^{41,42}. Due to this complexity, it is very difficult to accurately assess to what extent the chemokine axes are really redundant. Current understanding of the roles and dynamics of each chemokine in the complexity of an *in vivo* system is also incomplete. Therefore, the evolutionary, functional and regulative forces, that permitted expansion and radiation of diverse chemokine ligand and receptor lineages, remain largely unknown⁶⁵.

Homologies between the chemokine signalling axes of human and zebrafish

While most of the ligand-receptor interacting partners have been identified in human and mammalian models, functional roles in zebrafish are only beginning to be examined⁷⁰. At transcriptional level, several zebrafish chemokines have been found to be expressed in discrete areas/organs and tissues (homeostatic) and others were found to be highly responsive to inflammation (inflammatory), like in humans^{14,71,72,73,74}. In a recent review on the translational value of zebrafish and other teleosts to study chemokine functions⁷⁰, it was reported that only one CCL chemokine axis has been currently shown to have clear *in vivo* functional homologies with the human and mammalian system. In this study, the zebrafish Ccl25a gene was found essential for normal T cell lymphotaxis to the thymus⁷³.

While the CCL ligand family is more divergent between mammals and teleosts, the CXCL family is much better conserved. For example, 11 canonical CXCRs and 20 CXCLs can be currently identified in zebrafish, versus 7 CXCRs and 17 CXCLs described in humans¹⁴. Several true homologies between zebrafish and mammals (corroborated by sequence alignments, synteny studies and functional analyses) have been demonstrated for CXC axes. As described in this thesis (**Chapter 3**), the CXCR3/CXCL9-10-11 axis exists in zebrafish. On the ligand side, a cluster of seven CXCL11-like tandem-duplicated genes have been identified. Some of these genes are infection-inducible, similar to their mammalian counterpart, and we showed that two of the CXCL11-like chemokines encoded by these infection-inducible genes signal via an orthologue of CXCR3, the zebrafish receptor Cxcr3.2 (**Chapter 3-4**). The *CXCR3* gene is also duplicated in zebrafish and two paralogues of *cxcr3.2* exist, namely *cxcr3.1* and *cxcr3.3*. In zebrafish embryos and larvae the *cxcr3.2* and *cxcr3.3* genes are expressed by macrophages and neutrophils (**Chapter 3**), while the expression of *cxcr3.1* is very low in these cells. Preliminary data suggest that *cxcr3.1* is detectable at higher levels than the other two orthologues in *lck*⁺ cells (immature T cells)⁷⁵. Therefore, it is possible that the three isoforms regulate chemotaxis of different cell types and represent sub-functionalisation of the mammalian CXCR3 that is expressed by both myeloid and lymphoid leukocytes (T cells, natural killer cells, dendritic cells, macrophages and microglial cells). However, this remains speculative, since the expression patterns remain to be further examined in adult zebrafish. The function of Cxcr3.3 has not been yet elucidated. However, as described in **Chapter 5**, *cxcr3.3* genes exist exclusively in fish and in most cases they present peculiar sequence adaptations, including disruption of the E/DRY-motif consensus, that suggest atypical functions.

Two clear orthologous CXCL12/CXCR4 axes have also been reported in zebrafish and extensive functional studies have been performed for both systems^{72,76,77,78,79} (**Chapter 6**). Both CXCL12 and CXCR4 are in fact duplicated in fish (Cxcl12a/b and Cxcr4a/b). Cxcr4a and Cxcr4b are mostly expressed in separate tissues and cell types, which corroborates the hypothesis that after gene duplication the two receptors have divided functions between the two copies⁸⁰. It appears that Cxcr4a evokes more potent responses via Cxcl12b while Cxcr4b predominantly signals via Cxcl12a, however, cross-reactivity has also been described⁸¹. Importantly, it has been demonstrated that cross-communication between the zebrafish and human ligands and receptors takes place, supporting the translational value of the zebrafish model⁴⁰. The mammalian CXCL12 and either one or both of the zebrafish counterparts, Cxcl12a/b, are chemotactic for endothelial cells, haematopoietic progenitor cells, lymphocytes and several

other types of leukocytes. In agreement with their more homoeostatic function, these axes were implicated in the development of morphogenetic/organogenetic programs, such as the formation of blood and lymphatic vessels, the migration of germ cells and of the lateral line primordium, the organisation and patterning of the nervous system and the homing of haematopoietic stem cells^{72,76,79,82,83}. The axis (mostly the Cxcr4b/Cxcl12a paralogue axis) has also been shown to play a crucial function during inflammatory responses, for example, promoting the metastatic behaviour of CXCR4-positive cancer human-xenotransplants⁴⁰. Notably in both mammalian models and in zebrafish, this axis is regulated by the activity of the atypical chemokine receptor CXCR7/ACKR3, which acts as a Cxcl12 ligand scavenger to maintain an efficient formation of ligand gradient and facilitate cell migration⁷⁶.

At least 2 *cxcl8* genes were identified and characterised in zebrafish, which are both induced by inflammation, wounding and infection. Similar to the human CXCL8 (IL8), the zebrafish paralogues are potent neutrophil chemoattracts via the Cxcr2 receptor, although in contrast to humans, expression of Cxcr1 alone does not seem to permit any significant Cxcl8-dependent recruitment^{84,85,86,87}. In mammals, CXCR2 is highly promiscuous and can respond to a range of CXC-chemokines (**Table 1**). Therefore, it is likely that multiple ligands also exist for this receptor in zebrafish. As indicated in this thesis (**Chapter 7**), the zebrafish-specific chemokine Cxcl18b also partly requires signalling via Cxcr2, and not Cxcr1, to evoke optimal recruitment of neutrophils.

Chemokine function in mycobacterial diseases

Since chemokines and chemokine receptors direct cells to specific sites within the tissues, these molecules may be highly relevant for the progression of TB and granuloma formation, for example by directly controlling the cellular trafficking at the granuloma. There have been a variety of studies aimed at characterising expression of chemokines in response to *Mtb* *in vitro* and *in vivo*. *Mtb* is a strong inducer of chemokine expression. Following *Mtb* infection of human or mice macrophages *in vitro*, these cells exhibit rapid expression of many chemokines, detectable as early as 2 hours post-infection^{88,89}. Mammalian macrophages can produce the chemokines CCL2-3-4-5-7-12, CXCL2-9-10-11, and CX3CL1 with a response that, in many cases, depends on the pathogen virulence^{90,91,92,93,94,95,96,97}. The cell wall components of mycobacteria are considered crucial for chemokine induction, as the production of several chemokines is significantly reduced following exposure to wall-deficient mycobacteria⁹⁷. Macrophages are not the only cells inducing expression of chemokines upon *Mtb* infection. Chemokine expression is also induced in cells intrinsic to tissue surrounding the infection and other attracted leukocytes^{88,98}. Expression of CCL2 and CXCL5-8-9-10-11 was detected in human or murine epithelial cells in response to virulent strains of *M. tuberculosis*^{98,99,100,101}. Circulating granulocytes from TB patients produced high levels of CCL2-3 and CXCL1-8^{102,103}, bronchoalveolar lavage of patients displayed elevated levels of CCL2-5-7-12, CXCL8-10^{90,104,105} and plasma/serum level of CCL4-11-24-26 and CXCL10 were also found increased^{106,107,108}. Two other studies that profiled gene expression of *Mtb*-infected murine and non-human-primate lungs described overall upregulation of CXCL1-2-3-5-9-10-11-13 and CCL2-3-4-5-7-8-12-19-20^{98,109,110}. Recent studies also revealed that, in TB of the central nervous system (TB meningitis), microglia (specialised macrophages residing in the brain) can also produce several chemokines, including CCL2-5 and CXCL10¹¹¹. These studies indicate that a variety of cells participate in generating chemotactic/chemokinetic

cues in response to mycobacteria (**Figure 2**). Regarding the specificity of recruited cells, the chemokines reported are known to induce mobilisation of diverse types of cells (monocyte/macrophages, neutrophils, eosinophils, T cells, natural killer cells, dendritic cells, and others) according to the expression pattern of their receptor counterparts on target cells (**Table 1, Figure 2**)^{41,42,65}.

Regulation of the expression of signals inducing chemokine ligands still requires complete characterisation, as much in TB as in other physiological and pathological contexts. Chemokines are regulated by many cytokines and infection-related products. There is evidence for example that IFN γ , TNF α and *Mtb*-components can induce expression of chemokines (**Figure 2**). Expression of chemokines by macrophages appears to be highly influenced by TNF α production, which controls CCL2-5 and CXCL9-10-11¹¹². TNF α is produced by macrophages themselves in response to *Mtb* infection and blockade of TNF α signalling can attenuate chemokine production. Notably, CXCL9-10-11, which are generally regarded as IFN γ inducible chemokines, were also found to be induced by TNF α during mycobacterial infection^{112,113}. However, IFN γ itself is also found increased in TB patients and TNF α /IFN γ may synergise to induce these and other chemokines.

While expression of chemokines is increased by *Mtb* infection, the expression of chemokine receptors remains much more stable. However, transcriptional profiling of the murine lung revealed a general upregulation of CXCR3-6 and of CCR1-5-9^{98,110}. Clinically relevant might be also the fact that monocytes increase expression of CCR5 and CXCR4 (HIV co-receptors) in response to intracellular *Mtb*, which may result in increased infectivity of HIV in cases of *Mtb*/HIV co-infections^{114,115}. In another study, dendritic cells having taken up *Mtb* were found to upregulate CCR7, which induces their reverse migration from the infection focus to the draining lymph nodes (expressing its ligands CCL19-21) and permits a faster priming of the T cell-mediated adaptive immunity^{116,117}. Similarly, upon recruitment to the lung, T cells upregulate CXCR3 and CCR5 receptors. However, this upregulation maintains them at the infected locus, since high levels of CXCR3 and CCR5 ligands are produced at the lesion¹¹⁸.

Leprosy represents another relevant mycobacterial disease, caused by the species *Mycobacterium leprae*. The disease, which essentially affects the skin and the peripheral nervous system, is still endemic in several developing countries, where it represents a relevant cause of permanent disability¹¹⁹. Leprosy manifests itself in two main polarised forms, namely lepromatous leprosy and tuberculoid leprosy. Lepromatous leprosy is characterised by exuberant bacillary replication and failure of T-helper 1 activation (which is essential to counteract the pathogen). Conversely, tuberculoid leprosy, is characterised by reduced count of bacilli, expression of T-helper 1 cytokines and is generally restricted to delimited skin lesions. Compared to TB, leprosy still remains poorly understood, for several limitations, which include the scarce cultivability in axenic conditions^{120,121,122} and the limited availability of animal models^{123,124}. However, several studies have demonstrated that expression levels of different chemokines, including CXCL8-9-10 and CCL2-3-7, are sensitive to leprosy infection^{125,126,127,128,129}. Notably, recent studies have also demonstrated that the expression of several chemokine ligands and receptors can be specifically correlated with the different manifestations of leprosy.




Figure 2. Schematic representation of main chemokine ligands and receptors implicated in the response to mycobacterial infection. Infected macrophages (MΦ) and epithelial/endothelial cells (E) respond to microbe/damage-associated molecular patterns (M/DAMPs and red arrows) by producing cytokines and several chemokines, including interferon γ (IFN γ and blue arrows), tumour necrosis factor α (TNF α and green arrows), CCL2-3-4-5-7-12, CXCL1-2-5-8-9-10-11 and CX3CL1 (orange arrows). TNF α and IFN γ can act on other cells in the surrounding microenvironment (initially, mainly monocyte/macrophages, epithelial, endothelial, stromal cells and neutrophils) and stimulate the production of more chemokines. Different types of leukocytes express different kinds of chemokine receptors (blue names) and, depending on their ligand specificity, these receptors mediate chemotaxis of leukocytes to the initial infection focus. CXCL1-2-5-8 (orange names) are mostly chemotactic towards neutrophils (NΦ), while other chemokines, such as CCL2-3-4-5-7-11, CXCL9-10-11 and CX3CL1 (violet names) are chemotactic towards monocyte/macrophages (M(Φ)), dendritic cells (DCs), T, B and/or natural killer (T, B, NK) cells. Antigen presenting activity by M(Φ) and DCs, will support activation of adaptive B and T cell-mediated immune responses. Additionally activated NK and T cells will produce high levels of inflammatory cytokines (especially IFN γ), which will sustain further stimulation of the innate immune cells and additional release of inflammatory chemokines, which in turn will support supplementary infiltration of leukocytes. The atypical chemokine receptor D6/ACKR2 (grey) is also upregulated in the cells surrounding the infection site and exerts a scavenging activity against a wide spectrum of CC-cytokines, contributing to tight control of chemokine levels. Ultimately these forces permit aggregation of immune cells at the infection focus and contribute to the formation of tubercular granulomas.

In particular, it was identified that expression of CCL17 or CCL18 are either correlated to the lepromatous (CCL18) or to the tuberculoid (CCL17) form¹³⁰, and that the expression levels of both CXCR3 and its ligand CXCL10 can be used as indicator of leprosy type 1 reaction, a severe complication of leprosy, characterised by acute activation of the cell-mediated immune response with consequent inflammation and nerve damage¹³¹.

***In vivo* analysis of chemokine axes in murine TB models**

While large amounts of data are available from *in vitro* studies, the expression and relevance of chemokines have only been characterised to a limited degree *in vivo* and the real meaning of these molecules for the pathology of TB remains elusive and speculative in several cases (**Table 1**). Using knockout murine models it was shown that CCR2 and its ligand CCL2 are essential for migration of macrophages and dendritic cells to the lungs and to control infection^{92,94}. However, the susceptibility of these mice was found to be dose-dependent, since these animals were highly susceptible to a moderate or high dose of *Mtb* administered intravenously, but did not have increased bacterial loads compared to wildtypes (wt) when exposed to low dose aerosol or intravenous infection^{132,133}. Another controversial finding came from the study of a *CCR4* knockout (KO). In a model of acute *M. bovis*-BCG pulmonary infection, these animals exhibited higher susceptibility and increased bacterial burden. However, a follow-up study showed that *CCR4*^{-/-} mice were not more susceptible to low dose of *Mtb* infection^{110,134}. *CCR6* mutants also displayed increased susceptibility to *M. bovis*¹³⁵, but studies with *Mtb* are still necessary. The function of CCR5 signalling is also controversial; an initial study in *CCR5*^{-/-} mice reported no difference in the bacterial burden evoked by *Mtb*¹³⁶ and suggested that another receptor, such as CCR1 (which can bind the same ligands of CCR5, including CCL3-5), could compensate for CCR5 loss. Another study performed with *CCL5* mutant mice with low dose aerosol infection, revealed instead an increased bacterial burden early after infection. In this study, recruitment of immune cells via CCL5 was predominantly evoked by CCR5, suggesting that compensation may occur at later time points but that CCR5 and CCR1 are not entirely redundant early after infection¹³⁷. Furthermore, in the *CCR2*^{-/-} mice studies described above, it was reported that while the number of macrophages migrating to the infection is initially lower, new cohorts of monocyte/macrophages can still be recruited at later time points. It was hypothesised that the initial increased burden in these animals evokes a higher upregulation of other chemokines in a second phase, which still sustains infiltration of immune cells and formation of granulomas^{92,94}. These examples illustrate well the difficulties that can be experienced when addressing chemokine function *in vivo*.

A murine model was also employed to study the function of CXCR3 and its ligands CXCL9-10-11 during *Mtb* infection. A beneficial effect of *CXCR3* mutation was observed, with associated decrease in number, size and density of granulomas^{110,138}. Unexpectedly, the recruitment of T cells was increased, rather than diminished in *CXCR3* mutants, and the improved resistance of *CXCR3*-deficient mice was attributed to the function of CXCR3 in T-cell priming, rather than T-cell recruitment^{138,139}. Mice deficient for either *CXCR2* or its ligand *CXCL5* had reduced lung pathology in comparison to their wt counterparts, and this phenotype was linked to a reduced accumulation of neutrophils in the mutants and to decreased detrimental neutrophil-dependent inflammation¹⁰¹. CXCL8, one of the most potent chemoattractants for neutrophils in humans, which is abundantly expressed upon *Mtb* infection, is however absent in mice¹⁴⁰, which makes it difficult to exactly address the role of CXCR2-signalling using this model. Finally, a study of *Mtb* infection in a

Chapter 1

CX3CR1^{-/-} mouse model, showed that this mutation, while affecting infiltration of different leukocyte subsets, did not lead to protection or increased susceptibility to the pathogen, indicating that this receptor is dispensable¹⁴¹.

In few cases, specific chemokine axes have been shown to play unique, non-redundant, functions *in vivo*. For example, both receptor and ligand mutants of the *CXCR5/CXCL13* pair were shown to display decreased control of *Mtb*, linked to aberrant recruitment of a specific set of *CXCR5*⁺ lymphocytes^{114,115,142}. Similarly, the function of *CCR7* in dendritic cell reverse migration to the draining lymph nodes to prime the adaptive immune response was also found to be unique^{116,117}.

Regarding the relevance of atypical chemokine receptors for the response to mycobacterial infection, a recent study on *D6/ACKR2* KO mice showed that these animals are highly susceptible to low dose aerosol exposure to *Mtb*, owing to the increased number of monocyte/macrophages, dendritic cells and T cells infiltrated in the lesioned tissues. Blockade of a wide spectrum of chemokines with a cocktail of antibodies could rescue the phenotype of *D6* deficiency, which suggests that the activity of *D6/ACKR2* as a chemokine scavenger receptor is fundamental to maintain a balanced activation of leukocyte recruitment via chemokine signalling¹⁴³.

Despite the seemingly large redundancy in function exerted by chemokines in TB suggested by *in vitro* and *in vivo* models, human studies have still shown associations between mutations of multiple chemokine ligands (*CCL2-5*, *CXCL8-10*) and TB, indicating that besides the apparent redundancy, several chemokines may have defined roles to play in the pathogenesis of TB which we still do not entirely comprehend^{144,145,146,147,148,149,150,151}.

Taken together, it has been difficult to dissect the function of chemokine signalling in TB using the mice as an *in vivo* model. This situation may reflect, at least partly, the fact that mice do not represent an ideal model for TB, since *Mtb* is not a natural murine pathogen and infection in mice does not recapitulate the primate pathology. Additionally, the use of the murine-*Mtb* infection model leads to restrictions in terms of research opportunities. *In vivo* imaging and longitudinal studies in rodents require invasive and technically challenging procedures. Furthermore, work with the mouse model imposes severe ethical restrictions, since adult animals must be sacrificed for the experiments or for intra-uterine collection of embryos/foetuses. Moreover, research prospects are severely limited by the use of the *Mtb* pathogen, which represents a serious health risk for the researchers. To circumvent all these difficulties, in this thesis we have exploited the zebrafish-*Mm* natural infection model to evaluate the role of chemokines in mycobacterial infection. Using this surrogate system, we took advantage of its attractive optical and genetic tools to obtain a more dynamic representation of how the chemokine axes calibrate the immune response to mycobacteria.

Contribution of this thesis: a brief outline

In this thesis, we applied the zebrafish (*Danio rerio*)-*Mycobacterium marinum* natural infection model to obtain novel insights into the dynamic processes underlying host-pathogen interactions and, in particular, to dissect how chemokines, a class of small chemotactic proteins, orchestrate the response of immune cell types to mycobacterial

infection. Several chemokine axes are conserved from fish to mammals and the similarities between human-*M. tuberculosis* and zebrafish-*M. marinum* pathology are remarkable. These two aspects, together with the excellent possibilities for intravital imaging provided by zebrafish transparency, have enabled us to study more dynamically chemokine-dependent processes in an *in vivo* context and allowed us to dissect how these axes are implicated in defined spatio-temporal windows of mycobacterial disease.

The work presented in this thesis has helped to elucidate the function of several chemokine axes in mycobacterial disease, and contributed to interpret the translational impact of the zebrafish model for biomedical research. This chapter (**Chapter 1**) introduces the chemokine signalling system and the current knowledge of its role in tuberculosis from *in vitro* and murine studies. **Chapter 2** elaborates on the translational value of the zebrafish model and reviews how the application of this model has advanced our understanding of host-pathogen interactions for different infectious diseases. **Chapter 3** and **Chapter 4** dissect the function of the chemokine receptor Cxcr3.2 and of its ligands Cxcl11aa and Cxcl11af during the response to mycobacterial infection, expanding on the effects of this signalling system for both direct chemotaxis of macrophages and control of their intrinsic microbicidal competence against mycobacteria. **Chapter 5** describes the generation and initial characterisation of a *cxcr3.3* mutant zebrafish line and explores the possibility of atypical receptor crosstalk at play between Cxcr3.3 and Cxcr3.2. **Chapter 6** presents the importance of the homeostatic chemokine receptor Cxcr4b in order to sustain a pro-granuloma angiogenetic programme. In **Chapter 7** we characterised the function of the chemokine ligand Cxcl8b, a fish-specific inflammatory chemokine that, similarly to Cxcl8, exerts a neutrophil-specific chemotactic activity, at least partly via Cxcr2. Finally, **Chapter 8** summarises and contextualises the findings of this thesis in relation to the current scientific literature.

Due to the high redundancy of chemokines and to the practical limitations when attempting to study tuberculosis in a murine system, the use of a zebrafish platform to address chemokine signalling during mycobacterial pathogenesis promises to greatly contribute to our understanding of the actual *in vivo* significance of these multifaceted signalling systems.

REFERENCES

-
- 1 Grunwald DJ, Eisen JS. Headwaters of the zebrafish -- emergence of a new model vertebrate. *Nat Rev Genet.* 2002 Sep;3(9):717-24.
 - 2 Streisinger G, Walker C, Dower N, Knauber D, Singer F. Production of clones of homozygous diploid zebra fish (*Brachydanio rerio*). *Nature.* 1981 May 28;291(5813):293-6.
 - 3 Amsterdam A. Insertional mutagenesis in zebrafish. *Dev Dyn.* 2003 Nov;228(3):523-34.
 - 4 Busch-Nentwich E, Kettleborough R, Dooley CM, Scahill C, Sealy I, White R, Herd C, Mehroke S, Wali N, Carruthers S, Hall A, Collins J, Gibbons R, Pusztai Z, Clark R, Stemple DL. Sanger Institute Zebrafish Mutation Project mutant data submission. ZFIN Direct Data Submission. 2013.
 - 5 Meeker ND, Trede NS. Immunology and zebrafish: spawning new models of human disease. *Dev Comp Immunol.* 2008;32(7):745-57.
 - 6 Torraca V, Cui C, Boland R, Bebelman JP, van der Sar AM, Smit MJ, Siderius M, Spaink HP, Meijer AH. The CXCR3-CXCL11 signaling axis mediates macrophage recruitment and dissemination of mycobacterial infection. *Dis Model Mech.* 2015 Mar;8(3):253-69.
 - 7 Jao LE, Wente SR, Chen W. Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system. *Proc Natl Acad Sci U S A.* 2013 Aug 20;110(34):13904-9.

- 8 Hwang WY, Fu Y, Reyon D, Maeder ML, Tsai SQ, Sander JD, Peterson RT, Yeh JR, Joung JK. Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol*. 2013 Mar;31(3):227-9.
- 9 Auer TO, Del Bene F. CRISPR/Cas9 and TALEN-mediated knock-in approaches in zebrafish. *Methods*. 2014 Sep;69(2):142-50.
- 10 Rougeot J, Zakrzewska A, Kanwal Z, Jansen HJ, Spaink HP, Meijer AH. RNA sequencing of FACS-sorted immune cell populations from zebrafish infection models to identify cell specific responses to intracellular pathogens. *Methods Mol Biol*. 2014;1197:261-74.
- 11 North TE, Goessling W, Walkley CR, Lengerke C, Kopani KR, Lord AM, Weber GJ, Bowman TV, Jang IH, Grosser T, Fitzgerald GA, Daley GQ, Orkin SH, Zon LI. Prostaglandin E2 regulates vertebrate haematopoietic stem cell homeostasis. *Nature*. 2007 Jun 21;447(7147):1007-11.
- 12 Cutler C, Multani P, Robbins D, Kim HT, Le T, Hoggatt J, Pelus LM, Despons C, Chen YB, Rezner B, Armand P, Koreth J, Glotzbecker B, Ho VT, Alyea E, Isom M, Kao G, Armant M, Silberstein L, Hu P, Soiffer RJ, Scadden DT, Ritz J, Goessling W, North TE, Mendlein J, Ballen K, Zon LI, Antin JH, Shoemaker DD. Prostaglandin-modulated umbilical cord blood hematopoietic stem cell transplantation. *Blood*. 2013 Oct 24;122(17):3074-81.
- 13 Kanwal Z, Wiegertjes GF, Veneman WJ, Meijer AH, Spaink HP. Comparative studies of Toll-like receptor signalling using zebrafish. *Dev Comp Immunol*. 2014 Sep;46(1):35-52.
- 14 Nomiya H, Hieshima K, Osada N, Kato-Unoki Y, Otsuka-Ono K, Takegawa S, Izawa T, Yoshizawa A, Kikuchi Y, Tanase S, Miura R, Kusuda J, Nakao M, Yoshie O. Extensive expansion and diversification of the chemokine gene family in zebrafish: identification of a novel chemokine subfamily CX. *BMC Genomics*. 2008 May 15;9:222.
- 15 Rytkönen KT, Akbarzadeh A, Miandare HK, Kamei H, Duan C, Leder EH, Williams TA, Nikinmaa M. Subfunctionalization of cyprinid hypoxia-inducible factors for roles in development and oxygen sensing. *Evolution*. 2013 Mar;67(3):873-82.
- 16 Carvalho R, de Sonnevile J, Stockhammer OW, Savage ND, Veneman WJ, Ottenhoff TH, Dirks RP, Meijer AH, Spaink HP. A high-throughput screen for tuberculosis progression. *PLoS One*. 2011 Feb 16;6(2):e16779.
- 17 Takaki K, Cosma CL, Troll MA, Ramakrishnan L. An in vivo platform for rapid high-throughput antitubercular drug discovery. *Cell Rep*. 2012 Jul 26;2(1):175-84.
- 18 Ordas A, Raterink RJ, Cunningham F, Jansen HJ, Wiweger MI, Jong-Raadsen S, Bos S, Bates RH, Barros D, Meijer AH, Vreeken RJ, Ballell-Pages L, Dirks RP, Hankemeier T, Spaink HP. Testing tuberculosis drug efficacy in a zebrafish high-throughput translational medicine screen. *Antimicrob Agents Chemother*. 2015 Feb;59(2):753-62.
- 19 Hisano Y, Sakuma T, Nakade S, Ohga R, Ota S, Okamoto H, Yamamoto T, Kawahara A. Precise in-frame integration of exogenous DNA mediated by CRISPR/Cas9 system in zebrafish. *Sci Rep*. 2015 Mar 5;5:8841.
- 20 Ablain J, Zon LI. Tissue-specific gene targeting using CRISPR/Cas9. *Methods Cell Biol*. 2016;135:189-202.
- 21 Mizgirev I, Revskoy S. Generation of clonal zebrafish lines and transplantable hepatic tumors. *Nat Protoc*. 2010 Mar;5(3):383-94.
- 22 Ramakrishnan L. Revisiting the role of the granuloma in tuberculosis. *Nat Rev Immunol*. 2012 Apr 20;12(5):352-66.
- 23 Global tuberculosis report 2015, World Health Organisation (WHO).
- 24 Kulchavenya E. Extrapulmonary tuberculosis: are statistical reports accurate? *Ther Adv Infect Dis*. 2014 Apr;2(2):61-70.
- 25 Prouty MG, Correa NE, Barker LP, Jagadeeswaran P, Klose KE. Zebrafish-*Mycobacterium marinum* model for mycobacterial pathogenesis. *FEMS Microbiol Lett*. 2003 Aug 29;225(2):177-82.
- 26 Swaim LE, Connolly LE, Volkman HE, Humbert O, Born DE, Ramakrishnan L. *Mycobacterium marinum* infection of adult zebrafish causes caseating granulomatous tuberculosis and is moderated by adaptive immunity. *Infect Immun*. 2006 Nov;74(11):6108-17. Erratum in: *Infect Immun*. 2007 Mar;75(3):1540.
- 27 Parikka M, Hammarén MM, Harjula SK, Halfpenny NJ, Oksanen KE, Lahtinen MJ, Pajula ET, Iivanainen A, Pesu M, Rämetsä M. *Mycobacterium marinum* causes a latent infection that can be reactivated by gamma irradiation in adult zebrafish. *PLoS Pathog*. 2012 Sep;8(9):e1002944.
- 28 Ramakrishnan L. The zebrafish guide to tuberculosis immunity and treatment. *Cold Spring Harb Symp Quant Biol*. 2013;78:179-92.
- 29 Cronan MR, Tobin DM. Fit for consumption: zebrafish as a model for tuberculosis. *Dis Model Mech*. 2014 Jul;7(7):777-84.
- 30 van Leeuwen LM, van der Sar AM, Bitter W. Animal models of tuberculosis: zebrafish. *Cold Spring Harb Perspect Med*. 2014 Nov 20;5(3):a018580.
- 31 Meijer AH. Protection and pathology in TB: learning from the zebrafish model. *Semin Immunopathol*. 2016 Mar;38(2):261-73.

- 32 Cambier CJ, Takaki KK, Larson RP, Hernandez RE, Tobin DM, Urdahl KB, Cosma CL, Ramakrishnan L. Mycobacteria manipulate macrophage recruitment through coordinated use of membrane lipids. *Nature*. 2014 Jan 9;505(7482):218-22.
- 33 Oehlers SH, Cronan MR, Scott NR, Thomas MI, Okuda KS, Walton EM, Beerman RW, Crosier PS, Tobin DM. Interception of host angiogenic signalling limits mycobacterial growth. *Nature*. 2015 Jan 29;517(7536):612-5.
- 34 Gordon RJ, McGregor AL, Connor B. Chemokines direct neural progenitor cell migration following striatal cell loss. *Mol Cell Neurosci*. 2009 Jun;41(2):219-32.
- 35 Cyster JG, Ngo VN, Ekland EH, Gunn MD, Sedgwick JD, Ansel KM. Chemokines and B-cell homing to follicles. *Curr Top Microbiol Immunol*. 1999;246:87-92; discussion 93.
- 36 Mehrad B, Keane MP, Strieter RM. Chemokines as mediators of angiogenesis. *Thromb Haemost*. 2007 May;97(5):755-62.
- 37 Perlin JR, Talbot WS. Signals on the move: chemokine receptors and organogenesis in zebrafish. *Sci STKE*. 2007 Aug 21;2007(400):pe45.
- 38 Bondar C, Araya RE, Guzman L, Rua EC, Chopita N, Chirido FG. Role of CXCR3/CXCL10 axis in immune cell recruitment into the small intestine in celiac disease. *PLoS One*. 2014 Feb 20;9(2):e89068.
- 39 Monin L, Khader SA. Chemokines in tuberculosis: the good, the bad and the ugly. *Semin Immunol*. 2014 Dec;26(6):552-8.
- 40 Tulotta C, Stefanescu C, Beletkaia E, Bussmann J, Tarbashevich K, Schmidt T, Snaar-Jagalska BE. Inhibition of signaling between human CXCR4 and zebrafish ligands by the small molecule IT1t impairs the formation of triple-negative breast cancer early metastases in a zebrafish xenograft model. *Dis Model Mech*. 2016 Feb 1;9(2):141-53.
- 41 Lira SA, Furtado GC. The biology of chemokines and their receptors. *Immunol Res*. 2012 Dec;54(1-3):111-20.
- 42 Kunkel SL, Strieter RM, Lindley IJ, Westwick J. Chemokines: new ligands, receptors and activities. *Immunol Today*. 1995 Dec;16(12):559-61.
- 43 Kolakowski LF Jr. GCRDb: a G-protein-coupled receptor database. *Receptors Channels*. 1994;2(1):1-7.
- 44 Lappano R, Maggiolini M. G protein-coupled receptors: novel targets for drug discovery in cancer. *Nat Rev Drug Discov*. 2011 Jan;10(1):47-60.
- 45 Jacobson KA. New paradigms in GPCR drug discovery. *Biochem Pharmacol*. 2015 Dec 15;98(4):541-55.
- 46 Li J, Ning Y, Hedley W, Saunders B, Chen Y, Tindill N, Hannay T, Subramaniam S. The Molecule Pages database. *Nature*. 2002 Dec 12;420(6916):716-7.
- 47 Kuang Y, Wu Y, Jiang H, Wu D. Selective G protein coupling by C-C chemokine receptors. *J Biol Chem*. 1996 Feb 23;271(8):3975-8.
- 48 Curnock AP, Logan MK, Ward SG. Chemokine signalling: pivoting around multiple phosphoinositide 3-kinases. *Immunology*. 2002 Feb;105(2):125-36.
- 49 Jones GE, Allen WE, Ridley AJ. The Rho GTPases in macrophage motility and chemotaxis. *Cell Adhes Commun*. 1998;6(2-3):237-45.
- 50 Pettit EJ, Fay FS. Cytosolic free calcium and the cytoskeleton in the control of leukocyte chemotaxis. *Physiol Rev*. 1998 Oct;78(4):949-67.
- 51 Colvin RA, Means TK, Diefenbach TJ, Moita LF, Friday RP, Sever S, Campanella GS, Abraszinski T, Manice LA, Moita C, Andrews NW, Wu D, Hacohen N, Luster AD. Synaptotagmin-mediated vesicle fusion regulates cell migration. *Nat Immunol*. 2010 Jun;11(6):495-502.
- 52 Hall DA, Beresford IJ, Browning C, Giles H. Signalling by CXC-chemokine receptors 1 and 2 expressed in CHO cells: a comparison of calcium mobilization, inhibition of adenylyl cyclase and stimulation of GTPgammaS binding induced by IL-8 and GROalpha. *Br J Pharmacol*. 1999 Feb;126(3):810-8.
- 53 Hu XM, Liu YN, Zhang HL, Cao SB, Zhang T, Chen LP, Shen W. CXCL12/CXCR4 chemokine signaling in spinal glia induces pain hypersensitivity through MAPKs-mediated neuroinflammation in bone cancer rats. *J Neurochem*. 2015 Feb;132(4):452-63.
- 54 Vlahakis SR, Villasis-Keever A, Gomez T, Vanegas M, Vlahakis N, Paya CV. G protein-coupled chemokine receptors induce both survival and apoptotic signaling pathways. *J Immunol*. 2002 Nov 15;169(10):5546-54.
- 55 Locati M, Lamorte G, Luini W, Introna M, Bernasconi S, Mantovani A, Sozzani S. Inhibition of monocyte chemotaxis to C-C chemokines by antisense oligonucleotide for cytosolic phospholipase A2. *J Biol Chem*. 1996 Mar 15;271(11):6010-6.
- 56 Sallusto F, Mackay CR, Lanzavecchia A. The role of chemokine receptors in primary, effector, and memory immune responses. *Annu Rev Immunol*. 2000;18:593-620.
- 57 Ye RD. Regulation of nuclear factor kappaB activation by G-protein-coupled receptors. *J Leukoc Biol*. 2001 Dec;70(6):839-48.
- 58 Liu B, Hassan Z, Amisten S, King AJ, Bowe JE, Huang GC, Jones PM, Persaud SJ. The novel chemokine receptor, G-protein-coupled receptor 75, is expressed by islets and is coupled to stimulation of insulin secretion and improved glucose homeostasis. *Diabetologia*. 2013 Nov;56(11):2467-76.

- 59 Mellado M, Rodríguez-Frade JM, Aragay A, del Real G, Martín AM, Vila-Coro AJ, Serrano A, Mayor F Jr, Martínez-A C. The chemokine monocyte chemotactic protein 1 triggers Janus kinase 2 activation and tyrosine phosphorylation of the CCR2B receptor. *J Immunol*. 1998 Jul 15;161(2):805-13.
- 60 Steen A, Larsen O, Thiele S, Rosenkilde MM. Biased and G-protein-independent signaling of chemokine receptors. *Front Immunol*. 2014 Jun 23;5:277.
- 61 Signoret N, Hewlett L, Wavre S, Pelchen-Matthews A, Oppermann M, Marsh M. Agonist-induced endocytosis of CC chemokine receptor 5 is clathrin dependent. *Mol Biol Cell*. 2005 Feb;16(2):902-17.
- 62 Rajagopal S, Kim J, Ahn S, Craig S, Lam CM, Gerard NP, Gerard C, Lefkowitz RJ. Beta-arrestin- but not G protein-mediated signaling by the "decoy" receptor CXCR7. *Proc Natl Acad Sci U S A*. 2010 Jan 12;107(2):628-32.
- 63 Borroni EM, Cancellieri C, Vacchini A, Benureau Y, Lagane B, Bachelier F, Arenzana-Seisdedos F, Mizuno K, Mantovani A, Bonecchi R, Locati M. β -arrestin-dependent activation of the cofilin pathway is required for the scavenging activity of the atypical chemokine receptor D6. *Sci Signal*. 2013 Apr 30;6(273):ra30.1-11, S1-3.
- 64 Bacon K, Baggiolini M, Broxmeyer H, Horuk R, Lindley I, Mantovani A, Maysushima K, Murphy P, Nomiya H, Oppenheim J, Rot A, Schall T, Tsang M, Thorpe R, Van Damme J, Wadhwa M, Yoshie O, Zlotnik A, Zoon K; IUIS/WHO Subcommittee on Chemokine Nomenclature. Chemokine/chemokine receptor nomenclature. *J Interferon Cytokine Res*. 2002 Oct;22(10):1067-8.
- 65 Nomiya H, Osada N, Yoshie O. Systematic classification of vertebrate chemokines based on conserved synteny and evolutionary history. *Genes Cells*. 2013 Jan;18(1):1-16.
- 66 Clark-Lewis I, Schumacher C, Baggiolini M, Moser B. Structure-activity relationships of interleukin-8 determined using chemically synthesized analogs. Critical role of NH2-terminal residues and evidence for uncoupling of neutrophil chemotaxis, exocytosis, and receptor binding activities. *J Biol Chem*. 1991 Dec 5;266(34):23128-34.
- 67 Clark-Lewis I, Dewald B, Geiser T, Moser B, Baggiolini M. Platelet factor 4 binds to interleukin 8 receptors and activates neutrophils when its N terminus is modified with Glu-Leu-Arg. *Proc Natl Acad Sci U S A*. 1993 Apr 15;90(8):3574-7.
- 68 Cai Z, Gao C, Zhang Y, Xing K. Functional characterization of the ELR motif in piscine ELR+ CXC-like chemokine. *Mar Biotechnol (NY)*. 2009 Jul-Aug;11(4):505-12.
- 69 Graham GJ, Locati M, Mantovani A, Rot A, Thelen M. The biochemistry and biology of the atypical chemokine receptors. *Immunol Lett*. 2012 Jul 30;145(1-2):30-8.
- 70 Bird S, Tafalla C. Teleost Chemokines and Their Receptors. *Biology (Basel)*. 2015 Nov 11;4(4):756-84.
- 71 Rotman J, van Gils W, Butler D, Spaink HP, Meijer AH. Rapid screening of innate immune gene expression in zebrafish using reverse transcription - multiplex ligation-dependent probe amplification. *BMC Res Notes*. 2011 Jun 15;4:196.
- 72 Tamplin OJ, Durand EM, Carr LA, Childs SJ, Hagedorn EJ, Li P, Yzaguirre AD, Speck NA, Zon LI. Hematopoietic stem cell arrival triggers dynamic remodeling of the perivascular niche. *Cell*. 2015 Jan 15;160(1-2):241-52.
- 73 Hess I, Boehm T. Intravital imaging of thymopoiesis reveals dynamic lympho-epithelial interactions. *Immunity*. 2012 Feb 24;36(2):298-309.
- 74 Chen LC, Chen JY, Hour AL, Shiau CY, Hui CF, Wu JL. Molecular cloning and functional analysis of zebrafish (*Danio rerio*) chemokine genes. *Comp Biochem Physiol B Biochem Mol Biol*. 2008 Dec;151(4):400-9.
- 75 Cui C. Chemokine signaling in innate immunity of zebrafish embryos. Doctoral Thesis, Leiden University. 2012 Dec 20.
- 76 Donà E, Barry JD, Valentin G, Quirin C, Khmelinskii A, Kunze A, Durdu S, Newton LR, Fernandez-Minan A, Huber W, Knop M, Gilmour D. Directional tissue migration through a self-generated chemokine gradient. *Nature*. 2013 Nov 14;503(7475):285-9.
- 77 Harrison MR, Bussmann J, Huang Y, Zhao L, Osorio A, Burns CG, Burns CE, Sucov HM, Siekmann AF, Lien CL. Chemokine-guided angiogenesis directs coronary vasculature formation in zebrafish. *Dev Cell*. 2015 May 26;33(4):442-54.
- 78 Haas P, Gilmour D. Chemokine signaling mediates self-organizing tissue migration in the zebrafish lateral line. *Dev Cell*. 2006 May;10(5):673-80.
- 79 Diotel N, Vaillant C, Gueguen MM, Mironov S, Anglade I, Servili A, Pellegrini E, Kah O. Cxcr4 and Cxcl12 expression in radial glial cells of the brain of adult zebrafish. *J Comp Neurol*. 2010 Dec 15;518(24):4855-76.
- 80 Chong SW, Emelyanov A, Gong Z, Korzh V. Expression pattern of two zebrafish genes, cxcr4a and cxcr4b. *Mech Dev*. 2001 Dec;109(2):347-54.
- 81 Chong SW, Nguyen LM, Jiang YJ, Korzh V. The chemokine Sdf-1 and its receptor Cxcr4 are required for formation of muscle in zebrafish. *BMC Dev Biol*. 2007 May 22;7:54.
- 82 Bussmann J, Wolfe SA, Siekmann AF. Arterial-venous network formation during brain vascularization involves hemodynamic regulation of chemokine signaling. *Development*. 2011 May;138(9):1717-26.

- 83 Knaut H, Werz C, Geisler R, Nüsslein-Volhard C; Tübingen 2000 Screen Consortium. A zebrafish homologue of the chemokine receptor Cxcr4 is a germ-cell guidance receptor. *Nature*. 2003 Jan 16;421(6920):279-82.
- 84 Deng Q, Sarris M, Bennin DA, Green JM, Herbomel P, Huttenlocher A. Localized bacterial infection induces systemic activation of neutrophils through Cxcr2 signaling in zebrafish. *J Leukoc Biol*. 2013 May;93(5):761-9.
- 85 Sarris M, Masson JB, Maurin D, Van der Aa LM, Boudinot P, Lortat-Jacob H, Herbomel P. Inflammatory chemokines direct and restrict leukocyte migration within live tissues as glycan-bound gradients. *Curr Biol*. 2012 Dec 18;22(24):2375-82.
- 86 de Oliveira S, Reyes-Aldasoro CC, Candel S, Renshaw SA, Mulero V, Calado A. Cxcl8 (IL-8) mediates neutrophil recruitment and behavior in the zebrafish inflammatory response. *J Immunol*. 2013 Apr 15;190(8):4349-59.
- 87 de Oliveira S, Lopez-Muñoz A, Martínez-Navarro FJ, Galindo-Villegas J, Mulero V, Calado A. Cxcl8-11 and Cxcl8-12 are required in the zebrafish defense against *Salmonella Typhimurium*. *Dev Comp Immunol*. 2015 Mar;49(1):44-8.
- 88 Rhoades ER, Cooper AM, Orme IM. Chemokine response in mice infected with *Mycobacterium tuberculosis*. *Infect Immun*. 1995 Oct;63(10):3871-7.
- 89 Thuong NT, Dunstan SJ, Chau TT, Thorsson V, Simmons CP, Quyen NT, Thwaites GE, Thi Ngoc Lan N, Hibberd M, Teo YY, Seielstad M, Aderem A, Farrar JJ, Hawn TR. Identification of tuberculosis susceptibility genes with human macrophage gene expression profiles. *PLoS Pathog*. 2008 Dec;4(12):e1000229.
- 90 Sadek MI, Sada E, Toossi Z, Schwander SK, Rich EA. Chemokines induced by infection of mononuclear phagocytes with mycobacteria and present in lung alveoli during active pulmonary tuberculosis. *Am J Respir Cell Mol Biol*. 1998 Sep;19(3):513-21.
- 91 Saukkonen JJ, Bazydlo B, Thomas M, Strieter RM, Keane J, Kornfeld H. Beta-chemokines are induced by *Mycobacterium tuberculosis* and inhibit its growth. *Infect Immun*. 2002 Apr;70(4):1684-93.
- 92 Peters W, Scott HM, Chambers HF, Flynn JL, Charo IF, Ernst JD. Chemokine receptor 2 serves an early and essential role in resistance to *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A*. 2001 Jul 3;98(14):7958-63.
- 93 Lin Y, Gong J, Zhang M, Xue W, Barnes PF. Production of monocyte chemoattractant protein 1 in tuberculosis patients. *Infect Immun*. 1998 May;66(5):2319-22.
- 94 Scott HM, Flynn JL. *Mycobacterium tuberculosis* in chemokine receptor 2-deficient mice: influence of dose on disease progression. *Infect Immun*. 2002 Nov;70(11):5946-54.
- 95 Hingley-Wilson SM, Connell D, Pollock K, Hsu T, Tchilian E, Sykes A, Grass L, Potiphar L, Bremang S, Kon OM, Jacobs WR Jr, Lalvani A. ESX1-dependent fractalkine mediates chemotaxis and *Mycobacterium tuberculosis* infection in humans. *Tuberculosis (Edinb)*. 2014 May;94(3):262-70.
- 96 Guirado E, Schlesinger LS, Kaplan G. Macrophages in tuberculosis: friend or foe. *Semin Immunopathol*. 2013 Sep;35(5):563-83.
- 97 Indrigo J, Hunter RL Jr, Actor JK. Influence of trehalose 6,6'-dimycolate (TDM) during mycobacterial infection of bone marrow macrophages. *Microbiology*. 2002 Jul;148(Pt 7):1991-8.
- 98 Kang DD, Lin Y, Moreno JR, Randall TD, Khader SA. Profiling early lung immune responses in the mouse model of tuberculosis. *PLoS One*. 2011 Jan 13;6(1):e16161.
- 99 Sauty A, Dziejman M, Taha RA, Iarossi AS, Neote K, Garcia-Zepeda EA, Hamid Q, Luster AD. The T cell-specific CXC chemokines IP-10, Mig, and I-TAC are expressed by activated human bronchial epithelial cells. *J Immunol*. 1999 Mar 15;162(6):3549-58.
- 100 Wickremasinghe MI, Thomas LH, Friedland JS. Pulmonary epithelial cells are a source of IL-8 in the response to *Mycobacterium tuberculosis*: essential role of IL-1 from infected monocytes in a NF-kappa B-dependent network. *J Immunol*. 1999 Oct 1;163(7):3936-47.
- 101 Nouailles G, Dorhoi A, Koch M, Zerrahn J, Weiner J 3rd, Faé KC, Arrey F, Kuhlmann S, Bandermann S, Loewe D, Mollenkopf HJ, Vogelzang A, Meyer-Schwesinger C, Mittrücker HW, McEwen G, Kaufmann SH. CXCL5-secreting pulmonary epithelial cells drive destructive neutrophilic inflammation in tuberculosis. *J Clin Invest*. 2014 Mar;124(3):1268-82.
- 102 Hilda JN, Narasimhan M, Das SD. Neutrophils from pulmonary tuberculosis patients show augmented levels of chemokines MIP-1 α , IL-8 and MCP-1 which further increase upon in vitro infection with mycobacterial strains. *Hum Immunol*. 2014 Aug;75(8):914-22.
- 103 Riedel DD, Kaufmann SH. Chemokine secretion by human polymorphonuclear granulocytes after stimulation with *Mycobacterium tuberculosis* and lipoarabinomannan. *Infect Immun*. 1997 Nov;65(11):4620-3.
- 104 Miotto D, Christodouloupoulos P, Olivenstein R, Taha R, Cameron L, Tscipopoulos A, Tonnel AB, Fahy O, Lafitte JJ, Luster AD, Wallaert B, Mapp CE, Hamid Q. Expression of IFN-gamma-inducible protein; monocyte chemoattractant proteins 1, 3, and 4; and eotaxin in TH1- and TH2-mediated lung diseases. *J Allergy Clin Immunol*. 2001 Apr;107(4):664-70.
- 105 Kurashima K, Mukaida N, Fujimura M, Yasui M, Nakazumi Y, Matsuda T, Matsushima K. Elevated chemokine levels in bronchoalveolar lavage fluid of tuberculosis patients. *Am J Respir Crit Care Med*. 1997 Apr;155(4):1474-7.

- 106 Xiong W, Dong H, Wang J, Zou X, Wen Q, Luo W, Liu S, He J, Cai S, Ma L. Analysis of Plasma Cytokine and Chemokine Profiles in Patients with and without Tuberculosis by Liquid Array-Based Multiplexed Immunoassays. *PLoS One*. 2016 Feb 16;11(2):e0148885.
- 107 Sharifabadi AR, Hassanshahi G, Ghalebi SR, Arababadi MK, Khorramdelazad H, Zainodini N, Shabani Z. All eotaxins CCL11, CCL24 and CCL26 are increased but to various extents in pulmonary tuberculosis patients. *Clin Lab*. 2014;60(1):93-7.
- 108 Juffermans NP, Verbon A, van Deventer SJ, van Deutekom H, Belisle JT, Ellis ME, Speelman P, van der Poll T. Elevated chemokine concentrations in sera of human immunodeficiency virus (HIV)-seropositive and HIV-seronegative patients with tuberculosis: a possible role for mycobacterial lipoarabinomannan. *Infect Immun*. 1999 Aug;67(8):4295-7.
- 109 Mehra S, Pahar B, Dutta NK, Conerly CN, Philippi-Falkenstein K, Alvarez X, Kaushal D. Transcriptional reprogramming in nonhuman primate (rhesus macaque) tuberculosis granulomas. *PLoS One*. 2010 Aug 31;5(8):e12266.
- 110 Slight SR, Khader SA. Chemokines shape the immune responses to tuberculosis. *Cytokine Growth Factor Rev*. 2013 Apr;24(2):105-13.
- 111 Rock RB, Hu S, Gekker G, Sheng WS, May B, Kapur V, Peterson PK. Mycobacterium tuberculosis-induced cytokine and chemokine expression by human microglia and astrocytes: effects of dexamethasone. *J Infect Dis*. 2005 Dec 15;192(12):2054-8.
- 112 Algood HM, Lin PL, Flynn JL. Tumor necrosis factor and chemokine interactions in the formation and maintenance of granulomas in tuberculosis. *Clin Infect Dis*. 2005 Aug 1;41 Suppl 3:S189-93.
- 113 Groom JR, Luster AD. CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol*. 2011 Feb;89(2):207-15.
- 114 Mayanja-Kizza H, Wajja A, Wu M, Peters P, Nalugwa G, Mubiru F, Aung H, Vanham G, Hirsch C, Whalen C, Ellner J, Toossi Z. Activation of beta-chemokines and CCR5 in persons infected with human immunodeficiency virus type 1 and tuberculosis. *J Infect Dis*. 2001 Jun 15;183(12):1801-4.
- 115 Juffermans NP, Paxton WA, Dekkers PE, Verbon A, de Jonge E, Speelman P, van Deventer SJ, van der Poll T. Up-regulation of HIV coreceptors CXCR4 and CCR5 on CD4(+) T cells during human endotoxemia and after stimulation with (myco)bacterial antigens: the role of cytokines. *Blood*. 2000 Oct 15;96(8):2649-54.
- 116 Bhatt K, Hickman SP, Salgame P. Cutting edge: a new approach to modeling early lung immunity in murine tuberculosis. *J Immunol*. 2004 Mar 1;172(5):2748-51.
- 117 Olmos S, Stukes S, Ernst JD. Ectopic activation of Mycobacterium tuberculosis-specific CD4+ T cells in lungs of CCR7-/- mice. *J Immunol*. 2010 Jan 15;184(2):895-901.
- 118 Campbell JD, HayGlass KT. T cell chemokine receptor expression in human Th1- and Th2-associated diseases. *Arch Immunol Ther Exp (Warsz)*. 2000;48(6):451-6.
- 119 Chaptini C, Marshman G. Leprosy: a review on elimination, reducing the disease burden, and future research. *Lepr Rev*. 2015 Dec;86(4):307-15.
- 120 Biswas SK. Cultivation of Mycobacterium leprae in artificial culture medium. *Indian J Med Sci*. 1989 Jan;43(1):5-10.
- 121 Levy L, Ji B. The mouse foot-pad technique for cultivation of Mycobacterium leprae. *Lepr Rev*. 2006 Mar;77(1):5-24. Review. Erratum in: *Lepr Rev*. 2006 Jun;77(2):170
- 122 Singh S, Eldin C, Kowalczywska M, Raoult D. Axenic culture of fastidious and intracellular bacteria. *Trends Microbiol*. 2013 Feb;21(2):92-9.
- 123 Balamayooran G, Pena M, Sharma R, Truman RW. The armadillo as an animal model and reservoir host for Mycobacterium leprae. *Clin Dermatol*. 2015 Jan-Feb;33(1):108-15.
- 124 Adams LB, Pena MT, Sharma R, Hagge DA, Schurr E, Truman RW. Insights from animal models on the immunogenetics of leprosy: a review. *Mem Inst Oswaldo Cruz*. 2012 Dec;107 Suppl 1:197-208.
- 125 Bobosha K, Tjon Kon Fat EM, van den Eeden SJ, Bekele Y, van der Ploeg-van Schip JJ, de Dood CJ, Dijkman K, Franken KL, Wilson L, Aseffa A, Spencer JS, Ottenhoff TH, Corstjens PL, Geluk A. Field-evaluation of a new lateral flow assay for detection of cellular and humoral immunity against Mycobacterium leprae. *PLoS Negl Trop Dis*. 2014 May 8;8(5):e2845.
- 126 Geluk A, van Meijgaarden KE, Wilson L, Bobosha K, van der Ploeg-van Schip JJ, van den Eeden SJ, Quinten E, Dijkman K, Franken KL, Haisma EM, Haks MC, van Hees CL, Ottenhoff TH. Longitudinal immune responses and gene expression profiles in type 1 leprosy reactions. *J Clin Immunol*. 2014 Feb;34(2):245-55.
- 127 Geluk A, van der Ploeg-van Schip JJ, van Meijgaarden KE, Commandeur S, Drijfhout JW, Benckhuijsen WE, Franken KL, Naafs B, Ottenhoff TH. Enhancing sensitivity of detection of immune responses to Mycobacterium leprae peptides in whole-blood assays. *Clin Vaccine Immunol*. 2010 Jun;17(6):993-1004.
- 128 Medeiros MF, Rodrigues MM, Vital RT, da Costa Nery JA, Sales AM, de Andrea Hacker M, Ferreira H, Chimelli L, Sarno EN, Antunes SL. CXCL10, MCP-1, and other immunologic markers involved in neural leprosy. *Appl Immunohistochem Mol Morphol*. 2015 Mar;23(3):220-9.
- 129 Guerreiro LT, Robottom-Ferreira AB, Ribeiro-Alves M, Toledo-Pinto TG, Rosa Brito T, Rosa PS, Sandoval

- FG, Jardim MR, Antunes SG, Shannon EJ, Sarno EN, Pessolani MC, Williams DL, Moraes MO. Gene expression profiling specifies chemokine, mitochondrial and lipid metabolism signatures in leprosy. *PLoS One*. 2013 Jun 14;8(6):e64748.
- 130 Berrington WR, Kunwar CB, Neupane K, van den Eeden SJ, Vary JC Jr, Peterson GJ, Wells RD, Geluk A, Hagge DA, Hawn TR. Differential dermal expression of CCL17 and CCL18 in tuberculoid and lepromatous leprosy. *PLoS Negl Trop Dis*. 2014 Nov 20;8(11):e3263.
- 131 Sharma I, Singh A, Mishra AK, Singh LC, Ramesh V, Saxena S. Is CXCL10/CXCR3 axis overexpression a better indicator of leprosy type 1 reaction than inducible nitric oxide synthase? *Indian J Med Res*. 2015 Dec;142(6):681-9.
- 132 Lu B, Rutledge BJ, Gu L, Fiorillo J, Lukacs NW, Kunkel SL, North R, Gerard C, Rollins BJ. Abnormalities in monocyte recruitment and cytokine expression in monocyte chemoattractant protein 1-deficient mice. *J Exp Med*. 1998 Feb 16;187(4):601-8.
- 133 Kipnis A, Basaraba RJ, Orme IM, Cooper AM. Role of chemokine ligand 2 in the protective response to early murine pulmonary tuberculosis. *Immunology*. 2003 Aug;109(4):547-51.
- 134 Stolberg VR, Chiu BC, Schmidt BM, Kunkel SL, Sandor M, Chensue SW. CC chemokine receptor 4 contributes to innate NK and chronic stage T helper cell recall responses during *Mycobacterium bovis* infection. *Am J Pathol*. 2011 Jan;178(1):233-44.
- 135 Stolberg VR, Chiu BC, Martin BE, Shah SA, Sandor M, Chensue SW. Cysteine-cysteinyll chemokine receptor 6 mediates invariant natural killer T cell airway recruitment and innate stage resistance during mycobacterial infection. *J Innate Immun*. 2011;3(1):99-108.
- 136 Algood HM, Chan J, Flynn JL. Chemokines and tuberculosis. *Cytokine Growth Factor Rev*. 2003 Dec;14(6):467-77.
- 137 Vesosky B, Rottinghaus EK, Stromberg P, Turner J, Beamer G. CCL5 participates in early protection against *Mycobacterium tuberculosis*. *J Leukoc Biol*. 2010 Jun;87(6):1153-65.
- 138 Seiler P, Aichele P, Bandermann S, Hauser AE, Lu B, Gerard C, Ehlers S, Mollenkopf HJ, Kaufmann SH. Early granuloma formation after aerosol *Mycobacterium tuberculosis* infection is regulated by neutrophils via CXCR3-signaling chemokines. *Eur J Immunol*. 2003 Oct;33(10):2676-86.
- 139 Chakravarty SD, Xu J, Lu B, Gerard C, Flynn J, Chan J. The chemokine receptor CXCR3 attenuates the control of chronic *Mycobacterium tuberculosis* infection in BALB/c mice. *J Immunol*. 2007 Feb 1;178(3):1723-35.
- 140 Zlotnik A, Yoshie O, Nomiya H. The chemokine and chemokine receptor superfamilies and their molecular evolution. *Genome Biol*. 2006;7(12):243.
- 141 Hall JD, Kurtz SL, Rigel NW, Gunn BM, Taft-Benz S, Morrison JP, Fong AM, Patel DD, Braunstein M, Kawula TH. The impact of chemokine receptor CX3CR1 deficiency during respiratory infections with *Mycobacterium tuberculosis* or *Francisella tularensis*. *Clin Exp Immunol*. 2009 May;156(2):278-84.
- 142 Slight SR, Rangel-Moreno J, Gopal R, Lin Y, Fallert Junecko BA, Mehra S, Selman M, Becerril-Villanueva E, Baquera-Heredia J, Pavon L, Kaushal D, Reinhart TA, Randall TD, Khader SA. CXCR5⁺ T helper cells mediate protective immunity against tuberculosis. *J Clin Invest*. 2013 Feb;123(2):712-26.
- 143 Di Liberto D, Locati M, Caccamo N, Vecchi A, Meraviglia S, Salerno A, Sireci G, Nebuloni M, Caceres N, Cardona PJ, Dieli F, Mantovani A. Role of the chemokine decoy receptor D6 in balancing inflammation, immune activation, and antimicrobial resistance in *Mycobacterium tuberculosis* infection. *J Exp Med*. 2008 Sep 1;205(9):2075-84.
- 144 Gao Q, Du Q, Zhang H, Guo C, Lu S, Deng A, Tang M, Liu S, Wang Y, Huang J, Guo Q. Monocyte chemotactic protein-1 -2518 gene polymorphism and susceptibility to spinal tuberculosis. *Arch Med Res*. 2014 Feb;45(2):183-7.
- 145 Feng WX, Flores-Villanueva PO, Mokrousov I, Wu XR, Xiao J, Jiao WW, Sun L, Miao Q, Shen C, Shen D, Liu F, Jia ZW, Shen A. CCL2-2518 (A/G) polymorphisms and tuberculosis susceptibility: a meta-analysis. *Int J Tuberc Lung Dis*. 2012 Feb;16(2):150-6.
- 146 Chu SF, Tam CM, Wong HS, Kam KM, Lau YL, Chiang AK. Association between RANTES functional polymorphisms and tuberculosis in Hong Kong Chinese. *Genes Immun*. 2007 Sep;8(6):475-9.
- 147 Sánchez-Castañón M, Baquero IC, Sánchez-Velasco P, Fariñas MC, Ausín F, Leyva-Cobián F, Ocejo-Vinyals JG. Polymorphisms in CCL5 promoter are associated with pulmonary tuberculosis in northern Spain. *Int J Tuberc Lung Dis*. 2009 Apr;13(4):480-5.
- 148 Flores-Villanueva PO, Ruiz-Morales JA, Song CH, Flores LM, Jo EK, Montaña M, Barnes PF, Selman M, Granados J. A functional promoter polymorphism in monocyte chemoattractant protein-1 is associated with increased susceptibility to pulmonary tuberculosis. *J Exp Med*. 2005 Dec 19;202(12):1649-58.
- 149 Tang NL, Fan HP, Chang KC, Ching JK, Kong KP, Yew WW, Kam KM, Leung CC, Tam CM, Blackwell J, Chan CY. Genetic association between a chemokine gene CXCL-10 (IP-10, interferon gamma inducible protein 10) and susceptibility to tuberculosis. *Clin Chim Acta*. 2009 Aug;406(1-2):98-102.

150Lindenau JD, Guimarães LS, Friedrich DC, Hurtado AM, Hill KR, Salzano FM, Hutz MH. Cytokine gene polymorphisms are associated with susceptibility to tuberculosis in an Amerindian population. *Int J Tuberc Lung Dis.* 2014 Aug;18(8):952-7.

151 Ma X, Reich RA, Wright JA, Tooker HR, Teeter LD, Musser JM, Graviss EA. Association between interleukin-8 gene alleles and human susceptibility to tuberculosis disease. *J Infect Dis.* 2003 Aug 1;188(3):349-55.