



Universiteit
Leiden
The Netherlands

Pharmacogenetics of advanced colorectal cancer treatment

Pander, J.

Citation

Pander, J. (2011, June 29). *Pharmacogenetics of advanced colorectal cancer treatment*. Retrieved from <https://hdl.handle.net/1887/17746>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/17746>

Note: To cite this publication please use the final published version (if applicable).



6

**Letter to the editor regarding:
“Explorative study to identify novel candidate
genes related to oxaliplatin efficacy and
toxicity using a DNA repair array”**

Jan Pander • Hans Gelderblom • Tahar van der Straaten • Cornelis J.A. Punt
Henk-Jan Guchelaar

British Journal of Cancer 2010 Jun 8;102(12):1791-2

Sir,

We earlier reported in this journal results from an explorative pharmacogenetic study for the efficacy of second-line treatment of oxaliplatin combined with capecitabine of advanced colorectal cancer (ACC).¹ These results were obtained using a DNA repair array (Asper Biotech, Tartu, Estonia) to identify novel single nucleotide polymorphisms (SNPs) that are associated with progression-free survival (PFS) for oxaliplatin/capecitabine combination therapy.² After correction for multiple testing for five DNA repair pathways investigated, SNPs in the genes encoding ataxia telangiectasia mutated (*ATM* rs1801516) and excision repair cross-complementing group 5 (*ERCC5* rs1047768) were significantly associated with PFS in the final multivariate analysis.

Owing to the explorative nature of the study, we concluded that confirmation was required in a separate cohort of oxaliplatin/capecitabine-treated patients. We, therefore, tested the associations of the same SNPs in the *ATM* and *ERCC5* genes with PFS in patients treated in another cohort – the CAIRO2 study. Blood samples were available of 560 patients who were treated with oxaliplatin combined with capecitabine and bevacizumab, with or without cetuximab, as first-line treatment of ACC.³ Germline DNA was isolated from peripheral white blood cells by the standard manual salting-out method. We genotyped the *ATM* and *ERCC5* polymorphisms using a Taqman 7500 (Applied Biosystems, Foster City, CA, USA) with pre-designed assays according to the manufacturer's protocol. Negative controls (water) were included. The collection of blood samples for pharmacogenetic research was approved by the local institutional review boards of all participating centers, and all patients gave written informed consent.

The genotype frequencies in the CAIRO2 patients were not significantly different from the earlier study ($P=0.38$ and $P=0.68$ for *ATM* and *ERCC5*, respectively), and were in Hardy-Weinberg equilibrium. However, the frequency of *ATM* homozygote mutants was 1.6% in the CAIRO2 patients vs 4.4% in patients in the earlier study.

The results for the associations with PFS are shown in table 1. As opposed to our initial observation, the *ATM* and *ERCC5* polymorphisms were not significantly associated with PFS in the CAIRO2 patients.

Several reasons could underlie the lack of replication of association. First, our initial results¹ may have been false positive findings. Even though we had corrected for multiple testing, this approach may have been ineffective to correct for false positives. On the other hand, the frequency of *ATM* homozygote mutant patients in the CAIRO2 was lower than in the earlier study, which could have impacted the power to detect the association. However, the HR for PFS was 4.25 (95%CI 1.45 to 12.44; homozygote mutants vs wild-type) in our initial study, whereas it was 0.90 (95%CI, 0.37 to 2.18) in the CAIRO2 patients, indicating lack of association regardless of genotype frequency. Second, our initial findings were derived from patients receiving second-line therapy of oxaliplatin combined with capecitabine, while CAIRO2 concerns data from first-line

Table 1 Associations of *ATM* (rs1801516) and *ERCC5* (rs1047768) polymorphisms with PFS

	n	median PFS in months (95%CI)	Univariate HR (95%CI) [#]	P [#]	Multivariate HR (95%CI) ^{**}	P ^{**}
ATM rs1801516						
Wild-type	371	9.1 (8.3-10.4)	1	-	1	-
Heterozygote	127	12.4 (9.6-13.5)	0.88 (0.70-1.09)	.245	0.93 (0.75-1.17)	.543
Homozygote mutant	8	11.8 (7.2-∞) [†]	0.61 (0.27-1.36)	.225	0.94 (0.42-2.12)	.881
ERCC5 rs1047768						
Wild-type	180	10.6 (9.1-12.5)	1	-	1	-
Heterozygote	267	9.2 (8.2-10.6)	1.13 (0.93-1.39)	.227	1.15 (0.93-1.42)	.194
Homozygote mutant	77	10.1 (8.5-12.2)	0.96 (0.72-1.29)	.797	0.94 (0.69-1.28)	.689

[#] Hazard ratios (HR), 95% confidence intervals (95%CI) and P-values computed using a Cox proportional hazards model with the wild-type as reference

^{*} Covariates included in the multivariate model: age, gender, serum LDH (normal vs above normal) and treatment arm (oxaliplatin, capecitabine and bevacizumab vs oxaliplatin, capecitabine, bevacizumab and cetuximab)

[†] The upper limit of the 95%CI for PFS of the ATM homozygote mutants could not be estimated because of the low number of patients

therapy with the addition of bevacizumab and cetuximab also. We also recently also reported an opposite association of the *FCGR3A* Phe158Val polymorphism with PFS for cetuximab in the first-line setting for ACC compared with results from third-line settings.⁴ As the DNA repair array should theoretically be applicable to any platinum-containing regimen, this explanation is less likely for the present finding.

Finally, it is possible that the addition of cetuximab could have negatively influenced the efficacy of oxaliplatin in the cetuximab-arm in the CAIRO2 study^{5,6}, which may have obscured the associations when both treatment arms were combined for analysis. However, the outcome of our analysis did not change when we restricted this to patients treated without cetuximab in the CAIRO2 study (data not shown).

We, therefore, conclude that the *ATM* and *ERCC5* SNPs have no relevant impact on the PFS of oxaliplatin-based therapy for ACC. The negative result of this study underlines the importance of validating and reporting the findings from retrospective explorative studies.⁷

References

1. Kweekel DM, Antonini NF, Nortier JW, *et al.* Explorative study to identify novel candidate genes related to oxaliplatin efficacy and toxicity using a DNA repair array. *Br J Cancer* 2009;101:357-62.
2. Koopman M, Antonini NF, Douma J, *et al.* Sequential versus combination chemotherapy with capecitabine, irinotecan, and oxaliplatin in advanced colorectal cancer (CAIRO): a phase III randomised controlled trial. *Lancet* 2007;370:135-42.
3. Tol J, Koopman M, Cats A, *et al.* Chemotherapy, bevacizumab, and cetuximab in metastatic colorectal cancer. *N Engl J Med* 2009;360:563-72.
4. Pander J, Gelderblom H, Antonini NF, Tol J, van Krieken JH, van der Straaten T, *et al.* Correlation of *FCGR3A* and *EGFR* germline polymorphisms with the efficacy of cetuximab in *KRAS* wild-type metastatic colorectal cancer. *Eur J Cancer*. In press 2010.
5. Dahan L, Sadok A, Formento JL, Seitz JF, Kovacic H. Modulation of cellular redox state underlies antagonism between oxaliplatin and cetuximab in human colorectal cancer cell lines. *Br J Pharmacol* 2009;158:610-20.
6. Punt CJ, Tol J. More is less -- combining targeted therapies in metastatic colorectal cancer. *Nat Rev Clin Oncol* 2009;6:731-3.
7. Koopman M, Venderbosch S, Nagtegaal ID, van Krieken JH, Punt CJ. A review on the use of molecular markers of cytotoxic therapy for colorectal cancer, what have we learned? *Eur J Cancer* 2009;45:1935-49.