



Universiteit
Leiden
The Netherlands

The structure of flower visitation webs: how morphology and abundance affect interaction patterns between flowers and flower visitors

Stang, M.

Citation

Stang, M. (2007, October 30). *The structure of flower visitation webs: how morphology and abundance affect interaction patterns between flowers and flower visitors*. Retrieved from <https://hdl.handle.net/1887/12411>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12411>

Note: To cite this publication please use the final published version (if applicable).

Publications

- Schick, B. & Stang, M. (2000) Zur Blütenökologie und Blühphänologie silbergrasreicher, lockerer Sandrasen und ausgewählter Kontaktgesellschaften. *Ber. Naturw. Ges. Bayreuth* 24: 299-301.
- Stang, M. (2002) Grote verschillen op kleine schaal: bloembioëlogie en soortendiversiteit in een vochtige duinvallei. *Holland's Duinen* 40: 3-30.
- Stang, M., Klinkhamer, P.G.L. & van der Meijden, E. (2006) Size constraints and flower abundance determine the number of interactions in a plant-flower visitor web. *Oikos* 112: 111-121.
- Stang, M. (2006) Book review: Waser, N.M. & Ollerton, J. (editors) Plant-pollinator interactions: from specialization to generalization. *Entomologische Berichten* 66: 134-135.
- Stang, M., Klinkhamer, P.G.L. & van der Meijden, E. (2007) Asymmetric specialization and extinction risk in plant-flower visitor webs: a matter of morphology or abundance? *Oecologia* 151: 442-453.

