



Universiteit
Leiden
The Netherlands

Insulin and cellular stress induced glucose uptake in 3T3-L1 adipocytes

Bazuine, M.

Citation

Bazuine, M. (2005, March 10). *Insulin and cellular stress induced glucose uptake in 3T3-L1 adipocytes*. Retrieved from <https://hdl.handle.net/1887/2709>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/2709>

Note: To cite this publication please use the final published version (if applicable).

Chapter II

An introduction to 3T3-L1 adipocytes.

The adipocyte is a remarkable cell type in several aspects. For years the adipocyte has been viewed as a rather passive cell, simply a deposit site of excess energy in the form of lipids [1;2]. However, work with tissue-specific knock-out mice, the complex phenotype of patients with altered adipocyte function and the description of a range of proteins secreted by these cells, have established the adipocyte as a major regulator of whole body energy-homeostasis, influencing metabolic settings in key organs such as muscle, liver and brain [3-5]. Furthermore, the tight connection between adipocyte-mediated vascular remodelling and several types of cancer also identify adipose tissue as an important endocrine organ [6-8]. Aside from its endocrine role, the adipocyte serves to protect other organs from the deleterious effects of excessive intracellular triglyceride storage [9-11]. Thus, although adipose tissue accounts for only ~10% of whole body glucose uptake, an adipose tissue specific GLUT4 knock-out mouse displays glucose intolerance caused by a secondary insulin-resistance in muscle- and liver-cells [12].

Main effectors in this cross-talk are the “adipokines” $\text{TNF}\alpha$, adiponectin and leptin [13-15]. Increases in levels of $\text{TNF}\alpha$, as seen in the obese state, are associated with a deleterious impact on insulin-sensitivity in adipocytes, muscle and liver (Fig. 1). Conversely, adiponectin has a positive effect on insulin-sensitivity by stimulating fatty acid oxidation through the activation of AMPK and $\text{PPAR}\gamma$ [16-19]. The central role of adiponectin is illustrated by the adipocyte-specific insulin-receptor knock-out mouse. Although these adipocytes are no longer capable of insulin-induced glucose uptake, blood glucose levels are normal, due to an elevation in levels of adiponectin in these mice [20;21].

Another adipokine, acting in conjunction with adiponectin is the satiety hormone, leptin. This hormone regulates food intake through its effects on the hypothalamus [22], and mediates metabolic effects on peripheral tissues [23;24]. Adipocyte selective reduction of leptin receptors has profound effects on the regulation of metabolic genes, characterising an autoendocrine-loop in these cells [25]. Other functions of leptin involve regulation of AMPK, leading to fatty acid oxidation [26-28], the lipogenic transcription factor SREBP-1c [26] and PGC-1 α , a powerful inducer of mitochondrial biogenesis (Fig. 1)[29;30]. The involvement of adipokines in metabolic homeostasis is further illustrated by the occurrence of insulin resistance associated with lipodystrophy.

Fig. 1




Fig. 1 A schematic overview of a 3T3-L1 adipocyte : cellular organelles, main vesicle pathways and adipokine signalling.

A fully mature adipocyte is an endocrine cell involved in regulating whole body lipid and glucose homeostasis through the secretion of both stimulatory (adiponectin, leptin) and inhibitory (TNF- α , glucocorticoids (GC) and Free Fatty Acids) adipokines. Aside from regulating metabolic settings in target tissues the adipocyte is also tightly involved in adipogenesis through TNF- α , resistin, IGF-1 and GC-signalling. Autocrine factors derived from the pre-adipocyte involved in regulating differentiation are MCSF, TGF- β and Wnt10b.

Characteristic cell-components of the fully mature adipocyte are the caveolae and the lipid droplets, which are derived from the Endoplasmic Reticulum (ER). The Low Density Microsomal fraction (LDM) consists of several vesicular components involved in cellular trafficking. Main trafficking routes contributing to the LDM are endocytotic : 1. recycling of GLUT4 from the plasma-membrane, 2. clathrin-coated pits involved in recycling of receptors, and exocytotic : A. translocation of insulin-responsive GLUT4 Storage Vesicles (GSV) towards the plasma membrane, B. direct endosomal shuttling of GLUT4 containing vesicles (either from Golgi stacks or from the endosomal Tubulo-Vesicular Sorting (TVS) compartment), C. shuttling of GLUT4 containing vesicles towards the TVS and translocation of synthesised adipokines from either the ER or Golgi.


In mice models of lipodystrophy, injections with adiponectin and leptin ameliorate insulin-resistance accompanied by clearance of triglycerides in muscle and liver [26;31;32].

The 3T3-L1 adipogenic cell-line was established thirty years ago when Green and Meuth noted a high tendency in clones of Swiss 3T3 fibroblasts to undergo spontaneous adipogenic conversion [33-35].


Though immortalised 3T3-L1 cells are not transformed as is evidenced by their contact-inhibition (Fig. 2). At this stage cellular changes in the postmitotic adipoblasts are readily apparent with the cell flattening out, the nucleoli becoming visible (see Fig. 2B) and at a molecular level, the upregulation of the growth-arrest associated gene 2 [36]. Overriding contact-inhibition results in a fully transformed phenotype and loss of the ability to differentiate [37]. When fully arrested, cells are challenged with a potent adipogenic cocktail consisting of insulin, IBMX and dexamethasone (Table I)[38;39]. Whereby the phosphodiesterase-inhibitor IBMX can be replaced by PPAR γ agonists [40]. At this stage a number of crucial events take place : The medium becomes viscoelastic due to the excretion of highly crosslinked hyaluronic acid and the induction of metalloproteinases indicating an important outside-in signalling contribution [41-44]. The cells round up without losing the filipoda-connections with which they are linked to one-another (Fig. 2C and D). At this stage profound cell-morphological differences between lots of FCS become readily apparent, initiating the discrepancy in adipocytes differentiated under different batches of FCS (see Fig. 3A). Subsequently the cells undergo 2-4 rounds of clonal expansion and arrest in G1, whereas many other cells simply round up and enter apoptosis. Components of the p53-signalling pathway : Mdm-2, p21 and its family member p27 are tightly regulated at this stage [45-48]. The pocket-proteins, pRb, p130 and p107 are involved in regulating adipogenesis too : After a distinct switch to p107 during the clonal expansion stage the re-emergence of p130 as the main E2F-binding protein marks the final commitment of the cell to enter the G0 state (see Fig. 4)[49-52].

Remarkably, when these contact-inhibited cells are passaged into a new culture p130 will not re-emerge again which prevents 3T3-L1 cells from entering the differentiation programme a second time. The pRb protein meanwhile, regulates C/EBP β activity and drives differentiation towards white over brown adipose tissue [53-56]. Important components of the mitogenic response are the MAPK family members. Illustrating this in PC12-cells, studies of MAPK activation in response to either EGF or NGF demonstrated that the determination between mitogenesis or differentiation is highly dependent on the kinetics of MAPK signalling [57;58].


Fig. 2A




B




C




D




E



F



G



H




Fig. 2 Nomarski-photographs of differentiating adipocytes.

Panel A. growing 3T3-L1 fibroblasts, B. fibroblasts, flattened out at the growth arrested stage, C. and D. pre-adipocytes in Diff. I with the extended filipodic connections and their cytoplasmic components shrunk to barely more than the nucleus. E. early in Diff. II, flattened out cells and the start of lipid droplet formation. F. and G. maturation of the lipid droplets. H. fully mature 3T3-L1 adipocytes. Size (in micrometer) is indicated by a white bar in the photographs.

Fig. 3A effect of FCS on adipocyte differentiation




Fig. 3B development of glucose uptake during adipogenesis




Fig. 3 Analysis of insulin-induced glucose uptake.

Panel A. 3T3-L1 adipocytes differentiated using several batches of Foetal Calf Serum (FCS) demonstrating profound differences in basal levels of glucose uptake (white bars) and insulin-stimulated glucose uptake (black bars). Lot A and F, and lot D and E were obtained from the same supplier. Panel B. Development of insulin-responsiveness during adipogenesis. 3T3-L1 fibroblasts (confluent stage) have a slight response to insulin. The increase in the Dif. I response is mediated by a stress-induced increase in GLUT1 synthesis. During Diff. II the insulin responsive GLUT4 and associated vesicular compartments generate a profound increase in insulin-induced glucose uptake capacity, though at this stage the differentiating adipocytes are insulin-resistant as can be seen in the maturation stage. After completion of the maturation stage, the now fully mature adipocyte has overcome its initial insulin-resistance and downmodulated GLUT1-mediated glucose uptake leading to the profound response in insulin-induced glucose uptake over characteristically low basal levels observed.

During the initial stages of adipogenesis the induction of MAPK family members ERK-1/2 leads to the induction of PPAR γ and C/EBP α [59-63]. However, after this initial stage ERK signalling is terminated. Prolonged activation, such as induced by EGF-signalling, inhibits adipocyte differentiation through the inhibition of crucial adipogenic transcription factors (Fig. 4)[64-66]. Meanwhile p38 MAPK induces activation of C/EBP β [67-69], though similar to ERK-1/2, prolonged activation inhibits adipogenesis through the activity of CHOP [70-72]. Another key transcription factor in adipogenesis is CREB [73], which is crucial in preventing apoptosis through its inhibition of several pro-apoptotic genes such as ICE and by stimulating PKB expression [74]. Subsequently, the downregulation of pre-adipocyte factor-1 and the induction of C/EBP β and $-\delta$ induces the upregulation of PPAR γ and C/EBP α (Fig. 4)[40;75-79]. These latter two regulate the late-stage genes in adipogenesis, such as GLUT4, aP2 and adiponectin. Simultaneously the characteristic insulin-responsive microsomal-vesicular GLUT4 storage compartment is formed (Fig. 3B)[80]. To be precise, C/EBP α is not required for the generation of an “adipocyte” as such, but is crucial for conferring proper insulin-responsiveness on the cell. Thus in a C/EBP α knock-out mouse adipocytes are incapable of lipid accumulation [81-85]. On the other hand, an adipose-specific PPAR γ knock-out mice displays adipocyte hypocellularity and loss of leptin and adiponectin [86-88]. The insulin present in the cocktail induces the activation of PI-3’ kinase through the IGF I Receptor [89-91], regulating the FKHR-transcription factors, C/EBP α and SREBP1 (Fig. 4)[92-95]. The lipid- and cholesterol-metabolism genes regulated by SREBP1 mediate the synthesis of endogenous ligands for PPAR γ [96;97], illustrating autocrine signalling loops involved in adipogenesis (Fig. 1). Potent adipogenesis stimulating factors are Macrophage Colony-Stimulating Factor (MCSF), Insulin-like Growth Factor-1 (IGF-I) and Glucocorticoids (GC)[98-101]. The latter are not generated by the adipocyte as such. Rather, both primary pre-adipocytes and fully mature adipocytes express 11 β -hydroxysteroid dehydrogenase 1, which catalyses the conversion of inactive corticosterone to active cortisol (a glucocorticoid)[102-104]. Conversely, the aforementioned TNF α , resistin, Transforming Growth Factor- β (TGF β) and Wnt10b-signalling maintains adipocytes in an undifferentiated form [105-109]. Matter of factly, the Wnt-signalling components β -catenin and GSK-3 β are extensively downregulated during the first days of differentiation [75;110;111]. With the onset of C/EBP α and PPAR γ the pre-adipocyte matures as is visible by the formation of lipid droplets in the perinuclear region (see

Fig. 2F and 3B). These droplets are derived from the endoplasmic reticulum and covered by the adipocyte-specific perilipins (Fig. 1)[112;113]. PKA-mediated perilipin phosphorylation induces a conformational change of the perilipins allowing access to Hormone Sensitive Lipase and induces translocation of HSL towards the lipid-droplet [114-116]. PKA is acutely stimulated by lipolytic-hormones explaining the large cellular effects of these hormones on adipocytes [117;118]. Conversely, insulin inhibits lipolysis by activating phosphodiesterase-3, which leads to a loss of PKA activity [119]. Furthermore, insulin also induces the formation of an inhibitory complex between HSL and lipotransin [120]. Consequently, the presence of insulin in the Diff. II medium allows the lipid droplets to coalesce and expand until only a small number of large droplets is left, taking up roughly 70% of the cell-volume (Fig. 2G and H). Recent analysis of the protein profile found associated with these lipid droplets suggests that it is an important signalling compartment [121]. This is illustrated by the observation that when perilipins are ablated in knock-out mice, the mice become resistant to diet-induced obesity. Microarray analysis of these mice demonstrates a coordinated upregulation of genes involved in beta-oxidation, the Krebs cycle and the electron transport chain concomitant with a downregulation of genes involved in lipogenesis [122]. During adipogenesis cellular levels of mitochondria also increase, accompanied by qualitative changes in the mitochondrial composition (Fig. 1)[123;124]. In contrast to many continuous cell-lines, the 3T3-L1 adipocyte employs oxidative phosphorylation as a source of ATP [125]. Intriguingly, in response to insulin adipocytes also activate fatty acid oxidation in the mitochondria, even though the net effect of insulin is lipogenesis. Though this ‘futile cycle’ may seem a waste of energy, this cycle generates body heat and intermediates needed for the synthesis of other biochemical compounds [126].

Another cell-morphological feature of adipocytes is the presence of caveolae in the plasma-membrane (Fig. 1). According to the lipid-ordering hypothesis, membranes co-exist in two predominant forms : the liquid-disordered state, composed of phospholipids with relatively rapid lateral diffusion and the lipid-rafts, which are high in cholesterol- and sphingolipid-content resulting in a more rigid and confining environment [127;128].

Fig. 4




Fig. 4 Signalling pathways and stages involved in 3T3-L1 adipogenesis.

Mitotic cell stages involved in clonal expansion are indicated by their respective phases (Gap₁, Synthesis, Gap₂ and Mitosis), with the Restriction point involved in growth arrest and switch to the Gap₀ differentiation pathway. This stage is under control of the pocket proteins p107 and p130, IGF-I signalling (PI-3'kinase and PKB) and MAPK-signalling (ERK-1,-2 and p38). Apoptosis-induced cell loss occurs throughout the differentiation process, but is indicated in this picture as an alternative side-route of the cell-cycle.

Entry of the G₀ marks the entry of the commitment-stage dominated by C/EBPβ and -δ. From this stage onwards, continued MAPK- or GSK-3β signalling at this stage inhibits adipogenesis. C/EBPβ and -δ induce the main adipogenic transcription factors C/EBPα and PPARγ in conjunction with autocrine signalling. C/EBPα and PPARγ are interlocked in positive autoregulatory loops and mediate transcription of adipocyte-specific genes leading to the formation of a fully mature insulin-responsive 3T3-L1 adipocyte.

Caveolae are a specialised lipid raft characterised by the structural protein caveolin-1 forming the neck of these invaginations, thereby restricting random diffusion of the caveolar constituents [129-131]. At the plasma-membrane they form 50-100 nm omega-shaped invaginations morphologically distinct from clathrin coated pits [132;133]. In adipocytes a higher order organisation of the caveolae in “rosetta”-structures exists, though the precise reason for this clustering of caveolae remains unclear [134;135]. The two other members of the caveolin-family, Cav-2 and -3, also target exclusively to caveolae [136;137]. Whereas Cav-1 and -2 are coexpressed [138;139], Cav-3 expression is limited to muscle cells [136]. During adipogenesis, caveolae increase dramatically in number concomitant with an increase in caveolin-expression [140;141]. However, Cav-1 knock-out mice display a mild phenotype, such as exercise intolerance and decreased vascular tone, but no overt diabetes [142;143]. And treatment of adipocytes with the cholesterol chelating compounds nystatin and filipin has no effect on insulin-stimulated glucose uptake [144]. Although treatment with the more potent agent methyl- β -cyclodextrin inhibits IRS-1 activation, a total depletion of membrane-cholesterol also affects the organisation of the actin-cytoskeleton [145;146]. Yet, a direct interaction between the insulin receptor and caveolin is required for stabilisation of the insulin receptor [147-149]. And indeed, Cav-1 knock-out mice display a pronounced loss of the number of insulin receptors [150]. Furthermore, Cav-1 knock-out mice are lean, resistant to diet-induced obesity and display adipocyte abnormalities with attenuated serum leptin and adiponectin levels and loss of lipid homeostasis [151]. At face value, these mice resemble an adipocyte specific insulin-receptor knock-out (FIRKO) mouse [21]. There are some substantial differences though, such as a decrease in brown fat mass, an increase in plasma leptin and adiponectin and consequently a reduction in serum triglyceride levels in FIRKO mice with the opposite occurring in Cav-1 null mice. This is due to additional functions of the caveolae, such as its involvement in lipid homeostasis and signalling [152-154].

One of the hallmarks of a fully differentiated 3T3-L1 adipocyte is its marked insulin-induced glucose uptake, mediated by GLUT4 (Fig. 3B)[80;155]. In unstimulated cells GLUT4 is mainly localised in several intracellular vesicular compartments distinct from those employed by adipokines, demonstrating adipocytes maintain several insulin-responsive membrane compartments [156-159]. Among the intracellular structures harbouring GLUT4 are : a tubulo-vesicular endosomal recycling compartment [160-163], AP1/clathrin coated vesicles budding from either the TGN or endosomes, AP2 coated vesicles budding from the plasma-membrane, and a distinct population of GLUT4 Storage Vesicles (GSV)

harbouring a preponderance of GLUT4 and excluding general endosomal markers (Fig. 1)[161;164-167]. Of these LDM-vesicles (as they are collectively known) especially the GSV translocate rapidly towards the plasma-membrane in a PI-3'kinase dependent manner. However, endosomal ablation also causes a partial block of insulin-stimulated GLUT4 translocation, illustrating an direct involvement of the endosomal compartment as well [164;165;168]. This endosomal pathway is involved in GLUT4 translocation induced by cellular stress, exercise and GTP γ S [168-170].

With respect to the cytoskeleton in support of these structures, during adipogenesis the fibroblastic "stress-like" F-actin filaments disappear and are replaced by a cortical F-actin structure accompanied by a rearrangement of the cytoskeleton structures involved in GLUT4 translocation [171-176]. Furthermore, a novel type of actin filament, the so-called cav-actin (caveolae associated F-actin) originates in the cell, associated with the aforementioned rosetta-structures [177]. Recent data show either actin stabilising, or actin disrupting pharmacological agents severely inhibit insulin-induced glucose uptake suggesting the cytoskeleton is actively involved in regulating GLUT4 translocation, rather than acting passively as a barrier or a molecular railroad [178-187]. In conclusion, the process of 3T3-L1 adipogenesis highlights the complex molecular rearrangements implemented in a terminally differentiating cell. The re-routing of MAPK-signalling pathways, closing down of Wnt-signalling and enabling CAP-signalling occurs in intimate association with cell-morphological alterations such as the formation of caveolae, the cortical actin structure and insulin-responsive GLUT4 storage vesicles. In the fully differentiated adipocyte, a complex signalling interplay exists between these cellular structures embedding the insulin-signalling pathway and the secreted adipokines.

Table I Experimental set-up of 3T3-L1 adipogenesis

day	medium	comments
1	normal	Normal adipocyte-culturing medium consists of DMEM with 10% FCS. The FCS serum deployed throughout the procedure must have been tested for its adipogenic potential (see also Fig. 3A). Routinely cells are set up 1:20, though as high as 1:100 can be maintained.
4	normal	Usually cells are now roughly 70% confluent and have to be passaged into a new culture to prevent contact-inhibition. Up till passage 8 can be used, thereafter the cells rapidly lose adipogenic potential through the consequent “selection” of the fastest growing (transformed) cells with each passage.
7	normal	
10	normal	Usually the cells are now fully confluent and growth arrested (Fig. 2B). The cells are left in their contact-inhibited state for at least two days.
12	Diff. I	Differentiation I medium consists of 1.6 μ M insulin, 0.5 mM IBMX, 0.25 μ M dexamethasone and 10% FCS. The following day cells show their characteristically “stressed” appearance as depicted in Fig. 2C and D.
15	Diff. II	Differentiation II medium consists of 1.6 μ M insulin and 10% FCS. Addition of this medium should be applied with care as the stressed cells are but loosely attached at this stage. The following day cells show their “relaxed” appearance as depicted in Fig. 2E.
18	Diff. II	A second treatment with insulin. At these stages the medium becomes highly viscous and acidified, making it sometimes prudent to refresh the medium an additional time in between. Cells are as depicted in Fig. 2F, by eye the plate looks clustered-opaque due to the presence of lipid droplets in the cells.
21	normal	The cells need time to recover from their initial insulin-resistance, as can be observed in Fig. 3B.
23	normal	Due to the fact that adipocytes are metabolically more active, leading to medium acidification, and excrete (amongst others) TNF α , the medium has to be replenished more regularly than in their fibroblastic stage.
25	normal	
27	normal	At this moment the cells are fully mature (see Fig. 2G and H) and highly insulin-responsive (see Fig. 3B). Roughly 95% of the cells will have been converted into mature, lipid laden adipocytes.
30	normal	
33	end culture	From the start adipocytes are lost due to apoptosis and cell death. As the adipocytes are terminally differentiated, lost cells are not replenished. Non-converted (fibroblastic) cells however keep on dividing and will occupy any open place available. Furthermore, at this stage adipocytes are rapidly becoming insulin-resistance as a consequence of their secreted adipokines.

References

- [1] Fruhbeck,G., Gomez-Ambrosi,J., Muruzabal,F.J., & Burrell,M.A. (2001) The adipocyte: a model for integration of endocrine and metabolic signaling in energy metabolism regulation. *Am. J. Physiol Endocrinol. Metab*, **280**, E827-E847.
- [2] Flier,J.S. (1995) The adipocyte: storage depot or node on the energy information superhighway? *Cell*, **80**, 15-18.
- [3] Fortuno,A., Rodriguez,A., Gomez-Ambrosi,J., Fruhbeck,G., & Diez,J. (2003) Adipose tissue as an endocrine organ: role of leptin and adiponectin in the pathogenesis of cardiovascular diseases. *J. Physiol Biochem.*, **59**, 51-60.
- [4] Mora,S. & Pessin,J.E. (2002) An adipocentric view of signaling and intracellular trafficking. *Diabetes Metab Res. Rev.*, **18**, 345-356.
- [5] Trayhurn,P. & Beattie,J.H. (2001) Physiological role of adipose tissue: white adipose tissue as an endocrine and secretory organ. *Proc. Nutr. Soc.*, **60**, 329-339.
- [6] Kratchmarova,I., Kalume,D.E., Blagoev,B., Scherer,P.E., Podtelejnikov,A.V., Molina,H., Bickel,P.E., Andersen,J.S., Fernandez,M.M., Bunkenborg,J., Roepstorff,P., Kristiansen,K., Lodish,H.F., Mann,M., & Pandey,A. (2002) A proteomic approach for identification of secreted proteins during the differentiation of 3T3-L1 preadipocytes to adipocytes. *Mol Cell Proteomics.*, **1**, 213-222.
- [7] Fukumura,D., Ushiyama,A., Duda,D.G., Xu,L., Tam,J., Krishna,V., Chatterjee,K., Garkavtsev,I., & Jain,R.K. (2003) Paracrine regulation of angiogenesis and adipocyte differentiation during in vivo adipogenesis. *Circ. Res.*, **93**, e88-e97.
- [8] Wiseman,B.S. & Werb,Z. (2002) Stromal effects on mammary gland development and breast cancer. *Science*, **296**, 1046-1049.
- [9] Lam,T.K., Carpentier,A., Lewis,G.F., van de,W.G., Fantus,I.G., & Giacca,A. (2003) Mechanisms of the free fatty acid-induced increase in hepatic glucose production. *Am. J. Physiol Endocrinol. Metab*, **284**, E863-E873.
- [10] Unger,R.H. (2003) Minireview: weapons of lean body mass destruction: the role of ectopic lipids in the metabolic syndrome. *Endocrinology*, **144**, 5159-5165.
- [11] Shulman,G.I. (2000) Cellular mechanisms of insulin resistance. *J. Clin. Invest*, **106**, 171-176.
- [12] Abel,E.D., Peroni,O., Kim,J.K., Kim,Y.B., Boss,O., Hadro,E., Minnemann,T., Shulman,G.I., & Kahn,B.B. (2001) Adipose-selective targeting of the GLUT4 gene impairs insulin action in muscle and liver. *Nature*, **409**, 729-733.
- [13] Goldstein,B.J. (2002) Insulin resistance as the core defect in type 2 diabetes mellitus. *Am. J. Cardiol.*, **90**, 3G-10G.
- [14] Terauchi,Y. & Kadowaki,T. (2002) Insights into molecular pathogenesis of type 2 diabetes from knockout mouse models. *Endocr. J.*, **49**, 247-263.
- [15] Hotamisligil,G.S. (1999) The role of TNFalpha and TNF receptors in obesity and insulin resistance. *J. Intern. Med.*, **245**, 621-625.
- [16] ShklyaeV,S., Aslanidi,G., Tennant,M., Prima,V., Kohlbrenner,E., Kroutov,V., Campbell-Thompson,M., Crawford,J., Shek,E.W., Scarpace,P.J., & Zolotukhin,S. (2003) Sustained peripheral expression of transgene adiponectin offsets the development of diet-induced obesity in rats. *Proc. Natl. Acad. Sci. U. S. A*, **100**, 14217-14222.
- [17] Combs,T.P., Pajvani,U.B., Berg,A.H., Lin,Y., Jelicks,L.A., Laplante,M., Nawrocki,A.R., Rajala,M.W., Parlow,A.F., Cheeseboro,L., Ding,Y.Y., Russell,R.G., Lindemann,D., Hartley,A., Baker,G.R., Obici,S., Deshaies,Y., Ludgate,M., Rossetti,L., & Scherer,P.E. (2004) A transgenic mouse with a deletion in the collagenous domain of adiponectin displays elevated circulating adiponectin and improved insulin sensitivity. *Endocrinology*, **145**, 367-383.
- [18] Yamauchi,T., Kamon,J., Minokoshi,Y., Ito,Y., Waki,H., Uchida,S., Yamashita,S., Noda,M., Kita,S., Ueki,K., Eto,K., Akanuma,Y., Froguel,P., Foufelle,F., Ferre,P., Carling,D., Kimura,S., Nagai,R., Kahn,B.B., & Kadowaki,T. (2002) Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat. Med.*, **8**, 1288-1295.
- [19] Tomas,E., Tsao,T.S., Saha,A.K., Murrey,H.E., Zhang,C.C., Itani,S.I., Lodish,H.F., & Ruderman,N.B. (2002) Enhanced muscle fat oxidation and glucose transport by ACRP30 globular domain: acetyl-CoA carboxylase inhibition and AMP-activated protein kinase activation. *Proc. Natl. Acad. Sci. U. S. A*, **99**, 16309-16313.
- [20] Yamauchi,T., Kamon,J., Waki,H., Terauchi,Y., Kubota,N., Hara,K., Mori,Y., Ide,T., Murakami,K., Tsuboyama-Kasaoka,N., Ezaki,O., Akanuma,Y., Gavrilova,O., Vinson,C., Reitman,M.L., Kagechika,H., Shudo,K., Yoda,M., Nakano,Y., Tobe,K., Nagai,R., Kimura,S.,

- Tomita,M., Froguel,P., & Kadowaki,T. (2001) The fat-derived hormone adiponectin reverses insulin resistance associated with both lipotrophy and obesity. *Nat. Med.*, **7**, 941-946.
- [21] Bluher,M., Michael,M.D., Peroni,O.D., Ueki,K., Carter,N., Kahn,B.B., & Kahn,C.R. (2002) Adipose tissue selective insulin receptor knockout protects against obesity and obesity-related glucose intolerance. *Dev. Cell*, **3**, 25-38.
- [22] Bjorbaek,C. & Kahn,B.B. (2004) Leptin signaling in the central nervous system and the periphery. *Recent Prog. Horm. Res.*, **59**, 305-331.
- [23] Margetic,S., Gazzola,C., Pegg,G.G., & Hill,R.A. (2002) Leptin: a review of its peripheral actions and interactions. *Int. J. Obes. Relat Metab Disord.*, **26**, 1407-1433.
- [24] Harris,R.B. (2000) Leptin--much more than a satiety signal. *Annu. Rev. Nutr.*, **20:45-75.**, 45-75.
- [25] Huan,J.N., Li,J., Han,Y., Chen,K., Wu,N., & Zhao,A.Z. (2003) Adipocyte-selective reduction of the leptin receptors induced by antisense RNA leads to increased adiposity, dyslipidemia, and insulin resistance. *J. Biol. Chem.*, **278**, 45638-45650.
- [26] Shimomura,I., Hammer,R.E., Ikemoto,S., Brown,M.S., & Goldstein,J.L. (1999) Leptin reverses insulin resistance and diabetes mellitus in mice with congenital lipodystrophy. *Nature*, **401**, 73-76.
- [27] Friedman,J. (2002) Fat in all the wrong places. *Nature*, **415**, 268-269.
- [28] Minokoshi,Y., Kim,Y.B., Peroni,O.D., Fryer,L.G., Muller,C., Carling,D., & Kahn,B.B. (2002) Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature*, **415**, 339-343.
- [29] Puigserver,P., Wu,Z., Park,C.W., Graves,R., Wright,M., & Spiegelman,B.M. (1998) A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell*, **92**, 829-839.
- [30] Kakuma,T., Wang,Z.W., Pan,W., Unger,R.H., & Zhou,Y.T. (2000) Role of leptin in peroxisome proliferator-activated receptor gamma coactivator-1 expression. *Endocrinology*, **141**, 4576-4582.
- [31] Berg,A.H., Combs,T.P., Du,X., Brownlee,M., & Scherer,P.E. (2001) The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat. Med.*, **7**, 947-953.
- [32] Ebihara,K., Ogawa,Y., Masuzaki,H., Shintani,M., Miyanaga,F., Aizawa-Abe,M., Hayashi,T., Hosoda,K., Inoue,G., Yoshimasa,Y., Gavrilova,O., Reitman,M.L., & Nakao,K. (2001) Transgenic overexpression of leptin rescues insulin resistance and diabetes in a mouse model of lipotrophic diabetes. *Diabetes*, **50**, 1440-1448.
- [33] Green,H. & Meuth,M. (1974) An established pre-adipose cell line and its differentiation in culture. *Cell*, **3**, 127-133.
- [34] Green,H. & Kehinde,O. (1976) Spontaneous heritable changes leading to increased adipose conversion in 3T3 cells. *Cell*, **7**, 105-113.
- [35] Green,H. & Kehinde,O. (1975) An established preadipose cell line and its differentiation in culture. II. Factors affecting the adipose conversion. *Cell*, **5**, 19-27.
- [36] Shugart,E.C. & Umek,R.M. (1997) Dexamethasone signaling is required to establish the postmitotic state of adipocyte development. *Cell Growth Differ.*, **8**, 1091-1098.
- [37] Ho,I.C., Kim,J.H., Rooney,J.W., Spiegelman,B.M., & Glimcher,L.H. (1998) A potential role for the nuclear factor of activated T cells family of transcriptional regulatory proteins in adipogenesis. *Proc. Natl. Acad. Sci. U. S. A.*, **95**, 15537-15541.
- [38] Reed,B.C. & Lane,M.D. (1980) Insulin receptor synthesis and turnover in differentiating 3T3-L1 preadipocytes. *Proc. Natl. Acad. Sci. U. S. A.*, **77**, 285-289.
- [39] Reed,B.C., Kaufmann,S.H., Mackall,J.C., Student,A.K., & Lane,M.D. (1977) Alterations in insulin binding accompanying differentiation of 3T3-L1 preadipocytes. *Proc. Natl. Acad. Sci. U. S. A.*, **74**, 4876-4880.
- [40] Hamm,J.K., Park,B.H., & Farmer,S.R. (2001) A role for C/EBPbeta in regulating peroxisome proliferator-activated receptor gamma activity during adipogenesis in 3T3-L1 preadipocytes. *J. Biol. Chem.*, **276**, 18464-18471.
- [41] Gagnon,A.M., Chabot,J., Padasani,D., & Sorisky,A. (1998) Extracellular matrix induced by TGFbeta impairs insulin signal transduction in 3T3-L1 preadipose cells. *J. Cell Physiol*, **175**, 370-378.
- [42] Chavey,C., Mari,B., Monthouel,M.N., Bonnafous,S., Anglard,P., Van Obberghen,E., & Tartare-Deckert,S. (2003) Matrix metalloproteinases are differentially expressed in adipose tissue during obesity and modulate adipocyte differentiation. *J. Biol. Chem.*, **278**, 11888-11896.
- [43] Halbleib,M., Skurk,T., de Luca,C., von Heimburg,D., & Hauner,H. (2003) Tissue engineering of white adipose tissue using hyaluronic acid-based scaffolds. I: in vitro differentiation of human adipocyte precursor cells on scaffolds. *Biomaterials*, **24**, 3125-3132.

- [44] Calvo, J.C., Gandjbakhche, A.H., Nossal, R., Hascall, V.C., & Yanagishita, M. (1993) Rheological effects of the presence of hyaluronic acid in the extracellular media of differentiated 3T3-L1 preadipocyte cultures. *Arch. Biochem. Biophys.*, **302**, 468-475.
- [45] Berberich, S.J., Litteral, V., Mayo, L.D., Tabesh, D., & Morris, D. (1999) mdm-2 gene amplification in 3T3-L1 preadipocytes. *Differentiation*, **64**, 205-212.
- [46] Morrison, R.F. & Farmer, S.R. (1999) Role of PPARgamma in regulating a cascade expression of cyclin-dependent kinase inhibitors, p18(INK4c) and p21(Waf1/Cip1), during adipogenesis. *J. Biol. Chem.*, **274**, 17088-17097.
- [47] Reichert, M. & Eick, D. (1999) Analysis of cell cycle arrest in adipocyte differentiation. *Oncogene*, **18**, 459-466.
- [48] Phelps, D.E. & Xiong, Y. (1998) Regulation of cyclin-dependent kinase 4 during adipogenesis involves switching of cyclin D subunits and concurrent binding of p18INK4c and p27Kip1. *Cell Growth Differ.*, **9**, 595-610.
- [49] Liu, K., Guan, Y., MacNicol, M.C., MacNicol, A.M., & McGehee, R.E., Jr. (2002) Early expression of p107 is associated with 3T3-L1 adipocyte differentiation. *Mol Cell Endocrinol.*, **194**, 51-61.
- [50] Prince, A.M., May, J.S., Burton, G.R., Lyle, R.E., & McGehee, R.E., Jr. (2002) Proteasomal degradation of retinoblastoma-related p130 during adipocyte differentiation. *Biochem. Biophys. Res. Commun.*, **290**, 1066-1071.
- [51] May, J.S., Prince, A.M., Lyle, R.E., & McGehee, R.E., Jr. (2001) Antisense suppression of p107 inhibits 3T3-L1 adipocyte differentiation. *Biochem. Biophys. Res. Commun.*, **283**, 837-842.
- [52] Richon, V.M., Lyle, R.E., & McGehee, R.E., Jr. (1997) Regulation and expression of retinoblastoma proteins p107 and p130 during 3T3-L1 adipocyte differentiation. *J. Biol. Chem.*, **272**, 10117-10124.
- [53] Classon, M., Kennedy, B.K., Mulloy, R., & Harlow, E. (2000) Opposing roles of pRB and p107 in adipocyte differentiation. *Proc. Natl. Acad. Sci. U. S. A.*, **97**, 10826-10831.
- [54] Hansen, J.B., Jorgensen, C., Petersen, R.K., Hallenborg, P., De Matteis, R., Boye, H.A., Petrovic, N., Enerback, S., Nedergaard, J., Cinti, S., te, R.H., & Kristiansen, K. (2004) Retinoblastoma protein functions as a molecular switch determining white versus brown adipocyte differentiation. *Proc. Natl. Acad. Sci. U. S. A.*, **101**, 4112-4117.
- [55] Cole, K.A., Harmon, A.W., Harp, J.B., & Patel, Y.M. (2004) Rb regulates C/EBPbeta-DNA-binding activity during 3T3-L1 adipogenesis. *Am. J. Physiol Cell Physiol*, **286**, C349-C354.
- [56] Cowherd, R.M., Lyle, R.E., & McGehee, R.E., Jr. (1999) Molecular regulation of adipocyte differentiation. *Semin. Cell Dev. Biol.*, **10**, 3-10.
- [57] Marshall, C.J. (1995) Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation. *Cell*, **80**, 179-185.
- [58] Vaudry, D., Stork, P.J., Lazarovici, P., & Eiden, L.E. (2002) Signaling pathways for PC12 cell differentiation: making the right connections. *Science*, **296**, 1648-1649.
- [59] Prusty, D., Park, B.H., Davis, K.E., & Farmer, S.R. (2002) Activation of MEK/ERK signaling promotes adipogenesis by enhancing peroxisome proliferator-activated receptor gamma (PPARgamma) and C/EBPalpha gene expression during the differentiation of 3T3-L1 preadipocytes. *J. Biol. Chem.*, **277**, 46226-46232.
- [60] Sekimoto, H. & Boney, C.M. (2003) C-terminal Src kinase (CSK) modulates insulin-like growth factor-I signaling through Src in 3T3-L1 differentiation. *Endocrinology*, **144**, 2546-2552.
- [61] Belmonte, N., Phillips, B.W., Massiera, F., Villageois, P., Wdziekonski, B., Saint-Marc, P., Nichols, J., Aubert, J., Saeki, K., Yuo, A., Narumiya, S., Ailhaud, G., & Dani, C. (2001) Activation of extracellular signal-regulated kinases and CREB/ATF-1 mediate the expression of CCAAT/enhancer binding proteins beta and -delta in preadipocytes. *Mol Endocrinol.*, **15**, 2037-2049.
- [62] Tang, Q.Q. & Lane, M.D. (2000) Role of C/EBP homologous protein (CHOP-10) in the programmed activation of CCAAT/enhancer-binding protein-beta during adipogenesis. *Proc. Natl. Acad. Sci. U. S. A.*, **97**, 12446-12450.
- [63] Bost, F., Caron, L., Marchetti, I., Dani, C., Marchand-Brustel, Y., & Binetruy, B. (2002) Retinoic acid activation of the ERK pathway is required for embryonic stem cell commitment into the adipocyte lineage. *Biochem. J.*, **361**, 621-627.
- [64] Hu, E., Kim, J.B., Sarraf, P., & Spiegelman, B.M. (1996) Inhibition of adipogenesis through MAP kinase-mediated phosphorylation of PPARgamma. *Science*, **274**, 2100-2103.
- [65] Vassaux, G., Negrel, R., Ailhaud, G., & Gaillard, D. (1994) Proliferation and differentiation of rat adipose precursor cells in chemically defined medium: differential action of anti-adipogenic agents. *J. Cell Physiol*, **161**, 249-256.

- [66] Font,d.M., Porras,A., Ahn,N., & Santos,E. (1997) Mitogen-activated protein kinase activation is not necessary for, but antagonizes, 3T3-L1 adipocytic differentiation. *Mol Cell Biol.*, **17**, 6068-6075.
- [67] Zhang,J.W., Tang,Q.Q., Vinson,C., & Lane,M.D. (2004) Dominant-negative C/EBP disrupts mitotic clonal expansion and differentiation of 3T3-L1 preadipocytes. *Proc. Natl. Acad. Sci. U. S. A.*, **101**, 43-47.
- [68] Engelman,J.A., Lisanti,M.P., & Scherer,P.E. (1998) Specific inhibitors of p38 mitogen-activated protein kinase block 3T3-L1 adipogenesis. *J. Biol. Chem.*, **273**, 32111-32120.
- [69] Engelman,J.A., Berg,A.H., Lewis,R.Y., Lin,A., Lisanti,M.P., & Scherer,P.E. (1999) Constitutively active mitogen-activated protein kinase kinase 6 (MKK6) or salicylate induces spontaneous 3T3-L1 adipogenesis. *J. Biol. Chem.*, **274**, 35630-35638.
- [70] Maytin,E.V., Ubeda,M., Lin,J.C., & Habener,J.F. (2001) Stress-inducible transcription factor CHOP/gadd153 induces apoptosis in mammalian cells via p38 kinase-dependent and -independent mechanisms. *Exp. Cell Res.*, **267**, 193-204.
- [71] Brenner,B., Koppenhoefer,U., Weinstock,C., Linderkamp,O., Lang,F., & Gulbins,E. (1997) Fas- or ceramide-induced apoptosis is mediated by a Rac1-regulated activation of Jun N-terminal kinase/p38 kinases and GADD153. *J. Biol. Chem.*, **272**, 22173-22181.
- [72] Wang,X.Z. & Ron,D. (1996) Stress-induced phosphorylation and activation of the transcription factor CHOP (GADD153) by p38 MAP Kinase. *Science*, **272**, 1347-1349.
- [73] Reusch,J.E., Colton,L.A., & Klemm,D.J. (2000) CREB activation induces adipogenesis in 3T3-L1 cells. *Mol Cell Biol.*, **20**, 1008-1020.
- [74] Reusch,J.E. & Klemm,D.J. (2002) Inhibition of cAMP-response element-binding protein activity decreases protein kinase B/Akt expression in 3T3-L1 adipocytes and induces apoptosis. *J. Biol. Chem.*, **277**, 1426-1432.
- [75] Moldes,M., Zuo,Y., Morrison,R.F., Silva,D., Park,B.H., Liu,J., & Farmer,S.R. (2003) Peroxisome-proliferator-activated receptor gamma suppresses Wnt/beta-catenin signalling during adipogenesis. *Biochem. J.*, **376**, 607-613.
- [76] Smas,C.M., Chen,L., Zhao,L., Latasa,M.J., & Sul,H.S. (1999) Transcriptional repression of pref-1 by glucocorticoids promotes 3T3-L1 adipocyte differentiation. *J. Biol. Chem.*, **274**, 12632-12641.
- [77] Gregoire,F.M., Smas,C.M., & Sul,H.S. (1998) Understanding adipocyte differentiation. *Physiol Rev.*, **78**, 783-809.
- [78] Tanaka,T., Yoshida,N., Kishimoto,T., & Akira,S. (1997) Defective adipocyte differentiation in mice lacking the C/EBPbeta and/or C/EBPdelta gene. *EMBO J.*, **16**, 7432-7443.
- [79] Tang,Q.Q., Otto,T.C., & Lane,M.D. (2003) CCAAT/enhancer-binding protein beta is required for mitotic clonal expansion during adipogenesis. *Proc. Natl. Acad. Sci. U. S. A.*, **100**, 850-855.
- [80] El Jack,A.K., Kandror,K.V., & Pilch,P.F. (1999) The formation of an insulin-responsive vesicular cargo compartment is an early event in 3T3-L1 adipocyte differentiation. *Mol Biol. Cell*, **10**, 1581-1594.
- [81] Dixon,T.M., Daniel,K.W., Farmer,S.R., & Collins,S. (2001) CCAAT/enhancer-binding protein alpha is required for transcription of the beta 3-adrenergic receptor gene during adipogenesis. *J. Biol. Chem.*, **276**, 722-728.
- [82] Darlington,G.J., Ross,S.E., & MacDougald,O.A. (1998) The role of C/EBP genes in adipocyte differentiation. *J. Biol. Chem.*, **273**, 30057-30060.
- [83] Lin,F.T. & Lane,M.D. (1992) Antisense CCAAT/enhancer-binding protein RNA suppresses coordinate gene expression and triglyceride accumulation during differentiation of 3T3-L1 preadipocytes. *Genes Dev.*, **6**, 533-544.
- [84] Chen,S.S., Chen,J.F., Johnson,P.F., Muppala,V., & Lee,Y.H. (2000) C/EBPbeta, when expressed from the C/ebpalpha gene locus, can functionally replace C/EBPalpha in liver but not in adipose tissue. *Mol Cell Biol.*, **20**, 7292-7299.
- [85] Wang,N.D., Finegold,M.J., Bradley,A., Ou,C.N., Abdelsayed,S.V., Wilde,M.D., Taylor,L.R., Wilson,D.R., & Darlington,G.J. (1995) Impaired energy homeostasis in C/EBP alpha knockout mice. *Science*, **269**, 1108-1112.
- [86] Kadowaki,T., Hara,K., Yamauchi,T., Terauchi,Y., Tobe,K., & Nagai,R. (2003) Molecular mechanism of insulin resistance and obesity. *Exp. Biol. Med.*, **228**, 1111-1117.
- [87] He,W., Barak,Y., Hevener,A., Olson,P., Liao,D., Le,J., Nelson,M., Ong,E., Olefsky,J.M., & Evans,R.M. (2003) Adipose-specific peroxisome proliferator-activated receptor gamma knockout causes insulin resistance in fat and liver but not in muscle. *Proc. Natl. Acad. Sci. U. S. A.*, **100**, 15712-15717.

- [88] Koutnikova,H., Cock,T.A., Watanabe,M., Houten,S.M., Champy,M.F., Dierich,A., & Auwerx,J. (2003) Compensation by the muscle limits the metabolic consequences of lipodystrophy in PPAR gamma hypomorphic mice. *Proc. Natl. Acad. Sci. U. S. A.*, **100**, 14457-14462.
- [89] Fasshauer,M., Klein,J., Kriauciunas,K.M., Ueki,K., Benito,M., & Kahn,C.R. (2001) Essential role of insulin receptor substrate 1 in differentiation of brown adipocytes. *Mol Cell Biol.*, **21**, 319-329.
- [90] Sakaue,H., Ogawa,W., Matsumoto,M., Kuroda,S., Takata,M., Sugimoto,T., Spiegelman,B.M., & Kasuga,M. (1998) Posttranscriptional control of adipocyte differentiation through activation of phosphoinositide 3-kinase. *J. Biol. Chem.*, **273**, 28945-28952.
- [91] Smith,P.J., Wise,L.S., Berkowitz,R., Wan,C., & Rubin,C.S. (1988) Insulin-like growth factor-I is an essential regulator of the differentiation of 3T3-L1 adipocytes. *J. Biol. Chem.*, **263**, 9402-9408.
- [92] Nakae,J., Kitamura,T., Kitamura,Y., Biggs,W.H., III, Arden,K.C., & Accili,D. (2003) The forkhead transcription factor Foxo1 regulates adipocyte differentiation. *Dev. Cell*, **4**, 119-129.
- [93] Hemati,N., Ross,S.E., Erickson,R.L., Groblewski,G.E., & MacDougald,O.A. (1997) Signaling pathways through which insulin regulates CCAAT/enhancer binding protein alpha (C/EBPalpha) phosphorylation and gene expression in 3T3-L1 adipocytes. Correlation with GLUT4 gene expression. *J. Biol. Chem.*, **272**, 25913-25919.
- [94] Ribaux,P.G. & Iyendjian,P.B. (2003) Analysis of the role of protein kinase B (cAKT) in insulin-dependent induction of glucokinase and sterol regulatory element-binding protein 1 (SREBP1) mRNAs in hepatocytes. *Biochem. J.*, **376**, 697-705.
- [95] Sul,H.S., Latasa,M.J., Moon,Y., & Kim,K.H. (2000) Regulation of the fatty acid synthase promoter by insulin. *J. Nutr.*, **130**, 315S-320S.
- [96] Fajas,L., Schoonjans,K., Gelman,L., Kim,J.B., Najib,J., Martin,G., Fruchart,J.C., Briggs,M., Spiegelman,B.M., & Auwerx,J. (1999) Regulation of peroxisome proliferator-activated receptor gamma expression by adipocyte differentiation and determination factor 1/sterol regulatory element binding protein 1: implications for adipocyte differentiation and metabolism. *Mol Cell Biol.*, **19**, 5495-5503.
- [97] Kim,J.B., Wright,H.M., Wright,M., & Spiegelman,B.M. (1998) ADD1/SREBP1 activates PPARgamma through the production of endogenous ligand. *Proc. Natl. Acad. Sci. U. S. A.*, **95**, 4333-4337.
- [98] Nanbu-Wakao,R., Fujitani,Y., Masuho,Y., Muramatu,M., & Wakao,H. (2000) Prolactin enhances CCAAT enhancer-binding protein-beta (C/EBP beta) and peroxisome proliferator-activated receptor gamma (PPAR gamma) messenger RNA expression and stimulates adipogenic conversion of NIH-3T3 cells. *Mol Endocrinol.*, **14**, 307-316.
- [99] Wolf,G. (1999) The molecular mechanism of the stimulation of adipocyte differentiation by a glucocorticoid. *Nutr. Rev.*, **57**, 324-326.
- [100] Levine,J.A., Jensen,M.D., Eberhardt,N.L., & O'Brien,T. (1998) Adipocyte macrophage colony-stimulating factor is a mediator of adipose tissue growth. *J. Clin. Invest.*, **101**, 1557-1564.
- [101] Fischer-Posovszky,P., Tornqvist,H., Debatin,K.M., & Wabitsch,M. (2004) Inhibition of death-receptor mediated apoptosis in human adipocytes by the insulin-like growth factor I (IGF-I)/IGF-I receptor autocrine circuit. *Endocrinology*, **145**, 1849-1859.
- [102] Bujalska,I.J., Walker,E.A., Hewison,M., & Stewart,P.M. (2002) A switch in dehydrogenase to reductase activity of 11 beta-hydroxysteroid dehydrogenase type 1 upon differentiation of human omental adipose stromal cells. *J. Clin. Endocrinol. Metab.*, **87**, 1205-1210.
- [103] Belanger,C., Luu-The,V., Dupont,P., & Tchernof,A. (2002) Adipose tissue intracrinology: potential importance of local androgen/estrogen metabolism in the regulation of adiposity. *Horm. Metab Res.*, **34**, 737-745.
- [104] Napolitano,A., Voice,M.W., Edwards,C.R., Seckl,J.R., & Chapman,K.E. (1998) 11Beta-hydroxysteroid dehydrogenase 1 in adipocytes: expression is differentiation-dependent and hormonally regulated. *J. Steroid Biochem. Mol Biol.*, **64**, 251-260.
- [105] MacDougald,O.A. & Mandrup,S. (2002) Adipogenesis: forces that tip the scales. *Trends Endocrinol. Metab.*, **13**, 5-11.
- [106] Ross,S.E., Hemati,N., Longo,K.A., Bennett,C.N., Lucas,P.C., Erickson,R.L., & MacDougald,O.A. (2000) Inhibition of adipogenesis by Wnt signaling. *Science*, **289**, 950-953.
- [107] Valyasevi,R.W., Jyonouchi,S.C., Dutton,C.M., Munsakul,N., & Bahn,R.S. (2001) Effect of tumor necrosis factor-alpha, interferon-gamma, and transforming growth factor-beta on adipogenesis and expression of thyrotropin receptor in human orbital preadipocyte fibroblasts. *J. Clin. Endocrinol. Metab.*, **86**, 903-908.

- [108] Choy,L., Skillington,J., & Derynck,R. (2000) Roles of autocrine TGF-beta receptor and Smad signaling in adipocyte differentiation. *J. Cell Biol.*, **149**, 667-682.
- [109] Bortell,R., Owen,T.A., Ignatz,R., Stein,G.S., & Stein,J.L. (1994) TGF beta 1 prevents the down-regulation of type I procollagen, fibronectin, and TGF beta 1 gene expression associated with 3T3-L1 pre-adipocyte differentiation. *J. Cell Biochem.*, **54**, 256-263.
- [110] Gerhold,D.L., Liu,F., Jiang,G., Li,Z., Xu,J., Lu,M., Sachs,J.R., Bagchi,A., Fridman,A., Holder,D.J., Doebber,T.W., Berger,J., Elbrecht,A., Moller,D.E., & Zhang,B.B. (2002) Gene expression profile of adipocyte differentiation and its regulation by peroxisome proliferator-activated receptor-gamma agonists. *Endocrinology*, **143**, 2106-2118.
- [111] Brady,M.J., Bourbonnais,F.J., & Saltiel,A.R. (1998) The activation of glycogen synthase by insulin switches from kinase inhibition to phosphatase activation during adipogenesis in 3T3-L1 cells. *J. Biol. Chem.*, **273**, 14063-14066.
- [112] Greenberg,A.S., Egan,J.J., Wek,S.A., Moos,M.C., Jr., Londos,C., & Kimmel,A.R. (1993) Isolation of cDNAs for perilipins A and B: sequence and expression of lipid droplet-associated proteins of adipocytes. *Proc. Natl. Acad. Sci. U. S. A*, **90**, 12035-12039.
- [113] Greenberg,A.S., Egan,J.J., Wek,S.A., Garty,N.B., Blanchette-Mackie,E.J., & Londos,C. (1991) Perilipin, a major hormonally regulated adipocyte-specific phosphoprotein associated with the periphery of lipid storage droplets. *J. Biol. Chem.*, **266**, 11341-11346.
- [114] Sztalryd,C., Xu,G., Dorward,H., Tansey,J.T., Contreras,J.A., Kimmel,A.R., & Londos,C. (2003) Perilipin A is essential for the translocation of hormone-sensitive lipase during lipolytic activation. *J. Cell Biol.*, **161**, 1093-1103.
- [115] Harada,K., Shen,W.J., Patel,S., Natu,V., Wang,J., Osuga,J., Ishibashi,S., & Kraemer,F.B. (2003) Resistance to high-fat diet-induced obesity and altered expression of adipose-specific genes in HSL-deficient mice. *Am. J. Physiol Endocrinol. Metab*, **285**, E1182-E1195.
- [116] Egan,J.J., Greenberg,A.S., Chang,M.K., Wek,S.A., Moos,M.C., Jr., & Londos,C. (1992) Mechanism of hormone-stimulated lipolysis in adipocytes: translocation of hormone-sensitive lipase to the lipid storage droplet. *Proc. Natl. Acad. Sci. U. S. A*, **89**, 8537-8541.
- [117] Holm,C. (2003) Molecular mechanisms regulating hormone-sensitive lipase and lipolysis. *Biochem. Soc. Trans.*, **31**, 1120-1124.
- [118] Londos,C., Brasamle,D.L., Schultz,C.J., Adler-Wailes,D.C., Levin,D.M., Kimmel,A.R., & Rondinone,C.M. (1999) On the control of lipolysis in adipocytes. *Ann. N. Y. Acad. Sci.*, **892**, 155-168.
- [119] Wijkander,J., Landstrom,T.R., Manganiello,V., Belfrage,P., & Degerman,E. (1998) Insulin-induced phosphorylation and activation of phosphodiesterase 3B in rat adipocytes: possible role for protein kinase B but not mitogen-activated protein kinase or p70 S6 kinase. *Endocrinology*, **139**, 219-227.
- [120] Syu,L.J. & Saltiel,A.R. (1999) Lipotransin: a novel docking protein for hormone-sensitive lipase. *Mol Cell*, **4**, 109-115.
- [121] Liu,P., Ying,Y., Zhao,Y., Mundy,D.I., Zhu,M., & Anderson,R.G. (2004) Chinese hamster ovary K2 cell lipid droplets appear to be metabolic organelles involved in membrane traffic. *J. Biol. Chem.*, **279**, 3787-3792.
- [122] Castro-Chavez,F., Yechoor,V.K., Saha,P.K., Martinez-Botas,J., Wooten,E.C., Sharma,S., O'Connell,P., Taegtmeyer,H., & Chan,L. (2003) Coordinated upregulation of oxidative pathways and downregulation of lipid biosynthesis underlie obesity resistance in perilipin knockout mice: a microarray gene expression profile. *Diabetes*, **52**, 2666-2674.
- [123] Das,K., Lewis,R.Y., Combatsiaris,T.P., Lin,Y., Shapiro,L., Charron,M.J., & Scherer,P.E. (1999) Predominant expression of the mitochondrial dicarboxylate carrier in white adipose tissue. *Biochem. J.*, **344**, 313-320.
- [124] Lopez,J.M., Hegardt,F.G., & Haro,D. (2001) Differential expression of cytosolic and mitochondrial 3-hydroxy-3-methylglutaryl CoA synthases during adipocyte differentiation. *Mol Cell Biochem.*, **217**, 57-66.
- [125] Wilson-Fritch,L., Burkart,A., Bell,G., Mendelson,K., Leszyk,J., Nicoloso,S., Czech,M., & Corvera,S. (2003) Mitochondrial biogenesis and remodeling during adipogenesis and in response to the insulin sensitizer rosiglitazone. *Mol Cell Biol.*, **23**, 1085-1094.
- [126] Hardie,D.G., Carling,D., & Carlson,M. (1998) The AMP-activated/SNF1 protein kinase subfamily: metabolic sensors of the eukaryotic cell? *Annu. Rev. Biochem.*, **67**, 821-855.
- [127] Galbiati,F., Razani,B., & Lisanti,M.P. (2001) Emerging themes in lipid rafts and caveolae. *Cell*, **106**, 403-411.
- [128] Simons,K. & Ikonen,E. (1997) Functional rafts in cell membranes. *Nature*, **387**, 569-572.

- [129] Rothberg,K.G., Heuser,J.E., Donzell,W.C., Ying,Y.S., Glenney,J.R., & Anderson,R.G. (1992) Caveolin, a protein component of caveolae membrane coats. *Cell*, **68**, 673-682.
- [130] Smart,E.J., Graf,G.A., McNiven,M.A., Sessa,W.C., Engelman,J.A., Scherer,P.E., Okamoto,T., & Lisanti,M.P. (1999) Caveolins, liquid-ordered domains, and signal transduction. *Mol Cell Biol.*, **19**, 7289-7304.
- [131] Thorn,H., Stenkula,K.G., Karlsson,M., Ortegren,U., Nystrom,F.H., Gustavsson,J., & Stralfors,P. (2003) Cell surface orifices of caveolae and localization of caveolin to the necks of caveolae in adipocytes. *Mol Biol. Cell*, **14**, 3967-3976.
- [132] Razani,B. & Lisanti,M.P. (2001) Caveolin-deficient mice: insights into caveolar function human disease. *J. Clin. Invest*, **108**, 1553-1561.
- [133] Shaul,P.W. & Anderson,R.G. (1998) Role of plasmalemmal caveolae in signal transduction. *Am. J. Physiol*, **275**, L843-L851.
- [134] Parton,R.G., Molero,J.C., Floetenmeyer,M., Green,K.M., & James,D.E. (2002) Characterization of a distinct plasma membrane macrodomain in differentiated adipocytes. *J. Biol. Chem.*, **277**, 46769-46778.
- [135] Khan,A.H. & Pessin,J.E. (2002) Insulin regulation of glucose uptake: a complex interplay of intracellular signalling pathways. *Diabetologia*, **45**, 1475-1483.
- [136] Tang,Z., Scherer,P.E., Okamoto,T., Song,K., Chu,C., Kohtz,D.S., Nishimoto,I., Lodish,H.F., & Lisanti,M.P. (1996) Molecular cloning of caveolin-3, a novel member of the caveolin gene family expressed predominantly in muscle. *J. Biol. Chem.*, **271**, 2255-2261.
- [137] Scherer,P.E., Okamoto,T., Chun,M., Nishimoto,I., Lodish,H.F., & Lisanti,M.P. (1996) Identification, sequence, and expression of caveolin-2 defines a caveolin gene family. *Proc. Natl. Acad. Sci. U. S. A*, **93**, 131-135.
- [138] Scherer,P.E., Lewis,R.Y., Volonte,D., Engelman,J.A., Galbiati,F., Couet,J., Kohtz,D.S., van Donselaar,E., Peters,P., & Lisanti,M.P. (1997) Cell-type and tissue-specific expression of caveolin-2. Caveolins 1 and 2 co-localize and form a stable hetero-oligomeric complex in vivo. *J. Biol. Chem.*, **272**, 29337-29346.
- [139] Parolini,I., Sargiacomo,M., Galbiati,F., Rizzo,G., Grignani,F., Engelman,J.A., Okamoto,T., Ikezu,T., Scherer,P.E., Mora,R., Rodriguez-Boulan,E., Peschle,C., & Lisanti,M.P. (1999) Expression of caveolin-1 is required for the transport of caveolin-2 to the plasma membrane. Retention of caveolin-2 at the level of the golgi complex. *J. Biol. Chem.*, **274**, 25718-25725.
- [140] Scherer,P.E., Lisanti,M.P., Baldini,G., Sargiacomo,M., Mastick,C.C., & Lodish,H.F. (1994) Induction of caveolin during adipogenesis and association of GLUT4 with caveolin-rich vesicles. *J. Cell Biol.*, **127**, 1233-1243.
- [141] Kandror,K.V., Stephens,J.M., & Pilch,P.F. (1995) Expression and compartmentalization of caveolin in adipose cells: coordinate regulation with and structural segregation from GLUT4. *J. Cell Biol.*, **129**, 999-1006.
- [142] Razani,B., Engelman,J.A., Wang,X.B., Schubert,W., Zhang,X.L., Marks,C.B., Macaluso,F., Russell,R.G., Li,M., Pestell,R.G., Di Vizio,D., Hou,H., Jr., Kneitz,B., Lagaud,G., Christ,G.J., Edelmann,W., & Lisanti,M.P. (2001) Caveolin-1 null mice are viable but show evidence of hyperproliferative and vascular abnormalities. *J. Biol. Chem.*, **276**, 38121-38138.
- [143] Park,D.S., Woodman,S.E., Schubert,W., Cohen,A.W., Frank,P.G., Chandra,M., Shirani,J., Razani,B., Tang,B., Jelicks,L.A., Factor,S.M., Weiss,L.M., Tanowitz,H.B., & Lisanti,M.P. (2002) Caveolin-1/3 double-knockout mice are viable, but lack both muscle and non-muscle caveolae, and develop a severe cardiomyopathic phenotype. *Am. J. Pathol.*, **160**, 2207-2217.
- [144] Ros-Baro,A., Lopez-Iglesias,C., Peiro,S., Bellido,D., Palacin,M., Zorzano,A., & Camps,M. (2001) Lipid rafts are required for GLUT4 internalization in adipose cells. *Proc. Natl. Acad. Sci. U. S. A*, **98**, 12050-12055.
- [145] Kwik,J., Boyle,S., Fooksman,D., Margolis,L., Sheetz,M.P., & Edidin,M. (2003) Membrane cholesterol, lateral mobility, and the phosphatidylinositol 4,5-bisphosphate-dependent organization of cell actin. *Proc. Natl. Acad. Sci. U. S. A*, **100**, 13964-13969.
- [146] Parpal,S., Karlsson,M., Thorn,H., & Stralfors,P. (2001) Cholesterol depletion disrupts caveolae and insulin receptor signaling for metabolic control via insulin receptor substrate-1, but not for mitogen-activated protein kinase control. *J. Biol. Chem.*, **276**, 9670-9678.
- [147] Kimura,A., Mora,S., Shigematsu,S., Pessin,J.E., & Saltiel,A.R. (2002) The insulin receptor catalyzes the tyrosine phosphorylation of caveolin-1. *J. Biol. Chem.*, **277**, 30153-30158.
- [148] Nystrom,F.H., Chen,H., Cong,L.N., Li,Y., & Quon,M.J. (1999) Caveolin-1 interacts with the insulin receptor and can differentially modulate insulin signaling in transfected Cos-7 cells and rat adipose cells. *Mol Endocrinol.*, **13**, 2013-2024.

- [149] Yamamoto, M., Toya, Y., Schwencke, C., Lisanti, M.P., Myers, M.G., Jr., & Ishikawa, Y. (1998) Caveolin is an activator of insulin receptor signaling. *J. Biol. Chem.*, **273**, 26962-26968.
- [150] Cohen, A.W., Razani, B., Wang, X.B., Combs, T.P., Williams, T.M., Scherer, P.E., & Lisanti, M.P. (2003) Caveolin-1-deficient mice show insulin resistance and defective insulin receptor protein expression in adipose tissue. *Am. J. Physiol Cell Physiol*, **285**, C222-C235.
- [151] Razani, B., Combs, T.P., Wang, X.B., Frank, P.G., Park, D.S., Russell, R.G., Li, M., Tang, B., Jelicks, L.A., Scherer, P.E., & Lisanti, M.P. (2002) Caveolin-1-deficient mice are lean, resistant to diet-induced obesity, and show hypertriglyceridemia with adipocyte abnormalities. *J. Biol. Chem.*, **277**, 8635-8647.
- [152] Bickel, P.E. (2002) Lipid rafts and insulin signaling. *Am. J. Physiol Endocrinol. Metab*, **282**, E1-E10.
- [153] Saltiel, A.R. & Pessin, J.E. (2002) Insulin signaling pathways in time and space. *Trends Cell Biol.*, **12**, 65-71.
- [154] Ikonen, E. & Parton, R.G. (2000) Caveolins and cellular cholesterol balance. *Traffic.*, **1**, 212-217.
- [155] Simpson, F., Whitehead, J.P., & James, D.E. (2001) GLUT4--at the cross roads between membrane trafficking and signal transduction. *Traffic.*, **2**, 2-11.
- [156] Roh, C., Roduit, R., Thorens, B., Fried, S., & Kandror, K.V. (2001) Lipoprotein lipase and leptin are accumulated in different secretory compartments in rat adipocytes. *J. Biol. Chem.*, **276**, 35990-35994.
- [157] Bogan, J.S. & Lodish, H.F. (1999) Two compartments for insulin-stimulated exocytosis in 3T3-L1 adipocytes defined by endogenous ACRP30 and GLUT4. *J. Cell Biol.*, **146**, 609-620.
- [158] Barr, V.A., Malide, D., Zarnowski, M.J., Taylor, S.I., & Cushman, S.W. (1997) Insulin stimulates both leptin secretion and production by rat white adipose tissue. *Endocrinology*, **138**, 4463-4472.
- [159] Millar, C.A., Meerloo, T., Martin, S., Hickson, G.R., Shimwell, N.J., Wakelam, M.J., James, D.E., & Gould, G.W. (2000) Adipsin and the glucose transporter GLUT4 traffic to the cell surface via independent pathways in adipocytes. *Traffic.*, **1**, 141-151.
- [160] Martin, S., Slot, J.W., & James, D.E. (1999) GLUT4 trafficking in insulin-sensitive cells. A morphological review. *Cell Biochem. Biophys.*, **30**, 89-113.
- [161] Rea, S. & James, D.E. (1997) Moving GLUT4: the biogenesis and trafficking of GLUT4 storage vesicles. *Diabetes*, **46**, 1667-1677.
- [162] Bryant, N.J., Govers, R., & James, D.E. (2002) Regulated transport of the glucose transporter GLUT4. *Nat. Rev. Mol Cell Biol.*, **3**, 267-277.
- [163] Martin, S., Rice, J.E., Gould, G.W., Keller, S.R., Slot, J.W., & James, D.E. (1997) The glucose transporter GLUT4 and the aminopeptidase vp165 colocalise in tubulo-vesicular elements in adipocytes and cardiomyocytes. *J. Cell Sci.*, **110**, 2281-2291.
- [164] Martin, S., Millar, C.A., Lyttle, C.T., Meerloo, T., Marsh, B.J., Gould, G.W., & James, D.E. (2000) Effects of insulin on intracellular GLUT4 vesicles in adipocytes: evidence for a secretory mode of regulation. *J. Cell Sci.*, **113**, 3427-3438.
- [165] Livingstone, C., James, D.E., Rice, J.E., Hanpeter, D., & Gould, G.W. (1996) Compartment ablation analysis of the insulin-responsive glucose transporter (GLUT4) in 3T3-L1 adipocytes. *Biochem. J.*, **315**, 487-495.
- [166] Yeh, J.I., Verhey, K.J., & Birnbaum, M.J. (1995) Kinetic analysis of glucose transporter trafficking in fibroblasts and adipocytes. *Biochemistry*, **34**, 15523-15531.
- [167] Keller, S.R. (2003) The insulin-regulated aminopeptidase: a companion and regulator of GLUT4. *Front Biosci.*, **8**, s410-s420.
- [168] Millar, C.A., Shewan, A., Hickson, G.R., James, D.E., & Gould, G.W. (1999) Differential regulation of secretory compartments containing the insulin-responsive glucose transporter 4 in 3T3-L1 adipocytes. *Mol Biol. Cell*, **10**, 3675-3688.
- [169] Randhawa, V.K., Bilan, P.J., Khayat, Z.A., Daneman, N., Liu, Z., Ramlal, T., Volchuk, A., Peng, X.R., Coppola, T., Regazzi, R., Trimble, W.S., & Klip, A. (2000) VAMP2, but not VAMP3/cellubrevin, mediates insulin-dependent incorporation of GLUT4 into the plasma membrane of L6 myoblasts. *Mol Biol. Cell*, **11**, 2403-2417.
- [170] Ploug, T., van Deurs, B., Ai, H., Cushman, S.W., & Ralston, E. (1998) Analysis of GLUT4 distribution in whole skeletal muscle fibers: identification of distinct storage compartments that are recruited by insulin and muscle contractions. *J. Cell Biol.*, **142**, 1429-1446.
- [171] Olson, A.L., Eyster, C.A., Duggins, Q.S., & Knight, J.B. (2003) Insulin promotes formation of polymerized microtubules by a phosphatidylinositol 3-kinase-independent, actin-dependent pathway in 3T3-L1 adipocytes. *Endocrinology*, **144**, 5030-5039.

- [172] Patki, V., Buxton, J., Chawla, A., Lifshitz, L., Fogarty, K., Carrington, W., Tuft, R., & Corvera, S. (2001) Insulin action on GLUT4 traffic visualized in single 3T3-L1 adipocytes by using ultra-fast microscopy. *Mol Biol. Cell*, **12**, 129-141.
- [173] Emoto, M., Langille, S.E., & Czech, M.P. (2001) A role for kinesin in insulin-stimulated GLUT4 glucose transporter translocation in 3T3-L1 adipocytes. *J. Biol. Chem.*, **276**, 10677-10682.
- [174] Fletcher, L.M., Welsh, G.I., Oatey, P.B., & Tavaré, J.M. (2000) Role for the microtubule cytoskeleton in GLUT4 vesicle trafficking and in the regulation of insulin-stimulated glucose uptake. *Biochem. J.*, **352**, 267-276.
- [175] Franke, W.W., Hergt, M., & Grund, C. (1987) Rearrangement of the vimentin cytoskeleton during adipose conversion: formation of an intermediate filament cage around lipid globules. *Cell*, **49**, 131-141.
- [176] Kanzaki, M., Furukawa, M., Raab, W., & Pessin, J.E. (2004) Phosphatidylinositol-4, 5-bisphosphate (PI4,5P2) regulates adipocyte actin dynamics and GLUT4 vesicle recycling. *J. Biol. Chem.*, **279**, 30622-30633.
- [177] Kanzaki, M. & Pessin, J.E. (2002) Caveolin-associated filamentous actin (Cav-actin) defines a novel F-actin structure in adipocytes. *J. Biol. Chem.*, **277**, 25867-25869.
- [178] Wang, Q., Bilan, P.J., Tsakiridis, T., Hinek, A., & Klip, A. (1998) Actin filaments participate in the relocalization of phosphatidylinositol3-kinase to glucose transporter-containing compartments and in the stimulation of glucose uptake in 3T3-L1 adipocytes. *Biochem. J.*, **331**, 917-928.
- [179] Guilherme, A., Emoto, M., Buxton, J.M., Bose, S., Sabini, R., Theurkauf, W.E., Leszyk, J., & Czech, M.P. (2000) Perinuclear localization and insulin responsiveness of GLUT4 requires cytoskeletal integrity in 3T3-L1 adipocytes. *J. Biol. Chem.*, **275**, 38151-38159.
- [180] Kanzaki, M. & Pessin, J.E. (2001) Insulin-stimulated GLUT4 translocation in adipocytes is dependent upon cortical actin remodeling. *J. Biol. Chem.*, **276**, 42436-42444.
- [181] Omata, W., Shibata, H., Li, L., Takata, K., & Kojima, I. (2000) Actin filaments play a critical role in insulin-induced exocytotic recruitment but not in endocytosis of GLUT4 in isolated rat adipocytes. *Biochem. J.*, **346**, 321-328.
- [182] Tsakiridis, T., Vranic, M., & Klip, A. (1994) Disassembly of the actin network inhibits insulin-dependent stimulation of glucose transport and prevents recruitment of glucose transporters to the plasma membrane. *J. Biol. Chem.*, **269**, 29934-29942.
- [183] Paul, D.S., Harmon, A.W., Winston, C.P., & Patel, Y.M. (2003) Calpain facilitates GLUT4 vesicle translocation during insulin-stimulated glucose uptake in adipocytes. *Biochem. J.*, **376**, 625-632.
- [184] Kanzaki, M., Watson, R.T., Khan, A.H., & Pessin, J.E. (2001) Insulin stimulates actin comet tails on intracellular GLUT4-containing compartments in differentiated 3T3L1 adipocytes. *J. Biol. Chem.*, **276**, 49331-49336.
- [185] Vollenweider, P., Martin, S.S., Haruta, T., Morris, A.J., Nelson, J.G., Cormont, M., Marchand-Brustel, Y., Rose, D.W., & Olefsky, J.M. (1997) The small guanosine triphosphate-binding protein Rab4 is involved in insulin-induced GLUT4 translocation and actin filament rearrangement in 3T3-L1 cells. *Endocrinology*, **138**, 4941-4949.
- [186] Martin, S.S., Haruta, T., Morris, A.J., Klippel, A., Williams, L.T., & Olefsky, J.M. (1996) Activated phosphatidylinositol 3-kinase is sufficient to mediate actin rearrangement and GLUT4 translocation in 3T3-L1 adipocytes. *J. Biol. Chem.*, **271**, 17605-17608.
- [187] Kanzaki, M., Watson, R.T., Hou, J.C., Stamnes, M., Saltiel, A.R., & Pessin, J.E. (2002) Small GTP-binding protein TC10 differentially regulates two distinct populations of filamentous actin in 3T3L1 adipocytes. *Mol Biol. Cell*, **13**, 2334-2346.