



Universiteit  
Leiden  
The Netherlands

## **New insights in mechanism, diagnosis and treatment of myocardial infarction**

Bergheanu, S.C.

### **Citation**

Bergheanu, S. C. (2011, April 21). *New insights in mechanism, diagnosis and treatment of myocardial infarction*. Retrieved from <https://hdl.handle.net/1887/17588>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/17588>

**Note:** To cite this publication please use the final published version (if applicable).

# CHAPTER

GENETIC DETERMINANTS OF  
ADVERSE OUTCOME (RESTENOSIS,  
MALAPPOSITION AND  
THROMBOSIS) AFTER  
STENT IMPLANTATION

# 8

## SUMMARY

Despite its unequivocal superiority compared with balloon angioplasty, coronary stenting did not abolish the restenosis problem and even brought along a completely new type of pathology. Bare-metal stents still associate with an approximate 20-30% in-stent restenosis rate and the need for repeat revascularization. Drug-eluting stents (which unfortunately did not completely prevent restenosis either) sometimes determine late-acquired stent malapposition in a significant number of patients. This is followed occasionally by a very serious event – stent thrombosis. Patient comorbidities, stent design, procedural characteristics and antiplatelet therapy influence the risk of poststenting complications. Research in the recent years has also revealed that individual genetic profile plays an important role in adverse outcome after stent implantation. This manuscript reviews the evidence of genetic variations associated with stent restenosis, late-acquired stent malapposition and stent thrombosis.

8

**Keywords:** coronary disease, stent, restenosis, malapposition, thrombosis, genes

Sandrin C Bergheanu<sup>1,2,4</sup>, Douwe Pons<sup>1,4</sup>, Ioannis Karalis<sup>1</sup>, Orçun Özsoy<sup>1,2</sup>, Jeffrey JW Verschuren<sup>1</sup>, Mark M Ewing<sup>1,3,4</sup>, Paul HA Quax<sup>3,4</sup> and J Wouter Jukema<sup>1,4,5,6,†</sup>

<sup>1</sup>Department of Cardiology, <sup>2</sup>Department of Clinical Epidemiology, <sup>3</sup>Department of Surgery, <sup>4</sup>Eindhoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, <sup>5</sup>Durrer Institute for Cardiogenetic Research, Amsterdam and <sup>6</sup>Interuniversity Cardiology Institute the Netherlands (ICIN), Utrecht, The Netherlands.

*Interv Cardiol.* 2010;2:147-157.


The era of percutaneous coronary intervention (PCI) began with the first balloon angioplasty performed by Andreas Gruentzig in 1977 [1]. Although this technique provided impressive immediate results, mid and long term follow up was characterized by high restenosis rates and need for repeat revascularization [1,2]. Evolving our techniques, bare-metal prosthetic devices (stents) were designed to act as a barrier against intima growth and recoil, assuring long-time patency of the coronary vessel. In 1986 Sigwart and Puel implanted the first coronary stent in a human patient [3]. Superior to balloon angioplasty alone (32-42% restenosis rate), bare-metal stent (BMS) implantation remains however vulnerable to restenosis (22-32% of cases) [4,5,6] and often requires re-intervention. Drug-eluting stents (DES) were conceived as an answer to this problem. They, for the majority, consist of a metallic platform covered with a combination of polymer and cellular proliferation inhibitor. The antiproliferative agent is gradually released in the arterial wall at the site of stent deployment preventing restenosis. The first successful DES trials were with sirolimus stents and led to their approval for use in 2002 in Europe and 2003 in USA [7,8]. Currently, other DES based upon paclitaxel, everolimus, zotarolimus, biolimus and tacrolimus are available. DES have successfully achieved their task of preventing restenosis but the experience of the last years revealed an increased incidence of stent malapposition and stent thrombosis associated with their use [9]. The aim of this article is to briefly present incidence and mechanisms of 1) stent restenosis, 2) stent malapposition and 3) stent thrombosis and to focus on potential genetic factors related to these complications. The majority of available data is retrieved from candidate gene approach studies, limiting thus the results to specific pre-targeted pathophysiologic sequences. Further novel pharmacogenomic approaches such as GWAS (genome wide association studies) may be able to identify new genetic factors for a better prediction of outcome after coronary stent deployment.

## IN-STENT RESTENOSIS

In-stent restenosis (ISR) is defined angiographically when neo-formation tissue represents more than 50% of the lumen diameter at the site of the stented vessel (Figure 1).

The clinical confirmation of ISR is the recurrence of angina pectoris, which further requires intervention: TLR (target lesion revascularization) or TVR (target vessel revascularization). Although the severity of angiographic stenosis correlates with the need for TLR, half of the patients with angiographically confirmed ISR do not manifest clinical complains [6,10]. For this reason authors generally prefer to conduct their research in relation to angiographically documented ISR when an insight in the mechanism of restenosis is aimed, while studies comparing different stents are in relation to clinically-driven TLR or TVR.

ISR is the result of in-stent cellular proliferation and migration along with extracellular matrix accumulation [11]. Classic predictors of angiographic ISR (both in



**Figure 1.** In-stent restenosis. a) Angiographically documented in-stent restenosis; b) IVUS documented in-stent restenosis. 1 – neointima; 2 – stent contour; 3 – vessel contour.

BMS and DES) include diabetes, renal failure, lesion length, reference vessel diameter and post-intervention lumen area [12,13]. Inflammation plays a pivotal role in ISR and it is triggered by the vascular injury during the stent deployment and by the presence of stent struts within the vessel wall [14,15]. Together with inflammation, major contributors are smooth muscle cell migration and proliferation but the process of restenosis involves many different cell-types, among which platelets and endothelial cells, and is also characterized by thrombus formation and to a lesser extent by matrix remodelling.

## GENETIC FACTORS RELATED TO IN-STENT RESTENOSIS

### Genetic variations in thrombus formation

In principle, any vascular intervention initiates the formation of a thrombus. Initial studies have shown associations of only a few polymorphisms in the hemostatic system with the risk for adverse events following a PCI. These early reports showed significant associations of the PLA1/A2 polymorphism with acute stent-thrombosis and coronary restenosis [16,17]. However, other studies in this field could not confirm these associations [18,19]. On grounds of the hypothesis that carriers of the PLA2 allele have a more intense binding of fibrinogen and vitronectin and thus a higher risk of platelet-rich white thrombus formation, the PLA2 allele can be expected to lead to an increased risk for acute stent thrombosis. However, as platelet inhibition by IIb/IIIa and P2Y12 antagonists does reduce acute stent thrombosis, but not in-stent restenosis rates [20], thrombus formation is probably not a main player in the development of restenosis. This hypothesis is further confirmed by findings showing that especially the strong pro-thrombotic genetic risk factors for venous thrombosis do not increase the risk for restenosis [21]. Moreover, results from the GENDER study [21] have shown

that the Factor V Leiden polymorphism (a well-known prothrombotic risk factor) was even found to reduce the risk for restenosis after PCI. A total of 3104 consecutive patients with stable angina pectoris or non-STEMI, of whom 2309 (74.4%) received stents, were included [21]. The factor V (1691 G>A or factor V Leiden) amino acid substitution was associated with a decreased risk of TVR (HR=0.41, 95%CI 0.19–0.86). The Factor V allele, which is known to lead to increased activation of protein C, might therefore influence restenosis risk by mechanisms not involved in coagulation, but in processes that have a more prominent role in neointimal growth, such as inflammation. Even though in another study of the same patient sample, associations were found between P2Y12 receptor haplotypes and restenosis [22], fewer and smaller effects were present in the stented subgroup. The decrease of the effects in this group could be due to inhibition of this receptor by clopidogrel (although several studies [23–26] failed to demonstrate a functional role of the P2Y12 receptor polymorphism in patients receiving dual antiplatelet therapy). Therefore, the genetic variation in this receptor, and also in many other genes with a role in the hemostatic system, may have been more important at a time in which not every patient was receiving a stent and concomitant platelet inhibition.

The 4G/4G genotype of the PAI-1 4G/5G polymorphism determines higher PAI-1 levels in plasma [27–29] and tissue [30–32]. The PAI-1 4G allele was associated with an increased risk of restenosis after PCI in the GENDER study [21]. When compared to 5G/5G homozygotes, heterozygous patients were at higher risk for clinically-driven TVR (HR=1.46, 95%CI 1.05–2.03), whereas patients with the 4G/4G genotype had an even further increased risk (HR=1.69, 95%CI 1.19 – 2.41). Although one smaller study could not confirm this association [33], many reports found a positive correlation between post-PCI PAI-1 levels or activity and restenosis [34,35]. Nevertheless, PAI-1 has a diverse role in several processes involved in restenosis, also in inflammation and proliferation [36]. Even if the 4G allele would increase the risk for restenosis, this could be mediated by a mechanism not related to fibrinolysis inhibition. Taking these findings together, we suggest that coagulation is not a main determinant of the long-term process that leads to restenosis.

## Genetic variations in inflammatory factors

Early studies investigating the role of genetics in restenosis showed associations between variants in genes encoding cytokines [37] and selectins [38] – important mediators of inflammation – and suggested a role for inflammation in restenosis. One of these studies was performed by Kastrati et al.[37], and included 1850 consecutive stented patients. They demonstrated a protective effect of allele 2 of a polymorphism in exon 2 of the gene encoding the IL-1 receptor antagonist (IL-1ra), an anti-inflammatory interleukin, on both angiographic and clinical restenosis (OR=0.78, 95%CI 0.63–0.97 and OR=0.73, 95%CI 0.58–0.92, respectively). Monraats et al. have further established the important role of inflammatory genes in the development of restenosis. In the GENDER study, the rare alleles of the -260 C/T polymorphism in the

CD14 gene, the 117 Ile/Thr polymorphism in the colony stimulating factor 2 gene (also known as granulocyte-macrophage colony stimulating factor, GM-CSF) and the -1328 G/A polymorphism in the eotaxin gene were associated with decreased risk of TVR [39]. Eotaxin is a chemokine which selectively recruits eosinophils and was previously reported to be elevated in plasma of patients with advanced atherosclerosis. After coronary interventions, eotaxin levels increase and remain high for at least 24 hours but no longer than 3 month [40].

Furthermore, the variant alleles of two promoter polymorphisms in the Tumor Necrosis Factor alpha (TNF- $\alpha$ ) gene have been shown to protect against the development of restenosis [41]. Stented patients with the -238A/A genotype (HR=0.44, 95%CI 0.23-0.83) and patients with the -1031C/C genotype (HR=0.72, 95%CI 0.52-1.00) needed TVR less frequently. Several other inflammatory genes were shown to be involved in the process of restenosis in this cohort, among which interleukin 10 and caspase-1 (IL-1 $\beta$  converting enzyme) [42,43]. All these findings support the hypothesis that restenosis is largely (albeit not solely) determined by inflammation.

## Genes involved in smooth muscle cell proliferation

Stents specifically aiming to inhibit inflammation (dexamethasone eluting stents) were not proven as effective as stents inhibiting both inflammation and cell proliferation [44]. Despite the fact that restenosis is mainly determined by proliferation and migration of vascular smooth muscle cells (VSMCs), relatively few studies investigated genes involved in proliferation, such as cell-cycle regulatory genes. A recent important finding in this field by Van Tiel et al.[45] was an association between the -838 G/A polymorphism in the cyclin-dependent kinase inhibitor p27(kip1) (a key regulator of SMC proliferation) with ISR. Three polymorphisms concerning the p27(kip1) gene (-838C>A; -79C>T; +326G>T) were determined in a cohort of 715 patients undergoing coronary angioplasty and stent placement. Patients with the p27(kip1) -838AA genotype had a decreased risk of ISR (HR=0.28, 95%CI 0.10-0.77). This finding was replicated in another cohort study of 2309 patients (HR= 0.61, 95%CI 0.40-0.93). The -838 A allele corresponded to enhanced promoter activity which in turn may explain decreased SMC proliferation.


## Genetic variations in matrix metalloproteinases

Matrix metalloproteinases (MMPs) are Zn<sup>2+</sup>-requiring proteases capable of degrading a variety of extracellular matrix components. Due to their significance in vascular remodeling, MMPs are suspected to play an (important) role in the pathogenesis of atherosclerosis and restenosis [46]. Especially MMP2, MMP3 and MMP9 are potential players in the process of restenosis after PCI. MMP2 and MMP9 (the gelatinases) are produced by vascular VSMCs and degrade basement membrane components and other matrix proteins to allow migration and proliferation of vascular smooth muscle cells (VSMCs) [47]. They are upregulated and activated in VSMCs during intima formation in many different animal models for restenosis involving balloon

angioplasty [47]. An increase in MMP2 levels and activity was demonstrated in human coronary sinus blood samples 4 and 24 hours after elective coronary angioplasty [48]. This small study, in which only 21 of 47 patients were stented, also showed an association between MMP2 levels and restenosis. MMP3 (stromelysin-1) expression has been found to be related to plaque-instability in pathological studies [49]. MMP3 reduces the matrix content of the vascular wall and is therefore expected to protect against restenosis [49]. Functional studies have shown that the MMP3 -1612 5A/6A promoter polymorphism is associated with altered MMP3 expression. Carriers of the 6A/6A genotype were found to have a reduced MMP3 expression [50-53] and were at increased risk of developing restenosis in a subset of the REGRESS study, in which stents were not yet frequently used [54], and in two other studies with luminal narrowing after plain balloon angioplasty [55,56]. However, an association between the MMP3 5A/6A polymorphism could not be confirmed in a study which included 217 stented patients. Unpublished results from the GENDER study indeed show no association between this polymorphism and clinical restenosis in stented patients. Therefore, even though matrix formation is an important process in the development of restenosis, variations in genes involved in matrix remodeling were infrequently investigated or studies yielding negative results and were not published.

## STENT MALAPPPOSITION

Stent malapposition (SM), commonly detected by intravascular ultrasonography (IVUS), represents a separation of the stent struts from the intimal surface of the arterial wall (in the absence of a side branch) with evidence of blood speckles behind the struts [57] (Figure 2a).



**Figure 2.** Stent malapposition and thrombosis. a) Intravascular ultrasound documented stent malapposition; b) angiographically documented stent thrombosis. 1 – lumen contour behind stent struts; 2 – vessel contour; 3 – stent thrombosis.



SM may be acute (present immediately after implantation), persistent (present both immediately after implantation and at follow-up) or late-acquired (present only at follow-up). Acute and persistent SM are mainly procedure-driven while late-acquired stent malapposition (LASM) is a consequence of positive remodelling of the vessel wall and and/or of plaque volume decrease behind the stent (including clot lysis or plaque regression) [58-62]. The main repercussion of late SM (persistent or acquired) is stent thrombosis (ST) [9]. Independent predictors of LASM include lesion length, unstable angina, absence of diabetes and primary stenting in acute MI [63,59]. The risk of LASM in patients with DES is approximately 4 times higher compared to those with BMS [9]. This is due to the fact that in BMS, hypersensitivity to the metallic stent is mostly associated with restenosis, whereas in DES, the hypersensitivity to the metallic stent, the polymer or to the drug is associated with positive remodelling and excessive inflammation in the vessel wall [64].

## 8

### GENETIC FACTORS RELATED TO STENT MALAPPOSITION

We have previously investigated 7 polymorphisms (involved in inflammatory processes and related to restenosis) on the risk of LASM in SES patients [65]. In total, 104 STEMI patients from the MISSION! intervention study [62] were genotyped for the caspase-1 5352 G/A, eotaxin 1382 A/G, CD14 260 A/G, colony stimulating factor 2 1943 C/T, IL10 -1117 C/T, IL10 4251 C/T and the TNF- $\alpha$  1211 C/T polymorphisms. LASM occurred in 26/104 (25%) of patients. We found a significantly higher risk for LASM in patients carrying the caspase-1 (CASP1) 5352 A allele (RR= 2.32, 95% CI 1.22-4.42). In addition, mean neointimal growth was significantly lower in patients carrying this LASM risk allele (1.6 vs 4.1%,  $p=0.014$ ). The other 6 polymorphisms related to inflammation were not significantly related to the risk of LASM. Given the limited number of patients included in the study, similar reports are needed to confirm our findings. Moreover, a direct relation between the CASP1 5352 A allele and the risk of ST was not investigated. To our knowledge, no other studies yet scrutinized the role of genetic variations in LASM.

### STENT THROMBOSIS

Stent thrombosis (ST) (Figure 2b) is a complication which occurs in 0.8-2% of patients undergoing PCI and is associated with large MI and death [66]. ST is categorized into “acute” ST (within 24 hours from stent implantation), “subacute” ST (within 1 – 30 days from stent implantation), “late” ST (within 30 days – 1 year) and “very late” ST (> 1 year after stent implantation). Subacute and acute ST are classically related to procedure parameters such as stent underdeployment (acute SM) [67,68] or procedure related complications such as coronary dissections [69,70]. In contrast, (very) late ST appears to be an active phenomenon associated with late SM

(persistent or acquired) [9,71], stent type [9] duration of dual anti-platelet therapy [66] and inflammation [58]. Gene variations in the platelet aggregation pathway, responsiveness to clopidogrel or presence of inherited thrombophilic disorders were associated with both acute and late ST.

## GENETIC FACTORS RELATED TO STENT THROMBOSIS

### Platelet receptor gene polymorphism

Platelet aggregation involves the binding of fibrinogen to the glycoprotein (GP) IIb/IIIa receptor on the platelet surface. One polymorphism of the GP IIIa gene (PLA1/A2 or HPA – 1a/1b) has been related to the inherited risk of coronary thrombosis [72]. Of importance, the same polymorphism had no influence on the degree of myocardial salvage achieved in 133 acute MI patients undergoing coronary stenting and abciximab administration [73]. The PLA2 polymorphism is a substitution of cytosine for thymidine at position 1565 in exon 2. Walter et al. [74] investigated the association of PLA2 allele with acute and subacute stent thrombosis in 318 consecutive BMS patients stented for coronary dissection, acute occlusion or high residual restenosis after PTCA lesions in by-pass grafts, and restenotic lesions. They found that patients with the PLA2 allele had an increased risk of stent thrombosis compared with patients homozygous for PLA1 (OR=5.26, 95%CI 1.55-17.85). Kastrati et al. [75] confirmed these findings partially in their prospective study including 1759 patients with stable and unstable angina pectoris. No difference was seen at 30 days after stent placement in terms of ST or a composite end-point of death, MI or urgent revascularization between PLA1/A1 and PLA1/A2 carriers. However, the incidence of ST and the composite end-point were higher in the PLA2 homozygotes versus PLA1 homozygotes (8.7% vs. 1.7%,  $p=0.002$  and 13.0% vs. 5.4%,  $p=0.06$ , respectively).

More recently, Sucker et al. [76] assessed the relevance of prothrombotic platelet receptor polymorphisms for the onset of coronary stent thrombosis in 316 patients. They compared the prevalence of GP Iba, GP IIb, GP IIIa (including PLA1/A2) and GP Ia prothrombotic polymorphisms in patients with coronary stent thrombosis occurring in the first 6 months after stent implantation and healthy control subjects. Carriers of the above mentioned prothrombotic versions did not appear to be at any increased risk for stent thrombosis. Selection of patients (differences in number of elective and acute stent implantations) and the treatment of more complex coronary lesions in the latter study or the limited power might explain these discrepancies [76]. Angiolillo et al. [77] have investigated the differential platelet sensitivity between PLA1 homozygotes and PLA2 carriers in 38 patients undergoing coronary stent implantation and receiving a 300 mg clopidogrel loading dose. They have shown that PLA2 carriers have a lower inhibition of platelet reactivity following the standard clopidogrel loading dose, which might finally lead to stent thrombosis.

## Genetic variations in response to clopidogrel

In current practice, patients undergoing PCI and stent deployment are given 300-600 mg clopidogrel as a loading dose followed by 1 year dual anti-platelet therapy (aspirin 80-325 mg and clopidogrel 75 mg daily) and continued with life-long aspirin intake.

A good responsiveness to clopidogrel is therefore crucial in order to prevent thrombotic events after stent deployment.

Clopidogrel is an inactive prodrug which requires a two-step oxidation by the hepatic cytochrome P450 (CYP) enzymes to transform into an active metabolite which further inhibits the ADP P2Y<sub>12</sub> receptor producing the anti-aggregation effect. The genes encoding the CYP enzymes are polymorphic and several variants were related to a decreased catalytic activity and subsequent attenuated effect of the drug.

The CYP3A5 gene has a functional polymorphism which includes the expressor (\*1) and non-expressor (\*3) alleles [78,79]. Suh et al. [79] compared clinical outcome in 348 patients (with stable angina, unstable angina or non-STEMI) who had PCI with BMS implantation. Antiplatelet therapy consisted of aspirin (100-300 mg daily, prescribed indefinitely) and clopidogrel (75 mg daily after 300 mg loading dose) administered for at least 4 weeks after the procedure. Atherothrombotic events (a composite of cardiac death, MI and non-hemorrhagic stroke) occurred more frequently within 6 months after stent implantation among the patients with the non-expressor genotype than among those with the expressor genotype (14/193 vs. 3/155,  $p=0.023$ ). Moreover, the CYP3A5 polymorphism was a predictor of atherothrombotic events in clopidogrel users.

These findings are interesting especially since a number of studies (which did not aim at clinical end-points) found no association between the CYP3A5 variants and clopidogrel response and/or residual platelet aggregation (RPA) [80-82] nor did a number of studies with clinical end-points [83,84].

Trenk et al. [85] investigated whether the CYP2C19 681G>A \*2 polymorphism was associated with high (>14%) RPA on clopidogrel and whether high on-clopidogrel RPA affects clinical outcome after elective coronary stent placement. RPA was assessed in 797 consecutive patients after a 600 mg loading dose and after the first 75 mg maintenance dose of clopidogrel before discharge. Patients were followed-up for 1 year. Between the \*2 carriers and \*1/\*1 carriers (wild-type) the authors found significant ( $p<0.001$ ) differences in the proportion of patients with RPA>14%, both after loading (62.4% vs. 43.4%) and at pre-discharge (41.3% vs. 22.5%). RPA >14% at discharge was associated with a 3-fold increase (95%CI 1.4-6.8,  $p=0.004$ ) in the 1-year incidence of death and myocardial infarction. However, authors could not show a direct relation between the CYP2C19\*2 allele and clinical outcome.

This relation was demonstrated by Giusti et al. [86] in a subanalysis of the RECLOSE trial. The role of the CYP2C19\*2 polymorphism in the occurrence of DES ST (definite or probable) or the composite end-point of ST (definite or probable) and cardiac mortality within 6-month follow-up was assessed in 772 patients undergoing PCI and receiving either sirolimus or paclitaxel DES. Patients with ACS

and STEMI were included as well as patients with left main disease, chronic total occlusions, bifurcation lesions or diffuse disease. All patients received aspirin (325 mg) and a loading dose of clopidogrel 600 mg before the procedure followed by a maintenance dose of clopidogrel 75 mg and aspirin 325 mg daily. Patients with ST or ST and cardiac mortality end-point had a higher prevalence of the \*2 allele (54.1% vs. 31.3%;  $p=0.025$  and 51.7% vs. 31.2%;  $p=0.020$ , respectively). At multivariate logistic regression analysis, the CYP2C19\*2 allele was an independent risk factor for ST (OR=3.43, 95%CI 1.01-12.78,  $p=0.047$ ) and ST and cardiac mortality (OR=2.7, 95%CI 1.00-8.42,  $p=0.049$ ).

Mega et al. [83] reconfirmed these findings on long term assessment of patients from TRITON-TIMI 38 study. A number of 1389 patients treated with clopidogrel who underwent PCI and stenting were followed-up for 15 months. Patients were initially admitted with non-STEMI (71%) and STEMI (29%). They received a 300 mg clopidogrel loading dose, followed by 75 mg daily maintenance dose for up to 15 months. For the CYP2C19, the presence of at least one copy of the \*2 allele was associated with a higher rate of composite death from cardiovascular causes, non-fatal MI, non-fatal stroke (HR=1.42, 95%CI 0.98-2.05) and of definite/probable ST (HR=3.33, 95%CI 1.28-8.62) than did non-carriers.

Sibbing et al. [87] assessed the role of the mutant \*2 allele of the CYP2C19 polymorphism on the 30-day incidence of definite ST in 2485 consecutive patients undergoing coronary stent placement. There are a number of differences with regard to the previous study [83]: (1) STEMI patients were excluded, (2) the end-point was acute and subacute definite ST and (3) patients received 600 mg clopidogrel loading dose. DES were used in 25% and BMS in 75% of the patients. Of the patients studied, 73% were CYP2C19 wild-type homozygotes (\*1/\*1) and 27% carried at least one of the \*2 allele. The cumulative 30-day incidence of ST was significantly higher in CYP2C19\*2 allele carriers vs. wild-type homozygotes (1.5% vs. 0.4%, HR=3.81, 95%CI 1.45-10.02,  $P=0.006$ ). The risk of ST was highest (2.1%) in patients carrying the CYP2C19 \*2/\*2 genotype ( $p=0.002$ ).

Recently, Collet et al. [88] demonstrated the role of the CYP2C19\*2 allele in 259 young patients (aged <45 years) who survived a first MI and received clopidogrel treatment for at least a month. The primary endpoint was a composite of death, MI, and urgent coronary revascularization occurring during exposure to clopidogrel. The secondary endpoint was angiography-documented stent thrombosis. Median clopidogrel treatment duration was approximately one year. The primary endpoint occurred more frequently in carriers than in non-carriers (15 vs. 11 events; HR=3.69, 95%CI 1.69-8.05,  $P=0.0005$ ), as did stent thrombosis (8 vs. 4 events; HR=6.02, 95%CI 1.81-20.04,  $P=0.0009$ ). The effect of the CYP2C19\*2 genetic variant persisted from 6 months after clopidogrel initiation up to the end of follow-up (HR=3.00, 95%CI 1.27-7.10,  $p=0.009$ ). The CYP2C19\*2 genetic variant appeared the only independent predictor of cardiovascular events (HR=4.04, 95%CI 1.81-9.02,  $P=0.0006$ ).

In a study [84] of 2208 patients presenting with acute MI (among which 1535 underwent PCI), patients carrying any two CYP2C19 loss-of-function alleles (\*2, \*3, \*4, or \*5), had a higher rate of death from any cause, nonfatal stroke, or myocardial infarction during 1 year of follow-up than patients with none (21.5% vs. 13.3%; adjusted HR=1.98; 95%CI 1.10-3.58). Among the patients who underwent PCI during hospitalization, the rate of cardiovascular events among carriers of CYP2C19 loss-of-function alleles was 3.58 (95%CI 1.71-7.51).times higher than among those with none.

For the development of a risk score for better prediction of RPA, Geisler et al. [81] analyzed the CYP2C19\*2 genotype and previously identified non-genetic risk factors (age >65 years, type 2 diabetes mellitus, decreased left ventricular function, renal failure and acute coronary syndrome). They demonstrated a significant correlation of the non-genetic factors ( $\chi^2 = 5.32$ ;  $P = 0.021$ ) and CYP2C19\*2 ( $\chi^2 = 21.31$ ;  $P < 0.0001$ ) with high RPA, and the highest association for the combination of both ( $\chi^2 = 25.85$ ;  $P < 0.0001$ ). This was the first study to show that prediction of clopidogrel responsiveness may substantially be improved by adding CYP2C19\*2 genotype to non-genetic risk factors. The important influence of the CYP2C19\*2 genotype over platelet function and cardiovascular outcomes was recently confirmed by Shuldiner et al. [89] in the first GWAS paper identifying CYP2C19 as a candidate gene. In the Pharmacogenomics of Antiplatelet Intervention (PAPI) Study, clopidogrel was administered for 7 days to 429 healthy individuals and the response was measured by ex vivo platelet aggregometry. A GWAS was performed followed by genotyping the loss-of-function cytochrome CYP2C19\*2 variant. The relation between CYP2C19\*2 genotype and platelet aggregation was replicated in 227 clopidogrel-treated patients undergoing PCI ( $P = 0.02$ ). Patients with the CYP2C19\*2 variant were more likely (20.9% vs 10.0%) to have a cardiovascular ischemic event or death during 1 year of follow-up (HR=2.42, 95%CI 1.18-4.99,  $P = 0.02$ ).

## Factor V Leiden mutation

Factor V Leiden is the most common inherited thrombophilic disorder, resulting from a single mutation (1691 G>A) in the factor V gene. Individual heterozygous for this mutation are at increased risk for venous thrombosis, and in homozygous the risk becomes extremely high. Although conceivable, there is only one case report to document a possible relation between a factor V Leiden heterozygous patient and stent thrombosis (simultaneous occlusion of two stents, one in left anterior descending artery and one in the right coronary artery at 4 days after implantation in a patient receiving standard dual anti-platelet therapy) [90]. Further larger studies are therefore needed before factor V Leiden may be linked to ST.

## LIMITATIONS

Many studies have managed to identify genes and polymorphisms involved in the post-stenting outcome after scrutinizing various plausible pathophysiologic mechanisms. However, to predict an accurate scale of adverse effects, an interaction assessment between genetic, non genetic (traditional risk factors) as well as epigenetic factors is of extreme importance. This information remains momentarily scarce.

Also of importance, findings from certain studies cannot sometimes be confirmed by other studies. This is largely explained by variation in study settings and therefore the replication of findings in independent studies needs to be further emphasized.

The candidate gene approach used to date in the majority of investigations narrows the results to specific areas of interest.

## CONCLUSION

In-stent restenosis and stent thrombosis remain important limitations of the current PCI practice. Besides the procedure-related risk factors and medication, solid evidence shows that patient's own response to stent implantation influences the outcome. Individual genetic response involves inflammation, cellular proliferation, platelet receptors and drug metabolism pathways. A better understanding of the stent pathology has led to the identification of new important genes and genetic polymorphisms. They may help us better identify the vulnerable patients who need extraordinary therapeutic measures. Conversely, genetic-epidemiologic studies have identified genes which subsequently have revealed important pathophysiologic mechanisms.

## FUTURE PERSPECTIVES

The speed by which new genes are being related to stent pathology is matched by the speed of new developments in stent technology and medication. Novel pharmacogenomic approaches (e.g., GWAS, 1000 genome project) may help to identify unknown genetic factors for a better prediction of outcome after stent implantation [91].

It is however difficult to predict whether screening for established polymorphisms will prove, in the future, a cost-effective method for improved stent type selection or medication in the daily routine.

The classic stents appear to be rapidly being replaced by new and complex body-polymer-drug constructs that address most, if not all, of the current problems. The new generation of stents may appear capable of modulating local inflammation, to permit a good re-endothelization, to prevent stent thrombosis, to reduce the duration of anti-platelet medication and, if necessary, even to degrade after local healing is achieved.

New drugs such as prasugrel, ticagrelor and cangrelor seem to effectively inhibit platelet aggregation with little or no interindividual response variability. The combination of lessons learned from genetic and pathophysiologic studies, the newly available resources (e.g., stents, antiplatelet drugs and imaging) and refined implantation techniques will definitely improve PCI performances and extend its use.

## EXECUTIVE SUMMARY

Genetic variants associated with an increased or decreased risk of in-stent restenosis (ISR) and stent thrombosis (ST)

### In-stent restenosis (ISR)

- » Genetic variations in thrombus formation
  - Associated with decreased risk:
    - Factor V 1691G>A (factor V Leiden) amino acid substitution
  - Associated with increased risk:
    - 4G allele of the PAI-1 4G/5G polymorphism
- » Genetic variations in inflammatory factors
  - Associated with decreased risk:
    - \*2 allele of the IL-1ra gene
    - T/T genotype of the CD14-260 C/T polymorphism
    - Thr allele of the CSF2-117 Ile/Thr polymorphism
    - A allele of the CCL 11 (Eotaxin) 1328 G/A polymorphism
    - A/A genotype of the TNF -238 G/A polymorphism
    - C/C genotype of the TNF -1031 T/C polymorphism
  - Associated with increased risk:
    - A/A genotype of the IL-10 -2849 G/A polymorphism
    - A/A genotype of the IL-10 -1082 G/A polymorphism
    - G/G genotype of the IL-10 +4259 A/G polymorphism
    - A/A genotype of the Caspase-1 5352 G/A polymorphism
- » Genes involved in smooth muscle cell proliferation
  - Associated with decreased risk:
    - A/A genotype of the p27(kip1)-838G/A polymorphism

### Stent thrombosis (ST)

- » Platelet receptor gene polymorphism
  - Associated with increased risk:
    - PLA2 allele of the GP IIIa PLA1/A2
- » Genetic variations in response to clopidogrel
  - Associated with increased risk:
    - \*3 allele of the CYP3A5 gene (encodes hepatic cytochrome P450 CYP enzymes)
    - \*2 allele of the CYP2C19 gene (encodes hepatic cytochrome P450 CYP enzymes)

## FINANCIAL DISCLOSURE/ACKNOWLEDGEMENTS

JWJ is an established clinical investigator of the Netherlands Heart Foundation (2001D032) and project leader of the HP 7 – pharmacogenetic EU project HEALTH-F2-2007 223004 PHASE. Authors have no financial disclosures to report

## REFERENCE LIST

1. Gruntzig AR, Senning A, Siegenthaler WE: Nonoperative dilatation of coronary-artery stenosis: percutaneous transluminal coronary angioplasty. *N.Engl.J.Med.* 301(2), 61-68 (1979).
2. Holmes DR, Jr., Vlietstra RE, Smith HC et al: Restenosis after percutaneous transluminal coronary angioplasty (PTCA): a report from the PTCA Registry of the National Heart, Lung, and Blood Institute. *Am.J.Cardiol.* 53(12), 77C-81C (1984).
3. Sigwart U, Puel J, Mirkovitch V, Joffre F, Kappenberger L: Intravascular stents to prevent occlusion and restenosis after transluminal angioplasty. *N.Engl.J.Med.* 316(12), 701-706 (1987).
4. Fischman DL, Leon MB, Baim DS et al: A randomized comparison of coronary-stent placement and balloon angioplasty in the treatment of coronary artery disease. Stent Restenosis Study Investigators. *N.Engl.J.Med.* 331(8), 496-501 (1994).
5. Serruys PW, de Jaegere P, Kiemeneij F et al: A comparison of balloon-expandable-stent implantation with balloon angioplasty in patients with coronary artery disease. Benestent Study Group. *N.Engl.J.Med.* 331(8), 489-495 (1994).
6. Cutlip DE, Chhabra AG, Baim DS et al: Beyond restenosis: five-year clinical outcomes from second-generation coronary stent trials. *Circulation* 110(10), 1226-1230 (2004).
7. Moses JW, Leon MB, Popma JJ et al: Sirolimus-eluting stents versus standard stents in patients with stenosis in a native coronary artery. *N.Engl.J.Med.* 349(14), 1315-1323 (2003).
8. Serruys PW, Kutryk MJ, Ong AT: Coronary-artery stents. *N.Engl.J.Med.* 354(5), 483-495 (2006).
9. Hassan AK, Bergheanu SC, Stijnen T et al: Late stent malapposition risk is higher after drug-eluting stent compared with bare-metal stent implantation and associates with late stent thrombosis. *Eur.Heart J.* (2009).
10. Scott NA: Restenosis following implantation of bare metal coronary stents: pathophysiology and pathways involved in the vascular response to injury. *Adv Drug Deliv Rev.* 58(3), 358-376 (2006).
11. Hoffmann R, Mintz GS, Dussaillant GR et al: Patterns and mechanisms of in-stent restenosis. A serial intravascular ultrasound study. *Circulation* 94(6), 1247-1254 (1996).
12. Rathore S, Terashima M, Katoh O et al: Predictors of angiographic restenosis after drug eluting stents in the coronary arteries: contemporary practice in real world patients. *EuroIntervention.* 5(3), 349-354 (2009).
13. Bhargava B, Karthikeyan G, Abizaid AS, Mehran R: New approaches to preventing restenosis. *BMJ* 327(7409), 274-279 (2003).
14. Farb A, Sangiorgi G, Carter AJ et al: Pathology of acute and chronic coronary stenting in humans. *Circulation* 99(1), 44-52 (1999).
15. Farb A, Weber DK, Kolodgie FD, Burke AP, Virmani R: Morphological predictors of restenosis after coronary stenting in humans. *Circulation* 105(25), 2974-2980 (2002).
16. Kastrati A, Schomig A, Seyfarth M et al: PIA polymorphism of platelet glycoprotein IIIa and risk of restenosis after coronary stent placement. *Circulation* 99(8), 1005-1010 (1999).
17. Wheeler GL, Braden GA, Bray PF, Marciniak SJ, Mascelli MA, Sane DC: Reduced inhibition by abciximab in



- platelets with the PIA2 polymorphism. *Am.Heart J.* 143(1), 76-82 (2002).
18. Mamotte CD, van Bockxmeer FM, Taylor RR: Pla1/a2 polymorphism of glycoprotein IIIa and risk of coronary artery disease and restenosis following coronary angioplasty. *Am.J.Cardiol.* 82(1), 13-16 (1998).
  19. Volzke H, Grimm R, Robinson DM *et al*: Candidate genetic markers and the risk of restenosis after coronary angioplasty. *Clin.Sci.(Lond)* 106(1), 35-42 (2004).
  20. Acute platelet inhibition with abciximab does not reduce in-stent restenosis (ERASER study). The ERASER Investigators. *Circulation* 100(8), 799-806 (1999).
  21. Pons D, Monraats PS, de Maat MP *et al*: The influence of established genetic variation in the haemostatic system on clinical restenosis after percutaneous coronary interventions. *Thromb. Haemost.* 98(6), 1323-1328 (2007).
  22. Rudez G, Pons D, Leebeek F *et al*: Platelet receptor P2RY12 haplotypes predict restenosis after percutaneous coronary interventions. *Hum.Mutat.* 29(3), 375-380 (2008).
  23. Angiolillo DJ, Fernandez-Ortiz A, Bernardo E *et al*: Lack of association between the P2Y12 receptor gene polymorphism and platelet response to clopidogrel in patients with coronary artery disease. *Thromb.Res.* 116(6), 491-497 (2005).
  24. Giusti B, Gori AM, Marcucci R *et al*: Cytochrome P450 2C19 loss-of-function polymorphism, but not CYP3A4 IVS10 + 12G/A and P2Y12 T744C polymorphisms, is associated with response variability to dual antiplatelet treatment in high-risk vascular patients. *Pharmacogenet. Genomics* 17(12), 1057-1064 (2007).
  25. Lev EI, Patel RT, Guthikonda S, Lopez D, Bray PF, Kleiman NS: Genetic polymorphisms of the platelet receptors P2Y(12), P2Y(1) and GPIIb/IIIa and response to aspirin and clopidogrel. *Thromb.Res.* 119(3), 355-360 (2007).
  26. von Beckerath N, von Beckerath O, Koch W, Eichinger M, Schomig A, Kastrati A: P2Y12 gene H2 haplotype is not associated with increased adenosine diphosphate-induced platelet aggregation after initiation of clopidogrel therapy with a high loading dose. *Blood Coagul.Fibrinolysis* 16(3), 199-204 (2005).
  27. Kathiresan S, Gabriel SB, Yang Q *et al*: Comprehensive survey of common genetic variation at the plasminogen activator inhibitor-1 locus and relations to circulating plasminogen activator inhibitor-1 levels. *Circulation* 112(12), 1728-1735 (2005).
  28. Diamanti-Kandarakis E, Palioniko G, Alexandraki K, Bergiele A, Koutsouba T, Bartzis M: The prevalence of 4G5G polymorphism of plasminogen activator inhibitor-1 (PAI-1) gene in polycystic ovarian syndrome and its association with plasma PAI-1 levels. *Eur.J.Endocrinol.* 150(6), 793-798 (2004).
  29. Asselbergs FW, Williams SM, Hebert PR *et al*: The gender-specific role of polymorphisms from the fibrinolytic, renin-angiotensin, and bradykinin systems in determining plasma t-PA and PAI-1 levels. *Thromb.Haemost.* 96(4), 471-477 (2006).
  30. Castello R, Espana F, Vazquez C *et al*: Plasminogen activator inhibitor-1 4G/5G polymorphism in breast cancer patients and its association with tissue PAI-1 levels and tumor severity. *Thromb. Res.* 117(5), 487-492 (2006).
  31. Burzotta F, Iacoviello L, Di Castelnuovo A *et al*: 4G/5G PAI-1 promoter polymorphism and acute-phase levels of PAI-1 following coronary bypass surgery: a prospective study. *J.Thromb. Thrombolysis.* 16(3), 149-154 (2003).
  32. Festa A, D'Agostino R, Jr., Rich SS, Jenny NS, Tracy RP, Haffner SM: Promoter (4G/5G) plasminogen activator inhibitor-1 genotype and plasminogen activator inhibitor-1 levels in blacks, Hispanics, and non-Hispanic whites: the Insulin Resistance Atherosclerosis Study. *Circulation* 107(19), 2422-2427 (2003).
  33. Bottiger C, Koch W, Lahn C *et al*: 4G/5G polymorphism of the plasminogen activator inhibitor-1 gene and risk of restenosis after coronary artery stenting. *Am.Heart J.* 146(5), 855-861 (2003).
  34. Prisco D, Fedi S, Antonucci E *et al*: Postprocedural PAI-1 activity is a risk marker of subsequent clinical restenosis

- in patients both with and without stent implantation after elective balloon PTCA. *Thromb.Res.* 104(3), 181-186 (2001).
35. Ishiwata S, Tukada T, Nakanishi S, Nishiyama S, Seki A: Postangioplasty restenosis: platelet activation and the coagulation-fibrinolysis system as possible factors in the pathogenesis of restenosis. *Am.Heart J.* 133(4), 387-392 (1997).
  36. Hoekstra T, Geleijnse JM, Schouten EG, Klufft C: Plasminogen activator inhibitor-type 1: its plasma determinants and relation with cardiovascular risk. *Thromb. Haemost.* 91(5), 861-872 (2004).
  37. Kastrati A, Koch W, Berger PB *et al*: Protective role against restenosis from an interleukin-1 receptor antagonist gene polymorphism in patients treated with coronary stenting. *J.Am.Coll. Cardiol.* 36(7), 2168-2173 (2000).
  38. Rauchhaus M, Gross M, Schulz S *et al*: The E-selectin SER128ARG gene polymorphism and restenosis after successful coronary angioplasty. *Int.J.Cardiol.* 83(3), 249-257 (2002).
  39. Monraats PS, Pires NM, Agema WR *et al*: Genetic inflammatory factors predict restenosis after percutaneous coronary interventions. *Circulation* 112(16), 2417-2425 (2005).
  40. Economou E, Tousoulis D, Katinioti A *et al*: Chemokines in patients with ischaemic heart disease and the effect of coronary angioplasty. *Int.J.Cardiol.* 80(1), 55-60 (2001).
  41. Monraats PS, Pires NM, Schepers A *et al*: Tumor necrosis factor-alpha plays an important role in restenosis development. *FASEB J.* 19(14), 1998-2004 (2005).
  42. Monraats PS, Kurreeman FA, Pons D *et al*: Interleukin 10: a new risk marker for the development of restenosis after percutaneous coronary intervention. *Genes Immun.* 8(1), 44-50 (2007).
  43. Monraats PS, de Vries F, de Jong LW *et al*: Inflammation and apoptosis genes and the risk of restenosis after percutaneous coronary intervention. *Pharmacogenet. Genomics* 16(10), 747-754 (2006).
  44. Pesarini G, Ferrero V, Tomai F *et al*: Steroid-eluting stents in patients with acute coronary syndromes. Angiographic results of DESIRE: Dexamethasone-Eluting Stent Italian REgistry. *J.Invasive.Cardiol.* 21(3), 86-91 (2009).
  45. van Tiel CM, Bonta PI, Rittersma SZ *et al*: p27kip1-838C>A single nucleotide polymorphism is associated with restenosis risk after coronary stenting and modulates p27kip1 promoter activity. *Circulation* 120(8), 669-676 (2009).
  46. Agema WR, Jukema JW, Pimstone SN, Kastelein JJ: Genetic aspects of restenosis after percutaneous coronary interventions: towards more tailored therapy. *Eur.Heart J.* 22(22), 2058-2074 (2001).
  47. Newby AC: Matrix metalloproteinases regulate migration, proliferation, and death of vascular smooth muscle cells by degrading matrix and non-matrix substrates. *Cardiovasc.Res.* 69(3), 614-624 (2006).
  48. Hojo Y, Ikeda U, Katsuki T, Mizuno O, Fujikawa H, Shimada K: Matrix metalloproteinase expression in the coronary circulation induced by coronary angioplasty. *Atherosclerosis* 161(1), 185-192 (2002).
  49. Ye S: Influence of matrix metalloproteinase genotype on cardiovascular disease susceptibility and outcome. *Cardiovasc.Res.* 69(3), 636-645 (2006).
  50. Ye S, Eriksson P, Hamsten A, Kurkinen M, Humphries SE, Henney AM: Progression of coronary atherosclerosis is associated with a common genetic variant of the human stromelysin-1 promoter which results in reduced gene expression. *J.Biol.Chem.* 271(22), 13055-13060 (1996).
  51. Beyzade S, Zhang S, Wong YK, Day IN, Eriksson P, Ye S: Influences of matrix metalloproteinase-3 gene variation on extent of coronary atherosclerosis and risk of myocardial infarction. *J.Am.Coll. Cardiol.* 41(12), 2130-2137 (2003).
  52. Medley TL, Kingwell BA, Gatzka CD, Pillay P, Cole TJ: Matrix metalloproteinase-3 genotype contributes to age-related aortic stiffening through modulation of gene and protein expression. *Circ.Res.* 92(11), 1254-1261 (2003).
  53. Lichtinghagen R, Bahr MJ, Wehmeier M *et al*: Expression and coordinated regulation of matrix metalloproteinases in chronic hepatitis C and hepatitis C

- virus-induced liver cirrhosis. *Clin.Sci. (Lond)* 105(3), 373-382 (2003).
54. De Maat MP, Jukema JW, Ye S *et al*: Effect of the stromelysin-1 promoter on efficacy of pravastatin in coronary atherosclerosis and restenosis. *Am.J.Cardiol.* 83(6), 852-856 (1999).
  55. Humphries S, Bauters C, Meirhaeghe A, Luong L, Bertrand M, Amouyel P: The 5A6A polymorphism in the promoter of the stromelysin-1 (MMP3) gene as a risk factor for restenosis. *Eur.Heart J.* 23(9), 721-725 (2002).
  56. Hoppmann P, Koch W, Schomig A, Kastrati A: The 5A/6A polymorphism of the stromelysin-1 gene and restenosis after percutaneous coronary interventions. *Eur.Heart J.* 25(4), 335-341 (2004).
  57. Mintz GS, Nissen SE, Anderson WD *et al*: American College of Cardiology Clinical Expert Consensus Document on Standards for Acquisition, Measurement and Reporting of Intravascular Ultrasound Studies (IVUS). A report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J.Am.Coll. Cardiol.* 37(5), 1478-1492 (2001).
  58. Cook S, Wenaweser P, Togni M *et al*: Incomplete stent apposition and very late stent thrombosis after drug-eluting stent implantation. *Circulation* 115(18), 2426-2434 (2007).
  59. Hong MK, Mintz GS, Lee CW *et al*: Incidence, mechanism, predictors, and long-term prognosis of late stent malapposition after bare-metal stent implantation. *Circulation* 109(7), 881-886 (2004).
  60. Mintz GS, Shah VM, Weissman NJ: Regional remodeling as the cause of late stent malapposition. *Circulation* 107(21), 2660-2663 (2003).
  61. Shah VM, Mintz GS, Apple S, Weissman NJ: Background incidence of late malapposition after bare-metal stent implantation. *Circulation* 106(14), 1753-1755 (2002).
  62. van der Hoeven BL, Liem SS, Dijkstra J *et al*: Stent malapposition after sirolimus-eluting and bare-metal stent implantation in patients with ST-segment elevation myocardial infarction: acute and 9-month intravascular ultrasound results of the MISSION! intervention study. *JACC.Cardiovasc.Interv.* 1(2), 192-201 (2008).
  63. Tanabe K, Serruys PW, Degertekin M *et al*: Incomplete stent apposition after implantation of paclitaxel-eluting stents or bare metal stents: insights from the randomized TAXUS II trial. *Circulation* 111(7), 900-905 (2005).
  64. Virmani R, Farb A, Guagliumi G, Kolodgie FD: Drug-eluting stents: caution and concerns for long-term outcome. *Coron.Artery Dis.* 15(6), 313-318 (2004).
  65. Bergheanu SC, Pons D, van der Hoeven B *et al*: Inflammatory polymorphisms and the risk of late-acquired stent malapposition. *Journal of the American College of Cardiology* 53(10), A392-A392 (2009).
  66. Schulz S, Schuster T, Mehilli J *et al*: Stent thrombosis after drug-eluting stent implantation: incidence, timing, and relation to discontinuation of clopidogrel therapy over a 4-year period. *Eur.Heart J.* 30(22), 2714-2721 (2009).
  67. Cutlip DE, Leon MB, Ho KK *et al*: Acute and nine-month clinical outcomes after "suboptimal" coronary stenting: results from the STent Anti-thrombotic Regimen Study (STARS) registry. *J.Am. Coll. Cardiol.* 34(3), 698-706 (1999).
  68. Uren NG, Schwarzacher SP, Metz JA *et al*: Predictors and outcomes of stent thrombosis: an intravascular ultrasound registry. *Eur.Heart J.* 23(2), 124-132 (2002).
  69. Huber MS, Mooney JF, Madison J, Mooney MR: Use of a morphologic classification to predict clinical outcome after dissection from coronary angioplasty. *Am.J.Cardiol.* 68(5), 467-471 (1991).
  70. van Werkum JW, Heestermans AA, Zomer AC *et al*: Predictors of coronary stent thrombosis: the Dutch Stent Thrombosis Registry. *J.Am.Coll. Cardiol.* 53(16), 1399-1409 (2009).
  71. Bergheanu SC, van der Hoeven BL, Hassan AKM *et al*: Post-intervention IVUS is not predictive for very late in-stent thrombosis in drug-eluting stents. *Acta Cardiol* 64(5), 611-616 (2009).
  72. Weiss EJ, Bray PF, Tayback M *et al*: A polymorphism of a platelet glycoprotein receptor as an inherited risk factor for

- coronary thrombosis. *N.Engl.J.Med.* 334(17), 1090-1094 (1996).
73. Gorchakova O, Koch W, Mehilli J *et al*: PIA polymorphism of the glycoprotein IIIa and efficacy of reperfusion therapy in patients with acute myocardial infarction. *Thromb.Haemost.* 91(1), 141-145 (2004).
  74. Walter DH, Schachinger V, Elsner M, Dimmeler S, Zeiher AM: Platelet glycoprotein IIIa polymorphisms and risk of coronary stent thrombosis. *Lancet* 350(9086), 1217-1219 (1997).
  75. Kastrati A, Koch W, Gawaz M *et al*: PIA polymorphism of glycoprotein IIIa and risk of adverse events after coronary stent placement. *J.Am.Coll.Cardiol.* 36(1), 84-89 (2000).
  76. Sucker C, Scheffold N, Cyran J, Ghodsizad A, Scharf RE, Zotz RB: No evidence for involvement of prothrombotic platelet receptor polymorphisms in acute coronary stent thrombosis. *Int.J.Cardiol.* 123(3), 355-357 (2008).
  77. Angiolillo DJ, Fernandez-Ortiz A, Bernardo E *et al*: PIA polymorphism and platelet reactivity following clopidogrel loading dose in patients undergoing coronary stent implantation. *Blood Coagul.Fibrinolysis* 15(1), 89-93 (2004).
  78. Kuehl P, Zhang J, Lin Y *et al*: Sequence diversity in CYP3A promoters and characterization of the genetic basis of polymorphic CYP3A5 expression. *Nat. Genet.* 27(4), 383-391 (2001).
  79. Suh JW, Koo BK, Zhang SY *et al*: Increased risk of atherothrombotic events associated with cytochrome P450 3A5 polymorphism in patients taking clopidogrel. *CMAJ.* 174(12), 1715-1722 (2006).
  80. Brandt JT, Close SL, Iturria SJ *et al*: Common polymorphisms of CYP2C19 and CYP2C9 affect the pharmacokinetic and pharmacodynamic response to clopidogrel but not prasugrel. *J.Thromb. Haemost.* 5(12), 2429-2436 (2007).
  81. Geisler T, Schaeffeler E, Dippon J *et al*: CYP2C19 and nongenetic factors predict poor responsiveness to clopidogrel loading dose after coronary stent implantation. *Pharmacogenomics.* 9(9), 1251-1259 (2008).
  82. Smith SM, Judge HM, Peters G *et al*: Common sequence variations in the P2Y12 and CYP3A5 genes do not explain the variability in the inhibitory effects of clopidogrel therapy. *Platelets.* 17(4), 250-258 (2006).
  83. Mega JL, Close SL, Wiviott SD *et al*: Cytochrome p-450 polymorphisms and response to clopidogrel. *N.Engl.J.Med.* 360(4), 354-362 (2009).
  84. Simon T, Verstuyft C, Mary-Krause M *et al*: Genetic determinants of response to clopidogrel and cardiovascular events. *N.Engl.J.Med.* 360(4), 363-375 (2009).
  85. Trenk D, Hochholzer W, Fromm MF *et al*: Cytochrome P450 2C19 681G>A polymorphism and high on-clopidogrel platelet reactivity associated with adverse 1-year clinical outcome of elective percutaneous coronary intervention with drug-eluting or bare-metal stents. *J.Am.Coll.Cardiol.* 51(20), 1925-1934 (2008).
  86. Giusti B, Gori AM, Marcucci R *et al*: Relation of cytochrome P450 2C19 loss-of-function polymorphism to occurrence of drug-eluting coronary stent thrombosis. *Am.J.Cardiol.* 103(6), 806-811 (2009).
  87. Sibbing D, Stegherr J, Latz W *et al*: Cytochrome P450 2C19 loss-of-function polymorphism and stent thrombosis following percutaneous coronary intervention. *Eur.Heart J.* 30(8), 916-922 (2009).
  88. Collet JP, Hulot JS, Pena A *et al*: Cytochrome P450 2C19 polymorphism in young patients treated with clopidogrel after myocardial infarction: a cohort study. *Lancet* 373(9660), 309-317 (2009).
  89. Shuldiner AR, O'Connell JR, Bliden KP *et al*: Association of cytochrome P450 2C19 genotype with the antiplatelet effect and clinical efficacy of clopidogrel therapy. *JAMA* 302(8), 849-857 (2009).
  90. Eshtehardi P, Eslami M, Moayed DA: Simultaneous subacute coronary drug-eluting stent thrombosis in two different vessels of a patient with factor V Leiden mutation. *J.Cardiovasc.Med. (Hagerstown.)* 9(4), 410-413 (2008).
  91. Hayden EC: International genome project launched. *Nature* 451(7177), 378-379 (2008).

