

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/35607> holds various files of this Leiden University dissertation

Authors: Buiten, Maurits

Title: Heart and kidney : a dynamic relationship

Issue Date: 2015-10-01

4

SOLUBLE KLOTHO IS NOT INDEPENDENTLY ASSOCIATED WITH CARDIOVASCULAR DISEASE IN A POPULATION OF DIALYSIS PATIENTS

M.S. Buiten, M.K. de Bie, A. Bouma- de Krijger, B. van Dam,
F.W. Dekker, J.W. Jukema, T.J. Rabelink, J.I. Rotmans

Published in BMC Nephrol. 2014 Dec 11;15(1):197.

ABSTRACT

Background

Dialysis patients suffer from a high burden of cardiovascular disease (CVD). Partly this is due to progressive deterioration of calcium-phosphate homeostasis. Previous studies suggested that besides FGF-23, low levels of Klotho, a protein linked to aging, might constitute a key factor in this detrimental relationship. The purpose of the present study was to determine the relationship between serum Klotho (sKlotho) and the presence of CVD in dialysis patients.

Methods

Plasma levels of sKlotho were measured in a cohort of dialysis patients and related to left ventricular (LV) dysfunction (defined as a LV ejection fraction <45%) and LV mass using echocardiography. Coronary artery disease (CAD) and calcification score were assessed using computed tomography angiography. Abdominal aortic calcification score (AACscore) was measured by abdominal X-ray.

Results

We included 127 dialysis patients, 67 ± 7 years old, 76% male, 67% on hemodialysis, median sKlotho 460pg/mL (25th-75th percentile 350-620pg/mL). Patients with a low sKlotho (<460pg/mL) showed significantly more CAD (81% versus 61%; $p=0.02$) and LV dysfunction (19% versus 3%; $p<0.01$) than patients with a high sKlotho level. However, after adjusting for confounders, sKlotho was not independently associated with the presence of CVD or the AACscore.

Conclusions

In the present cohort of dialysis patients, sKlotho was not independently associated with CVD. However, patients with a low sKlotho level (<460pg/mL) did show CAD and LV dysfunction more frequently than patients with high sKlotho. Therefore, while sKlotho might be a marker for CVD in dialysis patients, the current data does not support a direct cardioprotective effect of sKlotho.

INTRODUCTION

Patients with chronic kidney disease (CKD) suffer from a high burden of cardiovascular disease (CVD)^{1,2}. Whereas part of this burden is related to traditional risk factors, CKD-associated disturbance in calcium-phosphate homeostasis play a crucial role as well³. Recently, Fibroblast Growth Factor-23 (FGF-23) and its co-receptor Klotho have emerged as pivotal players in calcium-phosphate homeostasis and they could be the missing link in the detrimental relationship between CKD and CVD⁴.

Membrane-bound Klotho is a co-receptor for FGF-23 vital for FGF-23's phosphaturic effect. In CKD serum FGF-23 levels rise in response to phosphate retention, while Klotho levels decrease⁵. Elevated FGF-23 levels are associated with worse cardiovascular outcome but it is unknown whether this is due to the toxic effect of FGF-23 itself or due to Klotho deficiency causing a state of FGF-23 resistance. Studies examining the relationship between low Klotho and outcome show conflicting results^{6,7}.

Klotho was first identified in mice, where mutations of the gene led to a syndrome resembling aging⁸. The Klotho protein exists in both a membrane bound and a soluble form (sKlotho). sKlotho is derived from the extracellular part of membranous Klotho and through alternative splicing of the Klotho gene and has a unique role in renal calcium and phosphate excretion, independent of FGF-23⁹⁻¹². Furthermore, sKlotho has been associated with protective effects on vascular calcification and oxidative stress in preclinical studies^{13,14}. Since the kidneys are the principal source of sKlotho, some studies have suggested that CKD might be a state of progressive Klotho deficiency^{5,16}.

Given the strong cardioprotective effects of Klotho demonstrated in preclinical studies, the current study aimed to assess the association between sKlotho and CVD in patients on chronic dialysis. It was hypothesized that low levels of sKlotho are associated with a larger burden of CVD in these patients.

METHODS

Population and design

For this study, all patients enrolled in the ICD-2 trial (ISRCTN 20479861) until January 2013 were analyzed. The ICD-2 trial is an ongoing randomized controlled trial designed to evaluate the effectiveness of an implantable cardioverter defibrillator in the prevention of sudden cardiac death in dialysis patients. The study protocol has been described previously¹⁷. The study population consists of 55-80 year old patients with a left ventricular ejection fraction (LVEF) of at least 35%. All patients provided written informed consent and the design of the trial was approved by the local ethics committee.

An extensive screening protocol was performed in all patients at the time of enrollment in the ICD-2 trial, including blood analysis, computed tomography angiography (CTA), transthoracic echocardiography (TTE) and lateral abdominal X-ray. Furthermore clinical

data on demographic characteristics, coexisting conditions, use of medications and information regarding the dialysis procedures were collected. Twenty-four-hour urinary samples were collected in all patients. Residual renal function was calculated as the mean of creatinine and urea clearance adjusted for body surface area (ml/min per 1.73m^2). The mean of both post- and pre-dialysis plasma samples (if available) were used to estimate mean plasma creatinine and urea concentrations ¹⁸. The residual renal function was considered zero in patients with an urinary output $<100\text{mL}/24\text{hours}$.

Laboratory tests

After collection, the blood samples were immediately centrifuged, separated in vials and stored at -60°C for future essays. Serum levels of creatinine, calcium and phosphate were measured at time of inclusion. Concentrations of plasma c-terminal FGF-23, soluble serum Klotho (sKlotho), serum 25-hydroxyvitamin D (25(OH)D) and plasma parathyroid hormone (PTH) were measured from the frozen samples. FGF-23 was measured using an assay designed to measure C-terminal FGF-23 (Immutopics, San Clemente, CA, USA)¹⁹. For the measurement of sKlotho an ELISA assay was used (IBL, International GmbH, Hamburg, Germany) based on the antibodies and substrates designed, used and described by Yamazaki and colleagues (intra- and inter-assay coefficients of variation $<5\%$ and 8% , respectively)²⁰. 25(OH)D levels were measured using a LC-MS/MS method ²¹.

Echocardiography

All patients underwent 2-dimensional TTE with commercially available ultrasound equipment (M3s probe, Vivid 7, GE Vingmed, Horton, Norway). The images were digitally stored for off-line analysis (EchoPAC version 110.0.0, GE Vingmed, Horten, Norway). Using Simpson's biplane method, left ventricular mass index (LVMI) and LVEF were assessed by a single independent observer. In accordance with the current guidelines, LV dysfunction was defined as a $\text{LVEF} < 45\%$ ²².

Multi Slice CT protocol

The CTA protocol used has been previously described ²³. Patients were scanned using a 64-slice CT scanner (Aquilion64, Toshiba Medical Systems, Otawara, Japan) or a 320-slice CT scanner (Aquilion ONE, Toshiba Medical Systems, Otawara Japan). In the presence of residual renal function, post and prehydration of the patients was performed in accordance with the nephrologist. The coronary artery calcium score (CACscore) was analyzed using the Agatston method ²⁴. The presence of significant coronary artery stenosis was defined as $\geq 50\%$ luminal narrowing. Coronary artery disease (CAD) was defined as the presence of at least one coronary artery with significant stenosis. If coronary artery bypass grafting or a percutaneous coronary intervention had been performed in the past, treated vessels were scored as occluded and the native vessels scored using CTA.

Quantification of abdominal aortic calcification

The extent of aortic calcification was calculated on a lateral X-ray of the abdomen. This X-ray was taken in a standing position using standard radiographic equipment. The abdominal aorta calcification (AAC) score was calculated using a previously validated grading system, in which the extent of calcific deposits in the abdominal aorta is graded on a per segment basis ²⁵.

Statistical analysis

Continuous data are presented as mean $\pm$ SD and compared using the 2-tailed Student's t-test. Categorical data are presented as numbers and percentages and compared using the Chi-square test. For the initial data presentation, the cohort was divided into two groups using median sKlotho as a cut-off.

The relationship between sKlotho and CVD was explored using regression models. For the AACscore, CACscore and LVMI linear regression models were used with unstandardized Betas and a 95% confidence interval (CI). To assist practical usability of the models, sKlotho was transformed from 1pg/mL to 100pg/mL, thus Beta's represent changes of 100pg/mL sKlotho. Otherwise, Betas and ORs would have been very small. For the associations between sKlotho and CAD and between sKlotho and LV-dysfunction, logistic regression models were used. Normality, linearity and homoscedasticity of the different variables in relation to sKlotho were checked. If residuals were not normally distributed, a logarithmic transformation (ln) was performed. In both the linear and logistic regression analysis, two models were used; model 1: crude model and model 2:t additionally adjusting for potential confounders: age, gender, dialysis type, dialysis vintage and residual renal function (mL/min per 1.73m²).

The correlations between the different parameters in the calcium-phosphate metabolism and sKlotho levels were evaluated using a Spearman rank correlation test. In a method comparable to the one used for sKlotho, the relationship between FGF-23 and CVD (AACscore, CACscore, LVMI, CAD and LV-dysfunction) was explored using regression models adjusting for potential confounders. To assess any combined effect of sKlotho and FGF-23, we compared patients with a high suspected calcification risk (low sKlotho and high FGF-23) to patients with a low suspected calcification risk (high Klotho and low FGF-23) based on the median of sKlotho and FGF-23.

All statistical analyses were performed using SPSS (version 20, IBM Corp, Amonk, NY, USA). A p-value <0.05 was considered statistically significant.

RESULTS

Patient characteristics

A total of 127 patients were included for the current study. The average age of the participating patients was 67 ± 7 years, 24% was female and 68% utilized hemodialysis (HD) as modality for renal replacement therapy. Median sKlotho was 460pg/mL (25th-75th percentile; 350-620pg/mL). The baseline characteristics are shown in table 1. The cause of end-stage renal disease (ESRD) was hypertension (24%), diabetes (18%), glomerulonephritis (9%), acute tubular necrosis (5%) and other (22%) or unknown causes (22%).

Table 1: Baseline characteristics in the total cohort and in two groups of patients with a low and high soluble Klotho concentration in serum

	Total cohort n=127	Klotho <460pg/mL n=63	Klotho >460pg/mL n=64
Age, years	67 ± 7	67 ± 7	67 ± 8
Gender, female	30 (24%)	10 (16%)	20 (31%)*
Pulse pressure, mmHg	59 ± 24	59 ± 24	59 ± 16
Hypertension	104 (82%)	53 (87%)	51 (80%)
Diabetes	43 (34%)	24 (40%)	19 (30%)
Dialysis Modality, PD	42 (33%)	8 (13%)	34 (53%)*
Dialysis vintage, years	2.3 ± 2	2.6 ± 2.7	2.0 ± 2.2
RRF, (ml/min/1.73m ²)	1.8 ± 1.6	1.7 ± 1.8	2.0 ± 1.4
Anuria	30 (24%)	20 (33%)	10 (16%)*
Phosphate (mmol/L)	1.5 ± 0.4	1.6 ± 0.4	1.5 ± 0.4
Calcium (mmol/L)	2.4 ± 0.2	2.4 ± 0.2	2.4 ± 0.2
FGF-23 (RefU/mL)	7247 ± 16815	8723 ± 21627	5794 ± 10044
25(OH)D (nmol/L)	87 ± 57	102 ± 65	$71 \pm 42^{**}$
PTH (pmol/L)	35 ± 34	42 ± 41	$28 \pm 22^{*}$
Creatinine (umol/L)	659 ± 202	642 ± 211	676 ± 195
CRP (mg/L)	14.3 ± 30.7	20.1 ± 41.7	$8.9 \pm 12.22^{*}$
β-blocker	71 (56%)	37 (60%)	34 (54%)
ACEi	32 (25%)	19 (31%)	13 (21%)
ARB	37 (29%)	13 (21%)	24 (38%)*
Statin	79 (62%)	43 (69%)	36 (57%)
NCPB	107 (84%)	54 (87%)	53 (84%)
CCPB	50 (39%)	25 (40%)	25 (40%)
Cinacalcet	15 (12%)	7 (12%)	8 (13%)

PD; Peritoneal dialysis, RRF; Residual Renal Function, FGF-23; Fibroblast Growth Factor 23. PTH; Parathyroid hormone, ACEi; Angiotensin Converting Enzyme inhibitor, ARB; Angiotensin receptor blocker, NCPB; Non calcium containing phosphate binder, CCPB; Calcium containing phosphate binder. *, $p < 0.05$, **, $p < 0.01$; ***, $p < 0.001$

After stratifying patients for plasma sKlotho levels into two groups, patients with a high sKlotho (>460 pg/mL) were more often female (31% versus 16%; $p=0.04$), more often had residual renal function (84% versus 67%; $p=0.04$), more frequently utilized peritoneal dialysis (PD; 53% versus 13%; $p<0.001$) and more often used an angiotensin receptor blocker (ARB; 38% versus 21%; $p=0.04$) than patients with low sKlotho. Patients with a low sKlotho had a higher plasma 25(OH)D (102nmol/L versus 71nmol/L; $p=0.002$), PTH (42pmol/L versus 28pmol/L; $p=0.02$) and CRP (20.1mg/L versus 8.9mg/L; $p=0.04$) than patients with high sKlotho (table 1). No further differences in baseline characteristics were found between the two groups.

sKlotho and markers for cardiovascular disease

In the 127 included patients, the average AACscore was 9 ± 6 , mean CACscore was 1134 ± 1316 and the average LVMI was 128 ± 42 g/m². A total of 76 (60%) patients suffered from CAD and 13 patients (10%) had LV dysfunction. As described in the Method section, only patients with LVEF $\geq$ 35% were included in the study. Patients with a high sKlotho (>460 pg/mL) had significantly less CAD (61% versus 81% of patients, $p=0.02$; figure 1) as well as less LV dysfunction (3% versus 19% of patients; $p=0.006$). The other markers of CVD were not significantly different between both groups.

The results of the multivariate regression modelling, with the various different markers of CVD as dependent variables are shown in table 2. In model 1, the crude association between sKlotho and markers of CVD was assessed. In these first models, sKlotho was not significantly associated with AACscore, CACscore, LVMI or the presence of CAD. However, a statistically significant crude association between sKlotho and the presence of LV-dysfunction was observed (OR: 0.58; 95% CI 0.37-0.92; $p=0.02$). In




Figure 1: Serum sKlotho and the number of significantly stenosed coronary arteries.

Table 2: Association between serum sKlotho and cardiovascular disease

AACscore	Beta	CI	p-value
Model 1. Crude association	0.33	-0.26-0.91	0.27
Model 2. (+ Confounders)*	0.58	-0.07-1.22	0.08
CACscore	Beta	CI	p-value
Model 1. Crude association	-0.06	-0.31-0.20	0.66
Model 2. (+Confounders)*	0.08	-0.19-0.36	0.55
LVMI	Beta	CI	p-value
Model 1. Crude association	-1.27	-5.58-3.04	0.56
Model 2. (+Confounders)*	1.55	-3.49-6.60	0.54
LV-dysfunction (LVEF<45%)	OR	CI	p-value
Model 1. Crude association	0.58	0.37-0.92	0.02
Model 2. (+Confounders)*	0.66	0.39-1.11	0.12
CAD (any vessel occluded)	OR	CI	p-value
Model 1. Crude association	0.82	0.65-1.05	0.11
Model 2. (+Confounders)*	0.91	0.66-1.23	0.52

Betas represent changes of 100pg/mL in sKlotho
AACscore; Abdominal aorta calcification score, CACscore; Coronary artery calcification score, LVMI; Left ventricular mass index, LV-dysfunction; Left ventricular dysfunction. CAD; Coronary artery disease.
Model 1; sKlotho. ***Model 2;** + age, gender, (ln)dialysis vintage, dialysis type, residual renal function.

model 2 the association between sKlotho and markers of CVD was assessed after adjusting for confounders (age, gender, (ln)dialysis vintage, dialysis type and residual renal function). After the adjustments made in model 2, sKlotho showed no significant association with any of these markers of CVD.

sKlotho and markers of the calcium-phosphate axis

The associations between the parameters of the calcium-phosphate axis are shown in table 3. Serum sKlotho level was inversely associated with plasma 25(OH)D ($r=-0.22$, $p=0.01$) and PTH ($r=-0.26$; $p<0.01$). FGF-23 showed a strong positive association with phosphate level ($r=0.51$; $p<0.001$) and PTH $r=0.29$; $p<0.001$). There was no crude association between serum FGF-23, PTH, 25(OH)D or phosphate and any of the investigated markers of CVD (data not shown). Furthermore, in the different multivariate regression models, FGF-23 was not independently associated with CVD after adjusting for confounders (although a LVEF<45% showed a trend toward an independent association with FGF-23, OR: 0.51 (0.26-1.01; $p=0.053$, Table 4.)

A total of 26 patients had a high median serum FGF-23 (>2625 RefU/L) combined with a low sKlotho (<460pg/mL), and 26 patients had a low FGF-23 (<2625 RefU/L) and high

Table 3: Spearman correlation coefficients of parameters of calcium-phosphate metabolism and sKlotho levels in serum

	Phosphate (mmol/L)	25(OH)D (nmol/L)	PTH (pmol/L)	FGF-23 (RefU/L)	sKlotho (pg/mL)
25(OH)D (nmol/L)	r=0.06 p=0.53				
PTH (pmol/L)	r=0.30 p<0.01	r=-0.01 p=0.90			
FGF-23 (RefU/L)	r=0.51 p<0.001	r=-0.03 p=0.76	r=0.29 P=0.001		
sKlotho (pg/mL)	r=-0.04 p=0.64	r=-0.22 p=0.01	r=-0.26 p<0.01	r=0.11 p=0.22	
Calcium (mmol/L)	r=0.04 p=0.67	r=0.05 p=0.55	r=-0.09 p=0.34	r=0.22 p=0.01	r=-0.04 p=0.66

PTH; Parathyroid hormone, FGF-23; Fibroblast Growth Factor 23. Indicated are correlation coefficients (r) with levels of significance (p)

Table 4: Association between serum FGF-23 and cardiovascular disease

AACscore	Beta	CI	p-value
Model 1. Crude association	0.72	-0.13-1.57	0.10
Model 2. (+ Confounders)*	0.49	-0.35-1.34	0.25
CACscore	Beta	CI	p-value
Model 1. Crude association	0.25	-0.13-0.63	0.20
Model 2. (+Confounders)*	0.001	-0.40-0.40	1.00
LVMI	Beta	CI	p-value
Model 1. Crude association	0.48	-5.79-6.75	0.88
Model 2. (+Confounders)*	-0.70	-7.40-6.00	0.84
LV-dysfunction (LVEF<45%)	OR	CI	p-value
Model 1. Crude association	0.68	0.40-1.16	0.16
Model 2. (+Confounders)*	0.51	0.26-1.01	0.053
CAD (any vessel occluded)	OR	CI	p-value
Model 1. Crude association	0.94	0.66-0.63	0.20
Model 2. (+Confounders)*	0.65	0.41-1.04	0.08

Betas represent changes in LnFGF-23

AACscore; Abdominal aorta calcification score, CACscore; Coronary artery calcification score, LVMI; Left ventricular mass index, LV-dysfunction; Left ventricular dysfunction. CAD; Coronary artery disease.

Model 1; sKlotho. ***Model 2;** + age, gender, (ln)dialysis vintage, dialysis type, residual urine production.

sKlotho level ($>460\text{pg/mL}$). Patients in the former group had a significantly higher CACscore (1242 ± 1037 versus 485 ± 656 ; $p=0.01$) as well as a trend towards lower LVEF ($52\pm7\%$ versus $55\pm6\%$; $p=0.07$) and more CAD (85% versus 60% ; $p=0.07$) than patients in the latter group.

DISCUSSION

Recently, sKlotho has emerged as a powerful player in calcium-phosphate homeostasis that is thought to contribute to the high burden of CVD in CKD patients ⁴. To the best of our knowledge, the present study is the first study that investigated the association between sKlotho and CVD in dialysis patients. Contrary to our hypothesis, while patients with a low sKlotho demonstrated LV dysfunction and CAD more often, there was no independent association between sKlotho and CVD in the present cohort.

sKlotho and cardiovascular disease

Several clinical studies have suggested that Klotho exerts strong cardioprotective effects. For instance, Klotho has been shown to protect against vascular calcifications in rodent models of CKD while in humans without CKD, higher sKlotho levels have been related to a lower incidence of mortality and CVD ^{8,9,26,27}. Moreover, low sKlotho levels have been associated with increased arterial stiffness in CKD patients ²⁸. Further support for a direct role of Klotho in vascular homeostasis comes from *in vitro* studies showing endogenous expression of Klotho in human vascular smooth muscle cells (VSMCs) ¹⁴. Interestingly, inhibition of sKlotho expression in aortic VSMCs resulted in accelerated calcification of these cells ¹⁴. However, the exact role of Klotho in the progression of CVD in dialysis patients remains to be elucidated.

In the present study, dialysis patients with a low sKlotho demonstrated LV-dysfunction and CAD more frequently. However, an independent association between sKlotho and CVD was not observed. There are several explanations for these findings.

First, disorders of mineral homeostasis, as well as CVD start to develop in the early stages of CKD ^(1,3). Therefore, patients with ESRD have been exposed to an environment predisposing to vascular calcifications for a prolonged period of time. In our study, we solely measured sKlotho levels at time of inclusion and as a consequence, the sKlotho levels in our study do not reflect the total 'burden' of Klotho deficiency in the dialysis patients' preceding years with CKD, before Klotho was measured. Thus, the association between sKlotho and vascular pathology might be diminished by the time patients have developed ESRD.

Second, the role of sKlotho in the development of atherosclerotic disease in dialysis patients might be overshadowed by the large amount of other pathophysiological stimuli for CVD prevalent in these patients, such as smoking, obesity, diabetes and hypertension among others. This might explain the disparity between data found in rodents with only Klotho deficiency and patients suffering from a wide variety of co-morbidities ^{8,9}.

A third explanation could be that sKlotho simply has no independent contribution to CVD in dialysis patients. Several very recent publications revealed similar results^{29,32}. In patients with CKD up to stage IV, sKlotho was not associated with AACscore nor was it predictive for cardiovascular outcomes^{29,30}. In addition, pre-clinical studies have provided for a limited role of Klotho in the vasculature since they showed a very low or absent vascular Klotho expression and Klotho-mediated effects in human and murine VSMCs^{31,32}.

Even though sKlotho does not appear to be independently associated with CVD in dialysis patients in the present study, CAD and LV dysfunction were more often present in patients with a low sKlotho level than in those with a high sKlotho level. Furthermore, patients with a combination of low serum sKlotho and high FGF-23 showed a significantly higher CACscore as well as a trend toward a lower LVEF and less CAD than patients with high sKlotho and low FGF-23. Therefore, sKlotho might be an effective biomarker of CVD. In this respect, it would be interesting to explore the association between urinary sKlotho and CVD, since recent studies have illustrated that urinary sKlotho might correlate better with renal function than plasma sKlotho³³.

Effect of treatment modalities on sKlotho level

In the investigated cohort two interesting observations were made regarding the association between treatment modalities and sKlotho levels. In line with a previous study that showed that losartan increased sKlotho levels, we observed that patients with a higher sKlotho more often used an ARB³⁴. It was observed in rats that administration of angiotensin-II reduced Klotho gene expression at both mRNA and protein level in the kidney, whereas ARB and renin inhibitors increased Klotho expression^{35,36}. These studies suggest a direct inhibitory effect of the renin-angiotensin-aldosterone system (RAAS) on Klotho expression³⁷.

In addition, we observed that patients with a higher sKlotho more often performed PD than patients with low sKlotho levels. This is an intriguing difference, since sKlotho is excreted in the peritoneal dialysate (median amount of 114 ng/day), while the molecular weight of sKlotho is too high (120-130 kDa) to allow clearance by hemodialysis³⁸. A possible explanation could be that the higher RRF in PD patients as compared to a lower RRF in HD patients, as observed in our cohort, reflects a healthier kidney and thus higher sKlotho levels.

Limitations

This study has several limitations. Data on dietary phosphate intake was not collected, which is a known determinant for both serum phosphate and FGF-23 levels. Furthermore, sKlotho was not measured at a fixed time and thus a circadian variation in the measured sKlotho, as described in previous studies, cannot be excluded³⁹. However, all blood samples in the ICD-2 trial were taken in the morning, somewhat decreasing this variability.

Conclusions

In the present cohort of dialysis patients, sKlotho level was not independently associated with cardiovascular disease. However, patients with a low sKlotho level (<460pg/mL) did show CAD and LV dysfunction more often than patients with a high sKlotho level. Therefore, while sKlotho might be a marker for CVD in dialysis patients, the current data does not support a direct cardioprotective effect of sKlotho.

REFERENCE LIST

- Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N.Engl.J.Med.* 2004; **13**:1296-1305.
- Boddana P, Caskey F, Casula A, Ansell D. UK Renal Registry 11th Annual Report (December 2008): Chapter 14 UK Renal Registry and international comparisons. *Nephron Clin.Pract.* 2009; c269-c276.
- KDIGO clinical practice guideline for the diagnosis, evaluation, prevention, and treatment of Chronic Kidney Disease-Mineral and Bone Disorder (CKD-MBD). *Kidney Int. Suppl* 2009; **113**:S1-130.
- Shimada T, Hasegawa H, Yamazaki Y, Muto T, Hino R, Takeuchi Y et al. FGF-23 is a potent regulator of vitamin D metabolism and phosphate homeostasis. *J.Bone Miner.Res.* 2004; **3**:429-435.
- Pavik I, Jaeger P, Ebner L, Wagner CA, Petzold K, Spichtig D et al. Secreted Klotho and FGF23 in chronic kidney disease Stage 1 to 5: a sequence suggested from a cross-sectional study. *Nephrol.Dial.Transplant.* 2013; **2**:352-359.
- Hu MC, Shi M, Zhang J, Quinones H, Griffith C, Kuro-o M et al. Klotho deficiency causes vascular calcification in chronic kidney disease. *J.Am.Soc.Nephrol.* 2011; **1**:124-136.
- Seiler S, Wen M, Roth HJ, Fehrenz M, Flugge F, Herath E et al. Plasma Klotho is not related to kidney function and does not predict adverse outcome in patients with chronic kidney disease. *Kidney Int.* 2013; **1**:121-128.
- Kuro-o M, Matsumura Y, Aizawa H, Kawaguchi H, Suga T, Utsugi T et al. Mutation of the mouse klotho gene leads to a syndrome resembling ageing. *Nature* 1997; **6655**:45-51.
- Imura A, Iwano A, Tohyama O, Tsuji Y, Nozaki K, Hashimoto N et al. Secreted Klotho protein in sera and CSF: implication for post-translational cleavage in release of Klotho protein from cell membrane. *FEBS Lett.* 2004; **1-3**:143-147.
- Matsumura Y, Aizawa H, Shiraki-Iida T, Nagai R, Kuro-o M, Nabeshima Y. Identification of the human klotho gene and its two transcripts encoding membrane and secreted klotho protein. *Biochem.Biophys. Res.Comm.* 1998; **3**:626-630.
- Chang Q, Hoefs S, van der Kemp AW, Topala CN, Bindels RJ, Hoenderop JG. The beta-glucuronidase klotho hydrolyzes and activates the TRPV5 channel. *Science* 2005; **5747**:490-493.
- Hu MC, Shi M, Zhang J, Pastor J, Nakatani T, Lanske B et al. Klotho: a novel phosphaturic substance acting as an autocrine enzyme in the renal proximal tubule. *FASEB J.* 2010; **9**:3438-3450.
- Yamamoto M, Clark JD, Pastor JV, Gurnani P, Nandi A, Kurosu H et al. Regulation of oxidative stress by the anti-aging hormone klotho. *J.Biol.Chem.* 2005; **45**:38029-38034.
- Lim K, Lu TS, Molostvov G, Lee C, Lam FT, Zehnder D et al. Vascular Klotho deficiency potentiates the development of human artery calcification and mediates resistance to fibroblast growth factor 23. *Circulation* 2012; **18**:2243-2255.
- Hu MC, Shiizaki K, Kuro-o M, Moe OW. Fibroblast growth factor 23 and Klotho: physiology and pathophysiology of an endocrine network of mineral metabolism. *Annu.Rev.Physiol* 2013; 503-533.
- Lindberg K, Amin R, Moe OW, Hu MC, Erben RG, Ostman WA et al. The kidney is the principal organ mediating Klotho effects. *J.Am.Soc.Nephrol.* 2014; **10**: 2169-75
- de Bie MK, Lekkerkerker JC, van Dam B, Gaasbeek A, van Buren M, Putter H et al. Prevention of sudden cardiac death: rationale and design of the Implantable Cardioverter Defibrillators in Dialysis patients (ICD2) Trial-a prospective pilot study. *Curr.Med.Res. Opin.* 2008; **8**:2151-2157.
- Fouque D, Vennegoor M, ter Wee P, Wanner C, Basci A, Canaud B et al. EBPG guideline on nutrition. *Nephrol.Dial.Transplant.* 2007; ii45-ii87.
- Heijlboer AC, Levitus M, Vervloet MG, Lips P, ter Wee PM, Dijkstelbloem HM et al. Determination of fibroblast growth factor 23. *Ann.Clin.Biochem.* 2009; **Pt 4**:338-340.
- Yamazaki Y, Imura A, Urakawa I, Shimada T, Murakami J, Aono Y et al. Establishment of sandwich ELISA for soluble alpha-Klotho measurement: Age-dependent change of soluble alpha-Klotho levels in healthy subjects. *Biochem.Biophys.Res.Comm.* 2010; **3**:513-518.

- 21 Heijboer AC, Blankenstein MA, Kema IP, Buijs MM. Accuracy of 6 routine 25-hydroxyvitamin D assays: influence of vitamin D binding protein concentration. *Clin.Chem.* 2012; **3**:543-548.
- 22 Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA et al. Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J.Am.Soc.Echocardiogr.* 2005; **12**:1440-1463.
- 23 de Bie MK, Buiten MS, Gaasbeek A, Boogers MJ, Roos CJ, Schuijf JD et al. CT coronary angiography is feasible for the assessment of coronary artery disease in chronic dialysis patients, despite high average calcium scores. *PLoS.One.* 2013; **7**:e67936
- 24 Agatston AS, Janowitz WR, Hildner FJ, Zusmer NR, Viamonte M, Jr., Detrano R. Quantification of coronary artery calcium using ultrafast computed tomography. *J.Am.Coll.Cardiol.* 1990; **4**:827-832.
- 25 Wilson PW, Kauppila LI, O'Donnell CJ, Kiel DP, Hannan M, Polak JM et al. Abdominal aortic calcific deposits are an important predictor of vascular morbidity and mortality. *Circulation* 2001; **11**:1529-1534.
- 26 Semba RD, Cappola AR, Sun K, Bandinelli S, Dalal M, Crasto C et al. Plasma klotho and cardiovascular disease in adults. *J.Am.Geriatr.Soc.* 2011; **9**:1596-1601.
- 27 Navarro-Gonzalez JF, Donate-Correa J, Muros de FM, Perez-Hernandez H, Martinez-Sanz R, Mora-Fernandez C. Reduced Klotho is associated with the presence and severity of coronary artery disease. *Heart* 2014; **100**:34-40.
- 28 Kitagawa M, Sugiyama H, Morinaga H, Inoue T, Takiue K, Ogawa A et al. A decreased level of serum soluble Klotho is an independent biomarker associated with arterial stiffness in patients with chronic kidney disease. *PLoS.One.* 2013; **2**:e56695
- 29 Seiler S, Rogacev KS, Roth HJ, Shafein P, Emrich I, Neuhaus S et al. Associations of FGF-23 and sKlotho with Cardiovascular Outcomes among Patients with CKD Stages 2-4. *Clin.J.Am.Soc.Nephrol.* 2014; **9**: 1049-58
- 30 Craver L, Dusso A, Martinez-Alonso M, Sarro F, Valdivielso JM, Fernandez E. A low fractional excretion of Phosphate/Fgf23 ratio is associated with severe abdominal Aortic calcification in stage 3 and 4 kidney disease patients. *BMC.Nephrol.* 2013; **14**: 221
- 31 Lindberg K, Olsson H, Amin R, Ponnusamy A, Goetz R, Taylor RF et al. Arterial klotho expression and FGF23 effects on vascular calcification and function. *PLoS.One.* 2013; **4**: e60658
- 32 Scialla JJ, Lau WL, Reilly MP, Isakova T, Yang HY, Crouthamel MH et al. Fibroblast growth factor 23 is not associated with and does not induce arterial calcification. *Kidney Int.* 2013; **6**:1159-1168.
- 33 Akimoto T, Yoshizawa H, Watanabe Y, Numata A, Yamazaki T, Takeshima E et al. Characteristics of urinary and serum soluble Klotho protein in patients with different degrees of chronic kidney disease. *BMC.Nephrol.* 2012; **13**: 155
- 34 Lim SC, Liu JJ, Subramaniam T, Sum CF. Elevated circulating alpha-klotho by angiotensin II receptor blocker losartan is associated with reduction of albuminuria in type 2 diabetic patients. *J.Renin.Angiotensin.Aldosterone.Syst.* 2013; **15**: 487-90
- 35 Mitani H, Ishizaka N, Aizawa T, Ohno M, Usui S, Suzuki T et al. In vivo klotho gene transfer ameliorates angiotensin II-induced renal damage. *Hypertension* 2002; **4**:838-843.
- 36 Sun CY, Chong WJ, Jian HZ, Hsu HH, Wu IW, Hsu HJ et al. Aliskiren reduced renal fibrosis in mice with chronic ischemic kidney injury--beyond the direct renin inhibition. *Hypertens.Res.* 2012; **3**:304-311.
- 37 de Borst MH, Vervloet MG, ter Wee PM, Navis G. Cross talk between the renin-angiotensin-aldosterone system and vitamin D-FGF-23-klotho in chronic kidney disease. *J.Am.Soc.Nephrol.* 2011; **9**:1603-1609.
- 38 Akimoto T, Shiizaki K, Sugase T, Watanabe Y, Yoshizawa H, Otani N et al. The relationship between the soluble Klotho protein and the residual renal function among peritoneal dialysis patients. *Clin.Exp.Nephrol.* 2012; **3**:442-447.
- 39 Carpenter TO, Insogna KL, Zhang JH, Ellis B, Nieman S, Simpson C et al. Circulating levels of soluble klotho and FGF23 in X-linked hypophosphatemia: circadian variance, effects of treatment, and relationship to parathyroid status. *J.Clin.Endocrinol.Metab* 2010; **11**:E352-E357.

