

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/20872> holds various files of this Leiden University dissertation.

Author: Lommen, Suzanne Theresia Esther

Title: Exploring and exploiting natural variation in the wings of a predatory ladybird beetle for biological control

Issue Date: 2013-05-16

References

- Abouheif E (2004) A framework for studying the evolution of gene networks underlying polyphenism: insights from winged and wingless ant castes. In: Hall BK (ed) Environment, development, and evolution. MIT Press, pp. 125-137
- Abouheif E, Wray GA (2002) Evolution of the gene network underlying wing polyphenism in ants. *Science* 297:249-252
- Adachi-Hagimori T, Shibao M, Tanaka H, Seko T, Miura K (2011) Control of *Myzus persicae* and *Lipaphis erysimi* (Hemiptera: Aphididae) by adults and larvae of a flightless strain of *Harmonia axyridis* (Coleoptera: Coccinellidae) on non-heading Brassica cultivars in the greenhouse. *BioControl* 56:207-213
- Agarwala BK, Dixon AFG (1992) Laboratory study of cannibalism and interspecific predation in ladybirds. *Ecol. Entomol.* 17:303-309
- Anbesse SA, Strauch O, Ehlers R-U (2012) Genetic improvement of the biological control nematode *Heterorhabditis bacteriophora* (Rhabditidomorpha: Heterorhabditidae): heterosis effect enhances desiccation but not heat tolerance. *Biocontrol Sci. Technol.* 22:1035-1045
- Arnaud L, Spinneux Y, Haubruge E (2003) Preliminary observations of sperm storage in *Adalia bipunctata* (Coleoptera : Coccinellidae): sperm size and number. *Appl. Entomol. Zool.* 38:301-304
- Atallah J, Dworkin I, Cheung U, Greene A, Ing B, Leung L, Larsen E (2004) The environmental and genetic regulation of *obake* expressivity: morphogenetic fields as evolvable systems. *Evol. Dev.* 6:114-122
- Bakker FM, Klein ME, Mesa NC, Braun AR (1993) Saturation deficit tolerance spectra of phytophagous mites and their phytoseiid predators on cassava. *Exp. Appl. Acarol.* 17:97-113
- Bänsch R (1966) On prey-seeking behaviour of aphidophagous insects In: Hodek I (ed) Ecology of aphidophagous insects. Academia/Junk, Prague/The Hague, pp. 123-128
- Bartelt RJ (2010) Volatile hydrocarbon pheromones from beetles. In: Blomquist GJ and Bagnères A (eds) Insect hydrocarbons. Biology, biochemistry, and chemical ecology. Cambridge University Press, Cambridge, UK, pp. 492
- Bartlett AC (1993) Maintaining genetic diversity in laboratory colonies of parasites and predators. In: Narang SK, Bartlett AC and Faust RM (eds) Applications of genetics to arthropods of biological control significance. CRC Press, Inc., Boca Raton, Florida, pp. 133-145
- Beermann A, Jay DG, Beeman RW, Hülskamp M, Tautz D, Jürgens G (2001) The *Short antennae* gene of *Tribolium* is required for limb development and encodes the orthologue of the *Drosophila* *Distal-less* protein. *Development* 128:287-297
- Bengtson SA, Hagen R (1977) Melanism in 2-spot ladybird *Adalia bipunctata* in relation to climate in Western Norway. *Oikos* 28:16-19
- Blackman RL (1965) Studies on specificity in Coccinellidae. *Ann. Appl. Biol.* 56:336-338
- Blackman RL, Eastop VF (1984) Aphids on the world's crops: an identification guide. Wiley-Interscience, Chichester
- Blackman RL, Eastop VF (1994) Aphids on the world's trees. An identification and information guide. CAB International, Wallingford
- Bolckmans KJF (2003) State of affairs and future directions of product quality assurance in Europe. In: Van Lenteren JC (ed) Quality control and production of biological control agents theory and testing procedures. CABI Publishing, Wallingford, UK, pp. 215-224

- Bonte M, Samih MA, De Clercq P (2010) Development and reproduction of *Adalia bipunctata* on factitious and artificial foods. *BioControl* 55:485-491
- Boto L (2010) Horizontal gene transfer in evolution: facts and challenges. *Proc. R. Soc. B* 277:819-827
- Bowsher JH, Wray GA, Abouheif E (2007) Growth and patterning are evolutionarily dissociated in the vestigial wing discs of workers of the red imported fire ant, *Solenopsis invicta*. *J. Exp. Zool. (Mol. Dev. Evol.)* 308B:769-776
- Braendle C, Baer CF, Felix MA (2012) Bias and evolution of the mutationally accessible phenotypic space in a developmental system. *Plos Genetics* 6
- Braendle C, Davis GK, Brisson JA, Stern DL (2006) Wing dimorphism in aphids. *Heredity* 97:192-199
- Brakefield PM (1984a) Ecological studies on the polymorphic ladybird *Adalia bipunctata* in The Netherlands. 1. Population biology and geographical variation of melanism. *J. Anim. Ecol.* 53:761-774
- Brakefield PM (1984b) Ecological studies on the polymorphic ladybird *Adalia bipunctata* in The Netherlands. 2. Population dynamics, differential timing of reproduction and thermal melanism. *J. Anim. Ecol.* 53:775-790
- Brakefield PM (1984c) Selection along clines in the ladybird *Adalia bipunctata* in The Netherlands - a general mating advantage to melanics and its consequences. *Heredity* 53:37-49
- Brakefield PM (1985) Polymorphic Mullerian mimicry and interactions with thermal melanism in ladybirds and a soldier beetle - a hypothesis. *Biol. J. Linnean Soc.* 26:243-267
- Brakefield PM (2006) Evo-devo and constraints on selection. *Trends Ecol. Evol.* 21:362-368
- Brakefield PM, de Jong PW (2011) A steep cline in ladybird melanism has decayed over 25 years: a genetic response to climate change? *Heredity* 107:574-578
- Brakefield PM, Gates J, Keys D, Kesbeke F, Wijngaarden P, Monteiro A, French V, Carroll SB (1996) Development, plasticity and evolution of butterfly eyespot patterns. *Nature* 384:236-242
- Brakefield PM, Willmer PG (1985) The basis of thermal melanism in the ladybird *Adalia bipunctata* - differences in reflectance and thermal properties between the morphs. *Heredity* 54:9-14
- Brisson JA, Davis GK, Stern DL (2007) Common genome-wide patterns of transcript accumulation underlying the wing polyphenisms and polymorphism in the pea aphid (*Acyrtosiphon pisum*). *Evol. Dev.* 9:338-346
- Brisson JA, Ishikawa A, Miura T (2010) Wing development genes of the pea aphid and differential gene expression between winged and unwinged morphs. *Ins. Mol. Biol.* 19:63-73
- Brown PMJ, Adriaens T, Bathon H, Cuppen J, Goldarazena A, Hagg T, Kenis M, Klausnitzer BEM, Kovar I, Loomans AJM, Majerus MEN, Nedved O, Pedersen J, Rabitsch W, Roy HE, Ternois V, Zakharov IA, Roy DB (2008) *Harmonia axyridis* in Europe: spread and distribution of a non-native coccinellid. *BioControl* 53:5-21
- Brunetti CR, Selegue JE, Monteiro A, French V, Brakefield PM, Carroll SB (2001) The generation and diversification of butterfly eyespot color patterns. *Curr. Biol.* 11:1578-1585
- Campbell G, Tomlinson A (1998) The roles of the homeobox genes *aristaless* and *Distal-less* in patterning the legs and wings of *Drosophila*. *Development* 125:4483-4493
- Capinera JL (2008) Encyclopedia of entomology. 2nd edn. Springer, pp. 4346
- Carroll SB, Gates J, Keys DN, Paddock SW, Panganiban GEF, Selegue JE, Williams JA (1994) Pattern formation and eyespot determination in butterfly wings. *Science* 265:109-114
- Carter CI, Nichols JFA (1986) The lime aphid, *Eucallipterus tiliæ*. In: Juniper BE and Southwood TRE (eds) *Insects and the plant surface*. Edward Arnold, London, pp. 342
- Chan CX, Beiko RG, Darling AE, Ragan MA (2009) Lateral transfer of genes and gene fragments in prokaryotes. *Genome Biol. Evol.* 1:429-438

- Chandler CH (2010) Cryptic intraspecific variation in sex determination in *Caenorhabditis elegans* revealed by mutations. *Heredity* 105:473-482
- Cock MJW, van Lenteren JC, Brodeur J, Barratt BIP, Bigler F, Bolckmans K, Consoli FL, Haas F, Mason PG, Parra JRP (2010) Do new access and benefit sharing procedures under the Convention on Biological Diversity threaten the future of biological control? *BioControl* 55:199-218
- Cohen SM (1993) Imaginal disc development. In: Bate M and Martinez Arias A (eds) *The development of Drosophila melanogaster*. Cold Spring Harbour Press, New York, pp. 747-843
- Crawley M (2007) *The R Book*. John Wiley & Sons Ltd, Chichester
- Creed ER (1975) Melanism in 2 spot ladybird - nature and intensity of selection. *Proc. R. Soc. Lond. B* 190:135-148
- Crowson R (1981) *The biology of the Coleoptera*. Academic Press, London
- Dahlsten DL, Zuparko RL, Hajek AE, Rowney DL, Dreistadt SH (1999) Long-term sampling of *Eucallipterus tiliiae* (Homoptera: Drepanosiphidae) and associated natural enemies in a northern California site. *Environ. Entomol.* 28:845-850
- Damen WGM, Saridakis T, Averof M (2002) Diverse adaptations of an ancestral gill: A common evolutionary origin for wings, breathing organs, and spinnerets. *Curr. Biol.* 12:1711-1716
- Danchin E, Charmantier A, Champagne FA, Mesoudi A, Pujol B, Blanchet S (2011) Beyond DNA: integrating inclusive inheritance into an extended theory of evolution. *Nature Rev. Genet.* 12:475-486
- Darlington P (1936) Variation and atrophy of flying wings of some carabid beetles (Coleoptera). *Ann. Entomol. Soc. Am.* 49:136-179
- Darwin CR (1859) *On the origin of species by means of natural selection, or the preservation of favoured races in the struggle for life*. John Murray, London
- Day KR, Docherty M, Leather SR, Kidd NAC (2006) The role of generalist insect predators and pathogens in suppressing green spruce aphid populations through direct mortality and mediation of aphid dropping behavior. *Biol. Control* 38:233-246
- De Clercq P, Bonte M, Van Speybroeck K, Bolckmans K, Deforce K (2005) Development and reproduction of *Adalia bipunctata* (Coleoptera: Coccinellidae) on eggs of *Ephestia kuehniella* (Lepidoptera: Phycitidae) and pollen. *Pest Manag. Sci.* 61:1129-1132
- De Clercq P, Mason PG, Babendreier D (2011) Benefits and risks of exotic biological control agents. *BioControl* 56:681-698
- De Jong PW, Brakefield PM, Geerinck BP (1998) The effect of female mating history on sperm precedence in the two-spot ladybird, *Adalia bipunctata* (Coleoptera, Coccinellidae). *Behav. Ecol.* 9:559-565
- De Jong PW, Gussekloo SWS, Brakefield PM (1996) Differences in thermal balance, body temperature and activity between non-melanistic and melanistic two-spot ladybird beetles (*Adalia bipunctata*) under controlled conditions. *J. Exp. Biol.* 199:2655-2666
- Dekkers JCM, Hospital F (2002) The use of molecular genetics in the improvement of agricultural populations. *Nature Rev. Genet.* 3:22-32
- Den Boer PJ, Van Huizen THP, Den Boer-Daanje W, Aukema B, Den Bieman CFM (1980) Wing polymorphism and dimorphism in ground beetles as stages in an evolutionary process (Coleoptera, Carabidae). *Entomol. Gen.* 6:107-134
- Desender K (1989) Heritability of wing development and body size in a carabid beetle, *Pogonus chalceus* Marsham, and its evolutionary significance. *Oecologia* 78:513-520

- Dib H, Simon S, Sauphanor B, Capowiez Y (2010) The role of natural enemies on the population dynamics of the rosy apple aphid, *Dysaphis plantaginea* Passerini (Hemiptera: Aphididae) in organic apple orchards in south-eastern France. *Biol. Control* 55:97-109
- Dill LM, Fraser AHG, Roitberg BD (1990) The economics of escape behaviour in the pea aphid, *Acyrtosiphon pisum*. *Oecologia* 83:473-478
- Dixon AFG (1971a) Role of aphids in wood formation. 1. Effect of sycamore aphid, *Drepanosiphum platanoides* (Schr) (Aphididae), on growth of sycamore, *Acer pseudoplatanus* (L). *J. Appl. Ecol.* 8:165-179
- Dixon AFG (1971b) Role of aphids in wood formation. 2. Effect of lime aphid, *Eucallipterus tiliæ* L. (Aphididae), on growth of lime, *Tilia x vulgaris* Hayne. *J. Appl. Ecol.* 8:393-399
- Dixon AFG (1971c) Role of intra-specific mechanisms and predation in regulating numbers of lime aphid, *Eucallipterus tiliæ* L. *Oecologia* 8:179-193
- Dixon AFG (1998) Aphid ecology. Chapman & Hall, London
- Dixon AFG (2000) Insect predator-prey dynamics. Ladybird beetles and biological control. Cambridge University Press, Cambridge
- Dixon AFG, Agarwala BK (1999) Ladybird-induced life-history changes in aphids. *Proc. R. Soc. Lond. B* 266:1549-1553
- Dixon AFG, Barlow ND (1979) Population regulation in the lime aphid. *Zool. J. Linn. Soc.* 67:225-237
- Dobzhansky T (1924) Die geographische und individuelle Variabilität von *Harmonia axyridis* Pall in ihren Wechselbeziehungen. *Biol. Zentralblatt* 44 401-421
- Dobzhansky T (1933) Geographical variation in lady-beetles. *Am. Nat.* 67:97-126
- Dolenska M, Nedved O, Vesely P, Tesarova M, Fuchs R (2009) What constitutes optical warning signals of ladybirds (Coleoptera: Coccinellidae) towards bird predators: colour, pattern or general look? *Biol. J. Linnean Soc.* 98:234-242
- Doumbia M, Hemptinne JL, Dixon AFG (1998) Assessment of patch quality by ladybirds: role of larval tracks. *Oecologia* 113:197-202
- Dreistadt SH, Dahlsten DL (1988) Tuliptree aphid honeydew management. *J. Arboriculture* 14:209-214
- Duveau F, Felix MA (2012) Role of pleiotropy in the evolution of a cryptic developmental variation in *Caenorhabditis elegans*. *PLoS Biology* 10
- Dworkin I, Kennerly E, Tack D, Hutchinson J, Brown J, Mahaffey J, Gibson G (2009) Genomic consequences of background effects on *scalloped* mutant expressivity in the wing of *Drosophila melanogaster*. *Genetics* 181:1065-1076
- Dybas HS (1978) Polymorphism in featherwing beetles, with a revision of genus *Ptinellodes* (Coleoptera Ptiliidae). *Ann. Entomol. Soc. Am.* 71:695-714
- Eilenberg J, Hajek AE, Lomer C (2001) Suggestions for unifying the terminology for biological control. *BioControl*:387-400
- Engel MS, Grimaldi DA (2004) New light shed on the oldest insect. *Nature* 427:627-630
- Ettifouri M, Ferran A (1993) Influence of larval rearing diet on the intensive searching behavior of *Harmonia axyridis* (Col, Coccinellidae) larvae. *Entomophaga* 38:51-59
- Evans EW (2004) Habitat displacement of North American ladybirds by an introduced species. *Ecology* 85:637-647
- Falconer DS, Mackay TFC (1996) Introduction to quantitative genetics Longman Group Ltd Harlow, UK

- Ferran A, Giuge L, Tourniaire R, Gambier J, Fournier D (1998) An artificial non-flying mutation to improve the efficiency of the ladybird *Harmonia axyridis* in biological control of aphids. *BioControl* 43:53-64
- Ferran A, Niknam H, Kabiri F, Picart JL, DeHercé C, Brun J, Iperiti G, Lapchin L (1996) The use of *Harmonia axyridis* larvae (Coleoptera: Coccinellidae) against *Macrosiphum rosae* (Hemiptera: Sternorrhyncha: Aphididae) on rose bushes. *Eur. J. Entomol.* 93:59-67
- Fonseca MG, Zarbin PHG (2009) Mating behaviour and evidence for sex-specific pheromones in *Hedypathes betulinus* (Coleoptera: Cerambycidae: Lamiinae). *J. Appl. Entomol.* 133:695-701
- Frankino WA, Zwaan BJ, Stern DL, Brakefield PM (2007) Internal and external constraints in the evolution of morphological allometries in a butterfly. *Evolution* 61:2958-2970
- Fukaya M, Akino T, Yasuda T, Yasui H, Wakamura S (2004) Visual and olfactory cues for mate orientation behaviour in male white-spotted longicorn beetle, *Anoplophora malasiaca*. *Entomol. Exp. Appl.* 111:111-115
- Fukaya M, Akino T, Yasui H, Yasuda T, Wakamura S, Yamamura K (2005) Effects of size and color of female models for male mate orientation in the white-spotted longicorn beetle *Anoplophora malasiaca* (Coleoptera: Cerambycidae). *Appl. Entomol. Zool.* 40:513-519
- Fukaya M, Yasuda T, Wakamura S, Honda H (1996) Reproductive biology of the yellow-spotted longicorn beetle, *Psacotheta hilaris* (Pascoe) (Coleoptera: Cerambycidae). 3. Identification of contact sex pheromone on female body surface. *J. Chem. Ecol.* 22:259-270
- Futuyma DJ (2005) *Evolution*. Sinauer Associates, Sunderland, USA
- Geiselhardt S, Otte T, Hilker M (2009) The role of cuticular hydrocarbons in male mating behavior of the mustard leaf beetle, *Phaedon cochleariae* (F.). *J. Chem. Ecol.* 35:1162-1171
- Gibson G, Dworkin I (2004) Uncovering cryptic genetic variation. *Nature Rev. Genet.* 5:681-690
- Gibson G, van Helden S (1997) Is function of the *Drosophila* homeotic gene *Ultrabithorax* canalized? *Genetics* 147:1155-1168
- Gibson G, Wemple M, van Helden S (1999) Potential variance affecting homeotic *Ultrabithorax* and *Antennapedia* phenotypes in *Drosophila melanogaster*. *Genetics* 151:1081-1091
- Gillespie JH (2001) Is the population size of a species relevant to its evolution? *Evolution* 55:2161-2169
- Gillespie JH (2006) Stochastic processes in evolution. In: Fox CW and Wolf JB (eds) *Evolutionary genetics. Concepts and case studies*. Oxford University Press, Oxford, pp. 65-79
- Ginzel MD (2010) Hydrocarbons as contact pheromones of longhorned beetles (Coleoptera: Cerambycidae). In: Blomquist GJ and Bagnères A (eds) *Insect hydrocarbons. Biology, biochemistry, and chemical ecology*. Cambridge University Press, Cambridge, UK, pp. 492
- Ginzel MD, Blomquist GJ, Millar JG, Hanks LM (2003) Role of contact pheromones in mate recognition in *Xylotrechus colonus*. *J. Chem. Ecol.* 29:533-545
- Ginzel MD, Hanks LM (2003) Contact pheromones as mate recognition cues of four species of longhorned beetles (Coleoptera: Cerambycidae). *J. Ins. Behav.* 16:181-187
- Glen DM, Barlow ND (1980) Interaction of a population of the black-kneed capsid, *Blepharidopterus angulatus*, and its prey, the lime aphid. *Ecol. Entomol.* 5:335-344
- Golden KL, Meinke LJ, Stanleysamuelson DW (1992) Cuticular hydrocarbon discrimination of *Diabrotica* (Coleoptera, Chrysomelidae) sibling species. *Ann. Entomol. Soc. Am.* 85:561-570
- Gotoh A, Sameshima S, Tsuji K, Matsumoto T, Miura T (2005) Apoptotic wing degeneration and formation of an altruism-regulating glandular appendage (gemma) in the ponerine ant *Diacamma* sp. from Japan (Hymenoptera, Formicidae, Ponerinae). *Dev. Genes Evol.* 215:69-77

- Gould F (2008) Broadening the application of evolutionarily based genetic pest management. *Evolution* 62:500-510
- Grimaldi DA, Engel MS (2005) *Evolution of the Insects*. Cambridge University Press, Cambridge
- Gurney B, Hussey NW (1970) Evaluation of some Coccinellid species for biological control of aphids in protected cropping. *Ann. Appl. Biol.* 65:451-8
- Haddrill PR, Shuker DM, Amos W, Majerus MEN, Mayes S (2008) Female multiple mating in wild and laboratory populations of the two-spot ladybird, *Adalia bipunctata*. *Mol. Ecol.* 17:3189-3197
- Haddrill PR, Shuker DM, Mayes S, Majerus MEN (2007) Temporal effects of multiple mating on components of fitness in the two-spot ladybird, *Adalia bipunctata* (Coleoptera : Coccinellidae). *Eur. J. Entomol.* 104:393-398
- Hajek AE (2004) *Natural enemies. An introduction to biological control*. Cambridge University Press, Cambridge, UK
- Haldane JBS (1919) The combination of linkage values, and the calculation of distances between the loci of linked factors. *J. Genet.* 8:299-309
- Hämäläinen M (1976) Rearing the univoltine ladybeetles, *Coccinella septempunctata* and *Adalia bipunctata* (Col., Coccinellidae), all year round in the laboratory. *A.. Agric. Fenn.* 15:66-71
- Hämäläinen M (1977) Control of aphids on sweet peppers, chrysanthemums and roses in small greenhouses using ladybeetles *Coccinella septempunctata* and *Adalia bipunctata* (Col, Coccinellidae). *A.. Agric. Fenn.* 16:117-131
- Hämäläinen M (1980) Evaluation of two native coccinellids for aphid control in glasshouses IOBC/WPRS Bulletin 59-63
- Hämäläinen M, Markkula M (1977) Cool storage of *Coccinella septempunctata* and *Adalia bipunctata* (COL, Coccinellidae) eggs for use in biological control in greenhouses. *A.. Agric. Fenn.* 16:132-136
- Hammond PM (1985) Dimorphism of wings, wing-folding and wing-toiletry devices in the ladybird, *Rhyzobius litura* (F) (Coleoptera, Coccinellidae), with a discussion of inter-population variation in this and other wing-dimorphic beetle species. *Biol. J. Linnean Soc.* 24:15-33
- Hayden EJ, Ferrada E, Wagner A (2011) Cryptic genetic variation promotes rapid evolutionary adaptation in an RNA enzyme. *Nature* 474:92-U120
- Heimbach U (1982) Zur Honigtauproduktion von zwei Callaphididen an Linde und Eiche. *Apidologie* 13:331-332
- Held LI (2002) *Imaginal discs. The genetic and cellular logic of pattern formation*. Cambridge University Press, Cambridge
- Hemptinne JL, Desprets A (1986) Pollen as a spring food for *Adalia bipunctata* L. In: Hodek I (ed) *Ecology of Aphidophaga*. Academia and W. Junk, Prague and Dordrecht pp. 29-35
- Hemptinne JL, Dixon AFG, Adam B (2001) Do males and females of the two-spot ladybird, *Adalia bipunctata* (L.), differ in when they mature sexually? *J. Ins. Behav.* 14:411-419
- Hemptinne JL, Dixon AFG, Coffin J (1992) Attack strategy of ladybird beetles (Coccinellidae) - factors shaping their numerical response. *Oecologia* 90:238-245
- Hemptinne JL, Dixon AFG, Lognay G (1996) Searching behaviour and mate recognition by males of the two-spot ladybird beetle, *Adalia bipunctata*. *Ecol. Entomol.* 21:165-170
- Hemptinne JL, Lognay G, Dixon AFG (1998) Mate recognition in the two-spot ladybird beetle, *Adalia bipunctata*: role of chemical and behavioural cues. *J. Ins. Phys.* 44:1163-1171
- Hemptinne JL, Naisse J (1987) Ecophysiology of the reproductive activity of *Adalia bipunctata* L. *Med.Fac. Landbouw., Rijksuniv. Gent* 52:225-233

- Hermisson J, Wagner GP (2004) The population genetic theory of hidden variation and genetic robustness. *Genetics* 168:2271-2284
- Hiltpold I, Baroni M, Toepfer S, Kuhlmann U, Turlings TCJ (2010) Selection of entomopathogenic nematodes for enhanced responsiveness to a volatile root signal helps to control a major root pest. *J. Exp. Biol.* 213:2417-2423
- Hodek I (1973) *Biology of Coccinellidae*. Dr. W. Junk N.V., The Hague
- Hodek I, Evans EW (2012) Food relationships. In: Hodek I, Van Emden HF and Honěk A (eds) *Ecology and behaviour of the ladybird beetles (Coccinellidae)*. Blackwell Publishing Ltd, Chichester, pp. 141-274
- Hodek I, Honěk A (1996) *Ecology of Coccinellidae*. Kluwer Academic Publishers, Dordrecht/Boston/London
- Hodek I, Ipert G, Hodková M (1993) Long-distance flights in Coccinellidae (Coleoptera). *Eur. J. Entomol.* 90:403-414
- Hodek I, Van Emden HF, Honěk A (2012) *Ecology and behaviour of the ladybird beetles (Coccinellidae)*. Blackwell Publishing Ltd., Chichester
- Hoekstra RF (2003) Adaptive recovery after fitness reduction: the role of population size. In: Van Lenteren JC (ed) *Quality control and production of biological control agents theory and testing procedures*. CABI Publishing, Wallingford, UK, pp. 89-92
- Hoffmann MP, Orfanedes MS, Pedersen LH, Kirkwyland JJ, Hoebeke ER, Ayyappath R (1997) Survey of lady beetles (Coleoptera: Coccinellidae) in sweet corn using yellow sticky cards. *J. Entomol. Sci.* 32:358-369
- Honěk A (1985) Habitat preferences of aphidophagous coccinellids (Coleoptera). *Entomophaga* 30:253-264
- Honěk A (1993) Intraspecific variation in body size and fecundity in insects: a general relationship. *Oikos* 66:483-492
- Hopper KR, Roush RT, Powell W (1993) Management of genetics of biological-control introductions. *Annu. Rev. Entomol.* 38:27-51
- Howard RW, Blomquist GJ (1982) Chemical ecology and biochemistry of insect hydrocarbons *Annu. Rev. Entomol.* 27:149-172
- Howard RW, Blomquist GJ (2005) Ecological, behavioral, and biochemical aspects of insect hydrocarbons. *Annu. Rev. Entomol.* 50:371-393
- Hoy MA (1985) Recent advantages in genetics and genetic-improvement of the Phytoseiidae. *Annu. Rev. Entomol.* 30:345-370
- Hoy MA (1986) Use of genetic improvement in biological control. *Agric. Ecosyst. Environ.* 15:109-119
- Hoy MA (2002) Genetic improvement of biocontrol agents. In: Pimentel D (ed) *Encyclopedia of pest management*. Marcel Dekker, Inc., New York, pp. 329-332
- Hoy MA, Cave FE (1986) Screening for thelyticia in the parahaploid Phytoseiid, *Metaseiulus occidentalis* (Nesbitt). *Exp. Appl. Acarol.* 2:273-276
- Hundertmark A (1935) Die Entwicklung der Flügel des Mehlkäfers *Tenebrio molitor*, mit besonderer Berücksichtigung der Häutungsvorgänge *Z. Morphol. Okol. Tiere* 30:506-543
- Hurst GDD, Sharpe RG, Broomfield AH, Walker LE, Majerus TMO, Zakharov IA, Majerus MEN (1995) Sexually-transmitted disease in a promiscuous insect, *Adalia bipunctata*. *Ecol. Entomol.* 20:230-236
- Ignoffo CM, Garcia C, Dickerson WA, Schmidt GT, Biever KD (1977) Imprisonment of entomophages to increase effectiveness - evaluation of a concept. *J. Econ. Entomol.* 70:292-294
- Iguchi M, Fukushima F, Miura K (2012) Control of *Aphis gossypii* and *Myzus persicae* (Hemiptera: Aphididae) by a flightless strain of *Harmonia axyridis* (Coleoptera: Coccinellidae) on green pepper plants in open fields. *Entomol. Sci.* 15:127-132

- Ikeda H, Nishikawa M, Sota T (2012) Loss of flight promotes beetle diversification. *Nat. Commun.* 3
- Irvin NA, Hoddle MS (2006) The effect of intraspecific competition on progeny sex ratio in *Gonatocerus* spp. for *Homalodisca coagulata* egg masses: Economic implications for mass rearing and biological control. *Biol. Control* 39:162-170
- Ives AR, Kareiva P, Perry R (1993) Response of a predator to variation in prey density at three hierarchical scales: lady beetles feeding on aphids. *Ecology* 74:1929-1938
- Jack J, DeLotto Y (1992) Effect of wing scalloping mutations on *cut* expression and sense organ differentiation in the *Drosophila* wing margin. *Genetics* 131:353-363
- Jackson DJ (1928) The inheritance of long and short wings in the weevil, *Sitona hispidula*, with a discussion of wing reduction among beetles. *Trans. Roy. Soc. Edinburgh*:665-735
- Jalali MA, Michaud JP (2012) Aphid-plant interactions affect the suitability of *Myzus* spp. as prey for the two spot ladybird, *Adalia bipunctata* (Coleoptera: Coccinellidae). *Eur. J. Entomol.* 109:345-352
- Jalali MA, Tirry L, Arbab A, De Clercq P (2010) Temperature-dependent development of the two-spotted ladybeetle, *Adalia bipunctata*, on the green peach aphid, *Myzus persicae*, and a factitious food under constant temperatures. *J. Ins. Sci.* 10
- Jalali MA, Tirry L, De Clercq P (2009) Food consumption and immature growth of *Adalia bipunctata* (Coleoptera: Coccinellidae) on a natural prey and a factitious food. *Eur. J. Entomol.* 106:193-198
- Jennions MD, Petrie M (2000) Why do females mate multiply? A review of the genetic benefits. *Biol. Rev. Cambridge Philosophic. Soc.* 75:21-64
- Jermy T, Butt BA (1991) Method for screening female sex-pheromone extracts of the Colorado potato beetle. *Entomol. Exp. Appl.* 59:75-78
- Jetter K, Paine TD (2004) Consumer preferences and willingness to pay for biological control in the urban landscape. *Biol. Control* 30:312-322
- Johannes F, Colot V, Jansen RC (2008) Epigenome dynamics: a quantitative genetics perspective. *Nature Rev. Genet.* 9:883-890
- Johnson MW, Tabashnik WJ (1993) Laboratory selection for pesticide resistance in natural enemies. In: Narang SK, Bartlett AC and Faust RM (eds) Applications of genetics to arthropods of biological control significance. CRC Press, Inc., Boca Raton, Florida, pp. 91-105
- Kalushkov P (1998) Ten aphid species (Sternorrhyncha : Aphididae) as prey for *Adalia bipunctata* (Coleoptera : Coccinellidae). *Eur. J. Entomol.* 95:343-349
- Kearns PWE, Tomlinson IPM, Odonald P, Veltman CJ (1990) Non-random mating in the two-spot ladybird (*Adalia bipunctata*): I. A reassessment of the evidence. *Heredity* 65:229-240
- Kehrli P, Wyss E (2001) Effects of augmentative releases of the coccinellid, *Adalia bipunctata*, and of insecticide treatments in autumn on the spring population of aphids of the genus *Dysaphis* in apple orchards. *Entomol. Exp. Appl.* 99:245-252
- Kieckhefer RW, Olson GA (1974) Dispersal of marked adult coccinellids from crops in South-Dakota. *J. Econ. Entomol.* 67:52-54
- Kindlmann P, Ameixa O, Dixon AFG (2011) Ecological effects of invasive alien species on native communities, with particular emphasis on the interactions between aphids and ladybirds. *BioControl* 56:469-476
- Kitami T, Nadeau JH (2002) Biochemical networking contributes more to genetic buffering in human and mouse metabolic pathways than does gene duplication. *Nature Genet.* 32:191-194
- Knell RJ, Webberley KM (2004) Sexually transmitted diseases of insects: distribution, evolution, ecology and host behaviour. *Biol. Rev. Cambridge Philosophic. Soc.* 79:557-581

- Koch RL (2003) The multicolored Asian lady beetle *Harmonia axyridis*. A review of its biology, uses in biological control, and non-target impacts. J. Ins. Sci. 3:1-16
- Koonin EV (2009) Darwinian evolution in the light of genomics. Nucleic Acids Res. 37:1011-1034
- Kuroda T, Miura K (2003) Comparison of the effectiveness of two methods for releasing *Harmonia axyridis* (Pallas) (Coleoptera : Coccinellidae) against *Aphis gossypii* Glover (Homoptera : Aphididae) on cucumbers in a greenhouse. Appl. Entomol. Zool. 38:271-274
- Kuwayama H, Yaginuma T, Yamashita O, Niimi T (2006) Germ-line transformation and RNAi of the ladybird beetle, *Harmonia axyridis*. Ins. Mol. Biol. 15:507-512
- Laland KN, Sterelny K, Odling-Smee J, Hoppitt W, Uller T (2011) Cause and effect in biology revisited: Is Mayr's proximate-ultimate dichotomy still useful? Science 334:1512-1516
- Le Rouzic A, Carlborg O (2008) Evolutionary potential of hidden genetic variation. Trends Ecol. Evol. 23:33-37
- Lelito JP, Domingue MJ, Fraser I, Mastro VC, Tumlinson JH, Baker TC (2011) Field investigation of mating behaviour of *Agrilus cyanescens* and *Agrilus subcinctus*. Can. Entomol. 143:370-379
- Lelito JP, Fraser I, Mastro VC, Tumlinson JH, Boroczky K, Baker TC (2007) Visually mediated 'paratrooper copulations' in the mating behavior of *Agrilus planipennis* (Coleoptera : Buprestidae), a highly destructive invasive pest of North American ash trees. J. Ins. Behav. 20:537-552
- Leppla NC (2003) Aspects of total quality control for the production of natural enemies. In: Van Lenteren JC (ed) Quality control and production of biological control agents theory and testing procedures. CABI Publishing, Wallingford, UK, pp. 19-24
- Lindsley DL, Zimm GG (1992) The genome of *Drosophila melanogaster*. Academic Press, London
- Lobbia S, Niitsu S, Fujiwara H (2003) Female-specific wing degeneration caused by ecdysteroid in the Tussock Moth, *Orgyia recens*: hormonal and developmental regulation of sexual dimorphism. J. Ins. Sci. 3:11
- Lommen STE, De Jong PW, Brakefield PM (2005) Phenotypic plasticity of elytron length in wingless two-spot ladybird beetles, *Adalia bipunctata* (Coleoptera: Coccinellidae). Eur. J. Entomol. 102:553-556
- Lommen STE, De Jong PW, Koops KG, Brakefield PM (2012) Genetic linkage between melanism and winglessness in the ladybird beetle *Adalia bipunctata*. Genetica 140: 229–233 [CHAPTER 4]
- Lommen STE, Holness TC, Van Kuik AJ, De Jong PW, Brakefield PM (2013) Releases of a natural flightless strain of the ladybird beetle *Adalia bipunctata* reduce aphid-born honeydew beneath urban lime trees. BioControl 58: 195-204 [CHAPTER 3]
- Lommen STE, Middendorp CW, Luijten CA, Van Schelt J, Brakefield PM, De Jong PW (2008) Natural flightless morphs of the ladybird beetle *Adalia bipunctata* improve biological control of aphids on single plants. Biol. Control 47:340-346 [CHAPTER 2]
- Lommen STE, Saenko SV, Tomoyasu Y, Brakefield PM (2009) Development of a wingless morph in the ladybird beetle, *Adalia bipunctata*. Evol. Dev. 11:278-289 [CHAPTER 6]
- Lorenz MW (2007) Oogenesis-flight syndrome in crickets: Age-dependent egg production, flight performance, and biochemical composition of the flight muscles in adult female *Gryllus bimaculatus*. J. Ins. Phys. 53:819-832
- Losey JE, Denno RF (1998) The escape response of pea aphids to foliar-foraging predators: factors affecting dropping behaviour. Ecol. Entomol. 23:53-61
- Lozier JD, Roderick GK, Mills NJ (2008) Evolutionarily significant units in natural enemies: Identifying regional populations of *Aphidius transcaspicus* (Hymenoptera : Braconidae) for use in biological control of mealy plum aphid. Biol. Control 46:532-541
- Luo SL, Zhuge PP, Wang MQ (2011) Mating behavior and contact pheromones of *Batocera horsfieldi* (Hope) (Coleoptera: Cerambycidae). Entomol. Sci. 14:359-363

- Lusis JJ (1961) On the biological meaning of colour polymorphism of lady-beetle *Adalia bipunctata* L. Latvijas Entomologs 4:3-29
- Lynch M (2007) The frailty of adaptive hypotheses for the origins of organismal complexity. Proc. Natl. Acad. Sci. U. S. A. 104:8597-8604
- Mackauer M (1976) Genetic problems in production of biological-control agents. Annu. Rev. Entomol. 21:369-385
- Majerus M, Odonald P, Weir J (1982) Evidence for preferential mating in *Adalia bipunctata*. Heredity 49:37-49
- Majerus MEN (1994) Ladybirds. Collins, London
- Majerus MEN (1998) Melanism : evolution in action. Oxford University Press, Oxford [etc.]
- Majerus MEN, Kearns PWE (1989) Ladybirds. Richmond Publishing Company, Slough
- Majerus MEN, Odonald P, Kearns PWE, Ireland H (1986) Genetics and evolution of female choice. Nature 321:164-167
- Majerus MEN, Strawson V, Roy HE (2006) The potential impacts of the arrival of the harlequin ladybird *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae), in Britain. Ecol. Entomol. 31:207-215
- Majerus MEN, Zakharov IA (2000) Does thermal melanism maintain melanic polymorphism in the two-spot ladybird *Adalia bipunctata*. (Coleoptera: Coccinellidae)? Zhurnal Obshchei Biologii 61:381-392
- Maki H (2002) Origins of spontaneous mutations: Specificity and directionality of base-substitution, frameshift, and sequence-substitution mutageneses. Annu. Rev. Genet. 36:279-303
- Mally CW (1916) On the selection and breeding of desirable strains of beneficial insects. S. Afr. J. Sci.:369-385
- Mann JA, Tatchell GM, Dupuch MJ, Harrington R, Clark SJ, McCartney HA (1995) Movement of apterous *Sitobion avenae* (Homoptera, Aphididae) in response to leaf disturbances caused by wind and rain. Ann. Appl. Biol. 126:417-427
- Marples NM, Brakefield PM, Cowie RJ (1989) Differences between the seven-spot and two-spot ladybird beetles (Coccinellidae) in their toxic effects on a bird predator. Ecol. Entomol. 14:79-84
- Marples NM, De Jong PW, Ottenheim MM, Verhoog MD, Brakefield PM (1993) The inheritance of a wingless character in the 2-spot ladybird (*Adalia bipunctata*). Entomol. Exp. Appl. 69:69-73
- Marples NM, Van Veelen W, Brakefield PM (1994) The relative importance of color, taste and smell in the protection of an aposematic insect *Coccinella septempunctata*. Anim. Behav. 48:967-974
- Martini X, Haccou P, Olivieri I, Hemptinne JL (2009) Evolution of cannibalism and female's response to oviposition-detering pheromone in aphidophagous predators. J. Anim. Ecol. 78:964-972
- Masel J (2006) Cryptic genetic variation is enriched for potential adaptations. Genetics 172:1985-1991
- Mawson CA (1948) Honeydew on lime trees. Nature 161:176-176
- Mayr E (1961) Cause and effect in biology - kinds of causes, predictability, and teleology are viewed by a practicing biologist. Science 134:1501-1506
- Mayr E (2002) What evolution is. Basic Books, New York
- McGuigan K, Sgro CM (2009) Evolutionary consequences of cryptic genetic variation. Trends Ecol. Evol. 24:305-311
- McLellan ID (1999) A revision of *Zelandoperla* Tillyard (Plecoptera: Gripopterygidae: Zelandoperlinae). N. Z. J. Zool. 26:199-219
- Michaud JP (2012) Coccinellids in biological control. In: Hodek I, Van Emden HF and Honěk A (eds) Ecology and behaviour of the ladybird beetles (Coccinellidae). Blackwell Publishing Ltd, Chichester, pp. 488-519

- Michie LJ, Mallard F, Majerus MEN, Jiggins FM (2010) Melanic through nature or nurture: genetic polymorphism and phenotypic plasticity in *Harmonia axyridis*. J. Evol. Biol. 23:1699-1707
- Michie LJ, Masson A, Ware RL, Jiggins FM (2011) Seasonal phenotypic plasticity: wild ladybirds are darker at cold temperatures. Evol. Ecol. 25:1259-1268
- Milloz J, Duvéau F, Nuez I, Felix MA (2008) Intraspecific evolution of the intercellular signaling network underlying a robust developmental system. Genes Dev. 22:3064-3075
- Mills NJ (1979) *Adalia bipunctata* (L.) as a generalist predator of aphids. University East Anglia, Norwich,
- Mills NJ (1981) Essential and alternative foods for some British Coccinellidae. Entomol. Gaz. 32:197-202
- Mills NJ (1982) Voracity, cannibalism and coccinellid predation. Ann. Appl. Biol. 101:144-148
- Ming QL, Lewis SM (2010) Mate recognition and sex differences in cuticular hydrocarbons of the diurnal firefly *Ellychnia corrusca* (Coleoptera: Lampyridae). Ann. Entomol. Soc. Am. 103:128-133
- Minks AK, Harrewijn P (1987) Aphids, their biology, natural enemies and control. Elsevier, Amsterdam
- Minoretti N, Weisser WW (2000) The impact of individual ladybirds (*Coccinella septempunctata*, Coleoptera: Coccinellidae) cell aphid colonies. Eur. J. Entomol. 97:475-479
- Mole S, Zera AJ (1993) Differential allocation of resources underlies the dispersal-reproduction trade-off in the wing-dimorphic cricket, *Gryllus rubens*. Oecologia 93:121-127
- Morjan CL, Rieseberg LH (2004) How species evolve collectively: implications of gene flow and selection for the spread of advantageous alleles. Mol. Ecol. 13:1341-1356
- Mutis A, Parra L, Palma R, Pardo F, Perich F, Quiroz A (2009) Evidence of contact pheromone use in mating behavior of the raspberry weevil (Coleoptera: Curculionidae). Environ. Entomol. 38:192-197
- Nachappa P, Margolies DC, Nechols JR, Morgan TJ (2010) Response of a complex foraging phenotype to artificial selection on its component traits. Evol. Ecol. 24:631-655
- Nadeau JH (2001) Modifier genes in mice and humans. Nature Rev. Genet. 2:165-174
- Nahmad M, Glass L, Abouheif E (2008) The dynamics of developmental system drift in the gene network underlying wing polyphenism in ants: a mathematical model. Evol. Dev. 10:360-374
- Narang SK, Tabashnik WJ, Faust RM (1993) Complexities of population genetic structure and implications for biological control programs. In: Narang SK, Bartlett AC and Faust RM (eds) Applications of genetics to arthropods of biological control significance. CRC Press, Inc., Boca Raton, Florida, pp. 19-52
- Nelson EH, Matthews CE, Rosenheim JA (2004) Predators reduce prey population growth by inducing changes in prey behavior. Ecol. Lett. 85:1853-1858
- Ng M, Diaz-Benjumea FJ, Cohen SM (1995) *nubbin* encodes a POU-domain protein required for proximal-distal patterning in the *Drosophila* wing. Development 121:589-599
- Ng M, Diaz-Benjumea FJ, Vincent J, Wu J, Cohen SM (1996) Specification of the wing by localized expression of wingless protein. Nat. Commun. 381:316-318
- Niimi T, Kuwayama H, Yaginuma T (2005) Larval RNAi applied to the analysis of postembryonic development in the ladybird beetle, *Harmonia axyridis*. J. Insect Biotechnol. Sericology 74:95-102
- Niitsu S (2001) Wing degeneration due to apoptosis in the female of the winter moth, *Nyssiodes lefuarius*. Entomol. Sci. 4:1-7
- Niitsu S (2003) Postembryonic development of the wing imaginal discs in the female wingless bagworm moth *Eumeta variegata* (Lepidoptera, Psychidae). J. Morphol. 257:164-170

- Nunney L (2003) Managing captive populations for release: a population-genetic perspective. In: Van Lenteren JC (ed) Quality control and production of biological control agents theory and testing procedures. CABI Publishing, Wallingford, UK, pp. 73-88
- Obyrcki JJ, Harwood JD, Kring TJ, O'Neil RJ (2009) Aphidophagy by Coccinellidae: Application of biological control in agroecosystems. *Biol. Control* 51:244-254
- Obyrcki JJ, Kring TJ (1998) Predaceous Coccinellidae in biological control. *Annu. Rev. Entomol.* 43:295-321
- Ode PJ, Hardy ICW (2008) Parasitoid sex ratios and biological control. In: Wajnberg E, Bernstein C and Van Alphen J (eds) Behavioral ecology of insect parasitoids. From theoretical approaches to field applications. Blackwell Publishing, Malden, Massachusetts, pp. 253-291
- Ohde T, Masumoto M, Morita-Miwa M, Matsuura H, Yoshioka H, Yaginuma T, Niimi T (2009) *Vestigial* and *scalloped* in the ladybird beetle: a conserved function in wing development and a novel function in pupal ecdysis. *Ins. Mol. Biol.* 18:571-581
- Olszak RW (1986) The effectiveness of three species of coccinellid adults in controlling small colonies of the green apple aphid (*Aphis pomi*). In: Hodek I (ed) Ecology of Aphidophaga. Academia/Junk, Prague/Dordrecht, pp. 381-384
- Omkar, Pervez A (2005) Ecology of two-spotted ladybird, *Adalia bipunctata*: a review. *J. Appl. Entomol.* 129:465-474
- Panganiban G, Sebring A, Nagy L, Carroll SB (1995) The development of crustacean limbs and the evolution of arthropods. *Science* 270:1363-1366
- Patel NH, Kornberg TB, Goodman CS (1989) Expression of *engrailed* during segmentation in grasshopper and crayfish. *Development* 107:201-212
- Perry JC, Rowe L (2008) Neither mating rate nor spermatophore feeding influences longevity in a ladybird beetle. *Ethology* 114:504-511
- Perry JC, Sharpe DMT, Rowe L (2009) Condition-dependent female remating resistance generates sexual selection on male size in a ladybird beetle. *Anim. Behav.* 77:743-748
- Peschke K (1978) Female sex-pheromone of a Staphylinid beetle, *Aleochara curtula*. *J. Ins. Phys.* 24:197-200
- Peschke K (1979) Tactile orientation by mating males of the staphylinid beetle, *Aleochara curtula*, relative to the setal fields of the female. *Phys. Entomol.* 4:155-159
- Peschke K (1987) Cuticular hydrocarbons regulate mate recognition, male aggression, and female choice of the rove beetle, *Aleochara curtula*. *J. Chem. Ecol.* 13:1993-2008
- Peterson MA, Dobler S, Larson EL, Juarez D, Schlarbaum T, Monsen KJ, Francke W (2007) Profiles of cuticular hydrocarbons mediate male mate choice and sexual isolation between hybridising *Chrysochus* (Coleoptera : Chrysomelidae). *Chemoecology* 17:87-96
- Polaczyk PJ, Gasperini R, Gibson G (1998) Naturally occurring genetic variation affects *Drosophila* photoreceptor determination. *Dev. Genes Evol.* 207:462-470
- Pope RD (1977) Brachyptery and wing polymorphism among Coccinellidae (Coleoptera). *Syst. Entomol.* 2:59-66
- Powell W, Pell JK (2007) Biological Control. In: Van Emden HF and Harrington R (eds) Aphids as crop pests. CAB International, pp. 469-513
- Pureswaran DS, Poland TM (2009) The role of olfactory cues in short-range mate finding by the emerald ash borer, *Agrilus planipennis* Fairmaire (Coleoptera: Buprestidae). *J. Ins. Behav.* 22:205-216
- Quennedey A, Quennedey B (1990) Morphogenesis of the Wing Anlagen in the mealworm beetle *Tenebrio molitor* during the last larval instar. *Tissue Cell* 22:721-740

- Quennedey A, Quennedey B (1999) Development of the wing discs of *Zophobas atratus* under natural and experimental conditions: occurrence of a gradual larval-pupal commitment in the epidermis of tenebrionid beetles. *Cell Tissue Res.* 296:619-634
- R Development Core Team (2007) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria
- Ribaut JM, Hoisington D (1998) Marker-assisted selection: new tools and strategies. *Trends Plant Sci.* 3:236-239
- Roff DA (1986) The evolution of wing dimorphism in insects. *Evolution* 40:1009-1020
- Roff DA (1990) The evolution of flightlessness in insects. *Ecol. Monogr.* 60:389-421
- Roff DA (1994a) The evolution of flightlessness: is history important? *Evol. Ecol.* 8:639-657
- Roff DA (1994b) Why is there so much genetic variation for wing dimorphism? *Res. Popul. Ecol.* 36:145-150
- Roff DA, Fairbairn DJ (1991) Wing dimorphisms and the evolution of migratory polymorphisms among the Insecta. *Am. Zool.* 31:243-251
- Roff DA, Mostow S, Fairbairn DJ (2002) The evolution of trade-offs: Testing predictions on response to selection and environmental variation. *Evolution* 56:84-95
- Rosenheim JA, Hoy MA (1988) Genetic-improvement of a parasitoid biological-control agent - artificial selection for insecticide resistance in *Aphytis melinus* (Hymenoptera, Aphelinidae). *J. Econ. Entomol.* 81:1539-1550
- Roy HE, Wajnberg E (2008) From biological control to invasion: the ladybird *Harmonia axyridis* as a model species. *BioControl* 53:1-4
- Rutherford SL (2000) From genotype to phenotype: buffering mechanisms and the storage of genetic information. *BioEssays* 22:1095-1105
- Rutherford SL (2003) Between genotype and phenotype: Protein chaperones and evolvability. *Nature Rev. Genet.* 4:263-274
- Rutherford SL, Lindquist S (1998) Hsp90 as a capacitor for morphological evolution. *Nature* 396:336-342
- Salame L, Glazer I, Chubinishvili MT, Chkhubianishvili T (2010) Genetic improvement of the desiccation tolerance and host-seeking ability of the entomopathogenic nematode *Steinernema feltiae*. *Phytoparasitica* 38:359-368
- Sameshima S, Miura T, Matsumoto T (2004) Wing disc development during caste differentiation in the ant *Pheidole megacephala* (Hymenoptera: Formicidae). *Evol. Dev.* 6:336-341
- Sangster TA, Salathia N, Lee HN, Watanabe E, Schellenberg K, Morneau K, Wang H, Undurraga S, Queitsch C, Lindquist S (2008a) HSP90-buffered genetic variation is common in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* 105:2969-2974
- Sangster TA, Salathia N, Undurraga S, Milo R, Schellenberg K, Lindquist S, Queitsch C (2008b) HSP90 affects the expression of genetic variation and developmental stability in quantitative traits. *Proc. Natl. Acad. Sci. U. S. A.* 105:2963-2968
- Scali V, Creed ER (1975) Influence of climate on melanism in two-spot ladybird, *Adalia bipunctata*, in central Italy. *Trans. Roy. Entomol. Soc. Lond.* 127:163-169
- Schlichting CD (2008) Hidden reaction norms, cryptic genetic variation, and evolvability. *Ann. N. Y. Acad. Sci.* 1133:187-203
- Secretariat of the International Plant Protection Convention, FAO (2005) Guidelines for the export, shipment, import and release of biological control agents and other beneficial organisms. Food and Agriculture Organization of the United Nations, Rome
- Seko T, Yamashita K, Miura K (2008) Residence period of a flightless strain of the ladybird beetle *Harmonia axyridis* Pallas (Coleoptera: Coccinellidae) in open fields. *Biol. Control* 47:194-198

- Shapiro DI, Glazer I, Segal D (1997) Genetic improvement of heat tolerance in *Heterorhabditis bacteriophora* through hybridization. *Biol. Control* 8:153-159
- Smith DS (1964) Structure and development of flightless Coleoptera - light and electron microscopic study of wings, thoracic exoskeleton, and rudimentary flight musculature. *J. Morphol.* 114:107-183
- Smith SG (1953) Chromosome numbers of Coleoptera. *Heredity* 7:31-48
- Sokal RR, Rohlf FJ (1995) *Biometry: The principles and practice of statistics in biological research*. W.H. Freeman, New York
- Sollars V, Lu XY, Xiao L, Wang XY, Garfinkel MD, Ruden DM (2003) Evidence for an epigenetic mechanism by which Hsp90 acts as a capacitor for morphological evolution. *Nature Genet.* 33:70-74
- Spofford JB (1956) The relation between expressivity and selection against eyeless in *Drosophila melanogaster*. *Genetics* 41:938-959
- Steiger S, Peschke K, Francke W, Muller JK (2007) The smell of parents: breeding status influences cuticular hydrocarbon pattern in the burying beetle *Nicrophorus vespilloides*. *Proc. R. Soc. B* 274:2211-2220
- Steiger S, Whitlow S, Peschke K, Muller JK (2009) Surface chemicals inform about sex and breeding status in the biparental burying beetle *Nicrophorus vespilloides*. *Ethology* 115:178-185
- Strauch O, Oestergaard J, Hollmer S, Ehlers RU (2004) Genetic improvement of the desiccation tolerance of the entomopathogenic nematode *Heterorhabditis bacteriophora* through selective breeding. *Biol. Control* 31:218-226
- Sugeno W, Hori M, Matsuda K (2006) Identification of the contact sex pheromone of *Gastrophysa atrocyanea* (Coleoptera : Chrysomelidae). *Appl. Entomol. Zoolog.* 41:269-276
- Tabone E, Bardon C, Desneux N, Wajnberg E (2010) Parasitism of different *Trichogramma* species and strains on *Plutella xylostella* L. on greenhouse cauliflower. *Journal of Pest Science* 83:251-256
- Tanigaki T, Yamaoka R, Sota T (2007) The role of cuticular hydrocarbons in mating and conspecific recognition in the closely related longicorn beetles *Pidonia grallatrix* and *P. takechii*. *Zool. Sci.* 24:39-45
- Tauber CA, Tauber MJ, Giffin JG (2007) Flightless Hawaiian Hemerobiidae (Neuroptera): Comparative morphology and biology of a brachypterous species, its macropterous relative and intermediate forms. *Eur. J. Entomol.* 104:787-800
- Tautz D (1992) Redundancies, development and the flow of information. *BioEssays* 14:263-266
- Thayer MK (1992) Discovery of sexual wing dimorphism in Staphylinidae (Coleoptera) - *Omalius flavidum*, and a discussion of wing dimorphism in insects. *Journal of the New York Entomological Society* 100:540-573
- Tian DC, Wang Q, Zhang PF, Araki H, Yang SH, Kreitman M, Nagylaki T, Hudson R, Bergelson J, Chen JQ (2008) Single-nucleotide mutation rate increases close to insertions/deletions in eukaryotes. *Nature* 455:105-108
- Timofeeff-Ressovsky NW (1940) Zur Analyse des Polymorphismus bei *Adalia bipunctata* L. *Biol. Zentralblatt* 60:130-137
- Tomanovic Z, Kavallieratos NG, Stary P, Petrovic-Obradovic O, Athanassiou CG, Stanisavljevic LZ (2008) Cereal aphids (Hemiptera : Aphidoidea) in Serbia: Seasonal dynamics and natural enemies. *Eur. J. Entomol.* 105:495-501
- Tomlinson IPM, Kearns PWE, Veltman CJ (1995) Nonrandom mating in the 2-spot ladybird (*Adalia bipunctata*) - the influence of weight on mating success. *Behavior Genetics* 25:467-474
- Tomoyasu Y, Denell RE (2004) Larval RNAi in *Tribolium* (Coleoptera) for analyzing adult development. *Dev. Genes Evol.* 214:575-578
- Tomoyasu Y, Wheeler SR, E. DR (2005) *Ultrabithorax* is required for membranous wing identity in the beetle *Tribolium castaneum*. *Nature* 433:643-647

- Tourniaire R, Ferran A, Gambier J, Giuge L, Bouffault F (1999) Locomotor behavior of flightless *Harmonia axyridis* Pallas (Col., Coccinellidae). *J. Ins. Behav.* 12:545-558
- Tourniaire R, Ferran A, Giuge L, Piotte C, Gambier J (2000) A natural flightless mutation in the ladybird, *Harmonia axyridis*. *Entomol. Exp. Appl.* 96:33-38
- Tower WL (1903) The origin and development of the wings of Coleoptera. *Zool. Jahr. Abt. Anat. Ontog.* Thiere:517-572
- Tribolium* Genome Sequence Consortium (2008) The genome of the model beetle and pest *Tribolium castaneum*. *Nature* 452:949-955
- Trouvé C, Ledee S, Ferran A, Brun J (1997) Biological control of the damson-hop aphid, *Phorodon humuli* (Hom.: Aphididae), using the ladybeetle *Harmonia axyridis* (Col.: Coccinellidae). *Entomophaga* 42:57-62
- True JR (2003) Insect melanism: the molecules matter. *Trends Ecol. Evol.* 18:640-647
- Ueno H, de Jong PW, Brakefield PM (2004) Genetic basis and fitness consequences of winglessness in the two-spot ladybird beetle, *Adalia bipunctata*. *Heredity* 93:283-289
- Van 't Hof AE, Edmonds N, Dalikova M, Marec F, Saccheri IJ (2011) Industrial melanism in British peppered moths has a singular and recent mutational origin. *Science* 332:958-960
- Van 't Hof AE, Saccheri IJ (2010) Industrial melanism in the peppered moth is not associated with genetic variation in canonical melanisation gene candidates. *Plos One* 5
- Van Driesche R, Hoddle M, Center T (2008) Control of pests and weeds by natural enemies: an introduction to biological control. Blackwell Publishing
- Van Houten YM, Van Stratum P, Bruin J, Veerman A (1995) Selection for non-diapause in *Amblyseius cucumeris* and *Amblyseius barkeri* and exploration of the effectiveness of selected strains for thrips control. *Entomol. Exp. Appl.* 77:289-295
- Van Lenteren JC (1988) Evaluating the effectiveness of natural enemies. In: Niemczyk E and Dixon AFG (eds) *Ecology and Effectiveness of Aphidophaga*. SPB Academic Publishing, Den Haag, pp. 175–181
- Van Lenteren JC (2003a) Commercial availability of biological control agents. In: Van Lenteren JC (ed) *Quality control and production of biological control agents theory and testing procedures*. CABI Publishing, Wallingford, UK, pp. 167-180
- Van Lenteren JC (2003b) Need for quality control of mass-produced biological control agents. In: Van Lenteren JC (ed) *Quality control and production of biological control agents theory and testing procedures*. CABI Publishing, Wallingford, UK, pp. 1-18
- Van Lenteren JC (2003c) Quality control and production of biological control agents. Theory and testing procedures. CABI Publishing, Wallingford, UK, pp. 327
- Van Lenteren JC (2012) The state of commercial augmentative biological control: plenty of natural enemies, but a frustrating lack of uptake. *BioControl* 57:1-20
- Van Lenteren JC, Cock MJW, Brodeur J, Barratt BIP, Bigler F, Bolckmans K, Haas F, Mason PG, Parra JRP (2011) Will the Convention on Biological Diversity put an end to biological control? *Rev. Bras. Entomol.* 55:1-5
- Van Lenteren JC, Hale A, Klapwijk JN, Van Schelt J, Steinberg S (2003) Guidelines for quality control of commercially produced natural enemies. In: Van Lenteren JC (ed) *Quality control and production of biological control agents theory and testing procedures*. CABI Publishing, Wallingford, UK, pp. 265-304
- Van Lenteren JC, Loomans AJM, Babendreier D, Bigler F (2008) *Harmonia axyridis*: an environmental risk assessment for Northwest Europe. *BioControl* 53:37-54
- Vink CJ, Barratt BIP, Phillips CB, Barton DM (2012) Moroccan specimens of *Microctonus aethiopoides* spice our understanding of genetic variation in this internationally important braconid parasitoid of adult weevils. *BioControl* 57:751-758

- Vogler AP, Timmermans M (2012) Speciation: don't fly and diversify? *Curr. Biol.* 22:R284-R286
- Waddington CH (1957) *The strategy of the genes*. Allen and Unwin, London
- Wagner DL, Liebherr JK (1992) Flightlessness in insects. *Trends Ecol. Evol.* 7:216-220
- Wajnberg E (1991) Quality control of mass-reared arthropods: A genetical and statistical approach. In: Bigler F (ed) *Proceedings of the 5th Workshop on Quality Control of mass-reared arthropods*, Wageningen, Netherlands, pp. 15-25
- Wajnberg E (2004) Measuring genetic variation in natural enemies used for biological control: Why and how? In: Ehler L, Sforza R and Mateille T (eds) *Genetics, Evolution and Biological Control*. CAB International, Wallingford, pp. 19-37
- Wajnberg E (2010) Genetics of the behavioral ecology of egg parasitoids. In: Consoli FL, Parra JRP and Zucchi RA (eds) *Egg parasitoids in agroecosystems with emphasis on Trichogramma*. Springer Dordrecht, pp. 149-156
- Wajnberg E, Curty C, Jervis M (2012) Intra-population genetic variation in the temporal pattern of egg maturation in a parasitoid wasp. *Plos One* 7
- Ward JP (1981) Mating behaviour and the mechanism of male orientation in the anobiid bread beetle, *Stegobium paniceum*. *Phys. Entomol.* 6:213-217
- Webberley KM, Hurst GDD, Buszko J, Majerus MEN (2002) Lack of parasite-mediated sexual selection in a ladybird/sexually transmitted disease system. *Anim. Behav.* 63:131-141
- Webberley KM, Hurst GDD, Husband RW, Schulenburg J, Sloggett JJ, Isham V, Buszko J, Majerus MEN (2004) Host reproduction and a sexually transmitted disease: causes and consequences of *Coccipolipus hippodamiae* distribution on coccinellid beetles. *J. Anim. Ecol.* 73:1-10
- Webberley KM, Tinsley MC, Sloggett JJ, Majerus MEN, Hurst GDD (2006) Spatial variation in the incidence of a sexually transmitted parasite of the ladybird beetle *Adalia bipunctata* (Coleoptera : Coccinellidae). *Eur. J. Entomol.* 103:793-797
- Weissenberger A, Brun J, Piotte C, Ferran A (1999) Comparison between the wild type and flightless type of the coccinellid *Harmonia axyridis* (Pallas) in the control of the damson hop aphid *Phorodon humuli* (Schränk). Conference: Fifth International Conference on Pests in Agriculture pp. 727-734
- Weisser WW, Braendle C, Minoretti N (1999) Predator-induced morphological shift in the pea aphid. *Proc. R. Soc. Lond. B* 266:1175-1181
- White EB, Debach P, Garber MJ (1970) Artificial selection for genetic adaptation to temperature extremes in *Aphytis lingnanensis* Compere (Hymenoptera-Aphelinidae). *Hilgardia* 40:161-192
- Wittkopp PJ, Beldade P (2009) Development and evolution of insect pigmentation: Genetic mechanisms and the potential consequences of pleiotropy. *Sem. Cell Dev. Biol.* 20:65-71
- Wittkopp PJ, Carroll SB, Kopp A (2003) Evolution in black and white: genetic control of pigment patterns in *Drosophila*. *Trends Genet.* 19:495-504
- Wratten SD (1973) Effectiveness of coccinellid beetle, *Adalia bipunctata* (L), as a predator of lime aphid, *Eucallipterus tiliae* L. *J. Anim. Ecol.* 42:785-802
- Wratten SD (1976) Searching by *Adalia bipunctata* L (Coleoptera-Coccinellidae) and escape behavior of its aphid and cicadellid prey on lime (*Tilia x vulgaris* Hayne). *Ecol. Entomol.* 1:139-142
- Wyss E (1995) The effect of weed strips on aphids and aphidophagous predators in an apple orchard *Entomol. Exp. Appl.* 75:43-49
- Wyss E, Villiger M, Hemptinne JL, Müller-Schärer H (1999a) Effects of augmentative releases of eggs and larvae of the ladybird beetle, *Adalia bipunctata*, on the abundance of the rosy apple aphid, *Dysaphis plantaginea*, in organic apple orchards. *Entomol. Exp. Appl.* 90:167-173

- Wyss E, Villiger M, Müller-Schärer H (1999b) The potential of three native insect predators to control the rosy apple aphid, *Dysaphis plantaginea*. *BioControl* 44:171-182
- Yasuda H, Dixon AFG (2002) Sexual size dimorphism in the two spot ladybird beetle *Adalia bipunctata*: developmental mechanism and its consequences for mating. *Ecol. Entomol.* 27:493-498
- Yasui H (2009) Chemical communication in mate location and recognition in the white-spotted longicorn beetle, *Anoplophora malasiaca* (Coleoptera: Cerambycidae). *Appl. Entomol. Zoolog.* 44:183-194
- Yasui H, Akino T, Yasuda T, Fukaya M, Ono H, Wakamura S (2003) Ketone components in the contact sex pheromone of the white-spotted longicorn beetle, *Anoplophora malasiaca*, and pheromonal activity of synthetic ketones. *Entomol. Exp. Appl.* 107:167-176
- Yasui H, Inouchi J, Wakamura S, Fukaya M, Akino T, Arakaki N (2012) Male mate searching behavior in the black chafer *Holotrichia loochooana loochooana* (Sawada) (Coleoptera: Scarabaeidae): response to pheromone lures of various colors and intensities. *J. Ethol.* 30:233-238
- Zakharov IA, Goryacheva, II (1998) A study of nonrandom mating in the polymorphic *Adalia bipunctata* populations of the Crimea. *Genetika* 34:1090-1093
- Zeh JA, Zeh DW (1997) The evolution of polyandry. 2. Post-copulatory defences against genetic incompatibility. *Proc. R. Soc. Lond. B* 264:69-75
- Zera AJ, Denno RF (1997) Physiology and ecology of dispersal polymorphism in insects. *Annu. Rev. Entomol.* 42:207-230
- Zera AJ, Harshman LG (2001) The physiology of life history trade-offs in animals. *Annu. Rev. Ecol. Syst.* 32:95-126
- Zera AJ, Mole S (1994) The physiological costs of flight capability in wing dimorphic crickets *Res. Popul. Ecol.* 36:151-156
- Zera AJ, Mole S, Rokke K (1994) Lipid, carbohydrate and nitrogen-content of long-winged and short-winged *Gryllus firmus*-implications for the physiological cost of flight capability. *J. Ins. Phys.* 40:1037-1044
- Zhang XS (2008) Increase in quantitative variation after exposure to environmental stresses and/or introduction of a major mutation: $G \times E$ interaction and epistasis or canalization? *Genetics* 180:687-695
- Zuparko RL, Dahlsten DL (1994) Host plant resistance and biological control for linden aphids. *J. Arboriculture* 20:278-281
- Zuparko RL, Dahlsten DL (1995) Parasitoid complex of *Eucallipterus tiliæ* (Homoptera, Drepanosiphidae) in Northern California. *Environ. Entomol.* 24:730-737
- Zuparko RL, Dahlsten DL (1996) New potential for classical biological control of *Eucallipterus tiliæ* (Homoptera: Drepanosiphidae). *Biol. Control* 6:407-408

Affiliations of all authors

	<i>At the time of the research</i>	<i>Currently</i>
Ellyn V. Bitume	Institute of Biology, Leiden University, The Netherlands	Evolutionary Ecology and Genetics, Université catholique de Louvain, Belgium
Paul M. Brakefield	Institute of Biology, Leiden University, The Netherlands	Zoological Department of Cambridge/ Zoological Museum of Cambridge, UK
Bardo A. Cornelder	Institute of Biology, Leiden University, The Netherlands	
Peter W. de Jong	Laboratory of Entomology, Wageningen University, The Netherlands	Laboratory of Entomology, Wageningen University, The Netherlands
Thomas C. Holness	Applied Plant Research - Flower Bulbs and Nursery Stock, Wageningen UR, The Netherlands	
Kees G. Koops	Institute of Biology, Leiden University, The Netherlands	Institute of Biology, Leiden University, The Netherlands
Suzanne T.E. Lommen	Institute of Biology, Leiden University, The Netherlands	Department of Biology, University of Fribourg, Switzerland
Carola A. Luijten	Institute of Biology, Leiden University, The Netherlands	
Cock W. Middendorp	Laboratory of Entomology, Wageningen University, The Netherlands	
Suzanne V. Saenko	Institute of Biology, Leiden University, The Netherlands	Department of Genetics & Evolution, University of Geneva, Switzerland
Yoshinori Tomoyasu	Division of Biology, Kansas State University, USA	Department of Zoology, Miami University, USA
Alfons J. van Kuik	Applied Plant Research - Flower Bulbs and Nursery Stock, Wageningen UR, The Netherlands	Applied Plant Research - Flower Bulbs and Nursery Stock, Wageningen UR, The Netherlands
Jeroen van Schelt	Koppert B.V. , The Netherlands	Koppert B.V. , The Netherlands