



Universiteit
Leiden
The Netherlands

Design and synthesis of NLR and TLR based ligand-antigen conjugates

Willems, M.M.J.H.P.

Citation

Willems, M. M. J. H. P. (2012, November 1). *Design and synthesis of NLR and TLR based ligand-antigen conjugates*. Retrieved from <https://hdl.handle.net/1887/20082>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/20082>

Note: To cite this publication please use the final published version (if applicable).

Design & Synthesis of NLR and TLR based ligand-antigen conjugates

A focus on NOD2, NOD1 and TLR2 ligand modifications

PROEFSCHRIFT

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. P. F. van der Heijden,
volgens besluit van het College voor Promoties
te verdedigen op donderdag 1 november 2012
klokke 16.15 uur

door

Marian Martha Johanna Hendrika Petronella Willems

Geboren te Venray in 1984

Promotiecommissie

Promotor : Prof. Dr. G. A. van der Marel

Co-promotores : Dr. J. D. C. Codée
Dr. D. V. Filippov

Overige leden : Prof. Dr. H. S. Overkleef
Prof. Dr. F.A. Ossendorp
Prof. Dr. J. Brouwer
Prof. Dr. J. Lugtenburg
Dr M.S.M. Timmer



This research is supported by Top Institute Pharma, The TLR Project: Exploitation of Toll-like receptors in Drug Discovery (D1-101-0)

The printing of this thesis was financially supported by the J. E. Jurriaanse stichting

Printed by Wöhrmann Print Service



In the middle of difficulties lie opportunities – Albert Einstein

Table of Content

List of abbreviations

Chapter 1

General introduction 9

Chapter 2

The synthesis and biological evaluation of covalent
NOD2 ligand-antigen conjugates 31

Chapter 3

Lipophilic NOD2 ligands as immunostimulatory agents
61

Chapter 4

Synthesis and biological evaluation of NOD2- and TLR2-
ligand-antigen bis-conjugates 83

Chapter 5

Synthesis and biological evaluation of NOD1-ligands
and ligand-antigen conjugates 107

Chapter 6

Urea-derived Pam₃CSK₄ derivatives as new agonists for
TLR1/2 127

Chapter 7

Summary and future prospects 143

Nederlandse samenvatting 157

Curriculum Vitae (Nederlands) 161

List of Abbreviations

Ac	acetyl	Fmoc	9H-fluorenylmethoxy-carbonyl
AcOH	acetic acid	GlcNAc	<i>N</i> -acetyl glucosamine
All	allyl	h	hour
APC	antigen presenting cell	HATU	2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl-uronium hexafluorophosphate methanaminium
aq	aqueous	HEK	human embryonic kidney
ar	aryl	HCTU	2-(6-chloro-1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate
BMDC cell	bone marrow dendritic cell	HOBt	1-hydroxybenzotriazole
BOP	benzotriazole-1-yl-oxy-tris-dimethylamino-phosphonium hexafluorophosphate	HPLC	high performance liquid chromatography
Boc	<i>tert</i> -butyloxycarbonyl	HRMS	high resolution mass spectrometry
Boc ₂ O	di- <i>tert</i> -butyl dicarbonate	Hz	hertz
Bu	butyl	<i>i</i>	iso
^t Bu	<i>tert</i> -butyl	IR	infrared
Bu ₃ SnH	tributyltin hydride	IL	interleukin
Calc.	calculated	<i>J</i>	coupling constant
C _q	quaternary carbon atom	Lev	levulinoyl
CSA	camphor sulfonic acid	LC/Ms	liquid chromatography mass spectrometry
d	doublet	LPS	lipopolysaccharide
δ	chemical shift	m	multiplet
DAP	diaminopimelic acid	M	molar/mass
DBU	1,8-diazabicycloundec-7-ene	MDP	<i>N</i> -acetylmuramyl-L-alanine-D-isoglutamine
DC	dendritic cell	Me	methyl
DCE	dichloroethane	MeCN	acetonitrile
DCM	dichloromethane	MeOH	methanol
dd	double doublet	MHC I	major histocompatibility complex class I
DIC	<i>N,N</i> -diisopropylcarbodiimide	min	minutes
DMAP	<i>N,N</i> -dimethylaminopyridine	mg	milligram
DEVA ₅ K	DEVSGLEQLSIIINFEKLAATAAK	MS	mass spectrometry
DiPEA	<i>N,N</i> -diisopropylethylamine	Mtt	methyl trityl
DMF	<i>N,N</i> -dimethylformamide	MurNAc	<i>N</i> -acetylmuramic acid
DMSO	dimethyl sulfoxide	<i>m/z</i>	mass to charge ratio
EDC	1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide	NaH	sodium hydride
eq	equivalent	NaOMe	sodium methoxide
Et	ethyl	NH ₄ HCO ₃	ammonium bicarbonate
EtOAc	ethylacetate	nM	nanomolar
Et ₃ N	triethylamine		
Et ₂ O	diethyl ether		

NMP	<i>N</i> -methyl-2-pyrrolidone	rt	room temperature
NMR	nuclear magnetic resonance	Rt	retention time
NLR	NOD-like receptor	s	singulet
NOD	Nucleotide-binding Oligomerization Domain	SAR	structure- activity relationship
NOD1-L	NOD1 ligand	SPPS	solid phase peptide synthesis
NOD2-L	NOD2 ligand	t	triplet
PAMPS	pathogen associated molecular patterns	TFA	trifluoro acetic acid
PE	petroleum ether	THF	tetrahydrofuran
PG	peptidoglycan	TIS	triisopropyl silane
Ph	phenyl	TLC	tin layer chromatography
ppm	parts per million	TMSOTf	trimethylsilyl trifluoromethanesulfonate
PyBOP	benzotriazol-1-yl-oxy-tris-pyrrolidinophosphonium hexafluorophosphate	TLR	Toll-like receptor
PRR	pattern recognition receptors	TLR2-L	TLR2 ligand
PRR-L	PRR ligand	μM	micromolar
Pd(PPh ₃) ₄	palladium tetrakis	UV	ultraviolet
q	quartet		
RP-HPLC	reversed phase high performance liquid chromatography		

Three-letter codes for amino acids are used following the rules as proposed by the IUPAC-IUB commission on Biochemical Nomenclature. *J.Biol.Chem.*, **1966**, *241*, 2491-2495.
