



Universiteit
Leiden

The Netherlands

**Borneo : a quantitative analysis of botanical richness,
endemicity and floristic regions based on herbarium
records**

Raes, N.

Citation

Raes, N. (2009, February 11). *Borneo : a quantitative analysis of botanical richness, endemicity and floristic regions based on herbarium records.*

Retrieved from <https://hdl.handle.net/1887/13470>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/13470>

Note: To cite this publication please use the final published version (if applicable).

Borneo

A quantitative analysis of botanical richness,
endemicity and floristic regions
based on herbarium records

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. P.F. van der Heijden,
volgens besluit van het College van Promoties
te verdedigen op woensdag 11 februari 2009
klokke 15.00 uur

door

NIELS RAES

Geboren te Bladel
in 1971

Promotiecommissie:

Promotor: Prof. dr. P. Baas

Copromotoren: Dr. H. ter Steege
Dr. M.C. Roos

Referent: Prof. dr. F.J.J.M. Bongers (Wageningen University)

Overige leden: Prof. dr. D.J. Mabberley
Prof. dr. E.F. Smets
Prof. dr. M.S.M. Sosef
Prof. dr. P.C. van Welzen



A quantitative analysis
of botanical richness,
endemicity and floristic regions
based on herbarium records

Niels Raes

NATIONAAL HERBARIUM NEDERLAND
Leiden University branch
2009

Raes, N.

Borneo: A quantitative analysis of botanical richness, endemism and floristic regions based on herbarium records.

ISBN: 978-90-8570-318-1

Printing: Wöhrmann Print Service

Graphic Design: Ed van Oosterhout

Eindredactie Nederlandse samenvatting: Koen-Machiel van de Wetering

FSC Mixed Sources: This book is printed on FSC certified paper.

Cover: The symbol used on the cover and throughout the thesis is a Dayak symbol that stands for a man figure. The precise meaning of the symbol depends on the context in which it is placed, but generally it depicts the continuity of (human) life.

Chapter 1: Partly published in *Blumea* 54(1), N. Raes and P.C. van Welzen, The demarcation and internal division of Flora Malesiana: 1857 – Present, ©2009, with permission from the Nationaal Herbarium Nederland, Leiden University branch.

Chapter 2: Reprinted from *Blumea* 54(1), N. Raes, J.B. Mols, L.P.M. Willemsse and E.F. Smets, Georeferencing specimens by combining digitized maps with SRTM digital elevation data and satellite images, ©2009, with permission from the Nationaal Herbarium Nederland, Leiden University branch.

Chapter 3: Reprinted from *Ecography* 30, N. Raes and H. ter Steege, A null-model for significance testing of presence-only species distribution models, ©2007, with permission from the Oikos Editorial Office.

Chapter 4: Reprinted from *Ecography* (accepted), N. Raes, M.C. Roos, J.W.F. Slik, E.E. van Loon and H. ter Steege, Botanical richness and endemism patterns of Borneo derived from species distribution models, ©2009, with permission from the Oikos Editorial Office.

Chapter 5: Submitted to *Journal of Biogeography*

Chapter 6: In preparation

Remainder of the thesis ©2009, Nationaal Herbarium Nederland, Leiden University branch.

No part of this publication, apart from bibliographic data and brief annotations in critical reviews, may be reproduced, re-recorded or published in any form, including print, photocopy, microform, electronic or electromagnetic record without written permission by the publishers.

Summary

Although it is widely recognized that Borneo harbours one of the world's most important biodiversity hotspots, the spatial patterns of botanical richness, endemism, 'centres of endemism', and Borneo's floristic regions, until now have largely been based on informal expert opinion. Recent digitization of the botanical collections of Borneo, housed at the National Herbarium of the Netherlands, has provided a database that allowed a quantitative, spatial analysis of the components of botanical diversity of Borneo.

The objectives of this study are to develop high-resolution spatial maps of the patterns of botanical richness, -endemism, 'centres of endemism', and the floristic regions of Borneo derived from species distribution models. The resulting maps are related to environmental conditions to explain the patterns of the different components of botanical diversity and the recognized floristic regions. Finally, we assess the extent to which areas of high botanical diversity and the different floristic regions remain forested today to guide conservation efforts of the threatened forests of Borneo.

We used 'Maxent' to develop species distribution models for species treated in 'Flora Malesiana' and represented by at least five records. The 2273 species distribution models were statistically tested with a method we developed for this purpose, resulting in 1439 significant models (63.3%), covering 8577 grid cells (5 arc-minute resolution, ca. 100 km²) of Borneo.

The 1439 significant models were superimposed to generate patterns of botanical richness, -endemism, and 'centres of endemism'. The highest botanical richness is predicted to occur in northern and northwestern Borneo. The northern Crocker Mountains range with Mount Kinabalu, and the high mountains of central East Kalimantan have the highest botanical endemism values. The 'centres of endemism' are found on Mount Kinabalu and the northern Crocker Range Mountains, the southern Müller Mountains, the east side of the Meratus Mountains, and the Sangkulirang peninsula. Areas of high botanical richness and -endemism are characterized by a relatively small range in annual temperature, but with seasonality within that range. Furthermore, these areas are least affected by the El Niño Southern Oscillation drought events. 'Centres of endemism' are characterized by ecological distinctiveness in altitude, edaphic conditions, annual precipitation, or a combination of these factors.

The 11 floristic regions of Borneo were recognized based on a presence/absence matrix derived from the 1439 significant species distribution models. This matrix was analysed using a hierarchical cluster analysis, and the resulting cluster dendrogram was pruned using indicator species analysis (ISA) to partition floristic regions. This method allowed the quantitative confirmation of the floristic distinctiveness and extent of montane rain forest, kerangas, peat swamp, and fresh water swamp forest. The lowland rain forest, previously recognized as one floristic region was divided in at least four (and possibly six) different floristic regions, viz. the lowlands of (i) Sabah and Sarawak, (ii) East Kalimantan, (iii) southern Borneo, and (iv) the 'Wet hill forest of Sarawak'. We could not distinguish, but do recognize, the 'Kinabalu highlands', mangroves, and forests on limestone and ultramafic rock due to the spatial resolution (100 km²) of our analysis.

Correction for the impact of deforestation, forest fire, and land use change on the 11 floristic regions and areas of high botanical diversity revealed that 36% of Borneo's total surface, and 57% of its most diverse areas has already been lost. Especially the most diverse lowland forests have been severely hit by deforestation. Even more worrying is the fact that deforestation has taken its toll in IUCN recognized protected areas. Only 0.6% of Borneo's most diverse areas have an IUCN protected status, of which 33% was already deforested by the year 2000. Most dramatic is the loss of 84% of East Kalimantan's protected lowland rain forests.

To safeguard Borneo's genetic botanical diversity we urge governments and policy makers to award the remaining forested extents of areas with the highest botanical diversity with a protection status, to enforce the protection of recognized protected areas, and to conserve significant parts of each of the 11 floristic regions.

Table of Contents

| | |
|---|-----|
| CHAPTER 1 | |
| General Introduction | 10 |
| <hr/> | |
| CHAPTER 2 | |
| Georeferencing specimens by combining digitized maps with SRTM digital elevation data and satellite images | 18 |
| <hr/> | |
| CHAPTER 3 | |
| A null-model for significance testing of presence-only species distribution models | 24 |
| <hr/> | |
| CHAPTER 4 | |
| Botanical richness and endemism patterns of Borneo derived from species distribution models | 38 |
| <hr/> | |
| CHAPTER 5 | |
| The floristic regions of Borneo inferred from species distribution models | 56 |
| <hr/> | |
| CHAPTER 6 | |
| Borneo's remaining forests – Where to from here? | 76 |
| <hr/> | |
| REFERENCES | 84 |
| <hr/> | |
| Appendix Figures | 94 |
| <hr/> | |
| Appendix Tables | 101 |
| <hr/> | |
| Samenvatting (In Dutch) | 132 |
| <hr/> | |
| Curriculum Vitae | 138 |
| <hr/> | |
| Acknowledgements | 140 |
| <hr/> | |