



Universiteit
Leiden
The Netherlands

Glucose metabolism in healthy ageing

Wijsman, C.A.

Citation

Wijsman, C. A. (2014, October 8). *Glucose metabolism in healthy ageing*. Retrieved from <https://hdl.handle.net/1887/29017>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/29017>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden




The handle <http://hdl.handle.net/1887/29017> holds various files of this Leiden University dissertation

Author: Wijsman, Carolien A.

Title: Glucose metabolism in healthy ageing

Issue Date: 2014-10-08

The background of the slide is white and features several decorative elements. There are five stylized flowers in a light gray color, scattered across the page. A thin, light gray wavy line curves across the lower half of the slide. The text is centered and uses a mix of bold and regular fonts.

CHAPTER 4

Responsiveness of the innate immune system and glucose concentrations in the oldest old

Wijsman CA, Mooijaart SP, Westendorp RG, Maier AB
Age (Dordr) 2012 Aug; 34(4):983-6

Abstract

Background: Patients with diabetes mellitus show increased risk of infectious disease as well as disturbances in innate immunity. In critical care settings, hyperglycemia is associated with increased risk of sepsis. It is unclear whether elevated glucose concentrations and innate immunity are associated in a non-clinical setting. We aimed to assess the association between glucose concentrations and innate immune response in the oldest old, who are at increased risk of both disturbed glucose metabolism as well as infectious disease.

Methods: This study was part of the Leiden 85-plus Study. In 562 subjects aged 85 years old of the general population, venous blood samples were taken for measurement of morning glucose, C-reactive protein and glycated hemoglobin (HbA1c). The innate immune response was assessed by performing ex vivo whole blood lipopolysaccharide (LPS) stimulation for production capacity of tumor necrosis factor alpha (TNF- α), interleukin 6 (IL-6), interleukin 1-beta (IL1- β), interleukin 10 (IL-10) and interleukin 1 receptor antagonist (IL-1Ra). Using linear regression analysis, cross-sectional analysis between glucose and cytokine production capacity were performed.

Results: We found a significant negative association between glucose concentrations, but not HbA1c, and cytokine response capacity in four out of five measured cytokines (all $p < 0.05$). Both glucose and HbA1c were positively associated with circulating levels of CRP.

Conclusion: Higher glucose concentrations in non-diabetic elderly associate with lower innate immune response. As elderly show increased vulnerability for disturbances in glucose metabolism as well as infectious disease, this relation could be of clinical significance.

Introduction

Patients with diabetes mellitus have a higher vulnerability for infectious diseases, which contributes to increased hospitalization and mortality (13). It has been hypothesized that elevated glucose concentrations negatively influence cytokine production and neutrophil function (1; 6).

In vitro studies testing the effect of different glucose concentrations on innate immunity have yielded conflicting results (reviewed in (7)). In healthy young subjects, induction of hyperglycemia and lipopolysaccharide (LPS) stimulation in vivo did not attenuate the cytokine response (8). In the critical care setting, detrimental effects of hyperglycemia have been shown in non-diabetic patients, where stress-induced hyperglycemia is related to higher risk of sepsis and increased morbidity and mortality (4).

In the oldest old both infectious disease and disturbances of glucose homeostasis are highly prevalent. In the present study we tested the relationship between innate immunity and glucose concentrations in non-acutely diseased community-dwelling 85-year olds without diabetes mellitus.

Methods

The Leiden 85-plus Study is a prospective, observational, population-based follow-up study of all inhabitants aged 85 years in Leiden, The Netherlands. Subjects were enrolled between 1997 and 1999. All participants were aged 85 years. There were no exclusion criteria. Of the 705 eligible subjects, 599 subjects were willing to participate (87%). At baseline, non-fasted venous blood samples were obtained in the morning (before 11 am) for all participants. Ex vivo whole blood LPS stimulation was performed in 562 subjects. Levels of tumor necrosis factor alpha (TNF- α), interleukin 6 (IL-6), interleukin 1-beta (IL1- β), interleukin 10 (IL-10) and interleukin 1 receptor antagonist (IL-1Ra) were determined 24 hours after incubation with LPS, as more extensively described earlier (16). Non-fasted plasma glucose concentrations, glycated hemoglobin (HbA1c) and C-reactive protein (CRP) levels were measured in 559 subjects. Subjects with known diabetes ($n=52$) or hyperglycemia (non-fasted glucose > 11 mmol/L, $n = 39$) were excluded from analysis. Analyses were done using linear regression with adjustment for gender using SPSS version 16.0.




Figure 1. Cytokine production after LPS-stimulation ex vivo by glucose concentrations (n=468). Cytokine data represent geometric means $\pm$ 95% confidence intervals, after linear regression with adjustment for gender.

Results

Figure 1 shows the relation between quartiles of glucose and cytokine levels in response to LPS-stimulation in 468 participants. After adjustment for gender, concentrations of non-fasted glucose correlated negatively with TNF- α , IL-6, IL-1 β , and IL-10 (all p for trend < 0.01), but not with IL-1Ra (p for trend = 0.73). Additionally excluding the subjects with glycated hemoglobin $> 6.5\%$ (n=20) did not alter results (data not shown). Glucose concentrations were positively associated with CRP (p for trend = 0.02).

To assess whether the association between innate immunity and glucose concentrations was affected by acute or long-term regulation of glucose metabolism, we repeated the analyses of stimulated cytokine levels dependent on HbA1c. No consistent linear association was found between HbA1c and cytokines levels in response to LPS-stimulation (data not shown). HbA1c did correlate positively with CRP levels (p = 0.001).

Discussion

We found a strong negative association between non-fasted glucose concentrations, but not HbA1c levels, and cytokine production upon LPS stimulation ex vivo in a community-dwelling, non-diabetic population of the oldest old. Both non-fasted glucose concentrations and HbA1c showed a positive correlation with CRP levels.

The observed relationship between non-fasted glucose and cytokine production capacity in non-diabetic subjects suggests that alterations in glucose metabolism, even in non-diabetics, may attenuate innate immune response. The lack of association between HbA1c and cytokine levels suggests that this effect might be driven by temporarily increases of glucose rather than chronic elevations. Earlier, hyperglycemic fluctuations, rather than continuous hyperglycemia, has been associated with negative outcome in a non-diabetic population cohort(2).

The relationship between higher glucose concentrations and low cytokine response could be of clinical significance. A low cytokine response has been independently associated with morbidity and mortality in our study population of the oldest old(15). Furthermore, low cytokine response has been associated with detrimental outcome in patients with sepsis at the intensive care unit (11; 17). Also in this acutely ill patient cohort, innate immune response related inversely to levels of inflammation, such as CRP (14).

A limitation of this study is that we did not measure insulin sensitivity. Insulin sensitivity has been associated with higher circulating inflammatory markers and is a prognostic factor for outcome of critically ill patients with myocardial infarction (10; 18). We were therefore not able to assess which mechanism of glucose metabolism is associated with innate immune function. Furthermore, as glucose concentrations were assessed at one time point only, we could also not assess glycemic variability accurately. Further studies with sensitive tools, such as continuous glucose monitoring systems, could better clarify this issue.

Importantly, our results were found in a non-diabetic population, suggesting that the blunted cytokine response is not due to specific factors related to diabetes mellitus, such as use of anti-diabetic medication, or other pathophysiological changes due to the disease, but suggest a direct relationship between glucose concentration and innate immune response. This association might be of clinical importance, as also in apparently healthy older individuals, disturbances in glucose and insulin metabolism are found such as impaired beta-cell function and decreased insulin action (3; 5). Moreover, the glycemic response to stress might be increased

with aging, which could further aggravate glycemic disturbances in clinical settings (9; 12).

Further studies in vivo including non-diabetic subjects are warranted to identify the mechanism behind the association between glucose concentrations and innate immune response and to investigate whether results can be extended to younger populations.

References

1. **Andreasen AS, Pedersen-Skovsgaard T, Berg RM, Svendsen KD, Feldt-Rasmussen B, Pedersen BK and Moller K.** Type 2 diabetes mellitus is associated with impaired cytokine response and adhesion molecule expression in human endotoxemia. *Intensive Care Med* 2010.
2. **Barr EL, Boyko EJ, Zimmet PZ, Wolfe R, Tonkin AM and Shaw JE.** Continuous relationships between non-diabetic hyperglycaemia and both cardiovascular disease and all-cause mortality: the Australian Diabetes, Obesity, and Lifestyle (AusDiab) study. *Diabetologia* 52: 415-424, 2009.
3. **Basu R, Breda E, Oberg AL, Powell CC, Dalla MC, Basu A, Vittone JL, Klee GG, Arora P, Jensen MD, Toffolo G, Cobelli C and Rizza RA.** Mechanisms of the age-associated deterioration in glucose tolerance: contribution of alterations in insulin secretion, action, and clearance. *Diabetes* 52: 1738-1748, 2003.
4. **Berghe Van den G., Wouters P, Weekers F, Verwaest C, Bruyninckx F, Schetz M, Vlasselaers D, Ferdinande P, Lauwers P and Bouillon R.** Intensive insulin therapy in the critically ill patients. *N Engl J Med* 345: 1359-1367, 2001.
5. **Chang AM, Smith MJ, Galecki AT, Bloem CJ and Halter JB.** Impaired beta-cell function in human aging: response to nicotinic acid-induced insulin resistance. *J Clin Endocrinol Metab* 91: 3303-3309, 2006.
6. **Delamaire M, Maugendre D, Moreno M, Le Goff MC, Allannic H and Genetet B.** Impaired leucocyte functions in diabetic patients. *Diabet Med* 14: 29-34, 1997.
7. **Geerlings SE and Hoepelman AI.** Immune dysfunction in patients with diabetes mellitus (DM). *FEMS Immunol Med Microbiol* 26: 259-265, 1999.
8. **Krogh-Madsen R, Moller K, Dela F, Kronborg G, Jauffred S and Pedersen BK.** Effect of hyperglycemia and hyperinsulinemia on the response of IL-6, TNF-alpha, and FFAs to low-dose endotoxemia in humans. *Am J Physiol Endocrinol Metab* 286: E766-E772, 2004.
9. **Lazzeri C, Valente S, Chiostrì M, Picariello C and Gensini GF.** Acute glucose dysmetabolism in the early phase of ST-elevation myocardial infarction: the age response. *Diab Vasc Dis Res* 7: 131-137, 2010.
10. **Lazzeri C, Valente S, Chiostrì M, Picariello C and Gensini GF.** Acute glucose dysmetabolism in the elderly with ST elevation myocardial infarction submitted to mechanical revascularization. *Int J Cardiol* 2011.
11. **Muller Kobold AC, Tulleken JE, Zijlstra JG, Sluiter W, Hermans J, Kallenberg CG and Tervaert JW.** Leukocyte activation in sepsis; correlations with disease state and mortality. *Intensive Care Med* 26: 883-892, 2000.
12. **Odio MR and Brodish A.** Effects of age on metabolic responses to acute and chronic stress. *Am J Physiol* 254: E617-E624, 1988.
13. **Shah BR and Hux JE.** Quantifying the risk of infectious diseases for people with diabetes. *Diabetes Care* 26: 510-513, 2003.
14. **Tschaikowsky K, Hedwig-Geissing M, Schiele A, Bremer F, Schywalsky M and Schuttler J.** Coincidence of pro- and anti-inflammatory responses in the early phase of severe sepsis: Longitudinal study of mononuclear histocompatibility leukocyte antigen-DR expression, procalcitonin, C-reactive protein, and changes in T-cell subsets in septic and postoperative patients. *Crit Care Med* 30: 1015-1023, 2002.

15. **van den Biggelaar AH, Huizinga TW, de Craen AJ, Gussekloo J, Heijmans BT, Frolich M and Westendorp RG.** Impaired innate immunity predicts frailty in old age. The Leiden 85-plus study. *Exp Gerontol* 39: 1407-1414, 2004.
16. **van der Linden MW, Huizinga TW, Stoeken DJ, Sturk A and Westendorp RG.** Determination of tumour necrosis factor-alpha and interleukin-10 production in a whole blood stimulation system: assessment of laboratory error and individual variation. *J Immunol Methods* 218: 63-71, 1998.
17. **Westendorp RG, Langermans JA, Huizinga TW, Elouali AH, Verweij CL, Boomsma DI and Vandenbroucke JP.** Genetic influence on cytokine production and fatal meningococcal disease. *Lancet* 349: 170-173, 1997.
18. **Zuliani G, Galvani M, Maggio M, Volpato S, Bandinelli S, Corsi AM, Lauretani F, Cherubini A, Guralnik JM, Fellin R and Ferrucci L.** Plasma soluble gp130 levels are increased in older subjects with metabolic syndrome. The role of insulin resistance. *Atherosclerosis* 213: 319-324, 2010.