



Universiteit
Leiden
The Netherlands

Unraveling mechanisms of vascular remodeling in arteriovenous fistulas for hemodialysis

Wong, C.Y.

Citation

Wong, C. Y. (2017, March 8). *Unraveling mechanisms of vascular remodeling in arteriovenous fistulas for hemodialysis*. Retrieved from <https://hdl.handle.net/1887/46406>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/46406>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/46406> holds various files of this Leiden University dissertation.

Author: Wong, C.Y.

Title: Unraveling mechanisms of vascular remodeling in arteriovenous fistulas for hemodialysis

Issue Date: 2017-03-08

Chapter 2

Arteriovenous access failure: More than just intimal hyperplasia?

Tonia C. Rothuizen, ChunYu Wong, Paul H.A. Quax, Anton Jan van Zonneveld,
Ton J. Rabelink and Joris I. Rotmans

Nephrol Dial Transplant. 2013 May;28(5):1085-92



Abstract

Hemodialysis vascular access patency is severely compromised by fistula non-maturation and access stenosis. Intimal hyperplasia (IH) is considered the culprit lesion in failed fistulas, resulting in luminal narrowing and stenosis. This review focuses on the biology and pathophysiology of fistula failure and highlights not only the classically associated intimal hyperplasia but also some relatively neglected but potentially important contributors such as inadequate outward remodeling. In addition, the complex process and fragile balance of successful fistula maturation might be partially hindered by pre-existent chronic kidney disease-mediated vasculopathy. Further unravelling the (patho)physiology of outward remodeling and intimal hyperplasia could contribute to novel therapies and enhance fistula patency.

Introduction

Patients with end-stage renal disease are largely dependent on dialysis as renal replacement therapy. For chronic hemodialysis, an adequately functioning high-flow vascular access is required. Arteriovenous fistulas (AVFs) are the preferred modality in view of the superior patency rates and fewer complications as compared to arteriovenous synthetic grafts¹⁻³. Nonetheless, the durability of AVFs is far from optimal with one year primary patency rates ranging from 60-65%^{4,5}. In fact, these numbers are too optimistic, as they frequently do not account for fistulas that failed to mature. Maturation failure contributes significantly to the dismal patency rates of AVFs as illustrated by a recent multi-center study which revealed that 60% of the AVFs were not suitable for dialysis between 4-5 months post surgery⁶, although, these numbers do vary between different types of AVFs⁷. According to the KDOQI guidelines, an AVF maturation is clinically considered successful if 6 weeks after surgery the fistula supports a flow of 600 mL/min, is located at a maximum of 6 mm from the surface and has a diameter greater than 6 mm². The exact underlying mechanisms responsible for maturation failure are however unknown, but impaired outward remodeling as well as intimal hyperplasia (IH) are both considered to contribute.

Thus far, most research on the pathophysiology of AVF failure focuses on IH. In contrast, the role of vascular outward remodeling in the setting of AVFs is often neglected. However, adequate outward remodeling could preserve luminal caliber and may therefore be valuable for successful fistula maturation. We postulate that the balance between outward expansion and potential luminal narrowing due to IH may ultimately determine fistula flow and patency (Fig.1). This review focuses on the pathophysiology of both AVF maturation failure and failure of already matured AVFs, and highlights some potential contributors thus far gaining relatively little attention, such as outward remodeling and the implications of chronic kidney disease (CKD) mediated vascular pathology.




Figure 1. Different modalities of the vascular remodeling response after fistula creation. Whereas a healthy vein has the potential for successful outward remodeling (top left), adequate maturation may be partially hindered by CKD-induced pre-existing vasculopathy such as IH (top right). As shown below, we postulate that the net resultant of IH and outward remodeling may determine ultimate luminal calibre. In case of IH formation, adequate outward remodeling could to a certain extent remain lumen calibre intact, thus providing a patent fistula. However, if IH outbalances outward remodeling this could result in stenosis and fistula failure.

Outward remodeling: an emerging concept in arteriovenous fistulas?

The connection of a low-pressure vein to the high-pressure arterial system results in a chain of vascular events that starts with an immediate increase of blood flow through the both the feeding artery and the draining vein^{4,8}. Directly after construction of the AVF, this rapid

increase in flow results in both passive vascular distension and in nitric oxide (NO) synthesis by endothelial cells with subsequent vascular smooth muscle cell (VSMC) relaxationⁿ⁹⁻¹¹ resulting in acute vasodilation. Concomitantly, the hemodynamic changes following AVF creation initiate a more structural vascular remodeling response leading not only to a further increase in arterial and venous calibre¹²⁻¹⁴ but also to thickening of especially the venous wall^{15,16}. Increased wall shear stress (WSS) and wall tension are the driving forces. WSS is the frictional force exerted by blood on the vessel wall and is mathematically defined by Poiseuille's formula: $4\eta Q/\pi r^3$ where η = blood viscosity, Q = flow and r = vessel radius. An increase in blood flow will provoke an adaptive response of the vessel in which the luminal diameter increases in attempt to reduce WSS to pre-AVF levels (5-10 dyn/cm²). Furthermore, due to the pressure increase in the venous outflow tract after fistula creation, the wall tension rises leading to another adaptive response culminating in medial thickening (i.e. venous arterialization). These phenomena are elegantly illustrated in a study of Corpataux et al¹², where hemodynamic changes in the venous part of AVFs in six patients were investigated using echo-tracking and Doppler-ultrasonography. Within the first week after fistula formation the flow increased to 539 mL/min accompanied by an almost threefold increase in WSS to 24.5 dyn/cm². The continuous increase in flow resulted in a progressive increment in venous luminal calibre from 2.4 mm preoperative to 6.6 mm after 12 weeks. Since WSS is inversely related to lumen size, the WSS gradually returned to a physiological range (10.4 dyn/cm² at 12 weeks). In addition, the venous wall thickness increased, demonstrated by an augmentation of cross sectional wall area. This adaptive response is also applicable for the arterial side of the AVF, where the increment of arterial flow results in an increase of arterial luminal diameter¹⁴, although to a lesser extent than the venous side¹². This change in diameter does not result in wall thickening but likely rather results in arterial wall remodeling (i.e. an increase in both internal and external diameter without an increase in wall cross sectional area)¹⁴.

On a biological level, these changes in WSS and wall tension are sensed by the endothelial cells, that function as mechanosensors and convert these hemodynamic stimuli to biochemical signals such as vasodilating agents (e.g. NO), growth factors that can control VSMC proliferation and migration and cellular adhesion molecules^{10;17-19}. Upregulation of proteases such as matrix metalloproteinases (MMPs) and cathepsins results in matrix degradation and restructuring of the vascular scaffold leading to luminal expansion^{15;20-22}. In addition, with an increased circumference of the vessel wall, it seems logical that some VSMC reorganization should occur as well to keep in pace with the expansion, as is also described in arterial setting²³. However, to date little is known about the role of VSMCs in outward remodeling in AVFs.




Figure 2. Potential mechanisms of the remodeling response upon fistula creation. Upon fistula creation, in response to the altered hemodynamic environment a plethora of structural changes may occur in the vessel wall; amongst others surgery induced endothelial denudation, matrix reorganization to accommodate outward expansion but possibly also VSMC proliferation and migration contributing to intimal hyperplasia formation. Physiological outward remodeling may to a certain extent compensate for detrimental intimal lesions.

As explained before, the adaptive outward remodeling response occurs both in the feeding artery and in the draining vein of the fistula^{12,14}. Nonetheless, the majority of the stenotic lesions in fistulas failing to mature are localized in the venous part, mostly in the juxta-anastomotic region^{24,25}. The latter observation could suggest that not only IH but also venous luminal expansion may be important for the preservation of the luminal calibre, thereby allowing the fistula to mature. We postulate that the net resultant of adaptive outward expansion and potential luminal narrowing by IH and thrombosis may ultimately determine luminal calibre, flow and long term AVF patency.

Intimal hyperplasia: adverse vascular response that hampers AVF function

Intimal hyperplasia is the pathologic lesion in AVFs that may result in stenosis and ultimately thrombosis. It is characterized primarily by α -smooth muscle actin (α -SMA) positive cells, extracellular matrix proteins and cytokines such as platelet-derived growth factor, transforming growth factor- β and endothelin within the intima and media of the vein²⁶⁻²⁸. The vast majority of the α -SMA positive cells in the intimal lesions exhibit a myofibroblasts or synthetic VSMC phenotype²⁹. These cells could either be differentiated fibroblasts that migrated from the adventitia and acquired α -SMA expression³⁰, or dedifferentiated medial VSMCs⁵. Furthermore, recent studies of non-AVF models suggest that a proportion of the α -SMA positive cells in IH lesions might originate from multipotent vascular stem cells from the bone marrow^{31,32,33}.

In most physiological conditions, high laminar shear stress triggers endothelial quiescence, endothelial alignment in parallel with the flow and secretion of anti-inflammatory and anti-coagulant substances^{17,18,34}, thus preventing IH. In contrast, low flow and WSS levels as well as oscillating flow patterns are involved in endothelial cell activation with increased expression of pro-coagulant and pro-inflammatory mediators that predispose for IH^{34,35}. Although AVFs merely express high flow profiles, recent studies revealed the coexistence of spot regions with low and oscillating flow and WSS levels in the venous part of the AVF, using a pulsatile computational fluid dynamics simulation³⁶. These spot regions corresponded with in previous studies documented IH prone regions in the juxta-anastomotic area of the AVF. Alterations in anastomosis angle could impact the flow rates and patterns, potentially influencing IH formation, with sharper angles (30°) generating favourable outcomes³⁷. This was further illustrated by Krishnamoorthy et al³⁸, demonstrating a correlation between AVF configuration, WSS pattern and the development of IH in a porcine AVF model. Thus, IH in an AVF setting is likely to be associated with an abnormal WSS profile. Nevertheless, IH has also been observed in veins prior to vascular access placement³⁹ and in saphenous veins⁴⁰, suggesting that an abnormal WSS profile is not the sole cause for development of IH. Epidemiological studies have identified diabetes mellitus, race, older age, peripheral vascular disease, female sex and in

some studies also cardiovascular disease as risk factors for maturation failure⁴¹⁻⁴³. Furthermore, individual variation in patency outcomes may also in part be explained by genetic susceptibility. Indeed, several single nucleotide polymorphisms are associated with poor functional outcome of vascular access conduits for hemodialysis^{44;45}.

Most studies on fistula failure focussed on the effect of several parameters on IH, as IH is considered to be the pathognomonical lesion in fistula failure. However, as mentioned above, luminal calibre can be preserved by adequate outward remodeling. Consequently, in addition to IH, impaired outward remodeling may be another important but relatively overlooked contributor to fistula failure. Despite its potential impact on fistula patency, relatively little is known about outward remodeling in AVFs. Future research aiming to gain more insight in the (patho)physiology of outward remodeling in fistulas might therefore contribute to new therapeutic strategies that could improve fistula patency.

Interestingly, some of the elementary factors in outward remodeling are also involved in the process of IH. Whereas outward remodeling in AVFs is related to matrix protease activity such as MMPs and cathepsins²⁰⁻²², it is shown that MMPs are also involved in IH formation in AVFs^{46;47} and cathepsins in IH formation in balloon-injured artery⁴⁸. The relative contribution of these proteases to expansive remodeling versus IH is not yet established, though the crucial role in outward remodeling suggests a more beneficial than detrimental effect, especially in the initial phase of AVF maturation. The latter suggestion is underscored by a recent report showing increased serum MMP-2 levels in patients with matured fistulas as compared to those with maturation failure²⁰. However, as a consequence of elastolytic protease activity the internal elastic lamina is fragmented^{21;49}. This disruption of the elastic lamina and loss of integrity of this structural barrier may allow migration of adventitial fibroblasts or medial VSMCs to the intima. Moreover, the elastin degradation products can act as chemo-attractants for VSMCs and fibroblasts and might direct them to the intimal region and support their proliferation⁵⁰⁻⁵². Thus, potentially the balance of this partially overlapping beneficial outward remodeling and detrimental IH may affect fistula patency outcomes.

Vascular pathology in chronic kidney disease

Noteworthy, this complex process and fragile balance of successful fistula maturation might be partially hindered by pre-existent vascular abnormalities often present in patients with CKD. Especially in these patients with elevated comorbidity burden, a tailored surgical technique^{53;54} and surgical expertise⁵⁴ are important in determining fistula patency outcomes. Furthermore, CKD itself is a well-known risk factor for cardiovascular morbidity⁵⁵. The increased prevalence of cardiovascular disease in CKD-patients is only partly explained by traditional risk factors such as hypertension, diabetes, dyslipidemia and increased age. Epidemiological studies revealed

that CKD is an independent risk factor for cardiovascular morbidity⁵⁶. The latter observations implicate a role for additional inimical stimuli in CKD-patients such as chronic inflammation, increased oxidative stress, uremic toxins and endothelial dysfunction, as is reviewed in more detail elsewhere⁵⁷⁻⁵⁹. Most studies on CKD mediated vasculopathy concentrate on the arterial system. A functional AVF requires both adequate venous and arterial maturation. Pre-existent arterial vasculopathy may therefore reduce patency. In addition, although not extensively studied, the detrimental effects of CKD on the arterial system may influence veins in a similar manner⁶⁰. Indeed, recent studies elegantly showed marked pre-existing IH in venous segments of patients with end-stage renal disease prior to vascular access surgery^{39;61;62}. Another potential contributor to vascular pathology in CKD is vascular calcification. Whereas calcification in the tunica intima is classically associated with atherosclerosis, calcification in the tunica media can occur independently of atherosclerotic plaques and is frequently observed in arteries of any size in CKD-patients⁶³, resulting in vascular stiffness^{64;65}. In arterial setting this is known to impair the vessel's ability to expand upon high flow stimulation⁶⁶. Interestingly, Lee et al⁶⁷ recently also demonstrated extensive calcification in the intima and media of venous segments that were harvested at the time of vascular access surgery. This might result, similar to the arterial setting, in reduced venous compliance, thus potentially limiting the utility of AVFs by inhibiting outward remodeling and AVF maturation. Clinical studies already showed that forearm venous distensibility (i.e. increase in luminal diameter upon inflation of an upper arm cuff) rather than baseline venous diameter predicts successful AVF maturation⁶⁸. Although a recent study revealed that medial calcification in the supplying artery of AVFs was not associated with maturation failure⁶⁹, future studies should explore the impact of venous calcification prior to access surgery on maturation failure. Indeed, the outward remodeling response is much more pronounced in the venous part of AVFs compared to the feeding artery¹².

Therapeutic strategies to improve vascular access patency

In order to create a proper basis for a successful AVF, pre-operative vein preservation and careful selection of suitable vessels for AVF creation should be performed routinely^{2;70}. In case of fistula non-maturation or stenosis a percutaneous transluminal angioplasty (PTA) and/or surgical revision is required^{2;70}. To date, there are no adequate therapies improving the remodeling process of the AVF in the intermediary period. Moreover, despite the good results of PTA on the short term, it often induces restenosis on the longer term⁷¹⁻⁷³. Therefore there is a strong clinical need for new therapeutic strategies to improve fistula patency.

Possibilities to encourage outward remodeling include the use of elastase, thereby reorganizing the extracellular matrix scaffold of the vessel and promoting rapid dilation. After successful results in an AVF rabbit model³⁵, the use of recombinant-elastase PRT-201 was recently

clinically evaluated in a randomized controlled trial in human AVFs⁷⁴. Perivascular delivery of PRT-201 appeared to be safe but no effect on primary patency was observed. A larger clinical trial is underway since this first study was not powered to assess efficacy. Given the potential detrimental effects of elastin degradation products on both vascular calcification⁷⁵ and VSMC proliferation^{51,52}, the overall effectiveness of this agent remains to be elucidated.

Potential candidates for targeting IH are agents that inhibit VSMC proliferation, such as paclitaxel and sirolimus, although with data obtained only in AVG setting^{76,77} their effect in AVFs remains to be established.

As mentioned above, outward remodeling and IH are two processes that are in part intertwined. Some factors that are involved in outward remodeling such as MMPs may in a later stage also facilitate IH formation. Therefore, the potential of a therapy directed to a factor contributing in both types of remodeling could be influenced by the time of application. Limiting IH in an early stage might also decelerate the outward remodeling response and vice versa. Enhancement of maturation might require a different intervention than prevention of AVF failure once the AVF is successfully used for hemodialysis. Therefore, time dependent delivery might be a suitable approach to tackle fistula failure. Fistulas are ideal targets for such therapy, due to their easy accessibility and the potential to use perivascular delivery methods. However, to create a successful intervention strategy, more insight in the course and (patho)physiology of vascular remodeling is warranted.

Incorporation of other disciplines in the field of vascular access might offer new perspectives. IH is studied extensively in the field of cardiology and vascular surgery and the process of outward remodeling in physiological situations such as pregnancy. Furthermore the technology in slow-release drug delivery systems is rapidly expanding. Another relatively new field with exciting developments is vascular tissue engineering. With vascular tissue engineering it is possible to create a diameter- and length-matched blood vessel free from valves and accessory vessels and has the unique potential to adjust a vessel to patient specific requirements. Importantly, tissue engineered blood vessels are free from pre-existing vascular disease. The potential of a tissue engineered blood vessel (TEBV) as arteriovenous graft was illustrated by the group of l'Heureux and McAllister. Using a so-called sheet-based method, a completely biological TEBV was developed without the use of synthetic material, thereby creating a TEBV that resembles a native vessel in both composition and structure. The TEBV was evaluated as AVG in ten patients resulting in a primary patency rate of 78% and 60% after 1 and 6 months respectively⁷⁸. With spectacular progresses in the field of vascular tissue engineering⁷⁹⁻⁸¹, the use of a TEBV might become a realistic alternative in the nearby future.

Conclusion

Upon AVF creation a complex cascade of remodeling events should occur. The net resultant of beneficial outward expansion, potential luminal narrowing by intimal hyperplasia and the possible interference of CKD induced vascular pathology may ultimately determine luminal calibre, flow and long term AVF patency. Due to the potential positive contribution to fistula maturation and its assumed role in luminal calibre preservation, we pledge for more research emphasis on the role of outward remodeling. Further unravelling the complex pathways that mediate both IH and outward remodeling processes after AVF creation could provide new targets and therapies to improve fistula patency.

Acknowledgements

Tonia Rothuizen and Joris Rotmans are supported by the research program of the BioMedical Materials institute, co-funded by the Dutch Ministry of Economic Affairs, Agriculture and Innovation. ChunYu Wong is sponsored by a grant from the Dutch Kidney Foundation (KJPB 08.0003).

Reference List

1. Allon M, Robbin ML. Increasing arteriovenous fistulas in hemodialysis patients: problems and solutions. *Kidney Int* 2002; 62: 1109-1124
2. Clinical practice guidelines for vascular access. *Am J Kidney Dis* 2006; 48 Suppl 1: S248-S273
3. Gibson KD, Gillen DL, Caps MT, Kohler TR, Sherrard DJ, Stehman-Breen CO. Vascular access survival and incidence of revisions: a comparison of prosthetic grafts, simple autogenous fistulas, and venous transposition fistulas from the United States Renal Data System Dialysis Morbidity and Mortality Study. *J Vasc Surg* 2001; 34: 694-700
4. Tordoir JH, Rooyens P, Dammers R, van der Sande FM, de HM, Yo TI. Prospective evaluation of failure modes in autogenous radiocephalic wrist access for haemodialysis. *Nephrol Dial Transplant* 2003; 18: 378-383
5. Roy-Chaudhury P, Spergel LM, Besarab A, Asif A, Ravani P. Biology of arteriovenous fistula failure. *J Nephrol* 2007; 20: 150-163
6. Dember LM, Beck GJ, Allon M *et al.* Effect of clopidogrel on early failure of arteriovenous fistulas for hemodialysis: a randomized controlled trial. *JAMA* 2008; 299: 2164-2171
7. Lazarides MK, Georgiadis GS, Antoniou GA, Stamos DN. A meta-analysis of dialysis access outcome in elderly patients. *J Vasc Surg* 2007; 45: 420-426
8. Saucy F, Haesler E, Haller C, Deglise S, Teta D, Corpataux JM. Is intra-operative blood flow predictive for early failure of radiocephalic arteriovenous fistula? *Nephrol Dial Transplant* 2010; 25: 862-867
9. Cooke JP, Rossitch E Jr, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88: 1663-1671
10. Tronc F, Wassef M, Esposito B, Henrion D, Glagov S, Tedgui A. Role of NO in flow-induced remodeling of the rabbit common carotid artery. *Arterioscler Thromb Vasc Biol* 1996; 16: 1256-1262
11. Moncada S, Higgs A. The L-arginine-nitric oxide pathway. *N Engl J Med* 1993; 329: 2002-2012
12. Corpataux JM, Haesler E, Silacci P, Ris HB, Hayoz D. Low-pressure environment and remodelling of the forearm vein in Brescia-Cimino haemodialysis access. *Nephrol Dial Transplant* 2002; 17: 1057-1062
13. Ene-Iordache B, Mosconi L, Antiga L *et al.* Radial artery remodeling in response to shear stress increase within arteriovenous fistula for hemodialysis access. *Endothelium* 2003; 10: 95-102
14. Girerd X, London G, Boutouyrie P, Mourad JJ, Safar M, Laurent S. Remodeling of the radial artery in response to a chronic increase in shear stress. *Hypertension* 1996; 27: 799-803
15. Chan CY, Chen YS, Ma MC, Chen CF. Remodeling of experimental arteriovenous fistula with increased matrix metalloproteinase expression in rats. *J Vasc Surg* 2007; 45: 804-811
16. Langer S, Heiss C, Paulus N *et al.* Functional and structural response of arterialized femoral veins in a rodent AV fistula model. *Nephrol Dial Transplant* 2009; 24: 2201-2206
17. Garanich JS, Pahakis M, Tarbell JM. Shear stress inhibits smooth muscle cell migration via nitric oxide-mediated downregulation of matrix metalloproteinase-2 activity. *Am J Physiol Heart Circ Physiol* 2005; 288: H2244-H2252
18. Frangos JA, Eskin SG, McIntire LV, Ives CL. Flow effects on prostacyclin production by cultured human endothelial cells. *Science* 1985; 227: 1477-1479

19. Gibbons GH, Dzau VJ. The emerging concept of vascular remodeling. *N Engl J Med* 1994; 330: 1431-1438
20. Lee ES, Shen Q, Pitts RL *et al.* Serum metalloproteinases MMP-2, MMP-9, and metalloproteinase tissue inhibitors in patients are associated with arteriovenous fistula maturation. *J Vasc Surg* 2011; 54: 454-459
21. Tronc F, Mallat Z, Lehoux S, Wassef M, Esposito B, Tedgui A. Role of matrix metalloproteinases in blood flow-induced arterial enlargement: interaction with NO. *Arterioscler Thromb Vasc Biol* 2000; 20: E120-E126
22. Chang CJ, Chen CC, Hsu LA *et al.* Degradation of the internal elastic laminae in vein grafts of rats with aortocaval fistulae: potential impact on graft vasculopathy. *Am J Pathol* 2009; 174: 1837-1846
23. De Mey JG, Schiffers PM, Hilgers RH, Sanders MM. Toward functional genomics of flow-induced outward remodeling of resistance arteries. *Am J Physiol Heart Circ Physiol* 2005; 288: H1022-H1027
24. Nassar GM, Nguyen B, Rhee E, Achkar K. Endovascular treatment of the "failing to mature" arteriovenous fistula. *Clin J Am Soc Nephrol* 2006; 1: 275-280
25. Sivanesan S, How TV, Bakran A. Sites of stenosis in AV fistulae for haemodialysis access. *Nephrol Dial Transplant* 1999; 14: 118-120
26. Rotmans JJ, Pasterkamp G, Verhagen HJ, Pattynama PM, Blankestijn PJ, Stroes ES. Hemodialysis access graft failure: time to revisit an unmet clinical need? *J Nephrol* 2005; 18: 9-20
27. Weiss MF, Scivittaro V, Anderson JM. Oxidative stress and increased expression of growth factors in lesions of failed hemodialysis access. *Am J Kidney Dis* 2001; 37: 970-980
28. Lee T, Roy-Chaudhury P. Advances and new frontiers in the pathophysiology of venous neointimal hyperplasia and dialysis access stenosis. *Adv Chronic Kidney Dis* 2009; 16: 329-338
29. Roy-Chaudhury P, Wang Y, Krishnamoorthy M *et al.* Cellular phenotypes in human stenotic lesions from haemodialysis vascular access. *Nephrol Dial Transplant* 2009; 24: 2786-2791
30. Li L, Terry CM, Blumenthal DK *et al.* Cellular and morphological changes during neointimal hyperplasia development in a porcine arteriovenous graft model. *Nephrol Dial Transplant* 2007; 22: 3139-3146
31. Tang Z, Wang A, Yuan F *et al.* Differentiation of multipotent vascular stem cells contributes to vascular diseases. *Nat Commun* 2012; 3: 875
32. Diao Y, Guthrie S, Xia SL *et al.* Long-term engraftment of bone marrow-derived cells in the intimal hyperplasia lesion of autologous vein grafts. *Am J Pathol* 2008; 172: 839-848
33. Caplice NM, Wang S, Tracz M *et al.* Neoangiogenesis and the presence of progenitor cells in the venous limb of an arteriovenous fistula in the rat. *Am J Physiol Renal Physiol* 2007; 293: F470-F475
34. Owens CD. Adaptive changes in autogenous vein grafts for arterial reconstruction: clinical implications. *J Vasc Surg* 2010; 51: 736-746
35. Lee T, Roy-Chaudhury P. Advances and new frontiers in the pathophysiology of venous neointimal hyperplasia and dialysis access stenosis. *Adv Chronic Kidney Dis* 2009; 16: 329-338
36. Ene-Iordache B, Remuzzi A. Disturbed flow in radial-cephalic arteriovenous fistulae for haemodialysis: low and oscillating shear stress locates the sites of stenosis. *Nephrol Dial Transplant* 2012; 27: 358-368

37. Ene-Iordache B, Cattaneo L, Dubini G, Remuzzi A. Effect of anastomosis angle on the localization of disturbed flow in 'side-to-end' fistulae for haemodialysis access. *Nephrol Dial Transplant* 2012;
38. Krishnamoorthy MK, Banerjee RK, Wang Y *et al.* Hemodynamic wall shear stress profiles influence the magnitude and pattern of stenosis in a pig AV fistula. *Kidney Int* 2008; 74: 1410-1419
39. Lee T, Chauhan V, Krishnamoorthy M *et al.* Severe venous neointimal hyperplasia prior to dialysis access surgery. *Nephrol Dial Transplant* 2011; 26: 2264-2270
40. Friedl R, Li J, Schumacher B *et al.* Intimal hyperplasia and expression of transforming growth factor-beta1 in saphenous veins and internal mammary arteries before coronary artery surgery. *Ann Thorac Surg* 2004; 78: 1312-1318
41. Hernandez T, Saudan P, Berney T, Merminod T, Bednarkiewicz M, Martin PY. Risk factors for early failure of native arteriovenous fistulas. *Nephron Clin Pract* 2005; 101: c39-c44
42. Lok CE, Allon M, Moist L, Oliver MJ, Shah H, Zimmerman D. Risk equation determining unsuccessful cannulation events and failure to maturation in arteriovenous fistulas (REDUCE FTM I). *J Am Soc Nephrol* 2006; 17: 3204-3212
43. Monroy-Cuadros M, Yilmaz S, Salazar-Banuelos A, Doig C. Risk factors associated with patency loss of hemodialysis vascular access within 6 months. *Clin J Am Soc Nephrol* 2010; 5: 1787-1792
44. Heine GH, Ulrich C, Sester U, Sester M, Kohler H, Girndt M. Transforming growth factor beta1 genotype polymorphisms determine AV fistula patency in hemodialysis patients. *Kidney Int* 2003; 64: 1101-1107
45. Lin CC, Yang WC, Lin SJ *et al.* Length polymorphism in heme oxygenase-1 is associated with arteriovenous fistula patency in hemodialysis patients. *Kidney Int* 2006; 69: 165-172
46. Tummers AM, Mountain DJ, Mix JW *et al.* Serum levels of matrix metalloproteinase-2 as a marker of intimal hyperplasia. *J Surg Res* 2010; 160: 9-13
47. Misra S, Fu AA, Anderson JL *et al.* The rat femoral arteriovenous fistula model: increased expression of matrix metalloproteinase-2 and -9 at the venous stenosis. *J Vasc Interv Radiol* 2008; 19: 587-594
48. Cheng XW, Kuzuya M, Sasaki T *et al.* Increased expression of elastolytic cysteine proteases, cathepsins S and K, in the neointima of balloon-injured rat carotid arteries. *Am J Pathol* 2004; 164: 243-251
49. Abeles D, Kwei S, Stavrakis G, Zhang Y, Wang ET, Garcia-Cardena G. Gene expression changes evoked in a venous segment exposed to arterial flow. *J Vasc Surg* 2006; 44: 863-870
50. Amabile PG, Wong H, Uy M *et al.* In vivo vascular engineering of vein grafts: directed migration of smooth muscle cells by perivascular release of elastase limits neointimal proliferation. *J Vasc Interv Radiol* 2002; 13: 709-715
51. Mochizuki S, Brassart B, Hinek A. Signaling pathways transduced through the elastin receptor facilitate proliferation of arterial smooth muscle cells. *J Biol Chem* 2002; 277: 44854-44863
52. Reddel CJ, Weiss AS, Burgess JK. Elastin in asthma. *Pulm Pharmacol Ther* 2012; 25: 144-153
53. Palmes D, Kebschull L, Schaefer RM, Pelster F, Konner K. Perforating vein fistula is superior to forearm fistula in elderly haemodialysis patients with diabetes and arterial hypertension. *Nephrol Dial Transplant* 2011; 26: 3309-3314

54. Konner K, Hulbert-Shearon TE, Roys EC, Port FK. Tailoring the initial vascular access for dialysis patients. *Kidney Int* 2002; 62: 329-338
55. Vanholder R, Massy Z, Argiles A, Spasovski G, Verbeke F, Lameire N. Chronic kidney disease as cause of cardiovascular morbidity and mortality. *Nephrol Dial Transplant* 2005; 20: 1048-1056
56. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004; 351: 1296-1305
57. Brunet P, Gondouin B, Duval-Sabatier A *et al.* Does uremia cause vascular dysfunction? *Kidney Blood Press Res* 2011; 34: 284-290
58. Gunthner T, Jankowski V, Kretschmer A *et al.* Endothelium and vascular smooth muscle cells in the context of uremia. *Semin Dial* 2009; 22: 428-432
59. Moody WE, Edwards NC, Madhani M *et al.* Endothelial dysfunction and cardiovascular disease in early-stage chronic kidney disease: cause or association? *Atherosclerosis* 2012; 223: 86-94
60. Kokubo T, Ishikawa N, Uchida H *et al.* CKD accelerates development of neointimal hyperplasia in arteriovenous fistulas. *J Am Soc Nephrol* 2009; 20: 1236-1245
61. Wali MA, Eid RA, Dewan M, Al-Homrany MA. Pre-existing histopathological changes in the cephalic vein of renal failure patients before arterio-venous fistula (AVF) construction. *Ann Thorac Cardiovasc Surg* 2006; 12: 341-348
62. Wasse H, Huang R, Naqvi N, Smith E, Wang D, Husain A. Inflammation, oxidation and venous neointimal hyperplasia precede vascular injury from AVF creation in CKD patients. *J Vasc Access* 2012; 13: 168-174
63. O'Neill WC, Lomashvili KA. Recent progress in the treatment of vascular calcification. *Kidney Int* 2010; 78: 1232-1239
64. Blacher J, Guerin AP, Pannier B, Marchais SJ, London GM. Arterial calcifications, arterial stiffness, and cardiovascular risk in end-stage renal disease. *Hypertension* 2001; 38: 938-942
65. Guerin AP, London GM, Marchais SJ, Metivier F. Arterial stiffening and vascular calcifications in end-stage renal disease. *Nephrol Dial Transplant* 2000; 15: 1014-1021
66. Huang PH, Chen LC, Leu HB *et al.* Enhanced coronary calcification determined by electron beam CT is strongly related to endothelial dysfunction in patients with suspected coronary artery disease. *Chest* 2005; 128: 810-815
67. Lee T, Safdar N, Mistry MJ *et al.* Preexisting Venous Calcification Prior to Dialysis Vascular Access Surgery. *Semin Dial* 2012;
68. van der Linden J, Lameris TW, van den Meiracker AH, de Smet AA, Blankestijn PJ, van den Dorpel MA. Forearm venous distensibility predicts successful arteriovenous fistula. *Am J Kidney Dis* 2006; 47: 1013-1019
69. Allon M, Litovsky S, Young CJ *et al.* Medial fibrosis, vascular calcification, intimal hyperplasia, and arteriovenous fistula maturation. *Am J Kidney Dis* 2011; 58: 437-443
70. Tordoir J, Canaud B, Haage P *et al.* EBPG on Vascular Access. *Nephrol Dial Transplant* 2007; 22 Suppl 2: ii88-117
71. Chang CJ, Ko PJ, Hsu LA *et al.* Highly increased cell proliferation activity in the restenotic hemodialysis vascular access after percutaneous transluminal angioplasty: implication in prevention of restenosis. *Am J Kidney Dis* 2004; 43: 74-84
72. Beathard GA. Percutaneous transvenous angioplasty in the treatment of vascular access stenosis. *Kidney Int* 1992; 42: 1390-1397

73. Lee T, Ullah A, Allon M *et al.* Decreased cumulative access survival in arteriovenous fistulas requiring interventions to promote maturation. *Clin J Am Soc Nephrol* 2011; 6: 575-581
74. Peden EK, Leeseer DB, Dixon BS *et al.* A multi-center, dose-escalation study of human type I pancreatic elastase (PRT-201) administered after arteriovenous fistula creation. *J Vasc Access* 2012; 0: 0
75. Pai AS, Giachelli CM. Matrix remodeling in vascular calcification associated with chronic kidney disease. *J Am Soc Nephrol* 2010; 21: 1637-1640
76. Paulson WD, Kipshidze N, Kipiani K *et al.* Safety and efficacy of local periadventitial delivery of sirolimus for improving hemodialysis graft patency: first human experience with a sirolimus-eluting collagen membrane (Coll-R). *Nephrol Dial Transplant* 2012; 27: 1219-1224
77. Kelly B, Melhem M, Zhang J *et al.* Perivascular paclitaxel wraps block arteriovenous graft stenosis in a pig model. *Nephrol Dial Transplant* 2006; 21: 2425-2431
78. McAllister TN, Maruszewski M, Garrido SA *et al.* Effectiveness of haemodialysis access with an autologous tissue-engineered vascular graft: a multicentre cohort study. *Lancet* 2009; 373: 1440-1446
79. Dahl SL, Kypson AP, Lawson JH *et al.* Readily available tissue-engineered vascular grafts. *Sci Transl Med* 2011; 3: 68ra9
80. Peck MK, Dusserre N, Zagalski K *et al.* New biological solutions for hemodialysis access. *J Vasc Access* 2011; 12: 185-192
81. Wu W, Allen RA, Wang Y. Fast-degrading elastomer enables rapid remodeling of a cell-free synthetic graft into a neoartery. *Nat Med* 2012; 18: 1148-1153

