



Universiteit
Leiden
The Netherlands

Allosteric modulation by sodium ions and amilorides of G protein-coupled receptors : a closer look at the sodium ion site of the adenosine A2a receptor and development of a mass spectrometry ligand binding assay for adenosine A1 and A2a receptors

Massink, A.

Citation

Massink, A. (2016, December 8). *Allosteric modulation by sodium ions and amilorides of G protein-coupled receptors : a closer look at the sodium ion site of the adenosine A2a receptor and development of a mass spectrometry ligand binding assay for adenosine A1 and A2a receptors*. Retrieved from <https://hdl.handle.net/1887/44707>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/44707>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/44707> holds various files of this Leiden University dissertation.

Author: Massink, A.

Title: Allosteric modulation by sodium ions and amilorides of G protein-coupled receptors : a closer look at the sodium ion site of the adenosine A2a receptor and development of a mass spectrometry ligand binding assay for adenosine A1 and A2a receptors

Issue Date: 2016-12-08

Stellingen

bij het proefschrift

Allosteric modulation by sodium ions and amilorides of G protein-coupled receptors

A closer look at the sodium ion site of the adenosine A_{2A} receptor

Development of a mass spectrometry ligand binding assay for adenosine A₁ and A_{2A} receptors

1. Understanding allosteric mechanisms of GPCRs on a molecular level is important for drug development.
Keov P, et al. Neuropharmacology, 2011. 60(1): 24-35.
2. The sodium ion site is a versatile allosteric site that can exert various allosteric effects, even if it is well conserved amongst Class A GPCRs.
Chapter 2, this thesis.
3. At physiological concentrations sodium ions are important regulators of GPCR activity; in particular adenosine A_{2A} receptor activity is down-regulated significantly.
Chapter 3, this thesis.
4. The sodium ion site has a vital role in the activation mechanism of the adenosine A_{2A} receptor and of GPCRs in general.
Chapter 4, this thesis.
5. The proximity of the allosteric sodium ion site and the orthosteric site allows the synthesis of a bitopic ligand binding at both sites, which may provide more information on allosteric mechanisms of GPCRs.
Chapter 5, this thesis; Guo D, et al. Br J Pharmacol, 2014. 171(23): 5295-312.
6. Computational biology is vital to understand biochemical processes on a molecular level but still needs a firm foundation of 'wet lab' work.
Chapters 3 and 4, this thesis.
7. Even if the "snapshot" structural information provided by X-ray crystallization of GPCRs has given a substantial boost to the field, the next step to better understand GPCR allosteric and activation mechanisms will be pushed by time-resolved methods that can follow structural dynamics at an atomic scale.
Isogai S, et al. Nature, 2016. 530(7589): 237-41; Deupi X. Biochim Biophys Acta, 2014. 1837(5): 674-82.
8. As well researched targets adenosine receptors are excellent model systems to evaluate new methods.
Fredholm BB, et al. Pharmacol Rev, 2011. 63(1): 1-34.
9. The ongoing development of label-free assays for GPCRs greatly expands the possibilities to study protein-ligand interactions.
Jacobson KA. Biochem Pharmacol, 2015. 98(4): 541-55.
10. Mass spectrometry provides an excellent label-free alternative for the quantification of ligand binding to radiolabeled and fluorescently labeled ligands.
Chapter 6, this thesis; Neiens P, et al. ChemMedChem, 2015. 10(11): 1924-31.
11. Research is the process of going up alleys to see if they're blind.
Barstow Bates.
12. Doing a PhD comes with interesting time-dilation effects: four years seem plenty of time in the beginning, but appear to be nothing in the end.