



Universiteit
Leiden
The Netherlands

Deciphering fermionic matter: from holography to field theory

Meszéna, B.

Citation

Meszéna, B. (2016, December 21). *Deciphering fermionic matter: from holography to field theory*. Casimir PhD Series. Retrieved from <https://hdl.handle.net/1887/45226>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/45226>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/45226> holds various files of this Leiden University dissertation.

Author: Meszéna, B.

Title: Deciphering fermionic matter: from holography to field theory

Issue Date: 2016-12-21

Deciphering Fermionic Matter: from Holography to Field Theory

PROEFSCHRIFT

TER VERKRIJGING VAN
DE GRAAD VAN DOCTOR AAN DE UNIVERSITEIT LEIDEN,
OP GEZAG VAN RECTOR MAGNIFICUS
PROF. MR. C.J.J.M. STOLKER,
VOLGENS BESLUIT VAN HET COLLEGE VOOR PROMOTIES
TE VERDEDIGEN OP WOENSDAG 21 DECEMBER 2016
KLOKKE 13.45 UUR

DOOR

Balazs Meszena

GEBOREN TE BOEDAPEST, HONGARIJE IN 1989

Promotor: Prof. dr K. E. Schalm
Promotiecommissie: Prof. dr S. S. Lee (McMaster University, Hamilton,
Canada)
Dr P. Strack (University of Cologne, Duitsland)
Prof. dr E. R. Eliel
Prof. dr ir T. H. Oosterkamp
Prof. dr J. Zaanen

Casimir PhD series, Delft-Leiden 2016-35

ISBN 978-90-8593-228-9

An electronic version of this thesis can be found at
<https://openaccess.leidenuniv.nl>

Cover design: Proefschriftmaken.nl — Uitgeverij BOXPress

Printed & Lay Out by: Proefschriftmaken.nl — Uitgeverij BOXPress

The research presented in this thesis was founded by Leiden University
through a Huygens fellowship.

Contents

1	Introduction	1
1.1	Phases of fermionic matter: Fermi and non-Fermi liquids	3
1.1.1	Fermi liquids	3
1.1.2	Conductivity of Fermi liquids	5
1.1.3	Strongly correlated fermions: Non-Fermi liquids	7
1.1.4	Ising-nematic transitions: quantum critical boson coupled to a Fermi surface	11
1.2	Non-perturbative methods	14
1.2.1	Large-N theories	14
1.2.2	Conformal field theories	20
1.3	AdS/CFT	25
1.3.1	AdS spacetime	27
1.3.2	The dictionary	30
1.3.3	Finite density and temperature	34
1.3.4	Holographic fermions	38
1.4	This thesis	44
2	Pairing induced superconductivity in holography	53
2.1	Introduction	53
2.2	Review of fermion spectra in the AdS dual: spin splitting	56
2.3	Self interacting fermions in AdS and a bulk BCS theory	61
2.3.1	Majorana interaction	61
2.3.2	Nambu-Gorkov formalism	61
2.3.3	Perturbative calculation of the scalar source	65
2.3.4	Analytical study of the non-dynamical scalar: double gap equation	66
2.4	Fermionic ordering in holography	69
2.4.1	Purely scalar holographic superconductor	71
2.4.2	Bose-Fermi competition	71
2.4.3	A dynamical BCS scalar and a BCS/BEC crossover	73
2.5	Conclusions	82
2.A	Green's functions and charge densities	83
2.B	Perturbative solution	85

2.B.1	Perturbative fermion spectrum, AdS-gap equation	85
2.B.2	Simplified Gap equation	86
2.C	Numerical methods	87
2.C.1	General strategy	87
3	Non-perturbative emergence of non-Fermi liquid behaviour in $d = 2$ quantum critical metals	95
3.1	Introduction	95
3.2	2+1 dimensional quantum critical metals in the patch approximation	102
3.2.1	The $N_f = 0$ quenched approximation and Landau damping	104
3.2.2	The exact fermion Green's function	105
3.3	The physics of the planar quenched quantum critical metal	107
3.4	Conclusions	115
3.A	Comparison with perturbation theory	116
3.B	Calculating the real-space fermion Green's function	117
3.C	Fourier transforming the fermion Green's function	119
3.D	Integrals of the spectral function	121
4	The Green's function of a $d = 2$ quantum critical metal at large k_F and small N_f	129
4.1	Introduction	129
4.2	A closed system of Schwinger-Dyson equations and large k_F Ward identities for the elementary quantum critical metal	133
4.2.1	The $k_F \rightarrow \infty$ limit of the Schwinger-Dyson equations and the fermion two-point function: formal connection with the quenched result	135
4.3	The $k_F \rightarrow \infty$ limit of the Schwinger-Dyson equations and the exact boson two-point function	137
4.3.1	Polarization/Landau damping contribution at one-loop	138
4.3.2	The solution to the $k_F \rightarrow \infty$ Schwinger-Dyson equation: Robustness of the one-loop result	139
4.3.3	Multiloop cancellation	140
4.4	The fermion two-point function in the limit $k_F \rightarrow \infty$ at small N_f	142
4.4.1	The exact fermionic two-point function in real space: an analytical form	142

4.4.2	The exact fermion two-point function in momentum space: Numerical method	153
4.5	The physics of 2+1 quantum critical metals at large k_F	155
4.6	Conclusion	160
4.A	Derivation of the Schwinger-Dyson equations and the Ward Identity	161
4.B	The Fourier transform of the fermion Green's function in the large $N_f k_F$ approximation	162
4.C	The discontinuous transition from the quenched to the Landau- damped regime	167
Samenvatting		179
Summary		181
List of Publications		183
Curriculum Vitæ		185
Acknowledgments		187

