



Universiteit
Leiden
The Netherlands

Coulomb-assisted braiding of Majorana fermions in a Josephson junction array

Heck, B. van; Akhmerov, A.R.; Hassler, F.; Burrello, M.; Beenakker, C.W.J.

Citation

Heck, B. van, Akhmerov, A. R., Hassler, F., Burrello, M., & Beenakker, C. W. J. (2012). Coulomb-assisted braiding of Majorana fermions in a Josephson junction array. *New Journal Of Physics*, 14, 035019. doi:10.1088/1367-2630/14/3/035019

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/50550>

Note: To cite this publication please use the final published version (if applicable).

Coulomb-assisted braiding of Majorana fermions in a Josephson junction array

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2012 New J. Phys. 14 035019

(<http://iopscience.iop.org/1367-2630/14/3/035019>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 132.229.211.17

This content was downloaded on 10/05/2017 at 12:45

Please note that [terms and conditions apply](#).

You may also be interested in:

[The top-transmon: a hybrid superconducting qubit for parity-protected quantum computation](#)

F Hassler, A R Akhmerov and C W J Beenakker

[Minimal circuit for a flux-controlled Majorana qubit in a quantum spin-Hall insulator](#)

B van Heck, T Hyart and C W J Beenakker

[Anyonic interferometry without anyons: how a flux qubit can read out a topological qubit](#)

F Hassler, A R Akhmerov, C-Y Hou et al.

[Strongly interacting Majorana modes in an array of Josephson junctions](#)

Fabian Hassler and Dirk Schuricht

[New directions in the pursuit of Majorana fermions in solid state systems](#)

Jason Alicea

[Introduction to topological superconductivity and Majorana fermions](#)

Martin Leijnse and Karsten Flensberg

[Flux-charge duality and topological quantum phase fluctuations in quasi-one-dimensional superconductors](#)

Andrew J Kerman

[Controllable coupling between flux qubit and nanomechanical resonator by magnetic field](#)

Fei Xue, Y D Wang, C P Sun et al.

[Superconducting qubit network with controllable nearest-neighbour coupling](#)

M Wallquist, J Lantz, V S Shumeiko et al.

Coulomb-assisted braiding of Majorana fermions in a Josephson junction array

**B van Heck^{1,3}, A R Akhmerov¹, F Hassler², M Burrello¹
and C W J Beenakker¹**

¹ Instituut-Lorentz, Universiteit Leiden, PO Box 9506, 2300 RA Leiden, The Netherlands

² Institute for Quantum Information, RWTH Aachen University, D-52056 Aachen, Germany

E-mail: bernard@lorentz.leidenuniv.nl

New Journal of Physics **14** (2012) 035019 (14pp)

Received 16 December 2011

Published 28 March 2012

Online at <http://www.njp.org/>

doi:10.1088/1367-2630/14/3/035019

Abstract. We show how to exchange (braid) Majorana fermions in a network of superconducting nanowires by control over Coulomb interactions rather than tunneling. Even though Majorana fermions are charge-neutral quasiparticles (equal to their own antiparticle), they have an effective long-range interaction through the even–odd electron number dependence of the superconducting ground state. The flux through a split Josephson junction controls this interaction via the ratio of Josephson and charging energies, with exponential sensitivity. By switching the interaction on and off in neighboring segments of a Josephson junction array, the non-Abelian braiding statistics can be realized without the need to control tunnel couplings by gate electrodes.

³ Author to whom any correspondence should be addressed.

Contents

1. Introduction	2
2. The Majorana–Coulomb Hamiltonian	3
2.1. A single island	3
2.2. Multiple islands	4
2.3. Tri-junction	5
3. Majorana braiding	7
4. Discussion	8
Acknowledgments	10
Appendix A. Derivation of the Majorana–Coulomb Hamiltonian	10
Appendix B. Calculation of the Berry phase of the braiding operation	12
References	13

1. Introduction

Non-Abelian anyons have a topological charge that provides a nonlocal encoding of quantum information [1]. In superconducting implementations [2, 3] the topological charge is equal to the electrical charge modulo $2e$, shared nonlocally by a pair of midgap states called Majorana fermions [4]. This mundane identification of topological and electrical charge by no means diminishes the relevance for quantum computation. In contrast, it provides a powerful way to manipulate the topological charge through the well-established sub- e charge sensitivity of superconducting electronics [5, 6].

Following this line of thought, three of us recently proposed a hybrid device called a *top-transmon*, which combines the adjustable charge sensitivity of a superconducting charge qubit (the *transmon* [7]) to read out and rotate a topological (*top*) qubit [8]. A universal quantum computer with highly favorable error threshold can be constructed [9] if these operations are supplemented by the pairwise exchange (braiding) of Majorana fermions, which is a non-Abelian operation on the degenerate ground state [10, 11].

Here we show how Majorana fermions can be braided by means of charge-sensitive superconducting electronics. (Braiding was not implemented in [8], nor in other studies of hybrid topological/nontopological qubits [12–16].) We exploit the fact that the charge sensitivity can be switched on and off *with exponential accuracy* by varying the magnetic flux through a split Josephson junction [7]. This provides a macroscopic handle on the Coulomb interaction of pairs of Majorana fermions, which makes it possible to transport and exchange them in a Josephson junction array.

We compare and contrast our approach with that of Sau, Clarke and Tewari, who showed (building on the work of Alicea *et al* [17]) how non-Abelian braiding statistics could be generated by switching on and off the tunnel coupling of adjacent pairs of Majorana fermions [18]. The tunnel coupling is controlled by a gate voltage, while we rely on Coulomb interaction controlled by a magnetic flux. This becomes an essential difference when electric fields are screened by the superconductor too strongly to be effective. (For an alternative non-electrical approach to braiding, see [19].)




Figure 1. Cooper pair box, consisting of a superconducting island (brown) connected to a bulk superconductor by a split Josephson junction (black, with the gauge-variant phase differences indicated). The island contains Majorana fermions (yellow) at the end points of a nanowire (gray). These are coupled by the Coulomb charging energy, tunable via the flux Φ through the Josephson junction.

The basic procedure can be explained quite simply, see section 3, once the mechanism of the Coulomb coupling is presented in section 2. We make use of two more involved pieces of theoretical analysis: one is the derivation of the low-energy Hamiltonian of the Coulomb-coupled Majorana fermions (using the results of [20, 21]) and the other is the calculation of the non-Abelian Berry phase [22] of the exchange operation. To streamline the paper, details of these two calculations are given in the [appendices](#).

2. The Majorana–Coulomb Hamiltonian

2.1. A single island

The basic building block of the Josephson junction array is the Cooper pair box [23, 24], see figure 1, consisting of a superconducting island (capacitance C) connected to a bulk (grounded) superconductor by a split Josephson junction enclosing a magnetic flux Φ . The Josephson energy E_J is a periodic function of Φ with period $\Phi_0 = h/2e$. If the two arms of the split junction are balanced, each with the same coupling energy E_0 , the Josephson energy

$$E_J = 2E_0 \cos(\pi \Phi / \Phi_0) \quad (1)$$

varies between 0 and $2E_0 > 0$ as a function of $|\Phi| < \Phi_0/2$.

When the island contains no Majorana fermions, its Hamiltonian has the usual form [25]

$$H = \frac{1}{2C} (Q + q_{\text{ind}})^2 - E_J \cos \phi, \quad (2)$$

in terms of the canonically conjugate phase ϕ and charge $Q = -2ei d/d\phi$ of the island. The offset q_{ind} accounts for charges on nearby gate electrodes. We have chosen a gauge such that the phase of the pair potential is zero on the bulk superconductor.

A segment of a semiconductor nanowire (typically InAs) on the superconducting island can have Majorana midgap states bound to the end points [2, 3]. For N segments there can be

$2N$ Majorana fermions on the island. They have identical creation and annihilation operators $\gamma_n = \gamma_n^\dagger$ satisfying

$$\gamma_n \gamma_m + \gamma_m \gamma_n = 2\delta_{nm}. \quad (3)$$

The topological charge of the island equals the fermion parity

$$\mathcal{P} = i^N \prod_{n=1}^{2N} \gamma_n. \quad (4)$$

The eigenvalues of \mathcal{P} are ± 1 , depending on whether there is an even or an odd number of electrons on the island.

The Majorana operators do not enter explicitly H , but affect the spectrum through a constraint on the eigenstates [20],

$$\Psi(\phi + 2\pi) = (-1)^{(1-\mathcal{P})/2} \Psi(\phi). \quad (5)$$

This ensures that the eigenvalues of Q are even multiples of e for $\mathcal{P} = 1$ and odd multiples for $\mathcal{P} = -1$. Since \mathcal{P} contains the product of all the Majorana operators on the island, the constraint (5) effectively couples distant Majorana fermions—without requiring any overlap of wave functions.

We operate the Cooper pair box in the regime where the Josephson energy E_J is large compared to the single-electron charging energy $E_C = e^2/2C$. The phase ϕ (modulo 2π) then has small zero-point fluctuations around the value $\phi_{\min} = 0$, which minimizes the energy of the Josephson junction, with occasional 2π quantum phase slips.

In appendix A we derive the effective low-energy Hamiltonian for $E_J \gg E_C$,

$$H_{\text{eff}} = -E_J + \sqrt{2E_C E_J} - U\mathcal{P}, \quad (6)$$

$$U = 16(E_C E_J^3 / 2\pi^2)^{1/4} e^{-\sqrt{8E_J/E_C}} \cos(\pi q_{\text{ind}}/e). \quad (7)$$

The energy minimum $-2E_0$ at ϕ_{\min} is increased by $\sqrt{2E_C E_J}$ due to zero-point fluctuations of the phase. This offset does not contain the Majorana operators, so it can be ignored. The term $-U\mathcal{P}$ due to quantum phase slips depends on the Majorana operators through the fermion parity. This term acquires a dynamics for multiple coupled islands, because then the fermion parity of each individual island is no longer conserved.

2.2. Multiple islands

We generalize the description to multiple superconducting islands, labeled $k = 1, 2, \dots$, each connected to a bulk superconductor by a split Josephson junction enclosing a flux Φ_k . (See figure 2.) The Josephson junctions contribute an energy

$$H_J = - \sum_k E_{J,k} \cos \phi_k, \quad E_{J,k} = 2E_0 \cos(\pi \Phi_k / \Phi_0). \quad (8)$$

We assume that the charging energy is dominated by the self-capacitance C of each island, so that it has the additive form

$$H_C = \sum_k \frac{1}{2C} (Q_k + q_{\text{ind},k})^2. \quad (9)$$




Figure 2. Two Cooper pair boxes, each containing a pair of Majorana fermions. Single electrons can tunnel between the superconducting islands via the overlapping Majoranas γ_{12} and γ_{21} . This tunnel coupling has a slow (cosine) dependence on the enclosed fluxes, while the Coulomb coupling between the Majoranas on the same island varies rapidly (exponentially).

While both E_0 and C may be different for different islands, we omit a possible k dependence for ease of notation. There may be additional fluxes enclosed by the regions between the islands, but we do not include them to simplify the expressions. None of these simplifications are essential for the operation of the device.

The set of Majoranas on the k th island is indicated by γ_{kn} with $n = 1, 2, \dots, 2N_k$. The fermion parities $\mathcal{P}_k = i^{N_k} \prod_n \gamma_{kn}$ of neighboring islands k and k' are coupled with strength E_M by the overlapping Majoranas γ_{kn} and $\gamma_{k'm}$. We denote the gauge-invariant phase difference [25] by $\theta_{kk'} = \phi_k - \phi_{k'} + (2\pi/\Phi_0) \int_{k \rightarrow k'} \mathbf{A} \cdot d\mathbf{l}$. The corresponding tunnel Hamiltonian [4]

$$H_{kk'} = \Gamma_{kk'} \cos(\theta_{kk'}/2), \quad \Gamma_{kk'} = iE_M \gamma_{kn} \gamma_{k'm} \quad (10)$$

is 4π -periodic in the gauge-invariant phase difference, as an expression of the fact that single electrons (rather than Cooper pairs) tunnel through the midgap state. For example, in the two-island geometry of figure 2, one has

$$H_{12} = iE_M \gamma_{12} \gamma_{21} \cos(\theta_{12}/2), \quad (11a)$$

$$\theta_{12} = \phi_1 - \phi_2 - \pi(\Phi_1 + \Phi_2)/\Phi_0. \quad (11b)$$

In appendix A, we derive the effective low-energy Hamiltonian in the regime $E_J \gg E_C, E_M$,

$$H_{\text{eff}} = \text{const} - \sum_k U_k \mathcal{P}_k + \sum_{k,k'} \Gamma_{kk'} \cos \alpha_{kk'}, \quad (12)$$

$$\alpha_{kk'} = \lim_{\phi_k, \phi_{k'} \rightarrow 0} \frac{1}{2} \theta_{kk'}. \quad (13)$$

The single sum couples Majoranas within an island, through an effective Coulomb energy U_k . The double sum couples Majoranas in neighboring islands by tunneling. Both the Coulomb and tunnel couplings depend on the fluxes through the Josephson junctions, but in an entirely different way: the tunnel coupling varies slowly $\propto \cos(\pi \Phi / \Phi_0)$ with the flux, while the Coulomb coupling varies rapidly $\propto \exp[-4\sqrt{(E_0/E_C) \cos(\pi \Phi / \Phi_0)}]$.

2.3. Tri-junction

Since \mathcal{P}_k and $\Gamma_{kk'}$ in the Majorana–Coulomb Hamiltonian (12) do not commute, the evolution of the eigenstates upon variation of the fluxes is nontrivial. As we will demonstrate, it can provide the non-Abelian braiding statistic that we are seeking.




Figure 3. Three Cooper pair boxes connected at a tri-junction via three overlapping Majorana fermions (which effectively produce a single zero-mode γ_0 at the center). This is the minimal setup required for the braiding of a pair of Majoranas, controlled by the fluxes through the three Josephson junctions to a bulk superconductor.

Similarly to earlier braiding proposals [17, 18], the minimal setup consists of three superconductors in a tri-junction. (See figure 3.) Each superconductor contains a pair of Majorana fermions γ_k, γ'_k , with a tunnel coupling between γ'_1, γ'_2 and γ'_3 . The Majorana–Coulomb Hamiltonian (12) takes the form

$$H_{\text{eff}} = iE_M(\gamma'_1\gamma'_2 \cos \alpha_{12} + \gamma'_2\gamma'_3 \cos \alpha_{23} + \gamma'_3\gamma'_1 \cos \alpha_{31}) - \sum_{k=1}^3 U_k i\gamma_k\gamma'_k, \quad (14)$$

with gauge-invariant phase differences

$$\alpha_{12} = -(\pi/2\Phi_0)(\Phi_1 + \Phi_2 + 2\Phi_3), \quad (15a)$$

$$\alpha_{23} = (\pi/2\Phi_0)(\Phi_2 + \Phi_3), \quad (15b)$$

$$\alpha_{31} = (\pi/2\Phi_0)(\Phi_1 + \Phi_3). \quad (15c)$$

As we vary $|\Phi_k|$ between 0 and $\Phi_{\text{max}} < \Phi_0/2$, the Coulomb coupling U_k varies between two (possibly k -dependent) values U_{min} and U_{max} . We require $U_{\text{max}} \gg U_{\text{min}}$, which is readily achievable because of the exponential flux sensitivity of the Coulomb coupling expressed by equations (1) and (7). We call the Coulomb couplings U_{max} and U_{min} *on* and *off*, respectively. We also take $U_{\text{max}} \ll E_M$, meaning that the Coulomb coupling is weaker than the tunnel coupling. This is not an essential assumption, but it allows us to reduce the six-Majorana problem to a four-Majorana problem, as we will now show.

Consider first the case when $U_k = 0$ for all k . Then the Hamiltonian (14) has four eigenvalues equal to zero: three of these represent the Majoranas γ_k far away from the junction, while the fourth Majorana,

$$\gamma_0 = \frac{1}{\sqrt{3}}(\gamma'_1 + \gamma'_2 + \gamma'_3), \quad (16)$$

is situated at the tri-junction. The tri-junction also contributes two nonzero eigenvalues $\pm \frac{1}{2}E_{\text{gap}}$, separated by the gap

$$E_{\text{gap}} = E_M \sqrt{\cos^2 \alpha_{12} + \cos^2 \alpha_{23} + \cos^2 \alpha_{31}}. \quad (17)$$

Table 1. The variation of the flux through the three Josephson junctions during the braiding operation, at time steps corresponding to the diagrams in figure 4. The flux Φ_3 is varied in the opposite direction as Φ_1, Φ_2 , to ensure that the coupling parameters $\Delta_k \propto \beta_k$ do not change sign during the operation.

Time	Φ_1	Φ_2	Φ_3
0	0	0	$-\Phi_{\max}$
	Φ_{\max}	0	$-\Phi_{\max}$
T	Φ_{\max}	0	0
	Φ_{\max}	Φ_{\max}	0
$2T$	0	Φ_{\max}	0
	0	Φ_{\max}	$-\Phi_{\max}$
$3T$	0	0	$-\Phi_{\max}$

For Φ_{\max} well below Φ_0 and $U_{\max} \ll E_M$ these two gapped modes can be ignored, and only the four Majoranas $\gamma_0, \gamma_1, \gamma_2, \gamma_3$ need to be retained.

The Hamiltonian H_{int} that describes the Coulomb interaction of these four Majoranas for nonzero U_k is given, to first order in U_k/E_M , by

$$H_{\text{int}} = \sum_{k=1}^3 \Delta_k i\gamma_0\gamma_k, \quad \Delta_k = -(2E_M/E_{\text{gap}})\beta_k U_k, \quad (18)$$

$$\beta_1 = \cos \alpha_{23}, \quad \beta_2 = \cos \alpha_{31}, \quad \beta_3 = \cos \alpha_{12}. \quad (19)$$

3. Majorana braiding

The Hamiltonian (18) describes four flux-tunable Coulomb-coupled Majorana fermions. Although the coupling studied by Sau *et al* [18] has an entirely different origin (gate-tunable tunnel coupling), their Hamiltonian has the same form. We can therefore directly adapt their braiding protocol to our control parameters.

We have three fluxes Φ_1, Φ_2 and Φ_3 to control the couplings. The braiding operation consists of three steps, see table 1 and figure 4. (Sau *et al* [18] had more steps, involving six rather than four Majoranas.) At the beginning and at the end of each step two of the couplings are *off* ($\Phi_k = 0$) and one coupling is *on* ($|\Phi_k| = \Phi_{\max}$). We denote by $\mathcal{O}_{kk'}$ the step of the operation that switches the coupling that is *on* from k to k' . This is done by first increasing $|\Phi_{k'}|$ from 0 to Φ_{\max} and then decreasing $|\Phi_k|$ from Φ_{\max} to 0, fixing the third flux at 0.

During the entire process the degeneracy of the ground state remains unchanged (twofold degenerate), which is a necessary condition for an adiabatic operation. If, instead, we would have first decreased $|\Phi_k|$ and then increased $|\Phi_{k'}|$, the ground state degeneracy would have switched from two to four at some point during the process, precluding adiabaticity.

We start from coupling 3 *on* and couplings 1,2 *off*. The braiding operation then consists, in sequence, of the three steps $\mathcal{O}_{31}, \mathcal{O}_{12}$, and \mathcal{O}_{23} . Note that each coupling Δ_k appears twice in the *on* state during the entire operation, both times with the same sign s_k .

The step $\mathcal{O}_{kk'}$ transfers the uncoupled Majorana at site k' to site k in a time T . The transfer is described in the Heisenberg representation by $\gamma_k(T) = \mathcal{U}^\dagger(T)\gamma_k\mathcal{U}(T)$. We calculate the unitary




Figure 4. Schematic representation of the three steps of the braiding operation. The four Majoranas of the tri-junction in figure 3 (the three outer Majoranas $\gamma_1, \gamma_2, \gamma_3$ and the effective central Majorana γ_0) are represented by circles and the Coulomb coupling is represented by lines (solid in the *on* state, dashed in the *off* state). White circles indicate Majoranas with a large Coulomb splitting and colored circles those with a vanishingly small Coulomb splitting. The small diagram above each arrow shows an intermediate stage, with one Majorana delocalized over three coupled sites. The three steps together exchange Majoranas 1 and 2, which is a non-Abelian braiding operation.

evolution operator $\mathcal{U}(T)$ in the adiabatic $T \rightarrow \infty$ limit in appendix B, by integrating over the Berry connection. In the limit $U_{\min} \rightarrow 0$ we recover the result of [18]:

$$\gamma_k(T) = -s_k s_{k'} \gamma_{k'}(0). \quad (20)$$

The result after the three steps is that the Majoranas at sites 1 and 2 are switched, with a difference in sign,

$$\gamma_1(3T) = -s_1 s_2 \gamma_2(0), \quad \gamma_2(3T) = s_1 s_2 \gamma_1(0). \quad (21)$$

The corresponding unitary time evolution operator,

$$\mathcal{U}(3T) = \frac{1}{\sqrt{2}}(1 + s_1 s_2 \gamma_1 \gamma_2) = \exp\left(\frac{\pi}{4} s_1 s_2 \gamma_1 \gamma_2\right), \quad (22)$$

has the usual form of an adiabatic braiding operation [11]. For a nonzero U_{\min} the coefficient $\pi/4$ in the exponent acquires corrections of order U_{\min}/U_{\max} , see appendix B.

If one repeats the entire braiding operation, Majoranas 1 and 2 have returned to their original positions but the final state differs from the initial state by a unitary operator $\mathcal{U}(3T)^2 = s_1 s_2 \gamma_1 \gamma_2$ and not just by a phase factor. That is the hallmark of non-Abelian statistics [10].

4. Discussion

In summary, we have proposed a way to perform non-Abelian braiding operations on Majorana fermions, by controlling their Coulomb coupling via the magnetic flux through a Josephson junction. Majorana fermions are themselves charge-neutral particles (because they are their own antiparticle), so one may ask how there can be any Coulomb coupling at all. The answer is that the state of a pair of Majorana fermions in a superconducting island depends on the parity of the number of electrons on that island, and it is this dependence on the electrical charge modulo $2e$ that provides an electromagnetic handle on the Majoranas.

The Coulomb coupling can be made exponentially small by passing Cooper pairs through a Josephson junction between the island and a bulk (grounded) superconductor. The control parameter is the flux Φ through the junction, so it is purely magnetic. This is the key difference




Figure 5. A Josephson junction array containing Majorana fermions. The magnetic flux through a split Josephson junction controls the Coulomb coupling on each superconducting island. This device allows one to perform the three types of operations on topological qubits needed for a universal quantum computer: readout, rotation and braiding. All operations are controlled magnetically; no gate voltages are needed.

from braiding by electrostatically controlled tunnel couplings of Majorana fermions [18]. Gate voltages tend to be screened quite efficiently by the superconductor, so magnetic control is advantageous. Another advantage is that the dependence of the Coulomb coupling on the flux is governed by macroscopic electrical properties (capacitance of the island and resistance of the Josephson junction). Tunnel couplings, in contrast, require microscopic input (separation of the Majorana fermions on the scale of the Fermi wavelength), so they tend to be more difficult to control. Gate electrodes are still needed to drive the nanowire into a topologically nontrivial state, but no control on the scale of the Fermi wavelength is required.

The braiding operation is called topologically protected, because it depends on the *off/on* sequence of the Coulomb couplings, and not on details of the timing of the sequence. As in any physical realization of a mathematical concept, there are sources of error. Non-adiabaticity of the operation is one source of error, studied in [26]. Low-lying sub-gap excitations in the superconducting island break the adiabatic evolution by transitions which change the fermion parity of the Majoranas.

Another source of error, studied in appendix B, is governed by the *off/on* ratio U_{\min}/U_{\max} of the Coulomb coupling. This ratio depends exponentially on the ratio of the charging energy E_C and the Josephson energy E_J of the junction to the bulk superconductor. A value $E_J/E_C \simeq 50$ is not unrealistic [7], corresponding to $U_{\min}/U_{\max} \simeq 10^{-5}$.

The sign of the Coulomb coupling in the *on* state can be arbitrary, as long as it does not change during the braiding operation. Since $U_{\max} \propto \cos(\pi q_{\text{ind}}/e)$, any change in the induced charge by $\pm e$ will spoil the operation. The time scale for this quasiparticle poisoning can be milliseconds [27], so this does not seem to present a serious obstacle.

A universal quantum computation using Majorana fermions requires, in addition to braiding, the capabilities for single-qubit rotation and readout of up to four Majoranas [1]. The combination of [8] with the present proposal provides a scheme for all three operations, based on the interface of a topological qubit and a superconducting charge qubit. (See figure 5.) This is not a topological quantum computer, since single-qubit rotations of Majorana fermions lack topological protection. But by including the topologically protected braiding operations, one can improve the tolerance for errors of the entire computation by orders of magnitude (error rates as large as 10% are permitted [9]).

It might be possible to carry over some of the ideas of this work to braiding of non-Abelian anyons in the fractional quantum Hall effect. The leading proposal in that context is braiding by interferometric measurements [28]. Braiding by control over the coupling of the anyons could provide a useful alternative.

Acknowledgments

This research was supported by the Dutch Science Foundation NWO/FOM and by an ERC Advanced Investigator Grant.

Appendix A. Derivation of the Majorana–Coulomb Hamiltonian

A.1. A Single island

Considering first a single island, we start from the Cooper pair box Hamiltonian (2) with the parity constraint (5) on the eigenstates. Following [21], it is convenient to remove the constraint by the unitary transformation

$$\tilde{H} = \Omega^\dagger H \Omega, \quad \Omega = \exp[i(1 - \mathcal{P})\phi/4]. \quad (\text{A.1})$$

The transformed wave function $\tilde{\Psi}(\phi) = \Omega^\dagger \Psi(\phi)$ is then 2π -periodic, without any constraint. The parity operator \mathcal{P} appears in the transformed Hamiltonian,

$$\tilde{H} = \frac{1}{2C} (Q + \frac{1}{2}e(1 - \mathcal{P}) + q_{\text{ind}})^2 - E_J \cos \phi. \quad (\text{A.2})$$

For a single junction the parity is conserved, so eigenstates of H are also eigenstates of \mathcal{P} and we may treat the operator \mathcal{P} as a number. Equation (A.2) is therefore the Hamiltonian of a Cooper pair box with effective induced charge $q_{\text{eff}} = q_{\text{ind}} + e(1 - \mathcal{P})/2$. The expression for the ground state energy in the Josephson regime $E_J \gg E_C$ is given in the literature [29, 30]:

$$E_{\text{ground}} = -E_J + \sqrt{2E_C E_J} - 16(E_C E_J^3 / 2\pi^2)^{1/4} e^{-\sqrt{8E_J/E_C}} \cos(\pi q_{\text{eff}}/e). \quad (\text{A.3})$$

The first term $-E_J$ is the minimal Josephson energy at $\phi_{\text{min}} = 0$. Zero-point motion, with Josephson plasma frequency $\omega_p = \sqrt{8E_C E_J}/\hbar$, adds the second term $\sqrt{2E_C C_J} = \frac{1}{2}\hbar\omega_p$. The third term is due to quantum phase slips with transition amplitudes $\tau_{\pm} \simeq \exp(\pm i\pi q_{\text{eff}}/e) \sqrt{\hbar\omega_p E_J} \exp(-\hbar\omega_p/E_J)$ by which ϕ increments by $\pm 2\pi$.

Using $\mathcal{P}^2 = 1$, the ground state energy equation (A.3) may be written in the form

$$E_{\text{ground}} = -E_J + \sqrt{2E_C E_J} - U\mathcal{P}, \quad (\text{A.4})$$

with U defined in equation (7). Higher levels are separated by an energy $\hbar\omega_p$, which is large compared to U for $E_J \gg E_C$. We may therefore identify $E_{\text{ground}} = H_{\text{eff}}$ with the effective low-energy Hamiltonian of a single island in the large E_J limit.

A.2. Multiple islands

We now turn to the case of multiple islands with tunnel coupling. To be definite, we take the geometry of two islands shown in figure 2. The full Hamiltonian is $H = H_1 + H_2 + H_{12}$, where H_1 and H_2 are two copies of the Cooper box Hamiltonian (2) and H_{12} is the tunnel coupling from equation (11).

To obtain 2π -periodicity in both phases ϕ_1 and ϕ_2 , we make the unitary transformation $\tilde{H} = \Omega^\dagger H \Omega$ with

$$\Omega = e^{i(1-\mathcal{P}_1)\phi_1/4} e^{i(1-\mathcal{P}_2)\phi_2/4}. \quad (\text{A.5})$$

The Cooper pair box Hamiltonians are transformed into

$$\tilde{H}_k = \frac{1}{2C} (Q_k + eq_k + q_{\text{ind},k})^2 - E_{J,k} \cos \phi_k, \quad (\text{A.6})$$

with $q_k = \frac{1}{2}(1 - \mathcal{P}_k)$. The tunnel coupling transforms into

$$\tilde{H}_{12} = \frac{1}{2} e^{-iq_1\phi_1} \Gamma_{12} e^{iq_2\phi_2} e^{i\pi(\Phi_1+\Phi_2)/2\Phi_0} + \text{H.c.}, \quad (\text{A.7})$$

where $\Gamma_{12} = iE_M \gamma_{12} \gamma_{21}$ and H.c. stands for the Hermitian conjugate. Since $e^{iq\phi} = \cos \phi + iq \sin \phi$, the transformed tunnel coupling \tilde{H}_{12} is 2π -periodic in ϕ_1 and ϕ_2 .

For $E_J \gg E_C$ the phases remain close to the value which minimizes the sum of the Josephson energies to the bulk superconductor and between the islands. To leading order in $E_M/E_J \ll 1$ this minimal energy is given by

$$\mathcal{E}_{\text{min}} = -E_{J,1} - E_{J,2} + \Gamma_{12} \cos[\pi(\Phi_1 + \Phi_2)/2\Phi_0] + \mathcal{O}(E_M^2/E_J). \quad (\text{A.8})$$

The Josephson coupling of the islands changes the plasma frequency $\omega_{p,k}$ for phase ϕ_k by a factor $1 + \mathcal{O}(E_M/E_J)$, so the zero-point motion energy is

$$\frac{1}{2} \hbar \omega_{p,k} = \sqrt{2E_C E_{J,k} + E_M} \times \mathcal{O}(E_C/E_J)^{1/2}. \quad (\text{A.9})$$

The transition amplitudes τ_{\pm} for quantum phase slips of phase ϕ_k are similarly affected,

$$\tau_{\pm,k} = -U_k \mathcal{P}_k + E_M e^{-\hbar\omega_{p,k}/E_{J,k}} \times \mathcal{O}(E_C/E_J)^{1/4}. \quad (\text{A.10})$$

These are the contributions to the effective Hamiltonian,

$$H_{\text{eff}} = \mathcal{E}_{\text{min}} + \sum_k \left(\frac{1}{2} \hbar \omega_{p,k} + \tau_{+,k} + \tau_{-,k} \right) \text{ for } E_J \gg E_C, E_M$$

$$H_{\text{eff}} = \left(-U_1 \mathcal{P}_1 - U_2 \mathcal{P}_2 + \Gamma_{12} \cos[\pi(\Phi_1 + \Phi_2)/2\Phi_0] \right) [1 + \mathcal{O}(E_M/E_J)] + \text{const.} \quad (\text{A.11})$$

Equation (12) in the main text generalizes this expression for two islands to an arbitrary number of coupled islands.

Appendix B. Calculation of the Berry phase of the braiding operation

We evaluate the unitary evolution operator \mathcal{U} of the braiding operation in the adiabatic limit. This amounts to a calculation of the non-Abelian Berry phase (integral of the Berry connection) of the cyclic variation of the interaction Hamiltonian $H_{\text{int}}(\Delta_1, \Delta_2, \Delta_3)$.

In the Fock basis $|00\rangle, |01\rangle, |10\rangle, |11\rangle$ the interaction Hamiltonian (18) of four Majorana fermions is given by the occupation number of the two fermionic operators $c_1 = (\gamma_1 - i\gamma_2)/2$ and $c_2 = (\gamma_0 - i\gamma_3)/2$. It takes the form

$$H_{\text{int}} = \begin{pmatrix} -\Delta_3 & 0 & 0 & -i\Delta_1 - \Delta_2 \\ 0 & \Delta_3 & -i\Delta_1 - \Delta_2 & 0 \\ 0 & i\Delta_1 - \Delta_2 & -\Delta_3 & 0 \\ i\Delta_1 - \Delta_2 & 0 & 0 & \Delta_3 \end{pmatrix}. \quad (\text{B.1})$$

The eigenvalues are doubly degenerate at energy $\pm\varepsilon = \pm\sqrt{\Delta_1^2 + \Delta_2^2 + \Delta_3^2}$ (up to a flux-dependent offset, which only contributes an overall phase factor to the evolution operator). The two degenerate ground states at $-\varepsilon$ are distinguished by an even (e) or odd (o) quasiparticle number,

$$|e\rangle = \sqrt{\frac{\varepsilon - \Delta_3}{2\varepsilon}} \begin{pmatrix} i \frac{\varepsilon + \Delta_3}{\Delta_1 + i\Delta_2} \\ 0 \\ 0 \\ 1 \end{pmatrix}, \quad (\text{B.2a})$$

$$|o\rangle = \sqrt{\frac{\varepsilon + \Delta_3}{2\varepsilon}} \begin{pmatrix} 0 \\ i \frac{\varepsilon - \Delta_3}{\Delta_1 + i\Delta_2} \\ 1 \\ 0 \end{pmatrix}. \quad (\text{B.2b})$$

This parameterization is smooth and continuous except along the line $\Delta_1 = \Delta_2 = 0$.

If we avoid this line the Berry connection can be readily evaluated. It consists of three anti-Hermitian 2×2 matrices \mathcal{A}_k ,

$$\mathcal{A}_k = \begin{pmatrix} \langle e | \frac{d}{d\Delta_k} | e \rangle & 0 \\ 0 & \langle o | \frac{d}{d\Delta_k} | o \rangle \end{pmatrix}. \quad (\text{B.3})$$

Off-diagonal terms in \mathcal{A}_k are zero because of global parity conservation. Explicitly, we have

$$\mathcal{A}_1 = \frac{\Delta_2}{\Delta_1^2 + \Delta_2^2} \begin{pmatrix} i \frac{\varepsilon + \Delta_3}{2\varepsilon} & 0 \\ 0 & i \frac{\varepsilon - \Delta_3}{2\varepsilon} \end{pmatrix}, \quad (\text{B.4})$$

$$\mathcal{A}_2 = \frac{-\Delta_1}{\Delta_1^2 + \Delta_2^2} \begin{pmatrix} i \frac{\varepsilon + \Delta_3}{2\varepsilon} & 0 \\ 0 & i \frac{\varepsilon - \Delta_3}{2\varepsilon} \end{pmatrix}, \quad (\text{B.5})$$

$$\mathcal{A}_3 = 0. \quad (\text{B.6})$$




Figure B.1. The braiding path in three-dimensional parameter space along which the Berry phase is evaluated. This path corresponds to the flux values in table 1, with couplings $\Delta_k = \Delta_{\min}$ for $\Phi_k = 0$ and $\Delta_k = \Delta_{\max}$ for $|\Phi_k| = \Phi_{\max}$. The ratio $\Delta_{\min}/\Delta_{\max}$ in the figure is exaggerated for clarity.

A closed path \mathcal{C} in parameter space has the Berry phase [22]

$$\mathcal{U} = \exp\left(-\oint_{\mathcal{C}} \sum_k A_k d\Delta_k\right). \quad (\text{B.7})$$

The path \mathcal{C} corresponding to the braiding operation in figure 4 and table 1 is shown in figure B.1. We take all couplings Δ_k positive, varying between a minimal value Δ_{\min} and a maximal value Δ_{\max} . The parametrization equation (B.2) is well defined along the entire contour.

The contour integral evaluates to

$$\mathcal{U} = \exp\left[-i\left(\frac{\pi}{4} - \epsilon\right)\sigma_z\right], \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (\text{B.8})$$

$$\epsilon = \frac{3}{\sqrt{2}} \frac{\Delta_{\min}}{\Delta_{\max}} + \mathcal{O}\left(\left(\frac{\Delta_{\min}}{\Delta_{\max}}\right)^2\right). \quad (\text{B.9})$$

The limit $\Delta_{\min}/\Delta_{\max} \rightarrow 0$ corresponds to the braiding operator (22) in the main text (with $s_1, s_2 > 0$ and $\sigma_z = 1 - 2c_1^\dagger c_1 = i\gamma_1 \gamma_2$).

References

- [1] Nayak C, Simon S, Stern A, Freedman M and Das Sarma S 2008 *Rev. Mod. Phys.* **80** 1083
- [2] Lutchyn R M, Sau J D and Das Sarma S 2010 *Phys. Rev. Lett.* **105** 077001
- [3] Oreg Y, Refael G and von Oppen F 2010 *Phys. Rev. Lett.* **105** 177002
- [4] Kitaev A Yu 2001 *Phys.—Usp.* **44** (Suppl.) 131

- [5] Averin D V and Nazarov Yu V 1992 *Phys. Rev. Lett.* **69** 1993
- [6] Devoret M H, Wallraff A and Martinis J M 2004 arXiv:cond-mat/0411174
- [7] Schreier J A *et al* 2008 *Phys. Rev. B* **77** 180502
- [8] Hassler F, Akhmerov A R and Beenakker C W J 2011 *New J. Phys.* **13** 095004
- [9] Bravyi S and Kitaev A Yu *Phys. Rev. A* **71** 022316
Bravyi S 2006 *Phys. Rev. A* **73** 042313
- [10] Read N and Green D 2000 *Phys. Rev. B* **61** 10267
- [11] Ivanov D 2001 *Phys. Rev. Lett.* **86** 268
- [12] Hassler F, Akhmerov A R, Hou C-Y and Beenakker C W J 2010 *New J. Phys.* **12** 125002
- [13] Sau J D, Tewari S and Das Sarma S 2010 *Phys. Rev. A* **82** 052322
- [14] Flensberg K 2011 *Phys. Rev. Lett.* **106** 090503
Leijnse M and Flensberg K 2011 *Phys. Rev. Lett.* **107** 210502
- [15] Jiang L, Kane C L and Preskill J 2011 *Phys. Rev. Lett.* **106** 130504
- [16] Bonderson P and Lutchyn R M 2011 *Phys. Rev. Lett.* **106** 130505
- [17] Alicea J, Oreg Y, Refael G, von Oppen F and Fisher M P A 2011 *Nature Phys.* **7** 412
- [18] Sau J D, Clarke D J and Tewari S 2011 *Phys. Rev. B* **84** 094505
- [19] Romito A, Alicea J, Refael G and von Oppen F 2011 arXiv:1110.6193
- [20] Fu L 2010 *Phys. Rev. Lett.* **104** 056402
Xu C and Fu L 2010 *Phys. Rev. B* **81** 134435
- [21] van Heck B, Hassler F, Akhmerov A R and Beenakker C W J 2011 *Phys. Rev. B* **4** 180502
- [22] Wilczek F and Zee A 1984 *Phys. Rev. Lett.* **52** 2111
- [23] Büttiker M 1987 *Phys. Rev. B* **36** 3548
- [24] Makhlin Yu, Schön G and Shnirman A 2001 *Rev. Mod. Phys.* **73** 357
- [25] Tinkham M 1996 *Introduction to Superconductivity* (New York: McGraw-Hill)
- [26] Cheng M, Galitski V and Das Sarma S 2011 *Phys. Rev. B* **84** 104529
- [27] de Visser P J, Baselmans J J A, Diener P, Yates S J C, Endo A and Klapwijk T M 2011 *Phys. Rev. Lett.* **106** 167004
- [28] Bonderson P, Freedman M and Nayak C 2008 *Phys. Rev. Lett.* **101** 010501
- [29] Koch J, Yu T M, Gambetta J, Houck A A, Schuster D I, Majer J, Blais A, Devoret M H, Girvin S M and Schoelkopf R J 2007 *Phys. Rev. A* **76** 042319
- [30] Nazarov Yu V and Blanter Ya M 2009 *Quantum Transport: Introduction to Nanoscience* (Cambridge: Cambridge University Press)