



Universiteit
Leiden
The Netherlands

When galaxy clusters collide : the impact of merger shocks on cluster gas and galaxy evolution

Stroe, A.

Citation

Stroe, A. (2015, September 2). *When galaxy clusters collide : the impact of merger shocks on cluster gas and galaxy evolution*. Retrieved from <https://hdl.handle.net/1887/34937>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/34937>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/34937> holds various files of this Leiden University dissertation

Author: Stroe, Andra

Title: When galaxy clusters collide : the impact of merger shocks on cluster gas and galaxy evolution

Issue Date: 2015-09-02

When Galaxy Clusters Collide: the impact
of merger shocks on cluster gas and galaxy evolution

When Galaxy Clusters Collide: the impact of merger shocks on cluster gas and galaxy evolution

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof.mr. C.J.J.M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op woensdag 2 september 2015
klokke 11.15 uur

door Andra Stroe
geboren te Constanța, Roemenië
in 1987

Promotiecommissie

Promotor: Prof. dr. H. J. A. Röttgering

Co-promotor: Dr. David Sobral

Overige leden: Prof. dr. Annalisa Bonafede (Hamburg Observatory)

Dr. Jarle Brinchmann

Prof. dr. Marijn Franx

Prof. dr. Jelle Kaastra

Prof. dr. Koen Kuijken

Prof. dr. Tom Oosterloo

(ASTRON, Dwingeloo;

Kapteyn Astronomical Institute)

Cover: Fractal art representing a star forming galaxy (created by Logan Apple, a.k.a. ThoughtWeaver on deviantart.com)

Bookmark: Fractal art reminiscent of particle acceleration (created by Tatyana Zabanova, a.k.a. tatasz on deviantart.com)

*In loving memory of my mother.
Thank you for being my best friend, my greatest
supporter and for always believing in me!*

Contents

Contents	i
1 Introduction	1
1.1 Evolution and composition of the Universe	1
1.2 Galaxy clusters	2
1.3 This thesis	3
1.4 Future prospects	4
2 Discovery of spectral curvature in the shock downstream region	7
2.1 Introduction	8
2.2 Observations & Data Reduction	11
2.2.1 WSRT observations	12
2.2.2 GMRT observations	12
2.3 Results	15
2.3.1 Radio continuum maps	16
2.3.2 Spectral index and spectral curvature	20
2.3.3 Colour-colour plots	25
2.4 Discussion	28
2.4.1 Northern relic	29
2.4.2 Other relics	34
2.4.3 Radio galaxies	36
2.5 Conclusion	37
2.6 Appendix	40
2.6.1 Optical counterparts of compact and diffuse sources	40
2.6.2 Spectral index and curvature error maps	40
3 Spectral age modelling of the ‘Sausage’ cluster radio relic	45
3.1 Introduction	46
3.1.1 Cluster radio relics	46
3.1.2 The ‘Sausage’ cluster	46
3.1.3 Main questions and aim of this paper	47
3.2 Observations and imaging	48
3.3 Spectral age fitting with BRATS	49
3.4 Impact of finite resolution	50
3.5 Discussion	53
3.5.1 Best-fitting model?	53
3.5.2 Injection index	54
3.5.3 Mach number	55
3.5.4 Shock and ageing speed from spectral modelling	56
3.5.5 Spectral age	58
3.5.6 Merger history	62

3.6	Conclusions	62
4	The highest-frequency detection of a radio relic	65
4.1	Introduction	66
4.2	Observations & Data Reduction	67
4.2.1	AMI observations	67
4.2.2	Imaging	67
4.3	Results	68
4.3.1	Radio morphologies	68
4.3.2	Integrated spectrum	70
4.4	Discussion	70
4.4.1	Northern relic	71
4.4.2	Diffuse extension	73
4.5	Conclusions	74
5	The ‘ultimate’ radio relic spectrum	75
5.1	Introduction	76
5.2	Observations and data reduction	77
5.2.1	Jansky VLA data	77
5.2.2	Effelsberg data	78
5.2.3	AMI data	83
5.2.4	CARMA data	83
5.3	Imaging and flux density measurements	83
5.3.1	Interferometric measurements	84
5.3.2	Total power measurements	88
5.4	Possible systematics decreasing high frequency flux densities	92
5.4.1	Mosaicking in the interferometric spectrum	92
5.4.2	Flux scale	93
5.4.3	Sunyaev-Zeldovich decrement	94
5.5	Why is the integrated relic spectrum steepening?	95
5.5.1	Non stationary shock conditions	95
5.5.2	Structure in the Mach number or magnetic field distribution	95
5.5.3	Seed population with energy distribution cut-off	96
5.5.4	Best scenarios?	98
5.6	Conclusions	98
5.7	Appendix	99
5.7.1	Areas used for measuring integrated flux densities for the interferometric spectrum	99
6	Hα luminosity function in merging clusters	101
6.1	Introduction	102
6.2	Observations & Data Reduction	104
6.2.1	Isaac Newton Telescope observations	104
6.2.2	Reduction	105
6.2.3	Photometry	105
6.2.4	Source extraction	108

6.2.5	Galactic dust extinction correction	108
6.2.6	Narrow band excess selection	109
6.3	Results	113
6.3.1	Selecting H α emitters	113
6.3.2	Removing [NII] contamination	116
6.3.3	H α luminosity	116
6.3.4	Completeness correction	116
6.3.5	Volume	119
6.3.6	Filter profile correction	119
6.3.7	Survey limits	119
6.3.8	H α luminosity function	121
6.4	Discussion	123
6.5	Conclusion	127
6.6	Appendix	128
6.6.1	H α fractions	128
7	The rise and fall of star-formation in $z \sim 0.2$ merging galaxy clusters	131
7.1	Introduction	132
7.1.1	Star-forming galaxies in clusters	132
7.1.2	Merging clusters with shocks	133
7.1.3	The ‘Sausage’ and ‘Toothbrush’ clusters	133
7.1.4	This paper	134
7.2	Data Reduction & Analysis	137
7.2.1	Observations & data processing	137
7.2.2	Spectroscopy	139
7.2.3	Photometric reduction and source extraction	140
7.3	Methods and Results	142
7.3.1	Narrow band emitter selection	142
7.3.2	Identifying H α emitters among the line emitters	143
7.3.3	Removing [NII] contamination	150
7.3.4	H α luminosity	150
7.3.5	Completeness correction	150
7.3.6	Volume and filter profile corrections	151
7.3.7	Survey limits	151
7.3.8	H α luminosity function and star formation rate	151
7.3.9	Stellar masses	153
7.3.10	The star formation rate density	153
7.4	Discussion	154
7.4.1	H α luminosity function for the clusters	154
7.4.2	Star formation rate density for the two clusters	161
7.4.3	Stellar masses of cluster galaxies	162
7.4.4	Cause for enhanced star formation in the ‘Sausage’ cluster	164
7.5	Conclusions	167
7.6	Appendix	169
7.6.1	Dust extinction screens	169

8	Boosted AGN and star-formation activity in CIZA J2242.8+5301	173
8.1	Introduction	174
8.2	Sample Selection, Observations & Data Reduction	176
8.2.1	The “Sausage” Merging Cluster	176
8.2.2	Narrow-band survey and the sample of $H\alpha$ candidates	176
8.2.3	Follow-up spectroscopy with Keck and WHT	178
8.2.4	Redshifts and Emission line Measurements	181
8.2.5	Multiband photometry and stellar masses	184
8.2.6	Completeness: Stellar Mass	185
8.2.7	Completeness: SFR-Stellar Mass	185
8.2.8	The comparison sample: DEIMOS+AF2 non-cluster $H\alpha$ emitters	185
8.3	Results	185
8.3.1	Nature of $H\alpha$ emitters: SF vs AGN	189
8.3.2	Morphologies	190
8.3.3	Electron densities and Ionisation Potential	191
8.3.4	Outflows	195
8.3.5	Metallicities	196
8.3.6	The Mass-Metallicity relation for the Sausage merging cluster	198
8.4	Discussion: Shock induced star-formation, cooling or turbulence?	199
8.5	Conclusions	201
9	Neutral hydrogen in merging galaxy cluster	203
9.1	Introduction	204
9.2	Observations & Data Reduction	206
9.2.1	HI data	206
9.2.2	Optical imaging and spectroscopy	209
9.2.3	Broad-band radio data	209
9.3	Results	211
9.3.1	Direct detections	211
9.3.2	HI stacking	212
9.3.3	Correcting for the over-subtracted continuum emission	215
9.3.4	Measuring the HI signal and its significance	216
9.3.5	HI masses	217
9.3.6	$H\alpha$ - HI correlation	220
9.3.7	$H\alpha$ - radio correlation	221
9.4	Discussion	223
9.4.1	HI & $H\alpha$ - tracing the gas that fuels future star formation episodes	223
9.4.2	Radio broad data - tracing the SN emission	225
9.4.3	Relationship to cluster merger state and shocks	228
9.5	Conclusions	228
9.6	Appendix	230
9.6.1	Radio fluxes and morphologies	230

10 A large narrow band Hα survey at $z \sim 0.2$	233
10.1 Introduction	234
10.2 Observations & Data Reduction	235
10.2.1 Narrow band H α observations	236
10.2.2 Narrow band data reduction	236
10.2.3 Optical and IR data	237
10.3 Methods and selecting the H α samples	238
10.3.1 Star removal	238
10.3.2 Selection of line emitters	241
10.3.3 Selection of H α candidates	243
10.4 H α luminosity function and star-formation rate density	245
10.4.1 Completeness, volume and filter profile corrections	245
10.4.2 Survey limits	245
10.4.3 H α luminosity function	247
10.4.4 Star formation rate density	248
10.4.5 Distribution of H α emitters	251
10.4.6 Quantifying cosmic variance	251
10.5 Clustering of H α emitters	254
10.6 Conclusions	259
10.7 Appendix	261
10.7.1 Survey completeness	261
10.7.2 Survey completeness	261
Bibliography	265
Bibliography	265
Samenvatting	281
Rezumat	285
Publications	289
Curriculum Vitae	291
Acknowledgements	293

