



Universiteit
Leiden
The Netherlands

WRKY transcription factors involved in salicylic acid-induced defense gene expression

Verk, M.C. van

Citation

Verk, M. C. van. (2010, June 15). *WRKY transcription factors involved in salicylic acid-induced defense gene expression*. Retrieved from <https://hdl.handle.net/1887/15688>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/15688>

Note: To cite this publication please use the final published version (if applicable).

Bibliography

8



BIBLIOGRAPHY

A

- Aarts, N., Metz, M., Holub, E., Staskawicz, B. J., Daniels, M. J. and Parker, J. E. (1998). Different requirements for *EDS1* and *NDR1* by disease resistance genes define at least two *R* gene-mediated signaling pathways in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America* 95, 10306–10311.
- Akiyama, K., Chikayama, E., Yuasa, H., Shimada, Y., Tohge, T., Shinozaki, K., Hirai, M.Y., Sakurai, T., Kikuchi, J., and Saito, K. (2008). PRIME: A Web site that assembles tools for metabolomics and transcriptomics. *In Silico Biol.* 8, 0027.
- Allen, M. D., Yamasaki, K., Ohme-Takagi, M., Tateno, M. and Suzuki, M. (1998). A novel mode of DNA recognition by a β -sheet revealed by the solution structure of the GCC-box binding domain in complex with DNA. *The EMBO Journal* 17, 5484–5496.
- Alvarez-Venegas, R., Al Abdallat, A., Guo, M., Alfano, J. R. and Avramova, Z. (2007). Epigenetic control of a transcription factor at the cross section of two antagonistic pathways. *Epigenetics* 2, 106–113.
- Anand, A., Uppalapati, S. R., Ryu, C. M., Allen, S. N., Kang, L., Tang, Y. and Mysore, K. S. (2008). Salicylic acid and systemic acquired resistance play a role in attenuating crown gall disease caused by *Agrobacterium tumefaciens*. *Plant Physiology* 146, 703–715.
- Andreasson, E., Jenkins, T., Brodersen, P., Thorgrimsen, S., Petersen, N. H. T., Zhu, S., Qiu, J.-L., Micheelsen, P., Rocher, A., Petersen, M., Newman, M.-A., Nielsen, H. B. et al. (2005). The MAP kinase substrate MKS1 is a regulator of plant defense responses. *The EMBO Journal* 24, 2579–2589.
- Aoki, K., Ogata, Y. and Shibata, D. (2007). Approaches for Extracting Practical Information from Gene Co-expression Networks in Plant Biology. *Plant Cell Physiol.* 48, 381–390.
- Asai, T., Tena, G., Plotnikova, J., Willmann, M. R., Chiu, W.-L., Gomez-Gomez, L., Boller, T., Ausubel, F. M. and Sheen, J. (2002). MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature* 415, 977–983.
- Attaran, E., Zeier, T. E., Griebel, T. and Zeier, J. (2009). Methyl salicylate production and jasmonate signaling are not essential for systemic acquired resistance in *Arabidopsis*. *The Plant Cell* 21, 954–971.
- Axelos, M., Curie, C., Mazzolini, L., Bardet, C. and Lescure, B. (1992). A protocol for transient gene expression in *Arabidopsis thaliana* protoplasts isolated from cell suspension cultures. *Plant Physiol. Biochem.* 30, 123–128.

B

- Bastow, R., Mylne, J. S., Lister, C., Lippman, Z., Martienssen, R. A. and Dean, C. (2004). Vernalization requires epigenetic silencing of *FLC* by histone methylation. *Nature* 427, 164–167.
- Benfey, P. N., Ren, L. and Chua, N. H. (1990). Tissue-specific expression from CaMV 35S enhancer subdomains in early stages of plant development. *The EMBO Journal* 9, 1677–1684.
- Berger, S. L. (2002). Histone modifications in transcriptional regulation. *Current Opinion in Genetics & Development* 12, 142–148.
- Binder, B. M., Walker, J. M., Gagne, J. M., Emborg, T. J., Hemmann, G., Bleecker, A. B. and Vierstra, R. D. (2007). The *Arabidopsis* EIN3 binding F-box proteins EBF1 and EBF2 have distinct but overlapping roles in ethylene signaling. *The Plant Cell* 19, 509–523.
- Birnbaum, K., Shasha, D. E., Wang, J. Y., Jung, J. W., Lambert, G. M., Galbraith, D. W. and Benfey, P. N. (2003). A gene expression map of the *Arabidopsis* root. *Science* 302, 1956–1960.
- Blanco, F., Salinas, P., Cecchini, N. M., Jordana, X., Van Hummelen, P., Alvarez, M. E. and Holuigue, L. (2009). Early genomic responses to salicylic acid in *Arabidopsis*. *Plant Molecular Biology* 70, 79–102.
- Bleecker, A. B., Estelle, M. A., Somerville, C. and Kende, H. (1988). Insensitivity to ethylene conferred by a dominant mutation in *Arabidopsis thaliana*. *Science* 241, 1086–1089.
- Bol, J. F., Linthorst, H. J. M. and Cornelissen, B. J. C. (1990). Plant pathogenesis related proteins induced by virus infection. *Annual Rev. Phytopathol* 28, 113–138.
- Boller, T. (2005). Peptide signalling in plant development and self/non-self perception. *Current Opinion Cell Biol.* 17, 116–122.
- Borate, B. R., Chesler, E. J., Langston, M. A., Saxton, A. M. and Voy, B. H. (2009). Comparison of threshold selection methods for microarray gene co-expression matrices. *BMC Res. Notes* 2, 240.
- Borevitz, J. O., Xia, Y., Blount, J., Dixon, R. A. and Lamb, C. (2000). Activation tagging identifies a conserved MYB regulator of phenylpropanoid biosynthesis. *The Plant Cell* 12, 2383–2393.
- Boter, M., Ruíz-Rivero, O., Abdeen, A. and Prat, S. (2004). Conserved MYC transcription factors play a key role in jasmonate signaling both in tomato and *Arabidopsis*. *Genes & Development* 18, 1577–1591.
- Brazma, A., Parkinson, H., Sarkans, U., Shojatalab, M., Vilo, J., Abeygunawardena, N., Holloway, E., Kapushesky, M., Kemmeren, P., Lara, G. G., Oezcimen, A., Rocca-Serra, P. and Sansone, S. A. (2003). ArrayExpress - a public repository for microarray gene expression data at the EBI. *Nucleic Acids Res.* 31, 68–71.
- Brederode, F. T., Linthorst, H. J. M. and Bol, J. F. (1991). Differential induction of acquired resistance and PR gene expression in tobacco by virus infection, ethephon treatment, UV light and wounding. *Plant Molecular Biology* 17, 1117–1125.
- Brodersen, P., Petersen, M., Bjørn Nielsen, H., Zhu, S., Newman, M.-A., Shokat, K. M., Rietz,

- S., Parker, J. and Mundy, J. (2006). Arabidopsis MAP kinase 4 regulates salicylic acid- and jasmonic acid/ethylene-dependent responses via EDS1 and PAD4. *The Plant Journal* 47, 532–546.
- Buchel, A. S., Molenkamp, R., Bol, J. F. and Linthorst, H. J. M. (1996). The PR-1a promoter contains a number of elements that bind GT-1-like nuclear factors with different affinity. *Plant Molecular Biology* 30, 493–504.
- Buchel, A. S., Brederode, F. T., Bol, J. F. and Linthorst, H. J. (1999). Mutation of GT-1 binding sites in the *PR-1a* promoter influences the level of inducible gene expression in vivo. *Plant Molecular Biology* 40, 387–396.
- Buck, M. J. and Atchley, W. R. (2003). Phylogenetic analysis of plant basic helix-loop-helix proteins. *Journal of Molecular Evolution* 56, 742–750.
- Bülöw, L., Schindler, M. and Hehl, R. (2007). PathoPlant: a platform for microarray expression data to analyze co-regulated genes involved in plant defense responses. *Nucleic Acids Res.* 34, 841–845.
- Butterbrodt, T., Thurow, C. and Gatz, C. (2006). Chromatin immunoprecipitation analysis of the tobacco *PR-1a*- and the truncated *CaMV 35S* promoter reveals differences in salicylic acid-dependent TGA factor binding and histone acetylation. *Plant Molecular Biology* 61, 665–674.
- Büttner, M. and Singh, K. B. (1997). *Arabidopsis thaliana* ethylene-responsive element binding protein (AtEBP), an ethylene-inducible, GCC box DNA-binding protein interacts with an ocs element binding protein. *Proceedings of the National Academy of Sciences of the United States of America* 94, 5961–5966.

C

- Cancel, J. D. and Larsen, P. B. (2002). Loss-of-function mutations in the ethylene receptor *ETR1* cause enhanced sensitivity and exaggerated response to ethylene in Arabidopsis. *Plant Physiology* 129, 1557–1567.
- Cao, H., Glazebrook, J., Clarke, J. D., Volko, S. and Dong, X. (1997). The Arabidopsis *NPR1* gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. *Cell* 88, 57–63.
- Celenza, J. L., Quiel, J. A., Smolen, G. A., Merrih, H., Silvestro, A. R., Normanly, J. and Bender, J. (2005). The Arabidopsis ATR1 Myb transcription factor controls indolic glucosinolate homeostasis. *Plant Physiology* 137, 253–262.
- Chakravarthy, S., Tuori, R. P., D’Ascenzo, M. D., Fobert, P. R., Després, C. and Martin, G. B. (2003). The tomato transcription factor Pti4 regulates defense-related gene expression via GCC box and non-GCC box *cis* elements. *The Plant Cell* 15, 3033–3050.
- Chang, C., Kwok, S. F., Bleecker, A. B. and Meyerowitz, E. M. (1993). *Arabidopsis* ethylene-response gene *ETR1*: Similarity of product to two-component regulators. *Science* 262, 539–544.
- Chang, C. (2003). Ethylene signaling: the MAPK module has finally landed. *Trends in Plant Science* 8, 365–368.
- Chang, S. and Pikaard, C. S. (2005). Transcript profiling in *Arabidopsis* reveals complex-re-

sponses to global inhibition of DNA methylation and histone deacetylation. *Journal of Biological Chemistry* 280, 796–804.

Chen, C. and Chen, Z. (2000). Isolation and characterization of two pathogen- and salicylic acid-induced genes encoding WRKY DNA-binding proteins from tobacco. *Plant Molecular Biology* 42, 387–396.

Chen, C. and Chen, Z. (2002). Potentiation of developmentally regulated plant defense response by AtWRKY18, a pathogen-induced Arabidopsis transcription factor. *Plant Physiology* 129, 706–716.

Cheong, Y. H., Moon, B. C., Kim, J. K., Kim, C. Y., Kim, M. C., Kim, I. H., Park, C. Y., Kim, J. C., Park, B. O., Koo, S. C., Yoon, H. W., Chung, W. S. et al. (2003). BWMK1, a rice mitogen-activated protein kinase, locates in the nucleus and mediates pathogenesis-related gene expression by activation of a transcription factor. *Plant Physiology* 132, 1961–1972.

Chinchilla, D., Zipfel, C., Robatzek, S., Kemmerling, B., Nürnberger, T., Jones, J. D. G., Felix, G. and Boller, T. (2007). A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature* 448, 497–500.

Chini, A., Fonseca, S., Fernández, G., Adie, B., Chico, J. M., Lorenzo, O., García-Casado, G., López-Vidriero, I., Lozano, F. M., Ponce, M. R., Micol, J. L. and Solano, R. (2007). The JAZ family of repressors is the missing link in jasmonate signalling. *Nature* 448, 666–671.

Ciolkowski, I., Wanke, D., Birkenbihl, R. P. and Somssich, I. E. (2008). Studies on DNA-binding selectivity of WRKY transcription factors lend structural clues into WRKY-domain function. *Plant Molecular Biology* 68, 81–92.

Christians, M. J., Gingerich, D. J., Hansen, M., Binder, B. M., Kieber, J. J. and Vierstra, R. D. (2009). The BTB ubiquitin ligases ETO1, EOL1, and EOL2 act collectively to regulate ethylene biosynthesis in Arabidopsis by controlling type-2 ACC synthase levels. *The Plant Journal* 57, 332–345.

Ciftci-Yilmaz, S., Morsy, M. R., Song, L., Coutu, A., Krizek, B. A., Lewis, M. W., Warren, D., Cushman, J., Connolly, E. L. and Mittler, R. (2007). The EAR-motif of the Cys2/His2-type zinc finger protein Zat7 plays a key role in the defense response of Arabidopsis to salinity stress. *Journal of Biological Chemistry* 282, 9260–9268.

Clapier, C. R. and Cairns, B. R. (2009). The biology of chromatin remodeling complexes. *Annual Rev. of Biochemistry* 78, 273–304.

Clark, K. L., Larsen, P. B., Wang, X. and Chang, C. (1998). Association of the *Arabidopsis* CTR1 Raf-like kinase with the ETR1 and ERS1 ethylene receptors. *Proceedings of the National Academy of Sciences of the United States of America* 95, 5401–5406.

Conrath, U., Beckers, G. J. M., Flors, V., García-Agustín, P., Jakab, G., Mauch, F., Newman, M.-A., Pieterse, C. M. J., Poinssot, B., Pozo, M. J., Pugin, A., Schaffrath, U. et al. (2006). Priming: getting ready for battle. *Molecular Plant-Microbe Interactions* 19, 1062–1071.

Cormack, R. S., Eulgem, T., Rushton, P. J., Köchner, P., Hahlbrock, K. and Somssich, I. E. (2002). Leucine zipper containing WRKY proteins widen the spectrum of immediate early elicitor-induced WRKY transcription factors in parsley. *Biochim Biophys Acta* 1576, 92–100.

- Cosma, M. P. (2002). Ordered recruitment: Gene-specific mechanism of transcription activation. *Molecular Cell* 10, 227–236.
- Crooks, G. E., Hon, G., Chandonia, J. M. and Brenner, S.E. (2004). WebLogo: A sequence logo generator. *Genome Research* 14, 1188–1190.
- Czechowski, T., Bari, R. P., Stitt, M., Scheible, W. and Udvardi, M. K. (2004). Real-time RT-PCR profiling of over 1400 *Arabidopsis* transcription factors: unprecedented sensitivity reveals novel root- and shoot-specific gene. *The Plant Journal* 38, 366–379.

D

- Daniel, X., Lacomme, C., Morel, J.-B. and Roby, D. (1999). A novel *myb* oncogene homologue in *Arabidopsis thaliana* related to hypersensitive cell death. *The Plant Journal* 20, 57–66.
- Delaney, T. P., Friedrich, L. and Ryals, J. A. (1995). *Arabidopsis* signal transduction mutant defective in chemically and biologically induced disease resistance. *Proceedings of the National Academy of Sciences of the United States of America* 92, 6602–6606.
- Dempsey, D. M. A., Shah, J. and Klessig, D. F. (1999). Salicylic acid and disease resistance in plants. *Crit. Rev. Plant Sci.* 18, 547–575.
- Deppmann, C. D., Alvania, R. S. and Taparowsky, E. J. (2006). Cross-species annotation of basic leucine zipper factor interactions: Insight into the evolution of closed interaction networks. *Molecular Biol. Evol.* 23, 1480–1492.
- Deslandes, L., Olivier, J., Theulières, F., Hirsch, J., Feng, D. X., Bittner-Eddy, P., Beynon, J. and Marco, Y. (2002). Resistance to *Ralstonia solanacearum* in *Arabidopsis thaliana* is conferred by the recessive *RRS1-R* gene, a member of a novel family of resistance genes. *Proceedings of the National Academy of Sciences of the United States of America* 99, 2404–2409.
- Deslandes, L., Olivier, J., Peeters, N., Feng, D. X., Khounloatham, M., Boucher, C., Somssich, I., Genin, S. and Marco, Y. (2003). Physical interaction between RRS1-R, a protein conferring resistance to bacterial wilt, and PopP2, a type III effector targeted to the plant nucleus. *Proceedings of the National Academy of Sciences of the United States of America* 100, 8024–8029.
- Després, C., De Long, C., Glaze, S., Liu, E. and Fobert, P. R. (2000). The *Arabidopsis* NPR1/NIM1 protein enhances the DNA binding activity of a subgroup of the TGA family of bZIP transcription factors. *The Plant Cell* 12, 279–290.
- Després, C., Chubak, C., Rochon, A., Clark, R., Bethune, T., Desveaux, D. and Fobert, P. R. (2003). The *Arabidopsis* NPR1 disease resistance protein is a novel cofactor that confers redox regulation of DNA binding activity to the basic domain/leucine zipper transcription factor TGA1. *The Plant Cell* 15, 2181–2191.
- De Pater, S., Greco, V., Pham, K., Memelink, J. and Kijne, J. (1996). Characterization of a zinc-dependent transcriptional activator from *Arabidopsis*. *Nucleic Acids Res* 24, 4624–4631.
- De Vos, M., Denekamp, M., Dicke, M., Vuylsteke, M., Van Loon, L. C., Smeekens, S. C. M. and Pieterse, C. M. J. (2006). The *Arabidopsis thaliana* transcription factor AtMYB102 functions in defense against the insect herbivore *Pieris rapae*. *Plant Signaling & Behavior* 1, 305–311.

- Devoto, A., Nieto-Rostro, M., Xie, D., Ellis, C., Harmston, R., Patrick, E., Davis, J., Sherratt, L., Coleman, M. and Turner, J. G. (2002). COI1 links jasmonate signalling and fertility to the SCF ubiquitin-ligase complex in *Arabidopsis*. *The Plant Journal* 32, 457–466.
- Djamei, A., Pitzschke, A., Nakagami, H., Rajh, I. and Hirt, H. (2007). Trojan horse strategy in *Agrobacterium* transformation: abusing MAPK defense signaling. *Science* 318, 453–456.
- Dombrecht, B., Xue, G. P., Sprague, S. J., Kirkegaard, J. A., Ross, J. J., Reid, J. B., Fitt, G. P., Sewelam, N., Schenk, P. M., Manners, J. M. and Kazana, K. (2007). MYC2 differentially modulates diverse jasmonate-dependent functions in *Arabidopsis*. *The Plant Cell* 19, 2225–2245.
- Dong, J., Chen, C. and Chen, Z. (2003). Expression profile of the *Arabidopsis* WRKY gene superfamily during plant defense response. *Plant Molecular Biology* 51, 21–37.
- Dong, X. (1998). SA, JA, ethylene, and disease resistance in plants. *Current Opinion in Plant Biology* 1, 316–323.
- Du, L., Ali, G. S., Simons, K. A., Hou, J., Yang, T., Reddy, A. S. N. and Poovaiah, B. W. (2009). Ca^{2+} /calmodulin regulates salicylic-acid-mediated plant immunity. *Nature* 457, 1154–1158.
- Duan, M. R., Nan, J., Liang, Y. H., Mao, P., Lu, L., Li, L., Wei, C., Lai, L., Li, Y. and Su, X. D. (2007). DNA binding mechanism revealed by high resolution crystal structure of *Arabidopsis thaliana* WRKY1 protein. *Nucleic Acids Res.* 35, 1145–1154.
- Durrant, W. D. and Dong, X. (2004). Systemic acquired resistance. *Annual Rev. Phytopathol.* 42, 185–209.

E

- Edgar, R., Domrachev, M. and Lash, A. E. (2002). Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res.* 30, 207–210.
- Ellis, C., Karafyllidis, I., Wasternack, C. and Turner, J. G. (2002). The *Arabidopsis* mutant *cev1* links cell wall signaling to jasmonate and ethylene responses. *The Plant Cell* 14, 1557–1566.
- Elo, A., Lyznik, A., Gonzalez, D. O., Kachman, S. D. and Mackenzie, S. A. (2003). Nuclear genes that encode mitochondrial proteins for DNA and RNA metabolism are clustered in the *Arabidopsis* genome. *The Plant Cell* 15, 1619–1631.
- Eulgem, T. (2005). Regulation of the *Arabidopsis* defense transcriptome. *Trends in Plant Science* 10, 71–78.
- Eulgem, T., Rushton, P. J., Schmelzer, E., Hahlbrock, K. and Somssich, I. E. (1999). Early nuclear events in plant defense signaling: rapid gene activation by WRKY transcription factors. *The EMBO Journal* 18, 4689–4699.
- Eulgem, T., Rushton, P. J., Robatzek, S. and Somssich, I. E. (2000). The WRKY superfamily of plant transcription factors. *Trends in Plant Science* 5, 199–206.
- Eulgem, T. and Somssich, I. E. (2007). Networks of WRKY transcription factors in defense signaling. *Current Opinion in Plant Biology* 10, 366–371.

F

- Fan, W. and Dong, X. (2002). In vivo interaction between NPR1 and transcription factor TGA2 leads to salicylic acid-mediated gene activation in *Arabidopsis*. *The Plant Cell* 14, 1377–1389.
- Felix, G., Duran, J. D., Volko, S. and Boller, T. (1999). Plants have a sensitive perception system for the most conserved domain of bacterial flagellin. *The Plant Journal* 18, 265–276.
- Feng, S., Ma, L., Wang, X., Xie, D., Dinesh-Kumar, S. P., Wei, N. and Deng, X. W. (2003). The COP9 signalosome interacts physically with SCF^{COI1} and modulates jasmonate responses. *The Plant Cell* 15, 1083–1094.
- Feys, B. J. F., Benedetti, C. E., Penfold, C. N. and Turner, J. G. (1994). *Arabidopsis* mutants selected for resistance to the phytotoxin coronatine are male sterile, insensitive to methyl jasmonate, and resistant to a bacterial pathogen. *The Plant Cell* 6, 751–759.
- Feys, B. J., Moisan, L. J., Newman, M.-A. and Parker, J. E. (2001). Direct interaction between the *Arabidopsis* disease resistance signaling proteins, EDS1 and PAD4. *The EMBO Journal* 20, 5400–5411.
- Fode, B., Siensen, T., Thurow, C., Weigel, R. and Gatz, C. (2008). The *Arabidopsis* GRAS Protein SCL14 Interacts with Class II TGA Transcription Factors and Is Essential for the Activation of Stress-Inducible Promoters. *The Plant Cell* 20, 3122–3135.
- Fujimoto, S. Y., Ohta, M., Usui, A., Shinshi, H. and Ohme-Takagi, M. (2000). *Arabidopsis* ethylene-responsive element binding factors act as transcriptional activators or repressors of GCC box-mediated gene expression. *The Plant Cell* 12, 393–404.

G

- Gaille, C., Kast, P. and Haas, D. (2002). Salicylate biosynthesis in *Pseudomonas aeruginosa*. Purification and characterization of PchB, a novel bifunctional enzyme displaying isochorismate pyruvate-lyase and chorismate mutase activities. *Journal of Biological Chemistry* 277, 21768–21775.
- Gao, Z., Chen, Y.-F., Randlett, M. D., Zhao, X.-C., Findell, J. L., Kieber, J. J. and Schaller, G. E. (2003). Localization of the Raf-like kinase CTR1 to the endoplasmic reticulum of *Arabidopsis* through participation in ethylene receptor signaling complexes. *Journal of Biological Chemistry* 278, 34725–34732.
- Glazebrook, J. (2005). Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annual Review of Phytopathology* 43, 205–227.
- Gómez-Gómez, L. (2004). Plant perception systems for pathogen recognition and defence. *Mol. Immunol.* 41, 1055–1062.
- Green, P. J., Kay, S. A., Lam, E. and Chua, N. H. (1989). In vitro DNA footprinting. In SB Gelvin, RA Schilperoort, eds, *Plant Molecular Biology Manual B11*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 1–22.

- Grüner, R. and Pfitzner, U. M. (1994). The upstream region of the gene for the pathogenesis-related protein 1a from tobacco responds to environmental as well as to developmental signals in transgenic plants. *European Journal of Biochemistry* 220, 247–255.
- Grüner, R., Strompen, G., Pfitzner, A. J. P. and Pfitzner, U. M. (2003). Salicylic acid and the hypersensitive response initiate distinct signal transduction pathways in tobacco that converge on the *as-1*-like element of the *PR-1a* promoter. *European Journal of Biochemistry* 270, 4876–4886.
- Gu, Y.-Q., Yang, C., Thara, V. K., Zhou, J. and Martin, G. B. (2000). *Pti4* is induced by ethylene and salicylic acid, and its product is phosphorylated by the Pto kinase. *The Plant Cell* 12, 771–785.
- Guan, K. L. and Dixon, J. E. (1991). Eukaryotic proteins expressed in *Escherichia coli*: an improved thrombin cleavage and purification procedure of fusion proteins with glutathione S-transferase. *Anal. Biochem.* 192, 262–267.
- Guo, H. and Ecker, J. R. (2003). Plant responses to ethylene gas are mediated by SCF^{EBF1/EBF2}-dependent proteolysis of EIN3 transcription factor. *Cell* 115, 667–677.
- Guo, A., He, K., Liu, D., Bai, S., Gu, X., Wei, L. and Luo, J. (2005). DATF: a database of *Arabidopsis* transcription factors. *Bioinformatics* 21, 2568–2569.
- Guzmán, P. and Ecker, J. R. (1990). Exploiting the triple response of *Arabidopsis* to identify ethylene-related mutants. *The Plant Cell* 2, 513–523.

H

- He, Y. and Amasino, R. M. (2005). Role of chromatin modification in flowering-time control. *Trends in Plant Science* 10, 30–35.
- He, Y., Doyle, M. R. and Amasino, R. M. (2004). PAF1-complex-mediated histone methylation of FLOWERING LOCUS C chromatin is required for the vernalization-responsive, winter-annual habit in *Arabidopsis*. *Genes & Development* 18, 2774–2784.
- He, P., Shan, L., Lin, N.-C., Martin, G. B., Kemmerling, B., Nürnberger, T. and Sheen, J. (2006). Specific bacterial suppressors of PAMP signaling upstream of MAPKKK in *Arabidopsis* innate immunity. *Cell* 125, 563–575.
- He, P., Shan, L. and Sheen, J. (2007). The use of protoplasts to study innate immune responses. *Methods Mol. Biol.* 354, 1–9.
- Higashi, K., Ishiga, Y., Inagaki, Y., Toyoda, K., Shiraishi, T. and Ichinose, Y. (2008). Modulation of defense signal transduction by flagellin-induced WRKY41 transcription factor in *Arabidopsis thaliana*. *Molecular Genetics and Genomics* 279, 303–312.
- Hirai, M. Y., Sugiyama, K., Sawada, Y., Tohge, T., Obayashi, T., Suzuki, A., Araki, R., Sakurai, N., Suzuki, H., Aoki, K., Goda, H., Nishizawa, O. I. et al. (2007). Omics-based identification of *Arabidopsis* Myb transcription factors regulating aliphatic glucosinolate biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* 104, 6478–6483.
- Howe, G. A. (2004). Jasmonates as signals in the wound response. *Journal of Plant Growth Regulation* 23, 223–237.

- Hruz, T., Laule, O., Szabo, G., Wessendrop, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W. and Zimmermann, P. (2008). Genevestigator V3: A reference expression database for the meta-analysis of transcriptomes. *Advances in Bioinformatics* 2008, Article ID 420747.
- Hua, J., Chang, C., Sun, Q. and Meyerowitz, E. M. (1995). Ethylene insensitivity conferred by *Arabidopsis* *ERS* gene. *Science* 269, 1712–1714.
- Hua, J., Sakai, H., Nourizadeh, S., Chen, Q. G., Bleecker, A. B., Ecker, J. R. and Meyerowitz, E. M. (1998). *EIN4* and *ERS2* are members of the putative ethylene receptor gene family in *Arabidopsis*. *The Plant Cell* 10, 1321–1332.
- Hunt, M. D., Nuenschwander, U. H., Delaney, T. P., Weymann, K. B., Friedrich, L. B., Lawton, K. A., Steiner, H. Y. and Ryals, J. A. (1996). Recent advances in systemic acquired resistance research – a review. *Gene* 179, 89–95.

I

- Ichimura, K., Mizoguchi, T., Yoshida, R., Yuasa, T. and Shinozaki, K. (2000). Various abiotic stresses rapidly activate *Arabidopsis* MAP kinases ATMPK4 and ATMPK6. *The Plant Journal* 24, 655–665.
- Ishiguro, S. and Nakamura, K. (1994). Characterization of a cDNA encoding a novel DNA-binding protein, SPF1, that recognizes SP8 sequences in the 5' upstream regions of genes coding for sporamin and β -amylase from sweet potato. *Molecular and General Genetics* 244, 563–571.
- Ishihara, T., Sekine, K.-T., Hase, S., Kanayama, Y., Seo, S., Ohashi, Y., Kusano, T., Shibata, D., Shah, J. and Takahashi, H. (2008). Overexpression of the *Arabidopsis thaliana* *EDS5* gene enhances resistance to viruses. *Plant Biology* 10, 451–461.

J

- Jagadeeswaran, G., Raina, S., Acharya, B. R., Maqbool, S. B., Mosher, S. L., Appel, H. M., Schultz, J. C., Klessig, D. F. and Raina, R. (2007). *Arabidopsis* *GH3-LIKE DEFENSE GENE 1* is required for accumulation of salicylic acid, activation of defense responses and resistance to *Pseudomonas syringae*. *The Plant Journal* 51, 234–246.
- Jakoby, M., Weisshaar, B., Dröge-Laser, W., Vicente-Carbajosa, J., Tiedemann, J., Kroj, T. and Parcy, F. (2002). bZIP transcription factors in *Arabidopsis*. *Trends in Plant Science*. 7, 106–111.
- Jefferson, R. (1987). Assaying chimeric genes in plants: the GUS gene fusion system. *Plant Molecular Biology Rep* 5, 387–405.
- Jofuku, K. D., Den Boer, B. G. W., Van Montagu, M. and Okamura, J. K. (1994). Control of *Arabidopsis* flower and seed development by the homeotic gene *APETALA2*. *The Plant Cell* 6, 1211–1225.
- Johnson, C., Boden, E. and Arias, J. (2003). Salicylic acid and NPR1 induce the recruitment of *trans*-activating TGA factors to a defense gene promoter in *Arabidopsis*. *The Plant Cell* 15, 1846–1858.

- Jones, J. D. G. and Dangl, J. L. (2006). The plant immune system. *Nature* 444, 323–329.
- Journot-Catalino, N., Somssich, I. E., Roby, D. and Kroj, T. (2006). The transcription factors WRKY11 and WRKY17 act as negative regulators of basal resistance in *Arabidopsis thaliana*. *The Plant Cell* 18, 3289–3302.
- Jung, H. W., Tschaplinski, T. J., Wang, L., Glazebrook, J. and Greenberg, J. T. (2009). Priming in systemic plant immunity. *Science* 324, 89–91.

K

- Kagaya, Y., Ohmiya, K. and Hattori, T. (1999). RAV1, a novel DNA-binding protein, binds to bipartite recognition sequence through two distinct DNA-binding domains uniquely found in higher plants. *Nucleic Acids Res.* 27, 470–478.
- Kalde, M., Barth, M., Somssich, I. E. and Lippok, B. (2003). Members of the *Arabidopsis* WRKY group III transcription factors are part of different plant defense signaling pathways. *Molecular Plant-Microbe Interactions* 16, 295–305.
- Kaminaka, H., Näke, C., Epple, P., Dittgen, J., Schütze, K., Chaban, C., Holt, B. F., III, Merkle, T., Schäfer, E., Harter, K. and Dangl, J. L. (2006). bZIP10-LSD1 antagonism modulates basal defense and cell death in *Arabidopsis* following infection. *The EMBO Journal* 25, 4400–4411.
- Katagiri, F., Lam, E. and Chua, N.-H. (1989). Two tobacco DNA-binding proteins with homology to the nuclear factor CREB. *Nature* 340, 727–730.
- Kegler, C., Lenk, I., Krawczyk, S., Scholz, R. and Gatz, C. (2004). Functional characterization of tobacco transcription factor TGA2.1. *Plant Molecular Biology* 55, 153–164.
- Kesarwani, M., Yoo, J. and Dong, X. (2007). Genetic interactions of TGA transcription factors in the regulation of pathogenesis-related genes and disease resistance in *Arabidopsis*. *Plant Physiology* 144, 336–346.
- Kieber, J. J., Rothenberg, M., Roman, G., Feldman, K. A. and Ecker, J. R. (1993). *CTR1*, a negative regulator of the ethylene response pathway in *Arabidopsis*, encodes a member of the Raf family of protein kinases. *Cell* 72, 427–441.
- Kim, C. J. and Zhang, S. (2004). Activation of a mitogen-activated protein kinase cascade induces WRKY family of transcription factors and defense genes in tobacco. *The Plant Journal* 38, 142–151.
- Kim, K. C., Fan, B. and Chen, Z. (2006). Pathogen-induced *Arabidopsis* WRKY7 is a transcriptional repressor and enhances plant susceptibility to *Pseudomonas syringae*. *Plant Physiology* 142, 1180–1192.
- Kim, K.-C., Lai, Z., Fan, B. and Chen, Z. (2008). *Arabidopsis* WRKY38 and WRKY62 transcription factors interact with histone deacetylase 19 in basal defense. *The Plant Cell* 20, 2357–2371.
- Kinkema, M., Fan, W. and Dong, X. (2000). Nuclear localization of NPR1 is required for activation of *PR* gene expression. *The Plant Cell* 12, 2339–2350.

- Konishi, M. and Yanagisawa, S. (2008). Ethylene signaling in *Arabidopsis* involves feedback regulation via the elaborate control of *EBF2* expression by EIN3. *The Plant Journal* 55, 821–831.
- Koornneef, A. and Pieterse, C. M. J. (2008). Cross talk in defense signaling. *Plant Physiology* 146, 839–844.
- Koornneef, A., Rindermann, K., Gatz, C. and Pieterse, C. M. J. (2008). Histone modifications do not play a major role in salicylate-mediated suppression of jasmonate-induced *PDF1.2* gene expression. *Communicative & Integrative Biology* 1, 143–145.
- Kuhlmann, M., Horvay, K., Strathmann, A., Heinekamp, T., Fischer, U., Böttner, S. and Dröge-Laser, W. (2003). The α -helical D1 domain of the tobacco bZIP transcription factor BZI-1 interacts with the ankyrin-repeat protein ANK1 and is important for BZI-1 function, both in auxin signaling and pathogen response. *Journal of Biological Chemistry* 278, 8786–8794.
- Kunze, G., Zipfel, C., Robatzek, S., Niehaus, K., Boller, T. and Felix, G. (2004). The N terminus of bacterial elongation factor Tu elicits innate immunity in *Arabidopsis* plants. *The Plant Cell* 16, 3496–3507.
- Kuo, M. H., Vom Baur, E., Struhl, K. and Allis, C. D. (2000). Gcn4 activator targets Gcn5 histone acetyltransferase to specific promoters independently of transcription. *Molecular Cell* 6, 1309–1320.
- Kosugu, S., Ohashi, Y., Nakajima, K. and Arai, Y. (1990). An improved assay for β -glucuronidase in transformed cells: methanol almost completely suppresses a putative endogenous β -glucuronidase activity. *Plant Science* 70, 133–140.

L

- Lebel, E., Heifetz, P., Thorne, L., Uknes, S., Ryals, J. and Ward, E. (1998). Functional analysis of regulatory sequences controlling PR-1 gene expression in *Arabidopsis*. *The Plant Journal* 16, 223–233.
- Le Henanff, G., Heitz, T., Mestre, P., Mutterer, J., Walter, B. and Chong, J. (2009). Characterization of *Vitis vinifera* NPR1 homologs involved in the regulation of pathogenesis-related gene expression. *BMC Plant Biol.* 9, 54.
- Lee, H.-I., León, J. and Raskin, I. (1995). Biosynthesis and metabolism of salicylic acid. *Proceedings of the National Academy of Sciences of the United States of America* 92, 4076–4079.
- Lee, J. H., Yoo, S. J., Park, S. H., Hwang, I., Lee, J. S. and Ahn, J. H. (2007). Role of SVP in the control of flowering time by ambient temperature in *Arabidopsis*. *Genes & Dev* 21, 397–402.
- Leon-Reyes, A., Spoel, S. H., De Lange, E. S., Abe, H., Kobayashi, M., Tsuda, S., Millenaar, F. F., Welschen, R. A. M., Ritsema, T. and Pieterse, C. M. J. (2009). Ethylene modulates the role of NONEXPRESSOR OF PATHOGENESIS-RELATED GENES1 in cross talk between salicylate and jasmonate signaling. *Plant Physiology* 149, 1797–1809.
- Li, X., Zhang, Y., Clarke, J. D., Li, Y. and Dong, X. (1999). Identification and cloning of a negative regulator of systemic acquired resistance, SNI1, through a screen for suppressors of *npr1-1*. *Cell* 98, 329–339.

- Li, J., Brader, G. and Palva, E. T. (2004). The WRKY70 transcription factor: A node of convergence for jasmonate-mediated and salicylate-mediated signals in plant defense. *The Plant Cell* 16, 319–331.
- Li, J., Brader, G., Kariola, T. and Palva, E. T. (2006). WRKY70 modulates the selection of signaling pathways in plant defense. *The Plant Journal* 46, 477–491.
- Li, B., Carey, M. and Workman, J. L. (2007). The role of chromatin during transcription. *Cell* 128, 707–719.
- Linthorst, H. J. M. (1991). Pathogenesis-related proteins of plants. *Crit. Rev. Plant Sci.* 10, 123–150.
- Linthorst, H. J. M., Meuwissen, R. L. J., Kauffman, S. and Bol, J. F. (1989). Constitutive expression of pathogenesis-related proteins PR-1, GRP, and PR-S in tobacco has no effect on virus infection. *The Plant Cell* 1, 285–291.
- Liu, Y. and Zhang, S. (2004). Phosphorylation of 1-aminocyclopropane-1-carboxylic acid synthase by MPK6, a stress-responsive mitogen-activated protein kinase, induces ethylene biosynthesis in *Arabidopsis*. *The Plant Cell* 16, 3386–3399.
- Liu, Y. L., Schiff, M., Marathe, R. and Dinesh-Kumar, S. P. (2002). Tobacco Rar1, EDS1 and NPR1/NIM1 like genes are required for N-mediated resistance to tobacco mosaic virus. *The Plant Journal* 30, 415–429.
- Liu, Y., Schiff, M. and Dinesh-Kumar, S. P. (2004). Involvement of MEK1 MAPKK, NTF6 MAPK, WRKY/MYB transcription factors, COI1 and CTR1 in N-mediated resistance to tobacco mosaic virus. *The Plant Journal* 38, 800–809.
- Lorenzo, O. and Solano, R. (2005). Molecular players regulating the jasmonate signalling network. *Current Opinion in Plant Biology* 8, 532–540.
- Lorenzo, O., Piqueras, R., Sánchez-Serrano, J. J. and Solano, R. (2003). ETHYLENE RESPONSE FACTOR1 integrates signals from ethylene and jasmonate pathways in plant defense. *The Plant Cell* 15, 165–178.
- Lorenzo, O., Chico, J. M., Sánchez-Serrano, J. J. and Solano, R. (2004). *JASMONATE-INSENSITIVE1* encodes a MYC transcription factor essential to discriminate between different jasmonate-regulated defense responses in *Arabidopsis*. *The Plant Cell* 16, 1938–1950.

M

- Ma, S., Gong, Q. and Bohnert, H. J. (2007). An *Arabidopsis* gene network based on the graphical Gaussian model. *Genome Res.* 17, 1614–1625.
- Mackey, D. and McFall, A. J. (2006). MAMPs and MIMPs: proposed classifications for inducers of innate immunity. *Molecular Microbiology* 61, 1365–1371.
- Maeo, K., Hayashi, S., Kojima-Suzuki, H., Morikami, A. and Nakamura, K. (2001). Role of conserved residues of the WRKY domain in the DNA-binding of tobacco WRKY family proteins. *Biosci. Biotechnol Biochem.* 65, 2428–2436.
- Malamy, J., Carr, J. P., Klessig, D. F. and Raskin, I. (1990). Salicylic acid: a likely endogenous signal

- in the resistance response of tobacco to viral infection. *Science* 250, 1002-1004.
- Maleck, K., Levine, A., Eulgem, T., Morgan, A., Schmid, J., Lawton, K. A., Dangel, J.L. and Dietrich, R. A. (2000). The transcriptome of *Arabidopsis thaliana* during systemic acquired resistance. *Nature Genetics* 26, 403-410.
- Manfield, I. W., Jen, C. H., Pinney, J. W., Michalopoulos, I., Bradford, J. R., Gilmartin, P. M. and Westhead, D. R. (2006). *Arabidopsis* Co-expression Tool (ACT): web server tools for microarray-based gene expression analysis. *Nucleic Acids Res.* 34, W504-509.
- Mao, P., Duan, M., Wei, C. and Li, Y. (2007). WRKY62 transcription factor acts downstream of cytosolic NPR1 and negatively regulates jasmonate-responsive gene expression. *Plant & Cell Physiology* 48, 833-842.
- Mauch-Mani, B. and Slusarenko, A. J. (1996). Production of salicylic acid precursors is a major function of phenylalanine ammonia-lyase in the resistance of *Arabidopsis* to *Peronospora parasitica*. *The Plant Cell* 8, 203-212.
- McConn, M. and Browse, J. (1996). The critical requirement for linolenic acid is pollen development, not photosynthesis, in an *Arabidopsis* mutant. *The Plant Cell* 8, 403-416.
- McGrath, K. C., Dombrecht, B., Manners, J. M., Schenk, P. M., Edgar, C. I., Maclean, D. J., Scheible, W. R., Udvardi, M. K. and Kazan, K. (2005). Repressor- and activator-type ethylene response factors functioning in jasmonate signaling and disease resistance identified via a genome-wide screen of *Arabidopsis* transcription factor gene expression. *Plant Physiology* 139, 949-959.
- Meijer, A. H., Verpoorte, R. and Hoge, J. H. C. (1993). Regulation of enzymes and genes involved in terpenoid indole alkaloid biosynthesis in *Catharanthus roseus*. *Journal of Plant Research Special Issue* 3, 145-164.
- Mellor, J. (2006). Dynamic nucleosomes and gene transcription. *Trends in Genetics* 22, 320-329.
- Mengiste, T., Chen, X., Salmeron, J. and Dietrich, R. (2003). The *BOTRYTIS SUSCEPTIBLE1* gene encodes an R2R3MYB transcription factor protein that is required for biotic and abiotic stress responses in *Arabidopsis*. *The Plant Cell* 15, 2551-2565.
- Menke, F. L. H., Champion, A., Kijne, J. W. and Memelink, J. (1999). A novel jasmonate- and elicitor-responsive element in the periwinkle secondary metabolite biosynthetic gene *Str* interacts with a jasmonate- and elicitor-inducible AP2-domain transcription factor, ORCA2. *The EMBO Journal* 18, 4455-4463.
- Mészáros, T., Helfer, A. and Bögre, L. (2007). The more we know, the less we understand? Complexity of MAP kinase signaling. *Plant Signal Behav.* 2, 30-32.
- Métraux, J. P., Signer, H., Ryals, J., Ward, E., Wyss-Benz, M., Gaudin, J., Raschdorf, K., Schmid, E., Blum, W. and Inverardi, B. (1990). Increase in Salicylic Acid at the Onset of Systemic Acquired Resistance in Cucumber. *Science* 250, 1004-1006.
- Miao, Y., Laun, T. M., Smykowski, A. and Zentgraf, U. (2007). *Arabidopsis* MEKK1 can take a short cut: it can directly interact with senescence-related WRKY53 transcription factor on the protein level and can bind to its promoter. *Plant Molecular Biology* 65, 63-76.
- Mitchell, R. E. and Young, H. (1978). Identification of a chlorosis-inducing toxin of *Pseudomonas glycinea* as coronatine. *Phytochemistry* 17, 2028-2029.

- Mosher, R. A., Durrant, W. E., Wang, D., Song, J. and Dong, X. (2006). A comprehensive structure–function analysis of *Arabidopsis* SNI1 defines essential regions and transcriptional repressor activity. *The Plant Cell* 18, 1750–1765.
- Mou, Z., Fan, W. and Dong, X. (2003). Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* 113, 935–944.
- Murre, C., Bain, G., Van Dijk, M. A., Engel, I., Furnari, B. A., Massari, M. E., Matthews, J. R., Quong, M. W., Rivera, R. R. and Stuiver, M. H. (1994). Structure and function of helix-loop-helix proteins. *Biochimica et Biophysica Acta* 1218, 129–135.

N

- Nakano, T., Suzuki, K., Fujimura, T. and Shinshi, H. (2006). Genome-wide analysis of the ERF gene family in *Arabidopsis* and rice. *Plant Physiology* 140, 411–432.
- Narlikar, G. J., Fan, H.-Y. and Kingston, R. E. (2002). Cooperation between complexes that regulate chromatin structure and transcription. *Cell* 108, 475–487.
- Navarro, L., Zipfel, C., Rowland, O., Keller, I., Robatzek, S., Boller, T. and Jones, J. D. G. (2004). The transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis. *Plant Physiology* 135, 1113–1128.
- Nawrath, C. and Métraux, J.-P. (1999). Salicylic acid induction-deficient mutants of *Arabidopsis* express *PR-2* and *PR-5* and accumulate high levels of camalexin after pathogen inoculation. *The Plant Cell* 11, 1393–1404.
- Nawrath, C., Heck, S., Parinshawong, N. and Métraux, J.-P. (2002). EDS5, an essential component of salicylic acid-dependent signaling for disease resistance in *Arabidopsis*, is a member of the MATE transporter family. *The Plant Cell* 14, 275–286.
- Ndamukong, I., Al Abdallat, A., Thurow, C., Fode, B., Zander, M., Weigel, R. and Gatz, C. (2007). SA-inducible *Arabidopsis* glutaredoxin interacts with TGA factors and suppresses JA-responsive *PDF1.2* transcription. *The Plant Journal* 50, 128–139.
- Ng, D. W.-K., Chandrasekharan, M. B. and Hall, T. C. (2006). Ordered histone modifications are associated with transcriptional poising and activation of the *phaseolin* promoter. *The Plant Cell* 18, 119–132.
- Niggeweg, R., Thurow, C., Kegler, C. and Gatz, C. (2000a). Tobacco transcription factor TGA2.2 is the main component of as-1-binding factor ASF-1 and is involved in salicylic acid- and auxin-inducible expression of as-1-containing target promoters. *Journal of Biological Chemistry* 275, 19897–19905.
- Niggeweg, R., Thurow, C., Weigel, R., Pfitzner, U. and Gatz, C. (2000b). Tobacco TGA factors differ with respect to interaction with NPR1, activation potential and DNA-binding properties. *Plant Molecular Biology* 42, 775–788.
- Nobuta, K., Okrent, R. A., Stoutemyer, M., Rodibaugh, N., Kempema, L., Wildermuth, M. C. and Innes, R. W. (2007). The GH3 acyl adenylase family member PBS3 regulates salicylic acid-dependent defense responses in *Arabidopsis*. *Plant Physiology* 144, 1144–1156.

O

- Obayashi, T., Kinoshita, K., Nakai, K., Shibaoka, M., Hayashi, S., Saeki, M., Shibata, D., Saito, K. and Ohta, H. (2007). ATTED-II: a database of co-expressed genes and cis elements for identifying co-regulated gene groups in Arabidopsis. *Nucleic Acids Res.*, 35, D863–D869.
- Obayashi, T., Hayashi, S., Shibaoka, M., Saeki, M., Ohta, H. and Kinoshita, K. (2008). COXPRESdb: a database of coexpressed gene networks in mammals. *Nucleic Acids Res.* 36, D77–82.
- Obayashi, T., Hayashi, S., Saeki, M., Ohta, H. and Kinoshita, K. (2009). ATTED-II provides coexpressed gene networks for Arabidopsis. *Nucleic Acids Res.* 37, D987–991.
- Ohme-Takagi, M. and Shinshi, H. (1995). Ethylene-inducible DNA binding proteins that interact with an ethylene-responsive element. *The Plant Cell* 7, 173–182.
- Okrent, R. A., Brooks, M. D. and Wildermuth, M. C. (2009). Arabidopsis GH3.12 (PBS3) conjugates amino acids to 4-substituted benzoates and is inhibited by salicylate. *Journal of Biological Chemistry* 284, 9742–9754.
- Olmedo, G., Guo, H., Gregory, B. D., Nourizadeh, S. D., Aguilar-Henonin, L., Li, H., An, F., Guzman, P. and Ecker, J. R. (2006). *ETHYLENE-INSENSITIVE5* encodes a 5'→3' exoribonuclease required for regulation of the EIN3-targeting F-box proteins EBF1/2. *Proceedings of the National Academy of Sciences of the United States of America* 103, 13286–13293.
- Oñate-Sánchez, L., Anderson, J. P., Young, J. and Singh, K. B. (2007). AtERF14, a member of the ERF family of transcription factors, plays a nonredundant role in plant defense. *Plant Physiology* 143, 400–409.
- Ouaked, F., Rozhon, W., Lecourieux, D. and Hirt, H. (2003). A MAPK pathway mediates ethylene signaling in plants. *The EMBO Journal* 22, 1282–1288.
- Ouwierkerk, B. F. O. and Meijer, A. H. (2001). Yeast one-hybrid screening for DNA-protein interactions. *In* F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, K. Struhl, eds, *Current Protocols in Molecular Biology*. John Wiley & Sons, Chichester, UK, pp 12.12.1–12.12.22.

P

- Pandey, S. P. and Somssich, I. E. (2009). The Role of WRKY Transcription Factors in Plant Immunity. *Plant Physiology* 150, 1648–1655.
- Park, S. W., Kaimoyo, E., Kumar, D., Mosher, S. and Klessig, D. F. (2007). Methyl salicylate is a critical mobile signal for plant systemic acquired resistance. *Science* 318, 113–116.
- Pascuzzi, P., Hamilton, D., Bodily, K. and Arias, J. (1998). Auxin-induced stress potentiates transactivation by a conserved basic/leucine-zipper factor. *Journal of Biological Chemistry* 273, 26631–26637.
- Persson, S., Wei, H., Milne, J., Page, G. P. and Somerville, C. R. (2005). Identification of genes

required for cellulose synthesis by regression analysis of public microarray data sets. *Proceedings of the National Academy of Sciences of the United States of America* 102, 8633–8638.

- Pieterse, C. M. J., Van Wees, S. C. M., Van Pelt, J. A., Knoester, M., Laan, R., Gerrits, N., Weisbeek, P. J. and Van Loon, L. C. (1998). A novel signaling pathway controlling induced systemic resistance in *Arabidopsis*. *The Plant Cell* 10, 1571–1580.
- Pontier, D., Miao, Z. H. and Lam, E. (2001). Trans-dominant suppression of plant TGA factors reveals their negative and positive roles in plant defense responses. *The Plant Journal* 27, 529–538.
- Pontier, D., Privat, I., Trifa, Y., Zhou, J. M., Klessig, D. F. and Lam, E. (2002). Differential regulation of TGA transcription factors by post-transcriptional control. *The Plant Journal* 32, 641–653.
- Potuschak, T., Lechner, E., Parmentier, Y., Yanagisawa, S., Grava, S., Koncz, C. and Genschik, P. (2003). EIN3-dependent regulation of plant ethylene hormone signaling by two *Arabidopsis* F box proteins: EBF1 and EBF2. *Cell* 115, 679–689.
- Pozo, M. J., Van der Ent, S., Van Loon, L. C. and Pieterse, C. M. J. (2008). Transcription factor MYC2 is involved in priming for enhanced defense during rhizobacteria-induced systemic resistance in *Arabidopsis thaliana*. *New Phytologist* 180, 511–523.
- Pré, M. R. (2006). “Functional Analysis of Jasmonate-Responsive AP2/ERF-Domain Transcription Factors in *Arabidopsis thaliana*”. Ph.D. thesis, University of Leiden, Leiden, The Netherlands.
- Pré, M., Atallah, M., Champion, A., De Vos, M., Pieterse, C. M. J. and Memelink, J. (2008). The AP2/ERF domain transcription factor ORA59 integrates jasmonic acid and ethylene signals in plant defense. *Plant Physiology* 147, 1347–1357.

Q

- Qin, X.-F., Holuigue, L., Horvath, D. M. and Chua, N.-H. (1994). Immediate early transcription activation by salicylic acid via the cauliflower mosaic virus *as-1* element. *The Plant Cell* 6, 863–874.
- Qiu, J.-L., Zhou, L., Yun, B.-W., Nielsen, H. B., Fiil, B. K., Petersen, K., MacKinlay, J., Loake, G. J., Mundy, J. and Morris, P. C. (2008). *Arabidopsis* mitogen-activated protein kinases MKK1 and MKK2 have overlapping functions in defense signaling mediated by MEKK1, MPK4, and MKS1. *Plant Physiology* 148, 212–222.

R

- Raffaele, S., Rivas, S. and Roby, D. (2006). An essential role for salicylic acid in AtMYB30-mediated control of the hypersensitive cell death program in *Arabidopsis*. *FEBS Letters* 580, 3498–3504.
- Rayapuram, C. and Baldwin, I. T. (2007). Increased SA in NPR1-silenced plants antagonizes JA and JA-dependent direct and indirect defenses in herbivore-attacked *Nicotiana attenuata* in na-

- ture. *The Plant Journal* 52, 700–715.
- Rea, S., Eisenhaber, F., O'Carroll, D., Strahl, B. D., Sun, Z.-W., Schmid, M., Opravil, S., Mechtler, K., Ponting, C. P., Allis, C. D. and Jenuwein, T. (2000). Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature* 406, 593–599.
- Reed, J. W. (2001). Roles and activities of Aux/IAA proteins in *Arabidopsis*. *TIPS* 6, 420–425.
- Ren, C.-M., Zhu, Q., Gao, B.-D., Ke, S.-Y., Yu, W.-C., Xie, D.-X. and Peng, W. (2008). Transcription factor WRKY70 displays important but no indispensable roles in jasmonate and salicylic acid signaling. *Journal of Integrative Plant Biology* 50, 630–637.
- Reinke, H. and Horz, W. (2003). Histones are first hyperacetylated and then lose contact with the activated *PHO5* promoter. *Molecular Cell* 11, 1599–1607.
- Reymond, P. and Farmer, E. E. (1998). Jasmonate and salicylate as global signals for defense gene expression. *Current Opinion in Plant Biology* 1, 404–411.
- Reymond, P., Bodenhausen, N., Van Poecke, R. M. P., Krishnamurthy, V., Dicke, M. and Farmer, E. E. (2004). A conserved transcript pattern in response to a specialist and a generalist herbivore. *The Plant Cell* 16, 3132–3147.
- Robatzek, S. and Somssich, I. E. (2002). Targets of AtWRKY6 regulation during plant senescence and pathogen defense. *Genes & Development* 16, 1139–1149.
- Rochon, A., Boyle, P., Wignes, T., Fobert, P. R. and Després, C. (2006). The coactivator function of *Arabidopsis* NPR1 requires the core of its BTB/POZ domain and the oxidation of C-terminal cysteines. *The Plant Cell* 18, 3670–3685.
- Rogers, E. E. and Ausubel, F. M. (1997). *Arabidopsis* enhanced disease susceptibility mutants exhibit enhanced susceptibility to several bacterial pathogens and alterations in *PR-I* gene expression. *The Plant Cell* 9, 305–316.
- Roman, G., Lubarsky, B., Kieber, J. J., Rothenberg, M. and Ecker, J. R. (1995). Genetic analysis of ethylene signal transduction in *Arabidopsis thaliana*: Five novel mutant loci integrated into a stress response pathway. *Genetics* 139, 1393–1409.
- Ross, A. F. (1961). Systemic acquired resistance induced by localized virus infections in plants. *Virology* 14, 340–358.
- Rowland, O. and Jones, J. D. G. (2001). Unraveling regulatory networks in plant defense using microarrays. *Genome Biology* 2, 1001.1–1001.3.
- Ruegger, M., Dewey, E., Gray, W. M., Hobbie, L., Turner, J. and Estelle, M. (1998). The TIR1 protein of *Arabidopsis* functions in auxin response and is related to human SKP2 and yeast Grr1p. *Genes & Development* 12, 198–207.
- Rushton, P. J., Torres, J. T., Parniske, M., Wernert, P., Hahlbrock, K. and Somssich, I. E. (1996). Interaction of elicitor-induced DNA binding proteins with elicitor response elements in the promoters of parsley PR1 genes. *The EMBO Journal* 15, 5690–5700.
- Ryals, J. A., Neuenschwander, U. H., Willits, M. G., Molina, A., Steiner, H.-Y. and Hunt, M. D. (1996). Systemic Acquired Resistance. *The Plant Cell* 8, 1809–1819.

- Sakai, H., Hua, J., Chen, Q. G., Chang, C., Medrano, L. J., Bleecker, A. B. and Meyerowitz, E. M. (1998). *ETR2* is an *ETR1*-like gene involved in ethylene signaling in *Arabidopsis*. Proceedings of the National Academy of Sciences of the United States of America 95, 5812–5817.
- Sánchez-Navarro, J., Miglino, R., Ragozzino, A. and Bol, J. F. (2001). Engineering alfalfa mosaic virus RNA 3 into an expression vector. Arch Virol 146, 923–939.
- Schindler, U., Menkens, A. E., Beckmann, H., Ecker, J. R. and Cashmore, A. R. (1992). Heterodimerization between light-regulated and ubiquitously expressed *Arabidopsis* GBF bZIP proteins. The EMBO Journal 11, 1261–1273.
- Schirawski, J., Planchais, S. and Haenni, A. L. (2000). An improved protocol for the preparation of protoplasts from an established *Arabidopsis thaliana* cell suspension culture and infection with RNA of a turnip yellow mosaic tymovirus: a simple and reliable method. Journal of Virology 86, 85–94.
- Seet, B. T., Dikic, I., Zhou, M.-M. and Pawson, T. (2006). Reading protein modifications with interaction domains. Nature Reviews Molecular Cell Biology 7, 473–483.
- Seo, S., Okamoto, M., Seto, H., Ishizuka, K., Sano, H. and Ohashi, Y. (1995). Tobacco MAP kinase: A possible mediator in wound signal transduction pathways. Science 270, 1988–1992.
- Seo, S., Sano, H. and Ohashi, Y. (1999). Jasmonate-based wound signal transduction requires activation of WIPK, a tobacco mitogen-activated protein kinase. The Plant Cell 11, 289–298.
- Seo, H. S., Song, J. T., Cheong, J.-J., Lee, Y.-H., Lee, Y.-W., Hwang, I., Lee, J. S. and Choi, Y. D. (2001). Jasmonic acid carboxyl methyltransferase: A key enzyme for jasmonate-regulated plant responses. Proceedings of the National Academy of Sciences of the United States of America 98, 4788–4793.
- Sessa, G., Meller, Y. and Fluhr, R. (1995). A GCC element and a G-box motif participate in ethylene-induced expression of the *PRB-1b* gene. Plant Molecular Biology 28, 145–153.
- Shen, Q. H., Saijo, Y., Mauch, S., Biskup, C., Bieri, S., Keller, B., Seki, H., Ülker, B., Somssich, I. E. and Schulze-Lefert, P. (2007). Nuclear activity of MLA immune receptors links isolate-specific and basal disease-resistance responses. Science 315, 1098–1103.
- Solano, R., Stepanova, A., Chao, Q. and Ecker, J. R. (1998). Nuclear events in ethylene signaling: a transcriptional cascade mediated by ETHYLENEINSENSITIVE3 and ETHYLENE-RESPONSE-FACTOR1. Genes & Development 12, 3703–3714.
- Spoel, S. H. and Dong, X. (2008). Making sense of hormone crosstalk during plant immune responses. Cell Host & Microbe 3, 348–351.
- Spoel, S. H., Mou, Z., Tada, Y., Spivey, N. W., Genschik, P. and Dong, X. (2009). Proteasome-mediated turnover of the transcription coactivator NPR1 plays dual roles in regulating plant immunity. Cell 137, 860–872.
- Srinivasasainagendra, V., Page, G. P., Mehta, T., Coulibaly, I. and Loraine, A. E. (2008). CressExpress: a tool for large-scale mining of expression data from *Arabidopsis*. Plant Physiology

147, 1004–1016.

- Staswick, P. E. and Tiryaki, I. (2004). The oxylipin signal jasmonic acid is activated by an enzyme that conjugates it to isoleucine in *Arabidopsis*. *The Plant Cell* 16, 2117–2127.
- Steinhauser, D., Usadel, B., Luedemann, A., Thimm, O. and Kopka, J. (2004). CSB.DB: a comprehensive systems-biology database. *Bioinformatics*, 20, 3647–3651.
- Stracke, R., Werber, M. and Weisshaar, B. (2001). The *R2R3-MYB* gene family in *Arabidopsis thaliana*. *Current Opinion in Plant Biology* 4, 447–456.
- Strahl, B. D. and Allis, C. D. (2000). The language of covalent histone modifications. *Nature* 403, 41–45.
- Strompen, G., Grüner, R. and Pfitzner, U. M. (1998). An *as-1*-like motif controls the level of expression of the gene for the pathogenesis-related protein 1a from tobacco. *Plant Molecular Biology* 37, 871–883.
- Suarez-Rodriguez, M. C., Adams-Phillips, L., Liu, Y., Wang, H., Su, S-H., Jester, P. J., Zhang, S., Bent, A. F. and Krysan, P. J. (2007). MEKK1 is required for flg22-induced MPK4 activation in *Arabidopsis* plants. *Plant Physiology* 143, 661–669.
- Sun, C., Palmqvist, S., Olsson, H., Borén, M., Ahlandsberg, S. and Jansson, C. (2003). A novel WRKY transcription factor, *SUSIBA2*, participates in sugar signaling in barley by binding to the sugar-responsive elements of the *iso1* promoter. *The Plant Cell* 15, 2076–2092.

T

- Takahashia, F., Yoshidab, R., Ichimurac, K., Mizoguchib, T., Seoe, S., Yonezawac, M., Maruyamaf, K., Yamaguchi-Shinozakif, K. and Shinozakia, K. (2007). The Mitogen-Activated Protein Kinase Cascade MKK3–MPK6 Is an Important Part of the Jasmonate Signal Transduction Pathway in *Arabidopsis*. *The Plant Cell* 19, 805–818.
- Tang, X., Frederick, R. D., Zhou, J., Halterman, D. A., Jia, Y. and Martin, G. B. (1996). Initiation of plant disease resistance by physical interaction of AvrPto and Pto kinase. *Science* 274, 2060–2063.
- Thatcher, L. F., Manners, J. M. and Kazan, K. (2009). *Fusarium oxysporum* hijacks COI1-mediated jasmonate signaling to promote disease development in *Arabidopsis*. *The Plant Journal* 58, 927–939.
- Thines, B., Katsir, L., Melotto, M., Niu, Y., Mandaokar, A., Liu, G., Nomura, K., He, S. Y., Howe, G. A. and Browse, J. (2007). JAZ repressor proteins are targets of the SCF^{COI1} complex during jasmonate signalling. *Nature* 448, 661–665.
- Thomma, B. P. H. J., Penninckx, I. A., Broekaert, W. F. and Cammue, B. P. (2001). The complexity of disease signaling in *Arabidopsis*. *Current Opinion Immunol* 13, 63–68.
- Thurrow, C., Schiermeyer, A., Krawczyk, S., Butterbrodt, T., Nickolov, K. and Gatz, C. (2005). Tobacco bZIP transcription factor TGA2.2 and related factor TGA2.1 have distinct roles in plant defense responses and plant development. *The Plant Journal* 44, 100–113.
- Tiryaki, I. and Staswick, P. E. (2002). An *Arabidopsis* mutant defective in jasmonate response is

allelic to the auxin-signaling mutant *axr1*. *Plant Physiology* 130, 887–894.

Toledo-Ortiz, G., Huq, E. and Quail, P. H. (2003). The *Arabidopsis* basic/helix-loop-helix transcription factor family. *The Plant Cell* 15, 1749–1770.

Töpfer, R., Matzeit, V., Gronenborn, B., Schell, J. and Steinbiss, H. H. (1987). A set of plant expression vectors for transcriptional and translational fusions. *Nucleic Acids Res.* 15, 5890.

Toufighi, K., Brady, S. M., Austin, R., Ly, E. and Provart, N. J. (2005). The Botany Array Resource: e-Northerns, Expression Angling, and promoter analyses. *The Plant Journal* 43, 153–163.

Tsuda, K., Sato, M., Glazebrook, J., Cohen, J. D. and Katagiri, F. (2008). Interplay between MAMP-triggered and SA-mediated defense responses. *The Plant Journal* 53, 763–775.

Tuch, B. B., Li, H. and Johnson, A. D. (2008). Evolution of Eukaryotic Transcription Circuits. *Science* 319, 1797–1799.

Turner, J. G., Ellis, C. and Devoto, A. (2002). The jasmonate signal pathway. *The Plant Cell* 14, S153–S164.

U

Uknes, S., Dincher, S., Friedrich, L., Negrotto, D., Williams, S., Thompson-Taylor, H., Potter, S., Ward, E. and Ryals, J. (1993). Regulation of pathogenesis-related protein-1a gene expression in tobacco. *The Plant Cell* 5, 159–169.

Ülker, B. and Somssich, I. E. (2004). WRKY transcription factors: from DNA binding towards biological function. *Current Opinion in Plant Biology* 7, 491–498.

V

Vailleau, F., Daniel, X., Tronchet, M., Montillet, J.-L., Triantaphylidès, C. and Roby, D. (2002). A R2R3-MYB gene, *AtMYB30*, acts as a positive regulator of the hypersensitive cell death program in plants in response to pathogen attack. *Proceedings of the National Academy of Sciences of the United States of America* 99, 10179–10184.

Van der Ent, S., Verhagen, B. W. M., Van Doorn, R., Bakker, D., Verlaan, M. G., Pel, M. J. C., Joosten, R. G., Proveniers, M. C. G., Van Loon, L. C., Ton, J. and Pieterse, C. M. J. (2008). *MYB72* is required in early signaling steps of rhizobacteria-induced systemic resistance in *Arabidopsis*. *Plant Physiology* 146, 1293–1304.

Van de Rhee, M. D. and Bol, J. F. (1993). Induction of the tobacco PR-1a gene by virus infection and salicylate treatment involves an interaction of multiple regulatory elements. *The Plant Journal* 3, 71–82.

Van de Rhee, M. D., van Kan, J. A. L., González Jaén, M. T. and Bol, J. F. (1990). Analysis of regulatory elements involved in the induction of two tobacco genes by salicylate treatment and virus infection. *The Plant Cell* 2, 357–366.

Van der Fits, L. and Memelink, J. (1997). Comparison of the activities of CaMV 35S and FMV 34S promoter derivatives in *Catharanthus roseus* cells transiently and stably transformed by par-

- ticle bombardment. *Plant Molecular Biology* 33, 943–946.
- Van der Fits, L. and Memelink, J. (2000). ORCA3, a jasmonate-responsive transcriptional regulator of plant primary and secondary metabolism. *Science* 289, 295–297.
- Van Loon, L. C. (1997). Induced resistance in plants and the role of pathogenesis-related proteins. *European Journal Plant Pathology* 103, 753–65.
- Van Loon, L. C. and van Strien, E. A. (1999). The families of pathogenesis-related proteins, their activities and comparative analysis of PR-1 type proteins. *Physiol. Mol. Plant. Pathol.* 55, 85–97.
- Van Verk, M. C., Pappaioannou, D., Neeleman, L., Bol, J. F. and Linthorst, H. J. M. (2008). A Novel WRKY transcription factor is required for induction of *PR-1a* gene expression by salicylic acid and bacterial elicitors. *Plant Physiology* 146, 1983–1995.
- Verberne, M. C., Verpoorte, R., Bol, J. F., Mercado-Blanco, J. and Linthorst, H. J. M. (2000). Overproduction of salicylic acid in plants by bacterial transgenes enhances pathogen resistance. *Nature Biotechnology* 18, 779–783.
- Verberne, M. C., Sansuk, K., Bol, J. F., Linthorst, H. J. M. and Verpoorte, R. (2007). Vitamin K₁ accumulation in tobacco plants overexpressing bacterial genes involved in the biosynthesis of salicylic acid. *Journal of Biotechnology* 128, 72–79.

W

- Walley, J. W., Rowe, H. C., Xiao, Y., Chehab, E. W., Kliebenstein, D. J., Wagner, D. and Dehesh, K. (2008). The chromatin remodeler SPLAYED regulates specific stress signaling pathways. *PLoS Pathogens* 4, e1000237.
- Wang, Z., Yang, P., Fan, B. and Chen, Z. (1998). An oligo selection procedure for identification of sequence-specific DNA-binding activities associated with the plant defense response. *The Plant Journal* 16, 515–522.
- Wang, K. L.-C., Yoshida, H., Lurin, C. and Ecker, J. R. (2004). Regulation of ethylene gas biosynthesis by the *Arabidopsis* ETO1 protein. *Nature* 428, 945–950.
- Wang, D., Amornsiripanitch, N. and Dong, X. (2006). A genomic approach to identify regulatory nodes in the transcriptional network of systemic acquired resistance in plants. *PLoS Pathogens* 2, e123.
- Wasternack, C. (2007). Jasmonates: An update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Annals of Botany* 100, 681–697.
- Weigel, C., Schmidt, A., Rückert, B., Lurz, R. and Messer, W. (1997). DnaA protein binding to individual DnaA boxes in the *Escherichia coli* replication origin, oriC. *EMBO J* 16, 6574–6583.
- White, R. F. (1979). Acetylsalicylic acid (aspirin) induces resistance to tobacco mosaic virus in tobacco. *Virology* 99, 410–412.
- Wildermuth, M. C., Dewdney, J., Wu, G. and Ausubel, F. M. (2001). Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* 414, 562–565.
- Wirthmueller, L., Zhang, Y., Jones, J. D. G. and Parker, J. E. (2007). Nuclear accumulation of the *Arabidopsis* immune receptor RPS4 is necessary for triggering EDS1-dependent defense. *Current*

Biology 17, 2023–2029.

Wu and Brand, L. (1994). Resonance energy transfer: methods and applications. *Anal. Biochem.* 218, 1–13.

Wu, K., Tian, L., Hollingworth, J., Brown, D. C. W. and Miki, B. (2002). Functional analysis of tomato *Pti4* in *Arabidopsis*. *Plant Physiology* 128, 30–37.

Wu, K., Zhang, L., Zhou, C., Yu, C.-W. and Chaikam, V. (2008). HDA6 is required for jasmonate response, senescence and flowering in *Arabidopsis*. *Journal of Experimental Botany* 59, 225–234.

X

Xiang, C., Miao, Z.-H. and Lam, E. (1996). Coordinated activation of *as-1*-type elements and a tobacco glutathion *S*-transferase gene by auxins, salicylic acid, methyl-jasmonate and hydrogen peroxide. *Plant Molecular Biology* 32, 415–426.

Xiang, C., Miao, Z. and Lam, E. (1997). DNA-binding properties, genomic organization and expression pattern of TGA6, a new member of the TGA family of bZIP transcription factors in *Arabidopsis thaliana*. *Plant Molecular Biology* 34, 403–415.

Xie, D.-X., Feys, B. F., James, S., Nieto-Rostro, M. and Turner, J. G. (1998). COI1: An *Arabidopsis* gene required for jasmonate regulated defense and fertility. *Science* 280, 1091–1094.

Xing, D.-H., Laia, Z.-B., Zhenga, Z.-Y., Vinoda, K. M., Fana, B.-F. and Chen, Z.-X. (2008). Stress- and Pathogen-Induced *Arabidopsis* WRKY48 is a Transcriptional Activator that Represses Plant Basal Defense. *Mol. Plant* 1, 459–470.

Xu, X., Chen, C., Fan, B. and Chen, Z. (2006). Physical and functional interactions between pathogen-induced *Arabidopsis* WRKY18, WRKY40, and WRKY60 transcription factors. *The Plant Cell* 18, 1310–1326.

Y

Yalpani, N., Leon, J., Lawton, M. A. and Raskin, I. (1991). Pathway of salicylic acid biosynthesis in healthy and virus-inoculated tobacco. *Plant Physiology* 103, 315–321.

Yamamoto, S., Nakano, T., Suzuki, K. and Shinshi, H. (2004). Elicitor-induced activation of transcription via W box-related cis-acting elements from a basic chitinase gene by WRKY transcription factors in tobacco. *Biochim. Biophys. Acta* 1679, 279–287.

Yamasaki, K., Kigawa, T., Inoue, M., Tateno, M., Yamasaki, T., Yabuki, T., Aoki, M., Seki, E., Matsuda, T., Tomo, Y., et al (2005). Solution structure of an *Arabidopsis* WRKY DNA binding domain. *The Plant Cell* 17, 944–956.

Yang, Y. and Klessig, D. F. (1996). Isolation and characterization of a tobacco mosaic virus-inducible myb oncogene homolog from tobacco. *Proceedings of the National Academy of Sciences of the United States of America* 93, 14972–14977.

Yang, Z., Tian, L., Latoszek-Green, M., Brown, D. and Wu, K. (2005). *Arabidopsis* ERF4 is a

- transcriptional repressor capable of modulating ethylene and abscisic acid responses. *Plant Molecular Biology* 58, 585–596.
- Yoo, J.-H., Park, C. Y., Kim, J. C., Heo, W. D., Cheong, M. S., Park, H. C., Kim, M. C., Moon, B. C., Choi, M. S., Kang, Y. H., Lee, J. H., Kim, H. S., Lee, S. M., Yoon, H. W., Lim, C. O., Yun, D.-J., Lee, S. Y., Chung, W. S. and Cho, M. J. (2005). Direct Interaction of a Divergent CaM Isoform and the Transcription Factor, MYB2, Enhances Salt Tolerance in Arabidopsis. *Journal of Biological Chemistry* 280, 3697–3706.
- Yoo, S.-D., Cho, Y.-H., Tena, G., Xiong, Y. and Sheen, J. (2008). Dual control of nuclear EIN3 by bifurcate MAPK cascades in C₂H₄ signalling. *Nature* 451, 789–795.

Z

- Zhang, B., Chen, W., Foley, R. C., Büttner, M. and Singh, K. B. (1995). Interactions between distinct types of DNA binding proteins enhance binding to ocs element promoter sequences. *The Plant Cell* 7, 2241–2252.
- Zhang, Y. and Wang, L. (2005). The WRKY transcription factor superfamily: its origin in eukaryotes and expansion in plants. *BMC Evolutionary Biol.* 5, 1.
- Zhang, Y., Fan, W., Kinkema, M., Li, X. and Dong, X. (1999). Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the *PR-1* gene. *Proceedings of the National Academy of Sciences of the United States of America* 96, 6523–6528.
- Zhang, Y., Tessaro, M. J., Lassner, M. and Li, X. (2003). Knockout analysis of Arabidopsis transcription factors *TGA2*, *TGA5*, and *TGA6* reveals their redundant and essential roles in systemic acquired resistance. *The Plant Cell* 15, 2647–2653.
- Zhao, J., Davis, L. C. and Verpoorte, R. (2005). Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnology Advances* 23, 283–333.
- Zhou, J., Tang, X. and Martin, G. B. (1997). The Pto kinase conferring resistance to tomato bacterial speck disease interacts with proteins that bind a *cis*-element of pathogenesis-related genes. *The EMBO Journal* 16, 3207–3218.
- Zhou, J.-M., Trifa, Y., Silva, H., Pontier, D., Lam, E., Shah, J. and Klessig, D. F. (2000). NPR1 differentially interacts with members of the TGA/OBF family of transcription factors that bind an element of the PR-1 gene required for induction by salicylic acid. *Mol. Plant Microbe Interact.* 13, 191–202.
- Zhou, C., Zhang, L., Duan, J., Miki, B. and Wu, K. (2005). *HISTONE DEACETYLASE19* is involved in jasmonic acid and ethylene signaling of pathogen response in Arabidopsis. *The Plant Cell* 17, 1196–1204.
- Zimmermann, P., Hirsch-Hoffmann, M., Hennig, L. and Gruissem, W. (2004). GENEVESTIGATOR: Arabidopsis Microarray Database and Analysis Toolbox. *Plant Physiology* 136, 2621–2632.
- Zimmermann, P., Hennig, L. and Gruissem, W. (2005). Gene expression analysis and network

discovery using Genevestigator. Trends in Plant Science 10, 407-409.

Zwicker, S., Mast, S., Stos, V., Pfitzner, A. J. P. and Pfitzner, U. M. (2007). Tobacco NIMIN2 proteins control PR gene induction through transient repression early in systemic acquired resistance. Molecular Plant Pathology 8, 385-400.