



Universiteit
Leiden
The Netherlands

Clinical pharmacology of the tyrosine kinase inhibitors imatinib and sunitinib

Erp, P.H. van

Citation

Erp, P. H. van. (2009, December 16). *Clinical pharmacology of the tyrosine kinase inhibitors imatinib and sunitinib*. Retrieved from <https://hdl.handle.net/1887/14515>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/14515>

Note: To cite this publication please use the final published version (if applicable).



General introduction and scope of the thesis

Cancer is the second leading cause of death worldwide, after cardiovascular disease, accounting for 7.9 million deaths; ~ 13% of all deaths in 2007. Additionally the incidence of cancer is increasing. The five most mortal types of cancer are; lung, stomach, liver, colorectal and esophageal cancer. Over 30% of cancer can be prevented by not using tobacco, having a healthy diet, being physically active and by preventing infections that may cause cancer¹. Once diagnosed, there are several different types of treatment ranging from resection (surgery), to radiation (radiotherapy), to systemic therapy used as adjuvant or palliative therapy. The conventional cytotoxic chemotherapeutic agents have a generic working profile that interact non-specifically with cellular DNA and/ or tubulin resulting in growth arrest of all fast growing cells. With the increased understanding of cancer biology, rational design of targeted drugs has started. Targeted drugs have antitumor activity in selected subgroups of tumors expressing proteins that are specific for the malignant phenotype². The clinical use of targeted therapy started with the development of monoclonal antibodies³. Five years later, the first tyrosine kinase inhibitor was approved for cancer treatment. Tyrosine kinase inhibitors are a class of targeted therapy that is designed to compete with adenosine-5'-triphosphate (ATP) for the ATP-binding pocket within the intracellular domain of wild type and/or mutated tyrosine kinase receptor and thereby blocks downstream signaling important for tumor growth. Imatinib is the first rationally designed tyrosine kinase inhibitor approved in 2001 for the treatment of three Philadelphia chromosome positive leukemia subtypes⁴. Since 2001, seven additional tyrosine kinase inhibitors have been approved, all rationally designed to be active against specific tyrosine kinases. These targeted drugs tend to have a better toxicity profile than traditional cytotoxic chemotherapy that interacts non-specifically resulting in more collateral, transient damage in healthy tissues⁵. With the introduction of tyrosine kinase inhibitors a new era of treating tumors has started⁶.

All tyrosine kinase inhibitors exhibit rather similar pharmacokinetic characteristics. They are all highly protein bound, have a long half life and a large volume of distribution, they are all primarily metabolized by cytochrome P450 (CYP) 3A, and predominantly excreted with the feces⁷⁻¹⁴. However, several pharmacokinetic aspects of these drugs are also unknown. For example, the absolute bioavailability for most tyrosine kinase inhibitors is unknown as is the clinical relevance of their interactions with (substrates for and/or inhibitors of) drug transporters on intestinal cells, hepatocytes, cancer cells and renal cells. Since these drugs are both substrates and inhibitors of their own metabolic pathways, the metabolism of these drugs at steady-state exposure is complex and unpredictable.

Therefore, the aim of this thesis is to further explore clinical pharmacological aspects of two tyrosine kinase inhibitors; imatinib and sunitinib, to better understand steady-state pharmacokinetics, clinical relevant interactions and genetic determinants that may predispose for specific side effects of these drugs.

Most information of the pharmacokinetic behavior of the tyrosine kinase inhibitors originates from preclinical studies. In addition, clinical studies have revealed important pharmacokinetic data of these drugs. An overview of the current knowledge on absorption, distribution, metabolism, elimination, drug transporter affinity and drug-drug interactions of all approved tyrosine kinase inhibitors as well as their similarities and differences will be presented in **chapter 2**.

Little information is available on the relevance of drug interactions at steady-state pharmacokinetics. According to the drug label of imatinib, CYP3A4 is the most important enzyme responsible for the metabolism. Since many clinically used drugs are known to inhibit or induce CYP3A4, imatinib is prone for drug-drug interactions. In **chapter 3** we will determine the effect of ritonavir, a potent CYP3A4 inhibitor, on the steady-state imatinib exposure (AUC). Multiple CYP enzymes, such as CYP3A4, 3A5, 2D6, 2C9, 2C19, 1A2, 1A1, are capable of metabolizing imatinib in *in vitro* experiments; however there are no data available on the influence of these minor enzymes on imatinib exposure¹⁵. Since we know that smoking has a pronounced effect on CYP mediated metabolism and hereby on erlotinib exposure a similar effect is hypothesized for imatinib. In **chapter 4**, the effect cigarette smoking on imatinib exposure will be studied¹⁶.

The exact absorption-site of imatinib in the intestines is unknown. Some patients with gastrointestinal stromal tumor (GIST) may not be able to take imatinib orally, due to tumor related gastro-intestinal obstruction. Therefore, in **chapter 5** we will study imatinib pharmacokinetics in a patient after using the rectal route of administration.

Sunitinib, like all tyrosine kinase inhibitors, shows large interpatient variability in drug exposure which might affect the clinical outcome with respect to both toxicity and efficacy. In clinical practice ~ 33% of the patients need a dose interruption or a dose reduction due to drug related toxicities¹⁷⁻¹⁹. We will explore the use of a noninvasive and harmless phenotypic probe (midazolam) to determine CYP3A4 activity and thereby predict the exposure to sunitinib before starting sunitinib therapy. The results of this study will be described in **chapter 6**. Most interaction studies are performed with a single dose of the drug of interest, whereas the metabolism at steady-state can be distinctively different due to auto-inhibition of the primary metabolic pathway²⁰. Some tyrosine kinase inhibitors (imatinib, dasatinib and nilotinib) appear to be both substrates and inhibitors of CYP3A4^{12,21,22}. The effect of steady-state sunitinib exposure on CYP3A4 activity is also described in **chapter 6**. Additionally, we will study the association between genetic variants in genes encoding enzymes, transporters and sunitinib targets and sunitinib induced toxicities (**chapter 7**).

Since the absolute bioavailability of sunitinib is unknown, the influence of intestinal CYP3A4 activity on sunitinib exposure is unpredictable. However, in the drug label of sunitinib there is a warning for co-administration of CYP3A4 inhibitors, such as ketoconazole, clarithromycin and indinavir, but also for grapefruit juice which is a potent inhibitor of intestinal CYP3A4.

The suggested effect of grapefruit juice on steady-state sunitinib exposure will be determined (**chapter 8**). A drug-drug interaction in two patients treated with mitotane and sunitinib will be presented in **chapter 9**. In **chapter 10** a possible explanation will be presented for the pronounced effect of grapefruit juice on intestinal but absent effect on hepatic CYP3A4 in healthy volunteers.

Finally the results from these studies will be put into perspective in the general discussion (**chapter 11**).

References

1. WHO. World Health Organization - Cancer. <http://www.who.int/cancer/en/> Accessed february 2009.
2. Carden CP, Banerji U, Kaye SB, Workman P, de Bono JS. From darkness to light with biomarkers in early clinical trials of cancer drugs. *Clin Pharmacol Ther* 2009; 85(2):131-133.
3. Houshmand P, Zlotnik A. Targeting tumor cells. *Curr Opin Cell Biol* 2003; 15(5):640-644.
4. Druker BJ, Talpaz M, Resta DJ et al. Efficacy and safety of a specific inhibitor of the BCR-ABL tyrosine kinase in chronic myeloid leukemia. *N Engl J Med* 2001; 344(14):1031-1037.
5. Faivre S, Demetri G, Sargent W, Raymond E. Molecular basis for sunitinib efficacy and future clinical development. *Nat Rev Drug Discov* 2007; 6(9):734-745.
6. Baker SD, Hu S. Pharmacokinetic considerations for new targeted therapies. *Clin Pharmacol Ther* 2009; 85(2):208-211.
7. Cohen MH, Williams GA, Sridhara R et al. United States Food and Drug Administration Drug Approval summary: Gefitinib (ZD1839; Iressa) tablets. *Clin Cancer Res* 2004; 10(4):1212-1218.
8. Cohen MH, Williams G, Johnson JR et al. Approval summary for imatinib mesylate capsules in the treatment of chronic myelogenous leukemia. *Clin Cancer Res* 2002; 8(5):935-942.
9. Johnson JR, Cohen M, Sridhara R et al. Approval summary for erlotinib for treatment of patients with locally advanced or metastatic non-small cell lung cancer after failure of at least one prior chemotherapy regimen. *Clin Cancer Res* 2005; 11(18):6414-6421.
10. Kane RC, Farrell AT, Saber H et al. Sorafenib for the treatment of advanced renal cell carcinoma. *Clin Cancer Res* 2006; 12(24):7271-7278.
11. Goodman VL, Rock EP, Dagher R et al. Approval summary: sunitinib for the treatment of imatinib refractory or intolerant gastrointestinal stromal tumors and advanced renal cell carcinoma. *Clin Cancer Res* 2007; 13(5):1367-1373.
12. Brave M, Goodman V, Kaminskas E et al. Sprycel for chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia resistant to or intolerant of imatinib mesylate. *Clin Cancer Res* 2008; 14(2):352-359.
13. Medina PJ, Goodin S. Lapatinib: a dual inhibitor of human epidermal growth factor receptor tyrosine kinases. *Clin Ther* 2008; 30(8):1426-1447.
14. Hazarika M, Jiang X, Liu Q et al. Tasigna for chronic and accelerated phase Philadelphia chromosome-positive chronic myelogenous leukemia resistant to or intolerant of imatinib. *Clin Cancer Res* 2008; 14(17):5325-5331.
15. Peng B, Lloyd P, Schran H. Clinical pharmacokinetics of imatinib. *Clin Pharmacokinet* 2005; 44(9):879-894.
16. Hamilton M, Wolf JL, Rusk J et al. Effects of smoking on the pharmacokinetics of erlotinib. *Clin Cancer Res* 2006; 12(7 Pt 1):2166-2171.
17. Demetri GD, van Oosterom AT, Garrett CR et al. Efficacy and safety of sunitinib in patients with advanced gastrointestinal stromal tumour after failure of imatinib: a randomised controlled trial. *Lancet* 2006; 368(9544):1329-1338.
18. Motzer RJ, Hutson TE, Tomczak P et al. Sunitinib versus interferon alfa in metastatic renal-cell carcinoma. *N Engl J Med* 2007; 356(2):115-124.
19. van der Veldt AAM, Boven E, Helgason HH et al. Predictive factors for severe toxicity of sunitinib in unselected patients with advanced renal cell cancer. *Br J Cancer* 2008; 99(2):259-265.
20. van Erp NP, Gelderblom H, Karlsson MO et al. Influence of CYP3A4 inhibition on the steady-state pharmacokinetics of imatinib. *Clin Cancer Res* 2007; 13(24):7394-7400.
21. O'Brien SG, Meinhardt P, Bond E et al. Effects of imatinib mesylate (STI571, Glivec) on the pharmacokinetics of simvastatin, a cytochrome p450 3A4 substrate, in patients with chronic myeloid leukaemia. *Br J Cancer* 2003; 89(10):1855-1859.
22. Tanaka C, Smith T, Kantarjian H et al. Clinical pharmacokinetics (PK) of AMN107, a novel inhibitor of Bcr-Abl, in healthy subjects and patients with imatinib resistant or intolerant chronic myelogenous leukemia (CML) or relapsed/refractory Ph+ acute lymphocytic leukemia (Ph+ALL). *J Clin Oncol*, 2006 ASCO Annual Meeting Proceedings Part I Vol. 24, No. 18S, 3095. 2006.