

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/32966> holds various files of this Leiden University dissertation.

Author: Visser, Erwin Lourens

Title: Neutrinos from the Milky Way

Issue Date: 2015-05-12

NEUTRINOS FROM THE MILKY WAY

NEUTRINOS FROM THE MILKY WAY

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. C.J.J.M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op dinsdag 12 mei 2015
klokke 16:15 uur

door

Erwin Lourens Visser
geboren te Hoorn
in 1987

Promotiecommissie:

Promotor: Prof. dr. M. de Jong
Co-promotor: Dr. D.F.E. Samtleben

Overige leden: Dr. A.J. Heijboer (Nikhef, Amsterdam)
Prof. dr. M. Kadler (Universität Würzburg, Würzburg, Germany)
Prof. dr. P.M. Kooijman (Universiteit van Amsterdam)
Prof. dr. E.R. Eliel
Prof. dr. J.W. van Holten

Casimir PhD series, Delft-Leiden 2015-11

ISBN 978-90-8593-219-2

First published in print format 2015

An electronic version of this thesis can be found at <https://openaccess.leidenuniv.nl>

© Erwin Lourens Visser 2015

The work described in this thesis is part of the research programme of the Foundation for Fundamental Research on Matter (FOM), which is part of the Netherlands Organisation for Scientific Research (NWO).

The cover shows a panoramic image of the Milky Way over Lake Tekapo, New Zealand, after sunset with the zodiacal light visible.

© Alex Cherney (www.terrastro.com).

This document was typeset using the typographical look-and-feel *classicthesis* developed by André Miede. The style was inspired by Robert Bringhurst's seminal book on typography "*The Elements of Typographic Style*". *classicthesis* is available for both L^AT_EX and LyX:

<http://code.google.com/p/classicthesis>

Printed in the Netherlands by IJskamp Drukkers B.V.

Aan mijn ouders

CONTENTS

1	INTRODUCTION	1
1.1	The advent of astroparticle physics	5
1.2	Thesis goals and structure	11
2	NEUTRINO FLUXES FROM COSMIC RAY INTERACTIONS IN THE MILKY WAY	15
2.1	Model ingredients	15
2.1.1	The Milky Way	16
2.1.2	The interstellar matter	19
2.1.3	The magnetic field	23
2.1.4	Cosmic ray flux	25
2.2	Theoretical models for the neutrino flux	33
2.2.1	Assumptions	34
2.2.2	Calculation of $\nu_\mu + \bar{\nu}_\mu$ fluxes	41
2.3	Calculation of neutrino fluxes from Fermi γ -ray flux	48
2.3.1	Photon flux measured by Fermi	49
2.3.2	Photon flux from π^0 -decays	52
2.3.3	Determination of pion fluxes	56
2.3.4	Obtained $\nu_\mu + \bar{\nu}_\mu$ fluxes	58
2.4	Signal flux comparisons	60
2.4.1	Atmospheric neutrinos	60
2.4.2	Signal compared to the background	62
2.4.3	The Mediterranean sea versus the South Pole	63
3	THE ANTARES NEUTRINO TELESCOPE	67
3.1	Neutrino signatures	67
3.1.1	Muon propagation	70
3.2	The ANTARES detector	73
3.2.1	The optical module	74
3.2.2	Detector layout	76
3.2.3	Data acquisition	78
3.2.4	The shore station	78
3.2.5	Calibration	79
4	SIMULATION, TRIGGERS AND RECONSTRUCTION	81
4.1	Simulation tools	81
4.1.1	Neutrino generation	82
4.1.2	Atmospheric muon generation	84

4.1.3	Propagation of muons, light and other secondaries	84
4.1.4	Detector simulation	85
4.1.5	MC productions	85
4.2	Triggering	86
4.2.1	The 3N trigger algorithm	87
4.2.2	The 2T3 trigger algorithm	89
4.2.3	The directional trigger algorithm	89
4.2.4	The TQ trigger algorithm	92
4.3	Reconstruction	96
4.3.1	BBFIT	97
4.3.2	AAFIT	101
4.3.3	GRIDFIT	106
4.3.4	Energy reconstruction	126
4.3.5	Shower reconstruction	128
5	CONSTRAINTS ON THE DIFFUSE GALACTIC NEUTRINO FLUX FROM ANTARES	129
5.1	Determining the optimal signal region size	130
5.1.1	Statistical tools	130
5.1.2	Construction of the background regions	135
5.1.3	Signal region optimisation	137
5.2	Checks on the background regions	141
5.2.1	Data selection	142
5.2.2	Effective visibility	142
5.2.3	Checking for systematic biases	146
5.2.4	Data-MC comparison	149
5.3	Event selection	151
5.3.1	Optimisation without R_{GF}	151
5.3.2	Optimisation without β	153
5.3.3	Full optimisation	154
5.3.4	Additional optimisation	156
5.4	ANTARES sensitivity	161
5.4.1	The cosmic neutrino flux measured by Ice-Cube	164
5.5	Results	166
6	DETECTION POTENTIAL OF KM3NET FOR THE DIFFUSE GALACTIC NEUTRINO FLUX	173
6.1	KM3NeT	173
6.1.1	Muon track reconstruction	176
6.2	Determining the optimal signal region size	178
6.2.1	Statistical tools	178
6.2.2	Signal region optimisation	180

6.3 KM3NeT sensitivity	185
6.4 KM3NeT discovery potential	188
7 CONCLUSIONS AND OUTLOOK	191
BIBLIOGRAPHY	197
SUMMARY	209
SAMENVATTING	213
ABOUT THE AUTHOR	217
ACKNOWLEDGEMENTS	219

