



Universiteit  
Leiden  
The Netherlands

## **Vulnerability to cocaine: role of stress hormones**

Jong, I.E.M. de

### **Citation**

Jong, I. E. M. de. (2007, October 17). *Vulnerability to cocaine: role of stress hormones*. Division of Medical Pharmacology of the Leiden/Amsterdam Center for Drug Research (LACDR) and Leiden University Medical Center (LUMC), Leiden University. Retrieved from <https://hdl.handle.net/1887/12382>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12382>

**Note:** To cite this publication please use the final published version (if applicable).

# 9

---

## References

---



- [1] Abercrombie ED, Keefe KA, DiFrischia DS, Zigmond MJ. Differential effect of stress on in vivo dopamine release in striatum, nucleus accumbens, and medial frontal cortex. *J Neurochem* 1989; 52(5): 1655-1658.
- [2] Abraham I, Juhasz G, Kekesi KA, Kovacs KJ. Corticosterone peak is responsible for stress-induced elevation of glutamate in the hippocampus. *Stress* 1998; 2(3): 171-181.
- [3] Ackerman JM, White FJ. A10 somatodendritic dopamine autoreceptor sensitivity following withdrawal from repeated cocaine treatment. *Neurosci Lett* 1990; 117(1-2): 181-187.
- [4] Ahima R, Krozowski Z, Harlan R. Type I corticosteroid receptor-like immunoreactivity in the rat CNS: distribution and regulation by corticosteroids. *J Comp Neurol* 1991; 313(3): 522-538.
- [5] Akimoto K, Hamamura T, Kazahaya Y, Akiyama K, Otsuki S. Enhanced extracellular dopamine level may be the fundamental neuropharmacological basis of cross-behavioral sensitization between methamphetamine and cocaine - an in vivo dialysis study in freely moving rats. *Brain Res* 1990; 507(2): 344-346.
- [6] Albertson DN, Schmidt CJ, Kapatos G, Bannon MJ. Distinctive profiles of gene expression in the human nucleus accumbens associated with cocaine and heroin abuse. *Neuropsychopharmacology* 2006; 31(10): 2304-2312.
- [7] Alexander GE, Crutcher MD. Functional architecture of basal ganglia circuits: neural substrates of parallel processing. *Trends Neurosci* 1990; 13(7): 266-271.
- [8] Alleweireldt AT, Hobbs RJ, Taylor AR, Neisewander JL. Effects of SCH-23390 infused into the amygdala or adjacent cortex and basal ganglia on cocaine seeking and self-administration in rats. *Neuropsychopharmacology* 2006; 31(2): 363-374.
- [9] Alleweireldt AT, Weber SM, Kirschner KF, Bullock BL, Neisewander JL. Blockade or stimulation of D1 dopamine receptors attenuates cue reinstatement of extinguished cocaine-seeking behavior in rats. *Psychopharmacology (Berl)* 2002; 159(3): 284-293.
- [10] Ammassari-Teule M, Passino E, Restivo L, de Marsanich B. Fear conditioning in C57BL/6 and DBA/2 mice: variability in nucleus accumbens function according to the strain predisposition to show contextual- or cue-based responding. *Eur J Neurosci* 2000; 12(12): 4467-4474.
- [11] Anderson SM, Bari AA, Pierce RC. Administration of the D1-like dopamine receptor antagonist SCH-23390 into the medial nucleus accumbens shell attenuates cocaine priming-induced reinstatement of drug-seeking behavior in rats. *Psychopharmacology (Berl)* 2003; 168(1-2): 132-138.
- [12] Anderson SM, Schmidt HD, Pierce RC. Administration of the D2 dopamine receptor antagonist sulpiride into the shell, but not the core, of the nucleus accumbens attenuates cocaine priming-induced reinstatement of drug seeking. *Neuropsychopharmacology* 2006; 31(7): 1452-1461.
- [13] Antelman SM, Eichler AJ, Black CA, Kocan D. Interchangeability of stress and amphetamine in sensitization. *Science* 1980; 207(4428): 329-331.
- [14] Aouizerate B, Ho A, Schluger JH, Perret G, Borg L, Le Moal M, Piazza PV, Kreek MJ. Glucocorticoid negative feedback in methadone-maintained former heroin addicts with ongoing cocaine dependence: dose-response to dexamethasone suppression. *Addict Biol* 2006; 11(1): 84-96.
- [15] Appelboom T. Consumption of coca in history. *Verh K Acad Geneesk Belg* 1991; 53(5): 497-505.

- [16] Auchus AP, Green RC, Nemeroff CB. Cortical and subcortical neuropeptides in Alzheimer's disease. *Neurobiol Aging* 1994; 15(4): 589-595.
- [17] Azar MR, Acar N, Erwin VG, Barbato GF, Morse AC, Heist CL, Jones BC. Distribution and clearance of cocaine in brain is influenced by genetics. *Pharmacol Biochem Behav* 1998; 59(3): 637-640.
- [18] Back SE, Brady KT, Jaanimagi U, Jackson JL. Cocaine dependence and PTSD: a pilot study of symptom interplay and treatment preferences. *Addict Behav* 2006; 31(2): 351-354.
- [19] Backstrom P, Hyytia P. Ionotropic and metabotropic glutamate receptor antagonism attenuates cue-induced cocaine seeking. *Neuropsychopharmacology* 2006; 31(4): 778-786.
- [20] Badiani A, Cabib S, Puglisi-Allegra S. Chronic stress induces strain-dependent sensitization to the behavioral effects of amphetamine in the mouse. *Pharmacol Biochem Behav* 1992; 43(1): 53-60.
- [21] Badiani A, Morano MI, Akil H, Robinson TE. Circulating adrenal hormones are not necessary for the development of sensitization to the psychomotor activating effects of amphetamine. *Brain Res* 1995; 673(1): 13-24.
- [22] Badiani A, Robinson TE. Drug-induced neurobehavioral plasticity: the role of environmental context. *Behav Pharmacol* 2004; 15(5-6): 327-339.
- [23] Bahi A, Boyer F, Bussard G, Dreyer JL. Silencing dopamine D3-receptors in the nucleus accumbens shell in vivo induces changes in cocaine-induced hyperlocomotion. *Eur J Neurosci* 2005; 21(12): 3415-3426.
- [24] Baker TB, Morse E, Sherman JE. The motivation to use drugs: a psychobiological analysis of urges. *Nebr Symp Motiv* 1986; 34: 257-323.
- [25] Balfour ME, Yu L, Coolen LM. Sexual behavior and sex-associated environmental cues activate the mesolimbic system in male rats. *Neuropsychopharmacology* 2004; 29(4): 718-730.
- [26] Banihashemi L, Rinaman L. Noradrenergic inputs to the bed nucleus of the stria terminalis and paraventricular nucleus of the hypothalamus underlie hypothalamic-pituitary-adrenal axis but not hypophagic or conditioned avoidance responses to systemic yohimbine. *J Neurosci* 2006; 26(44): 11442-11453.
- [27] Baptista MA, Martin-Fardon R, Weiss F. Preferential effects of the metabotropic glutamate 2/3 receptor agonist LY379268 on conditioned reinstatement versus primary reinforcement: comparison between cocaine and a potent conventional reinforcer. *J Neurosci* 2004; 24(20): 4723-4727.
- [28] Bari AA, Pierce RC. D1-like and D2 dopamine receptor antagonists administered into the shell subregion of the rat nucleus accumbens decrease cocaine, but not food, reinforcement. *Neuroscience* 2005; 135(3): 959-968.
- [29] Barr AM, Hofmann CE, Weinberg J, Phillips AG. Exposure to repeated, intermittent d-amphetamine induces sensitization of HPA axis to a subsequent stressor. *Neuropsychopharmacology* 2002; 26(3): 286-294.
- [30] Barrot M, Marinelli M, Abrous DN, Rouge-Pont F, Le Moal M, Piazza PV. The dopaminergic hyper-responsiveness of the shell of the nucleus accumbens is hormone-dependent. *Eur J Neurosci* 2000; 12(3): 973-979.

- [31] Baumann MH, Gendron TM, Becketts KM, Henningfield JE, Gorelick DA, Rothman RB. Effects of intravenous cocaine on plasma cortisol and prolactin in human cocaine abusers. *Biol Psychiatry* 1995; 38(11): 751-755.
- [32] Beitner-Johnson D, Guitart X, Nestler EJ. Dopaminergic brain reward regions of Lewis and Fischer rats display different levels of tyrosine hydroxylase and other morphine- and cocaine-regulated phosphoproteins. *Brain Res* 1991; 561(1): 147-150.
- [33] Beitner-Johnson D, Nestler EJ. Morphine and cocaine exert common chronic actions on tyrosine hydroxylase in dopaminergic brain reward regions. *J Neurochem* 1991; 57(1): 344-347.
- [34] Belanoff JK, Rothschild AJ, Cassidy F, DeBattista C, Baulieu EE, Schold C, Schatzberg AF. An open label trial of C-1073 (mifepristone) for psychotic major depression. *Biol Psychiatry* 2002; 52(5): 386-392.
- [35] Bell K, Kalivas PW. Context-specific cross-sensitization between systemic cocaine and intra-accumbens AMPA infusion in the rat. *Psychopharmacology (Berl)* 1996; 127(4): 377-383.
- [36] Ben Shahar O, Ahmed SH, Koob GF, Ettenberg A. The transition from controlled to compulsive drug use is associated with a loss of sensitization. *Brain Res* 2004; 995(1): 46-54.
- [37] Benjamin J, Li L, Patterson C, Greenberg BD, Murphy DL, Hamer DH. Population and familial association between the D4 dopamine receptor gene and measures of Novelty Seeking. *Nat Genet* 1996; 12(1): 81-84.
- [38] Berglind WJ, Case JM, Parker MP, Fuchs RA, See RE. Dopamine D1 or D2 receptor antagonism within the basolateral amygdala differentially alters the acquisition of cocaine-cue associations necessary for cue-induced reinstatement of cocaine-seeking. *Neuroscience* 2006; 137(2): 699-706.
- [39] Berhow MT, Hiroi N, Nestler EJ. Regulation of ERK (extracellular signal regulated kinase), part of the neurotrophin signal transduction cascade, in the rat mesolimbic dopamine system by chronic exposure to morphine or cocaine. *J Neurosci* 1996; 16(15): 4707-4715.
- [40] Berke JD, Paletzki RF, Aronson GJ, Hyman SE, Gerfen CR. A complex program of striatal gene expression induced by dopaminergic stimulation. *J Neurosci* 1998; 18(14): 5301-5310.
- [41] Berman SM, Noble EP. The D2 dopamine receptor (DRD2) gene and family stress; interactive effects on cognitive functions in children. *Behav Genet* 1997; 27(1): 33-43.
- [42] Berns GS. Something funny happened to reward. *Trends Cogn Sci* 2004; 8(5): 193-194.
- [43] Berrettini WH, Alexander R, Ferraro TN, Vogel WH. A study of oral morphine preference in inbred mouse strains. *Psychiatr Genet* 1994; 4(2): 81-86.
- [44] Bevins RA, Klebaur JE, Bardo MT. Individual differences in response to novelty, amphetamine-induced activity and drug discrimination in rats. *Behav Pharmacol* 1997; 8(2-3): 113-123.
- [45] Bhattacharyya AK, Pradhan SN. Interactions between motor activity and stereotypy in cocaine-treated rats. *Psychopharmacology (Berl)* 1979; 63(3): 311-312.
- [46] Bibb JA, Chen J, Taylor JR, Svenningsson P, Nishi A, Snyder GL, Yan Z, Sagawa ZK, Ouimet CC, Nairn AC, Nestler EJ, Greengard P. Effects of chronic exposure to cocaine are regulated by the neuronal protein Cdk5. *Nature* 2001; 410(6826): 376-380.

- [47] Biron D, Dauphin C, Di Paolo T. Effects of adrenalectomy and glucocorticoids on rat brain dopamine receptors. *Neuroendocrinology* 1992; 55(4): 468-476.
- [48] Blaha CD, Yang CR, Floresco SB, Barr AM, Phillips AG. Stimulation of the ventral subiculum of the hippocampus evokes glutamate receptor-mediated changes in dopamine efflux in the rat nucleus accumbens. *Eur J Neurosci* 1997; 9(5): 902-911.
- [49] Blanc G, Trovero F, Vezina P, Herve D, Godeheu AM, Glowinski J, Tassin JP. Blockade of prefronto-cortical alpha 1-adrenergic receptors prevents locomotor hyperactivity induced by subcortical D-amphetamine injection. *Eur J Neurosci* 1994; 6(3): 293-298.
- [50] Bonci A, Williams JT. A common mechanism mediates long-term changes in synaptic transmission after chronic cocaine and morphine. *Neuron* 1996; 16(3): 631-639.
- [51] Borowsky B, Kuhn CM. Chronic cocaine administration sensitizes behavioral but not neuroendocrine responses. *Brain Res* 1991; 543(2): 301-306.
- [52] Borowsky B, Kuhn CM. Monoamine mediation of cocaine-induced hypothalamo-pituitary-adrenal activation. *J Pharmacol Exp Ther* 1991; 256(1): 204-210.
- [53] Borrell J, De Kloet ER, Bohus B. Corticosterone decreases the efficacy of adrenaline to affect passive avoidance retention of adrenalectomized rats. *Life Sci* 1984; 34(1): 99-104.
- [54] Borrell J, De Kloet ER, Versteeg DH, Bohus B. Inhibitory avoidance deficit following short-term adrenalectomy in the rat: the role of adrenal catecholamines. *Behav Neural Biol* 1983; 39(2): 241-258.
- [55] Borski RJ. Nongenomic membrane actions of glucocorticoids in vertebrates. *Trends Endocrinol Metab* 2000; 11(10): 427-436.
- [56] Boudreau AC, Wolf ME. Behavioral sensitization to cocaine is associated with increased AMPA receptor surface expression in the nucleus accumbens. *J Neurosci* 2005; 25(40): 9144-9151.
- [57] Bouthenet ML, Souil E, Martres MP, Sokoloff P, Giros B, Schwartz JC. Localization of dopamine D3 receptor mRNA in the rat brain using in situ hybridization histochemistry: comparison with dopamine D2 receptor mRNA. *Brain Res* 1991; 564(2): 203-219.
- [58] Brake WG, Zhang TY, Diorio J, Meaney MJ, Gratton A. Influence of early postnatal rearing conditions on mesocorticolimbic dopamine and behavioural responses to psychostimulants and stressors in adult rats. *Eur J Neurosci* 2004; 19(7): 1863-1874.
- [59] Brebner K, Ahn S, Phillips AG. Attenuation of d-amphetamine self-administration by baclofen in the rat: behavioral and neurochemical correlates. *Psychopharmacology (Berl)* 2005; 177(4): 409-417.
- [60] Brebner K, Phelan R, Roberts DC. Effect of baclofen on cocaine self-administration in rats reinforced under fixed-ratio 1 and progressive-ratio schedules. *Psychopharmacologia* 2000; 148(3): 314-321.
- [61] Brebner K, Childress AR, Roberts DCS. A potential role for GABAB agonists in the treatment of psychostimulant addiction. *Alcohol Alcoholism* 2002; 37(5): 478-484.
- [62] Budygin EA, John CE, Mateo Y, Jones SR. Lack of cocaine effect on dopamine clearance in the core and shell of the nucleus accumbens of dopamine transporter knock-out mice. *J Neurosci* 2002; 22(10): RC222.

- [63] Buijs RM, Kalsbeek A, van der Woude TP, van Heerikhuizen JJ, Shinn S. Suprachiasmatic nucleus lesion increases corticosterone secretion. *AJP - Legacy* 1993; 264(6 Pt 2): R1186-R1192.
- [64] Bunzow JR, Van Tol HH, Grandy DK, Albert P, Salon J, Christie M, Machida CA, Neve KA, Civelli O. Cloning and expression of a rat D2 dopamine receptor cDNA. *Nature* 1988; 336(6201): 783-787.
- [65] Cabib S. Strain-dependent behavioural sensitization to amphetamine: role of environmental influences. *Behav Pharmacol* 1993; 4(4): 367-374.
- [66] Cabib S, Algeri S, Perego C, Puglisi-Allegra S. Behavioral and biochemical changes monitored in two inbred strains of mice during exploration of an unfamiliar environment. *Physiol Behav* 1990; 47(4): 749-753.
- [67] Cabib S, Bonaventura N. Parallel strain-dependent susceptibility to environmentally-induced stereotypies and stress-induced behavioral sensitization in mice. *Physiol Behav* 1997; 61(4): 499-506.
- [68] Cabib S, Castellano C, Patacchioli FR, Cigliana G, Angelucci L, Puglisi-Allegra S. Opposite strain-dependent effects of post-training corticosterone in a passive avoidance task in mice: role of dopamine. *Brain Res* 1996; 729(1): 110-118.
- [69] Cabib S, Giardino L, Calza L, Zanni M, Mele A, Puglisi-Allegra S. Stress promotes major changes in dopamine receptor densities within the mesoaccumbens and nigrostriatal systems. *Neuroscience* 1998; 84(1): 193-200.
- [70] Cabib S, Kempf E, Schlee C, Oliverio A, Puglisi-Allegra S. Effects of immobilization stress on dopamine and its metabolites in different brain areas of the mouse: role of genotype and stress duration. *Brain Res* 1988; 441(1-2): 153-160.
- [71] Cabib S, Orsini C, Le Moal M, Piazza PV. Abolition and reversal of strain differences in behavioral responses to drugs of abuse after a brief experience. *Science* 2000; 289(5478): 463-465.
- [72] Cabib S, Puglisi-Allegra S. Genotype-dependent effects of chronic stress on apomorphine-induced alterations of striatal and mesolimbic dopamine metabolism. *Brain Res* 1991; 542(1): 91-96.
- [73] Cabib S, Puglisi-Allegra S, Ventura R. The contribution of comparative studies in inbred strains of mice to the understanding of the hyperactive phenotype. *Behav Brain Res* 2002; 130(1-2): 103-109.
- [74] Cadoni C, Solinas M, Di Chiara G. Psychostimulant sensitization: differential changes in accumbal shell and core dopamine. *Eur J Pharmacol* 2000; 388(1): 69-76.
- [75] Cador M, Bijiou Y, Cailhol S, Stinus L. Amphetamine-induced behavioral sensitization: implication of a glutamatergic medial prefrontal cortex-ventral tegmental area innervation. *Neuroscience* 1999; 94(3): 705-721.
- [76] Cador M, Cole BJ, Koob GF, Stinus L, Le Moal M. Central administration of corticotropin releasing factor induces long-term sensitization to D-amphetamine. *Brain Res* 1993; 606(2): 181-186.
- [77] Cador M, Dulluc J, Mormede P. Modulation of the locomotor response to amphetamine by corticosterone. *Neuroscience* 1993; 56(4): 981-988.
- [78] Caine SB, Heinrichs SC, Coffin VL, Koob GF. Effects of the dopamine D-1 antagonist SCH 23390 microinjected into the accumbens, amygdala or striatum on cocaine self-administration in the rat. *Brain Res* 1995; 692(1-2): 47-56.

- [79] Caine SB, Koob GF. Effects of mesolimbic dopamine depletion on responding maintained by cocaine and food. *J Exp Anal Behav* 1994; 61(2): 213-221.
- [80] Caldji C, Diorio J, Meaney MJ. Variations in maternal care in infancy regulate the development of stress reactivity. *Biol Psychiatry* 2000; 48(12): 1164-1174.
- [81] Caldji C, Francis D, Sharma S, Plotsky PM, Meaney MJ. The effects of early rearing environment on the development of GABAA and central benzodiazepine receptor levels and novelty-induced fearfulness in the rat. *Neuropsychopharmacology* 2000; 22(3): 219-229.
- [82] Camp DM, Browman KE, Robinson TE. The effects of methamphetamine and cocaine on motor behavior and extracellular dopamine in the ventral striatum of Lewis versus Fischer 344 rats. *Brain Res* 1994; 668(1-2): 180-193.
- [83] Campbell J, Spear LP. Effects of early handling on amphetamine-induced locomotor activation and conditioned place preference in the adult rat. *Psychopharmacology (Berl)* 1999; 143(2): 183-189.
- [84] Campbell UC, Carroll ME. Effects of ketoconazole on the acquisition of intravenous cocaine self-administration under different feeding conditions in rats. *Psychopharmacology (Berl)* 2001; 154(3): 311-318.
- [85] Campbell UC, Lac ST, Carroll ME. Effects of baclofen on maintenance and reinstatement of intravenous cocaine self-administration in rats. *Psychopharmacologia* 1999; 143(2): 209-214.
- [86] Cannon WB, de la Paz D. Emotional stimulation of adrenal secretion. *AJP - Legacy* 1911; 28(1): 64-70.
- [87] Capper-Loup C, Canales JJ, Kadaba N, Graybiel AM. Concurrent activation of dopamine D1 and D2 receptors is required to evoke neural and behavioral phenotypes of cocaine sensitization. *J Neurosci* 2002; 22(14): 6218-6227.
- [88] Capriles N, Rodaros D, Sorge RE, Stewart J. A role for the prefrontal cortex in stress- and cocaine-induced reinstatement of cocaine seeking in rats. *Psychopharmacology (Berl)* 2003; 168(1-2): 66-74.
- [89] Carelli RM. Nucleus accumbens cell firing and rapid dopamine signaling during goal-directed behaviors in rats. *Neuropharmacology* 2004; 47(suppl 1): 180-189.
- [90] Carlezon WA, Jr., Nestler EJ. Elevated levels of GluR1 in the midbrain: a trigger for sensitization to drugs of abuse? *Trends Neurosci* 2002; 25(12): 610-615.
- [91] Carney JM, Landrum RW, Cheng MS, Seale TW. Establishment of chronic intravenous drug self-administration in the C57BL/6J mouse. *Neuroreport* 1991; 2(8): 477-480.
- [92] Carroll ME. The role of food deprivation in the maintenance and reinstatement of cocaine-seeking behavior in rats. *Drug Alcohol Depend* 1985; 16(2): 95-109.
- [93] Carroll ME, Lac ST, Asencio M, Kragh R. Intravenous cocaine self-administration in rats is reduced by dietary L-tryptophan. *Psychopharmacology (Berl)* 1990; 100(3): 293-300.
- [94] Cartmell LW, Aufderhide A, Weems C. Cocaine metabolites in pre-Columbian mummy hair. *J Okla State Med Assoc* 1991; 84(1): 11-12.
- [95] Castellano C, Puglisi-Allegra S. Strain-dependent modulation of memory by stress in mice. *Behav Neural Biol* 1983; 38(1): 133-138.

- [96] Catalano M, Nobile M, Novelli E, Nothen MM, Smeraldi E. Distribution of a novel mutation in the first exon of the human dopamine D4 receptor gene in psychotic patients. *Biol Psychiatry* 1993; 34(7): 459-464.
- [97] Cervo L, Cocco A, Petrella C, Heidbreder CA. Selective antagonism at dopamine D3 receptors attenuates cocaine-seeking behaviour in the rat. *Int J Neuropsychopharmacol* 2007; 10(2): 167-181.
- [98] Chao HM, Choo PH, McEwen BS. Glucocorticoid and mineralocorticoid receptor mRNA expression in rat brain. *Neuroendocrinology* 1989; 50(4): 365-371.
- [99] Chen AC, LaForge KS, Ho A, McHugh PF, Kellogg S, Bell K, Schluger RP, Leal SM, Kreek MJ. Potentially functional polymorphism in the promoter region of prodynorphin gene may be associated with protection against cocaine dependence or abuse. *Am J Med Genet* 2002; 114(4): 429-435.
- [100] Chen J, Kelz MB, Hope BT, Nakabeppu Y, Nestler EJ. Chronic Fos-related antigens: stable variants of deltaFosB induced in brain by chronic treatments. *J Neurosci* 1997; 17(13): 4933-4941.
- [101] Chen J, Nye HE, Kelz MB, Hiroi N, Nakabeppu Y, Hope BT, Nestler EJ. Regulation of delta FosB and FosB-like proteins by electroconvulsive seizure and cocaine treatments. *Mol Pharmacol* 1995; 48(5): 880-889.
- [102] Chen R, Tilley MR, Wei H, Zhou F, Zhou FM, Ching S, Quan N, Stephens RL, Hill ER, Nottoli T, Han DD, Gu HH. Abolished cocaine reward in mice with a cocaine-insensitive dopamine transporter. *Proc Natl Acad Sci U S A* 2006; 103(24): 9333-9338.
- [103] Chen YZ, Hua SY, Wang CA, Wu LG, Gu Q, Xing BR. An electrophysiological study on the membrane receptor-mediated action of glucocorticoids in mammalian neurons. *Neuroendocrinology* 1991; 53(suppl 1): 25-30.
- [104] Chevrette J, Stellar JR, Hesse GW, Markou A. Both the shell of the nucleus accumbens and the central nucleus of the amygdala support amphetamine self-administration in rats. *Pharmacol Biochem Behav* 2002; 71(3): 501-507.
- [105] Chiamulera C, Epping-Jordan MP, Zocchi A, Marcon C, Cottiny C, Tacconi S, Corsi M, Orzi F, Conquet F. Reinforcing and locomotor stimulant effects of cocaine are absent in mGluR5 null mutant mice. *Nat Neurosci* 2001; 4(9): 873-874.
- [106] Childress AR, Mozley PD, McElgin W, Fitzgerald J, Reivich M, O'Brien CP. Limbic activation during cue-induced cocaine craving. *Am J Psychiatry* 1999; 156(1): 11-18.
- [107] Chio CL, Hess GF, Graham RS, Huff RM. A second molecular form of D2 dopamine receptor in rat and bovine caudate nucleus. *Nature* 1990; 343(6255): 266-269.
- [108] Chiu J, Kalant H, Le DA. Vasopressin opposes locomotor stimulation by ethanol, cocaine and amphetamine in mice. *Eur J Pharmacol* 1998; 355(1): 11-17.
- [109] Chiueh CC, Kopin IJ. Centrally mediated release by cocaine of endogenous epinephrine and norepinephrine from the sympathoadrenal medullary system of unanesthetized rats. *J Pharmacol Exp Ther* 1978; 205(1): 148-154.
- [110] Cho K, Little HJ. Effects of corticosterone on excitatory amino acid responses in dopamine-sensitive neurons in the ventral tegmental area. *Neuroscience* 1999; 88(3): 837-845.

- [111] Chocyk A, Czyrak A, Wedzony K. Acute and repeated cocaine induces alterations in FosB/delta FosB expression in the paraventricular nucleus of the hypothalamus. *Brain Res* 2006; 1090(1): 58-68.
- [112] Christie MJ, Summers RJ, Stephenson JA, Cook CJ, Beart PM. Excitatory amino acid projections to the nucleus accumbens septi in the rat: a retrograde transport study utilizing D[3H]aspartate and [3H]GABA. *Neuroscience* 1987; 22(2): 425-439.
- [113] Chuluyan HE, Saphier D, Rohn WM, Dunn AJ. Noradrenergic innervation of the hypothalamus participates in adrenocortical responses to interleukin-1. *Neuroendocrinology* 1992; 56(1): 106-111.
- [114] Ciamei A, Cestari V, Castellano C. Strain-dependent interactions between MK-801 and cocaine on retention of C57BL/6 and DBA/2 mice tested in a one-trial inhibitory avoidance task: involvement of dopaminergic mechanisms. *Neurobiol Learn Mem* 2000; 73(2): 188-194.
- [115] Ciccocioppo R, Sanna PP, Weiss F. Cocaine-predictive stimulus induces drug-seeking behavior and neural activation in limbic brain regions after multiple months of abstinence: reversal by D(1) antagonists. *Proc Natl Acad Sci U S A* 2001; 98(4): 1976-1981.
- [116] Coirini H, Magarinos AM, De Nicola AF, Rainbow TC, McEwen BS. Further studies of brain aldosterone binding sites employing new mineralocorticoid and glucocorticoid receptor markers in vitro. *Brain Res* 1985; 361(1-2): 212-216.
- [117] Colby CR, Whisler K, Steffen C, Nestler EJ, Self DW. Striatal cell type-specific overexpression of delta FosB enhances incentive for cocaine. *J Neurosci* 2003; 23(6): 2488-2493.
- [118] Cole BJ, Koob GF. Low doses of corticotropin-releasing factor potentiate amphetamine-induced stereotyped behavior. *Psychopharmacology (Berl)* 1989; 99(1): 27-33.
- [119] Cole BJ, Cador M, Stinus L, Rivier J, Vale W, Koob GF, Le Moal M. Central administration of a CRF antagonist blocks the development of stress-induced behavioral sensitization. *Brain Res* 1990; 512(2): 343-346.
- [120] Comings DE, Gonzalez N, Wu S, Saucier G, Johnson P, Verde R, MacMurray JP. Homozygosity at the dopamine DRD3 receptor gene in cocaine dependence. *Mol Psychiatry* 1999; 4(5): 484-487.
- [121] Comings DE, Muhleman D, Ahn C, Gysin R, Flanagan SD. The dopamine D2 receptor gene: a genetic risk factor in substance abuse. *Drug Alcohol Depend* 1994; 34(3): 175-180.
- [122] Comings DE, Muhleman D, Gade R, Johnson P, Verde R, Saucier G, MacMurray J. Cannabinoid receptor gene (CNR1): association with i.v. drug use. *Mol Psychiatry* 1997; 2(2): 161-168.
- [123] Comings DE, Muhleman D, Gysin R. Dopamine D2 receptor (DRD2) gene and susceptibility to posttraumatic stress disorder: a study and replication. *Biol Psychiatry* 1996; 40(5): 368-372.
- [124] Conversi D, Orsini C, Cabib S. Distinct patterns of Fos expression induced by systemic amphetamine in the striatal complex of C57BL/6JCo and DBA/2JCo inbred strains of mice. *Brain Res* 2004; 1025(1-2): 59-66.
- [125] Coolen MW, van Loo KM, van Bakel NN, Pulford DJ, Serneels L, De Strooper B, Ellenbroek BA, Cools AR, Martens GJ. Gene dosage effect on gamma-secretase component Aph-1b in a rat model for neurodevelopmental disorders. *Neuron* 2005; 45(4): 497-503.

- [126] Cools AR, Brachten R, Heeren D, Willemsen A, Ellenbroek B. Search after neurobiological profile of individual-specific features of wistar rats. *Brain Res Bull* 1990; 24(1): 49-69.
- [127] Cools AR, Ellenbroek BA, Gingras MA, Engbersen A, Heeren D. Differences in vulnerability and susceptibility to dexamphetamine in Nijmegen high and low responders to novelty: a dose-effect analysis of spatio-temporal programming of behaviour. *Psychopharmacology (Berl)* 1997; 132(2): 181-187.
- [128] Cornish JL, Duffy P, Kalivas PW. A role for nucleus accumbens glutamate transmission in the relapse to cocaine-seeking behavior. *Neuroscience* 1999; 93(4): 1359-1367.
- [129] Cornish JL, Kalivas PW. Glutamate transmission in the nucleus accumbens mediates relapse in cocaine addiction. *J Neurosci* 2000; 20(15): 89RC.
- [130] Cornish JL, Kalivas PW. Repeated cocaine administration into the rat ventral tegmental area produces behavioral sensitization to a systemic cocaine challenge. *Behav Brain Res* 2001; 126(1-2): 205-209.
- [131] Covington HE, III, Miczek KA. Repeated social-defeat stress, cocaine or morphine. Effects on behavioral sensitization and intravenous cocaine self-administration "binges". *Psychopharmacology (Berl)* 2001; 158(4): 388-398.
- [132] Crabbe JC. Genetic contributions to addiction. *Annu Rev Psychol* 2002; 53(1): 435-462.
- [133] Cubells JF, Kranzler HR, McCance-Katz E, Anderson GM, Malison RT, Price LH, Gelernter J. A haplotype at the DBH locus, associated with low plasma dopamine beta-hydroxylase activity, also associates with cocaine-induced paranoia. *Mol Psychiatry* 2000; 5(1): 56-63.
- [134] Cunningham ET, Jr., Bohn MC, Sawchenko PE. Organization of adrenergic inputs to the paraventricular and supraoptic nuclei of the hypothalamus in the rat. *J Comp Neurol* 1990; 292(4): 651-667.
- [135] Cunningham ET, Jr., Sawchenko PE. Anatomical specificity of noradrenergic inputs to the paraventricular and supraoptic nuclei of the rat hypothalamus. *J Comp Neurol* 1988; 274(1): 60-76.
- [136] Czyrak A, Wedzony K, Michalska B, Fijal K, Dziedzicka-Wasylewska M, Mackowiak M. The corticosterone synthesis inhibitor metyrapone decreases dopamine D1 receptors in the rat brain. *Neuroscience* 1997; 79(2): 489-495.
- [137] Dahl JP, Kampman KM, Oslin DW, Weller AE, Lohoff FW, Ferraro TN, O'Brien CP, Berrettini WH. Association of a polymorphism in the Homer1 gene with cocaine dependence in an African American population. *Psychiat Genet* 2005; 15(4): 277-283.
- [138] Dahlstroem A, Fuxe K. Evidence for the existence of monoamine-containing neurons in the central nervous system. I. Demonstration of monoamines in the cell bodies of brain stem neurons. *Acta Physiol Scand* 1964; 62(suppl 232): 1-55.
- [139] Dal Toso R, Sommer B, Ewert M, Herb A, Pritchett DB, Bach A, Shivers BD, Seeburg PH. The dopamine D2 receptor: two molecular forms generated by alternative splicing. *EMBO J* 1989; 8(13): 4025-4034.
- [140] Damianopoulos EN, Carey RJ. Evidence for N-methyl-aspartate receptor mediation of cocaine induced corticosterone release and cocaine conditioned stimulant effects. *Behav Brain Res* 1995; 68(2): 219-228.

- [141] Darracq L, Blanc G, Glowinski J, Tassin JP. Importance of the noradrenaline-dopamine coupling in the locomotor activating effects of D-amphetamine. *J Neurosci* 1998; 18(7): 2729-2739.
- [142] Das G. Cocaine and the cardiovascular system. *Can J Cardiol* 1990; 6(9): 411-415.
- [143] Davis WM, Smith SG, Khalsa JH. Noradrenergic role in the self-administration of morphine or amphetamine. *Pharmacol Biochem Behav* 1975; 3(3): 477-484.
- [144] De Jong IEM, De Kloet ER. Glucocorticoids and vulnerability to psychostimulant drugs: toward substrate and mechanism. *Ann N Y Acad Sci* 2004; 1018: 192-198.
- [145] de Jong IEM, Oitzl MS, de Kloet ER. Adrenalectomy prevents behavioural sensitisation of mice to cocaine in a genotype-dependent manner. *Behav Brain Res* 2007; 177(2): 329-339.
- [146] De Kloet ER, Joels M, Holsboer F. Stress and the brain: from adaptation to disease. *Nat Rev Neurosci* 2005; 6(6): 463-475.
- [147] De Kloet ER, Oitzl MS, Joels M. Functional implications of brain corticosteroid receptor diversity. *Cell Mol Neurobiol* 1993; 13(4): 433-455.
- [148] De Kloet ER, Reul JM. Feedback action and tonic influence of corticosteroids on brain function: a concept arising from the heterogeneity of brain receptor systems. *Psychoneuroendocrinology* 1987; 12(2): 83-105.
- [149] De Kloet ER, Vreugdenhil E, Oitzl MS, Joels M. Brain corticosteroid receptor balance in health and disease. *Endocr Rev* 1998; 19(3): 269-301.
- [150] De Souza EB. Corticotropin-releasing factor receptors: physiology, pharmacology, biochemistry and role in central nervous system and immune disorders. *Psychoneuroendocrinology* 1995; 20(8): 789-819.
- [151] De Vries TJ, Cools AR, Shippenberg TS. Infusion of a D-1 receptor agonist into the nucleus accumbens enhances cocaine-induced behavioural sensitization. *Neuroreport* 1998; 9(8): 1763-1768.
- [152] De Vries TJ, Schoffelmeer AN, Binnekade R, Raaso H, Vanderschuren LJ. Relapse to cocaine- and heroin-seeking behavior mediated by dopamine D2 receptors is time-dependent and associated with behavioral sensitization. *Neuropsychopharmacology* 2002; 26(1): 18-26.
- [153] De Vries TJ, Schoffelmeer AN, Tjon GH, Nestby P, Mulder AH, Vanderschuren LJ. Mifepristone prevents the expression of long-term behavioural sensitization to amphetamine. *Eur J Pharmacol* 1996; 307(2): R3-R4.
- [154] de Vries TJ, Schoffelmeer ANM, Binnekade R, Mulder AH, Vanderschuren LJM. Drug-induced reinstatement of heroin- and cocaine-seeking behaviour following long-term extinction is associated with expression of behavioural sensitization. *Eur J Neurosci* 1998; 10(11): 3565-3571.
- [155] de Vry J, Donselaar I, Van Ree JM. Effects of desglycinamide<sup>9</sup>, (Arg<sup>8</sup>) vasopressin and vasopressin antiserum on the acquisition of intravenous cocaine self-administration in the rat. *Life Sci* 1988; 42(26): 2709-2715.
- [156] de Waele JP, Gianoulakis C. Enhanced activity of the brain beta-endorphin system by free-choice ethanol drinking in C57BL/6 but not DBA/2 mice. *Eur J Pharmacol* 1994; 258(1-2): 119-129.
- [157] De Waele JP, Gianoulakis C. Characterization of the mu and delta opioid receptors in the brain

- of the C57BL/6 and DBA/2 mice, selected for their differences in voluntary ethanol consumption. *Alcohol Clin Exp Res* 1997; 21(4): 754-762.
- [158] De Wit H, Uhlenhuth EH, Johanson CE. Individual differences in the reinforcing and subjective effects of amphetamine and diazepam. *Drug Alcohol Depend* 1986; 16(4): 341-360.
- [159] Dearth A, Gingrich JA, Falardeau P, Fremeau RT, Jr., Bates MD, Caron MG. Molecular cloning and expression of the gene for a human D1 dopamine receptor. *Nature* 1990; 347(6288): 72-76.
- [160] DeBattista C, Belanoff J, Glass S, Khan A, Horne RL, Blasey C, Carpenter LL, Alva G. Mifepristone versus placebo in the treatment of psychosis in patients with psychotic major depression. *Biol Psychiatry* 2006; 60(12): 1343-1349.
- [161] Der-Avakian A, Bland ST, Schmid MJ, Watkins LR, Spencer RL, Maier SF. The role of glucocorticoids in the uncontrollable stress-induced potentiation of nucleus accumbens shell dopamine and conditioned place preference responses to morphine. *Psychoneuroendocrinology* 2006; 31(5): 653-663.
- [162] Deroche V, Le Moal M, Piazza PV. Cocaine self-administration increases the incentive motivational properties of the drug in rats. *Eur J Neurosci* 1999; 11(8): 2731-2736.
- [163] Deroche V, Marinelli M, Le Moal M, Piazza PV. Glucocorticoids and behavioral effects of psychostimulants. II: cocaine intravenous self-administration and reinstatement depend on glucocorticoid levels. *J Pharmacol Exp Ther* 1997; 281(3): 1401-1407.
- [164] Deroche V, Marinelli M, Maccari S, Le Moal M, Simon H, Piazza PV. Stress-induced sensitization and glucocorticoids. I. Sensitization of dopamine-dependent locomotor effects of amphetamine and morphine depends on stress-induced corticosterone secretion. *J Neurosci* 1995; 15(11): 7181-7188.
- [165] Deroche V, Piazza PV, Maccari S, Le Moal M, Simon H. Repeated corticosterone administration sensitizes the locomotor response to amphetamine. *Brain Res* 1992; 584(1-2): 309-313.
- [166] Deroche V, Piazza PV, Casolini P, Le Moal M, Simon H. Sensitization to the psychomotor effects of amphetamine and morphine induced by food restriction depends on corticosterone secretion. *Brain Res* 1993; 611(2): 352-356.
- [167] Deroche-Gamonet V, Belin D, Piazza PV. Evidence for addiction-like behavior in the rat. *Science* 2004; 305(5686): 1014-1017.
- [168] Deroche-Gamonet V, Sillaber I, Aouizerate B, Izawa R, Jaber M, Ghozland S, Kellendonk C, Le Moal M, Spanagel R, Schutz G, Tronche F, Piazza PV. The glucocorticoid receptor as a potential target to reduce cocaine abuse. *J Neurosci* 2003; 23(11): 4785-4790.
- [169] DeVries AC, Taymans SE, Sundstrom JM, Pert A. Conditioned release of corticosterone by contextual stimuli associated with cocaine is mediated by corticotropin-releasing factor. *Brain Res* 1998; 786(1-2): 39-46.
- [170] Di Chiara G, Imperato A. Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci U S A* 1988; 85(14): 5274-5278.
- [171] Di Ciano P, Coury A, Depoortere RY, Egilmez Y, Lane JD, Emmett-Oglesby MW, Lepiane FG, Phillips AG, Blaha CD. Comparison of changes in extracellular dopamine concentrations in the

- nucleus accumbens during intravenous self-administration of cocaine or d-amphetamine. *Behav Pharmacol* 1995; 6(4): 311-322.
- [172] Di Ciano P, Everitt BJ. Dissociable effects of antagonism of NMDA and AMPA/KA receptors in the nucleus accumbens core and shell on cocaine-seeking behavior. *Neuropsychopharmacology* 2001; 25(3): 341-360.
- [173] Di Ciano P, Everitt BJ. The GABA(B) receptor agonist baclofen attenuates cocaine- and heroin-seeking behavior by rats. *Neuropsychopharmacology* 2003; 28(3): 510-518.
- [174] Di Paolo T, Rouillard C, Morissette M, Levesque D, Bedard PJ. Endocrine and neurochemical actions of cocaine. *Can J Physiol Pharmacol* 1989; 67(9): 1177-1181.
- [175] Di S, Malcher-Lopes R, Marcheselli VL, Bazan NG, Tasker JG. Rapid glucocorticoid-mediated endocannabinoid release and opposing regulation of glutamate and gamma-aminobutyric acid inputs to hypothalamic magnocellular neurons. *Endocrinology* 2005; 146(10): 4292-4301.
- [176] Di S, Malcher-Lopes R, Halmos KC, Tasker JG. Nongenomic glucocorticoid inhibition via endocannabinoid release in the hypothalamus: a fast feedback mechanism. *J Neurosci* 2003; 23(12): 4850-4857.
- [177] Diaz J, Levesque D, Griffon N, Lammers CH, Martres MP, Sokoloff P, Schwartz JC. Opposing roles for dopamine D2 and D3 receptors on neurotensin mRNA expression in nucleus accumbens. *Eur J Neurosci* 1994; 6(8): 1384-1387.
- [178] Diaz J, Levesque D, Lammers CH, Griffon N, Martres M-P, Schwartz J-C, Sokoloff P. Phenotypical characterization of neurons expressing the dopamine D3 receptor in the rat brain. *Neuroscience* 1995; 65(3): 731-745.
- [179] Dietz DM, Tapocik J, Gaval-Cruz M, Kabbaj M. Dopamine transporter, but not tyrosine hydroxylase, may be implicated in determining individual differences in behavioral sensitization to amphetamine. *Physiol Behav* 2005; 86(3): 347-355.
- [180] Doherty MD, Gratton A. Effects of medial prefrontal cortical injections of GABA receptor agonists and antagonists on the local and nucleus accumbens dopamine responses to stress. *Synapse* 1999; 32(4): 288-300.
- [181] Douglass J, McKinzie AA, Couceyro P. PCR differential display identifies a rat brain mRNA that is transcriptionally regulated by cocaine and amphetamine. *J Neurosci* 1995; 15(3 Pt 2): 2471-2481.
- [182] Drouin C, Blanc G, Villegier AS, Glowinski J, Tassin JP. Critical role of alpha1-adrenergic receptors in acute and sensitized locomotor effects of D-amphetamine, cocaine, and GBR 12783: influence of preexposure conditions and pharmacological characteristics. *Synapse* 2002; 43(1): 51-61.
- [183] Drouin C, Darracq L, Trovero F, Blanc G, Glowinski J, Cotecchia S, Tassin JP. Alpha1b-adrenergic receptors control locomotor and rewarding effects of psychostimulants and opiates. *J Neurosci* 2002; 22(7): 2873-2884.
- [184] Drouin J, Sun YL, Chamberland M, Gauthier Y, De Lean A, Nemer M, Schmidt TJ. Novel glucocorticoid receptor complex with DNA element of the hormone-repressed POMC gene. *The EMBO Journal* 1993; 12(1): 145-156.

- [185] Du C, Yu M, Volkow ND, Koretsky AP, Fowler JS, Benveniste H. Cocaine increases the intracellular calcium concentration in brain independently of its cerebrovascular effects. *J Neurosci* 2006; 26(45): 11522-11531.
- [186] Dunn AJ. Stress-related changes in cerebral catecholamine and indoleamine metabolism: lack of effect of adrenalectomy and corticosterone. *J Neurochem* 1988; 51(2): 406-412.
- [187] Dunn JM, Inderwies BR, Licata SC, Pierce RC. Repeated administration of AMPA or a metabotropic glutamate receptor agonist into the rat ventral tegmental area augments the subsequent behavioral hyperactivity induced by cocaine. *Psychopharmacology (Berl)* 2005; 179(1): 172-180.
- [188] Ebstein RP, Novick O, Umansky R, Priel B, Osher Y, Blaine D, Bennett ER, Nemanov L, Katz M, Belmaker RH. Dopamine D4 receptor (D4DR) exon III polymorphism associated with the human personality trait of Novelty Seeking. *Nat Genet* 1996; 12(1): 78-80.
- [189] Edwards CR, Stewart PM, Burt D, Brett L, McIntyre MA, Sutanto WS, De Kloet ER, Monder C. Localisation of 11 beta-hydroxysteroid dehydrogenase - tissue specific protector of the mineralocorticoid receptor. *Lancet* 1988; 2(8618): 986-989.
- [190] Ellenbroek BA. Animal models in the genomic era: possibilities and limitations with special emphasis on schizophrenia. *Behav Pharmacol* 2003; 14(5-6): 409-417.
- [191] Ellenbroek BA, van der Kam EL, van der Elst MCJ, Cools AR. Individual differences in drug dependence in rats: the role of genetic factors and life events. *Eur J Pharmacol* 2005; 526(1-3): 251-258.
- [192] Elman I, Lukas SE, Karlsgodt KH, Gasic GP, Breiter HC. Acute cortisol administration triggers craving in individuals with cocaine dependence. *Psychopharmacol Bull* 2003; 37(3): 84-89.
- [193] Erb S, Hitchcott PK, Rajabi H, Mueller D, Shaham Y, Stewart J. Alpha-2 adrenergic receptor agonists block stress-induced reinstatement of cocaine seeking. *Neuropsychopharmacology* 2000; 23(2): 138-150.
- [194] Erb S, Shaham Y, Stewart J. Stress reinstates cocaine-seeking behavior after prolonged extinction and a drug-free period. *Psychopharmacology (Berl)* 1996; 128(4): 408-412.
- [195] Erb S, Shaham Y, Stewart J. The role of corticotropin-releasing factor and corticosterone in stress- and cocaine-induced relapse to cocaine seeking in rats. *J Neurosci* 1998; 18(14): 5529-5536.
- [196] Erb S, Brown ZJ. A role for corticotropin-releasing factor in the long-term expression of behavioral sensitization to cocaine. *Behav Brain Res* 2006; 172(2): 360-364.
- [197] Erb S, Stewart J. A role for the bed nucleus of the stria terminalis, but not the amygdala, in the effects of corticotropin-releasing factor on stress-induced reinstatement of cocaine seeking. *J Neurosci* 1999; 19(20): 35RC.
- [198] Erbllich J, Lerman C, Self DW, Diaz GA, Bovbjerg DH. Stress-induced cigarette craving: effects of the DRD2 TaqI RFLP and SLC6A3 VNTR polymorphisms. *Pharmacogenomics J* 2004; 4(2): 102-109.
- [199] Erwin VG, Womer DE, Campbell AD, Jones BC. Pharmacogenetics of cocaine: II. Mesocorticolimbic and striatal dopamine and cocaine receptors in C57BL and DBA mice. *Pharmacogenetics* 1993; 3(4): 189-196.
- [200] Escorihuela RM, Fernandez-Teruel A, Gil L, Aguilar R, Tobena A, Driscoll P. Inbred Roman

- high- and low-avoidance rats: differences in anxiety, novelty-seeking, and shuttlebox behaviors. *Physiol Behav* 1999; 67(1): 19-26.
- [201] Everitt BJ, Cardinal RN, Parkinson JA, Robbins TW. Appetitive behavior: impact of amygdala-dependent mechanisms of emotional learning. *Ann N Y Acad Sci* 2003; 985(1): 233-250.
- [202] Everitt BJ, Wolf ME. Psychomotor stimulant addiction: a neural systems perspective. *J Neurosci* 2002; 22(9): 3312-3320.
- [203] Exner M, Clark D. Behaviour in the novel environment predicts responsiveness to d-amphetamine in the rat: a multivariate approach. *Behav Pharmacol* 1993; 4(1): 47-56.
- [204] Fadda P, Scherma M, Fresu A, Collu M, Fratta W. Baclofen antagonizes nicotine-, cocaine-, and morphine-induced dopamine release in the nucleus accumbens of rat. *Synapse* 2003; 50(1): 1-6.
- [205] Falck RS, Wang J, Siegal HA, Carlson RG. The prevalence of psychiatric disorder among a community sample of crack cocaine users: an exploratory study with practical implications. *J Nerv Ment Dis* 2004; 192(7): 503-507.
- [206] Feldman S, Weidenfeld J. Involvement of amygdalar alpha adrenoceptors in hypothalamo-pituitary-adrenocortical responses. *Neuroreport* 1996; 7(18): 3055-3057.
- [207] Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, Koss MP, Marks JS. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the adverse childhood experiences (ACE) study. *Am J Prev Med* 1998; 14(4): 245-258.
- [208] Fierens E. Archeological and artistic sources of coca consumption in pre-hispanic America. *Verh K Acad Geneeskd Belg* 1991; 53(5): 463-485.
- [209] Filip M, Frankowska M, Zaniewska M, Golda A, Przegalinski E. The serotonergic system and its role in cocaine addiction. *Pharmacol Rep* 2005; 57(6): 685-700.
- [210] Filip M, Siwanowicz J. Implication of the nucleus accumbens shell, but not core, in the acute and sensitizing effects of cocaine in rats. *Pol J Pharmacol* 2001; 53(5): 459-466.
- [211] Filip M, Thomas ML, Cunningham KA. Dopamine D5 receptors in nucleus accumbens contribute to the detection of cocaine in rats. *J Neurosci* 2000; 20(19): RC98.
- [212] Fink JS, Reis DJ. Genetic variations in midbrain dopamine cell number: parallel with differences in responses to dopaminergic agonists and in naturalistic behaviors mediated by central dopaminergic systems. *Brain Res* 1981; 222(2): 335-349.
- [213] Fletcher PJ, Korth KM, Chambers JW. Depletion of brain serotonin following intra-raphé injections of 5,7-dihydroxytryptamine does not alter d-amphetamine self-administration across different schedule and access conditions. *Psychopharmacology (Berl)* 1999; 146(2): 185-193.
- [214] Flores G, Wood GK, Barbeau D, Quirion R, Srivastava LK. Lewis and Fischer rats: a comparison of dopamine transporter and receptors levels. *Brain Res* 1998; 814(1-2): 34-40.
- [215] Floresco SB, Yang CR, Phillips AG, Blaha CD. Basolateral amygdala stimulation evokes glutamate receptor-dependent dopamine efflux in the nucleus accumbens of the anaesthetized rat. *Eur J Neurosci* 1998; 10(4): 1241-1251.
- [216] Floresco SB, Todd CL, Grace AA. Glutamatergic afferents from the hippocampus to the nucleus

- accumbens regulate activity of ventral tegmental area dopamine neurons. *J Neurosci* 2001; 21(13): 4915-4922.
- [217] Follenius M, Simon C, Brandenberger G, Lenzi P. Ultradian plasma corticotropin and cortisol rhythms: time-series analyses. *J Endocrinol Invest* 1987; 10(3): 261-266.
- [218] Forray MI, Gysling K. Role of noradrenergic projections to the bed nucleus of the stria terminalis in the regulation of the hypothalamic-pituitary-adrenal axis. *Brain Res Brain Res Rev* 2004; 47(1-3): 145-160.
- [219] Francis DD, Meaney MJ. Maternal care and the development of stress responses. *Curr Opin Neurobiol* 1999; 9(1): 128-134.
- [220] Franklin KB, Paxinos G. The mouse brain in stereotaxic coordinates. Academic Press Inc, San Diego: 1997.
- [221] Freeman WM, Brebner K, Lynch WJ, Robertson DJ, Roberts DC, Vrana KE. Cocaine-responsive gene expression changes in rat hippocampus. *Neuroscience* 2001; 108(3): 371-380.
- [222] Fremeau RT, Jr., Duncan GE, Fornaretto M, Dearth A, Gingrich JA. Localization of D1 dopamine receptor mRNA in brain supports a role in cognitive, affective, and neuroendocrine aspects of dopaminergic neurotransmission. *Proc Natl Acad Sci U S A* 1991; 88(9): 3772-3776.
- [223] Funder JW, Pearce PT, Smith R, Smith AI. Mineralocorticoid action: target tissue specificity is enzyme, not receptor, mediated. *Science* 1988; 242(4878): 583-585.
- [224] Fuxe K, Hokfelt T. Further evidence for the existence of tubero-infundibular dopamine neurons. *Acta Physiol Scand* 1966; 66(1): 245-246.
- [225] Fuxe K, Wikstrom AC, Okret S, Agnati LF, Harfstrand A, Yu ZY, Granholm L, Zoli M, Vale W, Gustafsson. Mapping of glucocorticoid receptor immunoreactive neurons in the rat tel- and diencephalon using a monoclonal antibody against rat liver glucocorticoid receptor. *Endocrinology* 1985; 117(5): 1803-1812.
- [226] Gaillet S, Alonso G, Le Borgne R, Barbanel G, Malaval F, Assenmacher I, Szafarczyk A. Effects of discrete lesions in the ventral noradrenergic ascending bundle on the corticotropic stress response depend on the site of the lesion and on the plasma levels of adrenal steroids. *Neuroendocrinology* 1993; 58(4): 408-419.
- [227] Gal K, Gyertyan I. Dopamine D3 as well as D2 receptor ligands attenuate the cue-induced cocaine-seeking in a relapse model in rats. *Drug Alcohol Depend* 2006; 81(1): 63-70.
- [228] Gamaro GD, Denardin JD, Jr., Michalowski MB, Catelli D, Correa JB, Xavier MH, Dalmaz C. Epinephrine effects on memory are not dependent on hepatic glucose release. *Neurobiol Learn Mem* 1997; 68(3): 221-229.
- [229] Garavan H, Pankiewicz J, Bloom A, Cho JK, Sperry L, Ross TJ, Salmeron BJ, Risinger R, Kelley D, Stein EA. Cue-Induced cocaine craving: neuroanatomical specificity for drug users and drug stimuli. *Am J Psychiat* 2000; 157(11): 1789-1798.
- [230] Gelernter J, Kranzler HR, Satel SL, Rao PA. Genetic association between dopamine transporter protein alleles and cocaine-induced paranoia. *Neuropsychopharmacology* 1994; 11(3): 195-200.
- [231] Gelernter J, Kranzler H, Satel SL. No association between D2 dopamine receptor (DRD2) alleles

- or haplotypes and cocaine dependence or severity of cocaine dependence in European- and African-Americans. *Biol Psychiatry* 1999; 45(3): 340-345.
- [232] George FR, Goldberg SR. Genetic approaches to the analysis of addiction processes. *Trends Pharmacol Sci* 1989; 10(2): 78-83.
- [233] Gerfen CR, Engber TM, Mahan LC, Susel Z, Chase TN, Monsma FJ, Jr., Sibley DR. D1 and D2 dopamine receptor-regulated gene expression of striatonigral and striatopallidal neurons. *Science* 1990; 250(4986): 1429-1432.
- [234] Gerrits MA, Petromilli P, Westenberg HG, Di Chiara G, Van Ree JM. Decrease in basal dopamine levels in the nucleus accumbens shell during daily drug-seeking behaviour in rats. *Brain Res* 2002; 924(2): 141-150.
- [235] Gerrits MAFM, Van Ree JM. Effect of nucleus accumbens dopamine depletion on motivational aspects involved in initiation of cocaine and heroin self-administration in rats. *Brain Res* 1996; 713(1-2): 114-124.
- [236] Giorgi O, Corda MG, Carboni G, Frau V, Valentini V, Di Chiara G. Effects of cocaine and morphine in rats from two psychogenetically selected lines: a behavioral and brain dialysis study. *Behav Genet* 1997; 27(6): 537-546.
- [237] Giorgi O, Piras G, Lecca D, Corda MG. Behavioural effects of acute and repeated cocaine treatments: a comparative study in sensitisation-prone RHA rats and their sensitisation-resistant RLA counterparts. *Psychopharmacology (Berl)* 2005; 180(3): 530-538.
- [238] Giros B, Sokoloff P, Martres MP, Riou JF, Emorine LJ, Schwartz JC. Alternative splicing directs the expression of two D2 dopamine receptor isoforms. *Nature* 1989; 342(6252): 923-926.
- [239] Glenthøj BY, Hemmingsen R. Dopaminergic sensitization: implications for the pathogenesis of schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 1997; 21(1): 23-46.
- [240] Goeders NE. The HPA axis and cocaine reinforcement. *Psychoneuroendocrinology* 2002; 27(1-2): 13-33.
- [241] Goeders NE. Stress and cocaine addiction. *J Pharmacol Exp Ther* 2002; 301(3): 785-789.
- [242] Goeders NE, Dworkin SI, Smith JE. Neuropharmacological assessment of cocaine self-administration into the medial prefrontal cortex. *Pharmacol Biochem Behav* 1986; 24(5): 1429-1440.
- [243] Goeders NE, Guerin GF. Pretreatment with corticosterone or dexamethasone fails to affect ongoing cocaine self-administration in rats. *Neuroscience abstracts* 2006; 25: 1872.
- [244] Goeders NE, Guerin GF. Non-contingent electric footshock facilitates the acquisition of intravenous cocaine self-administration in rats. *Psychopharmacology (Berl)* 1994; 114(1): 63-70.
- [245] Goeders NE, Guerin GF. Effects of surgical and pharmacological adrenalectomy on the initiation and maintenance of intravenous cocaine self-administration in rats. *Brain Res* 1996; 722(1-2): 145-152.
- [246] Goeders NE, Peltier RL, Guerin GF. Ketoconazole reduces low dose cocaine self-administration in rats. *Drug Alcohol Depend* 1998; 53(1): 67-77.
- [247] Goeders NE, Smith JE. Cortical dopaminergic involvement in cocaine reinforcement. *Science* 1983; 221(4612): 773-775.
- [248] Goeders NE, Smith JE. Reinforcing properties of cocaine in the medial prefrontal cortex: primary

- action on presynaptic dopaminergic terminals. *Pharmacol Biochem Behav* 1986; 25(1): 191-199.
- [249] Goeders NE. The impact of stress on addiction. *Eur Neuropsychopharmacol* 2003; 13(6): 435-441.
- [250] Goeders NE, Guerin GF. Effects of the CRH receptor antagonist CP-154,526 on intravenous cocaine self-administration in rats. *Neuropsychopharmacology* 2000; 23(5): 577-586.
- [251] Gold PE, van Buskirk R. Posttraining brain norepinephrine concentrations: correlation with retention performance of avoidance training and with peripheral epinephrine modulation of memory processing. *Behav Biol* 1978; 23(4): 509-520.
- [252] Gold PE, Van Buskirk RB. Facilitation of time-dependent memory processes with posttrial epinephrine injections. *Behav Biol* 1975; 13(2): 145-153.
- [253] Gold PE, Vogt J, Hall JL. Glucose effects on memory: behavioral and pharmacological characteristics. *Behav Neural Biol* 1986; 46(2): 145-155.
- [254] Goldman D, Oroszi G, Ducci F. The genetics of addictions: uncovering the genes. *Nat Rev Genet* 2005; 6(7): 521-532.
- [255] Gomez F, de Kloet ER, Armario A. Glucocorticoid negative feedback on the HPA axis in five inbred rat strains. *AJP - Reg Integr Comp Physiol* 1998; 274(2): R420-R427.
- [256] Gorelova N, Yang CR. The course of neural projection from the prefrontal cortex to the nucleus accumbens in the rat. *Neuroscience* 1997; 76(3): 689-706.
- [257] Grahame NJ, Cunningham CL. Genetic differences in intravenous cocaine self-administration between C57BL/6J and DBA/2J mice. *Psychopharmacology (Berl)* 1995; 122(3): 281-291.
- [258] Grandy DK, Zhang Y, Bouvier C, Zhou Q, Johnson RA, Allen L, Buck K, Bunzow JR, Salon J, Civelli O. Multiple human D5 dopamine receptor genes: a functional receptor and two pseudo-genes. *Proc Natl Acad Sci U S A* 1991; 88(20): 9175-9179.
- [259] Graybiel AM, Aosaki T, Flaherty AW, Kimura M. The basal ganglia and adaptive motor control. *Science* 1994; 265(5180): 1826-1831.
- [260] Graybiel AM, Hirsch EC, Agid Y. The nigrostriatal system in Parkinson's disease. *Adv Neurol* 1990; 53: 17-29.
- [261] Greenberg DA, Prichard DC, Snyder SH. Alpha-noradrenergic receptor binding in mammalian brain: differential labeling of agonist and antagonist states. *Life Sci* 1976; 19(1): 69-76.
- [262] Gritti A, Bove D, Di Sarno AM, D'Addio AA, Chiapparato S, Bove RM. Stereotyped movements in a group of autistic children. *Funct Neurol* 2003; 18(2): 89-94.
- [263] Groenewegen HJ, Russchen FT. Organization of the efferent projections of the nucleus accumbens to pallidal, hypothalamic, and mesencephalic structures: a tracing and immunohistochemical study in the cat. *J Comp Neurol* 1984; 223(3): 347-367.
- [264] Groenewegen HJ, Vermeulen-Van der Zee E, te KA, Witter MP. Organization of the projections from the subiculum to the ventral striatum in the rat. A study using anterograde transport of Phaseolus vulgaris leucoagglutinin. *Neuroscience* 1987; 23(1): 103-120.
- [265] Grootendorst J, De Kloet ER, Vossen C, Dalm S, Oitzl MS. Repeated exposure to rats has persis-

- tent genotype-dependent effects on learning and locomotor activity of apolipoprotein E knockout and C57Bl/6 mice. *Behav Brain Res* 2001; 125(1-2): 249-259.
- [266] Grootendorst J, Kempes MM, Lucassen PJ, Dalm S, De Kloet ER, Oitzl MS. Differential effect of corticosterone on spatial learning abilities in apolipoprotein E knockout and C57Bl/6 mice. *Brain Res* 2002; 953(1-2): 281-285.
- [267] Guindalini C, Howard M, Haddley K, Laranjeira R, Collier D, Ammar N, Craig I, O'Gara C, Bubb V, Greenwood T, Kelsoe J, Asherson P, Murray RM, Castelo A, Quinn JP, Vallada H, Breen G. A dopamine transporter gene functional variant associated with cocaine abuse in a Brazilian sample. *Proc Natl Acad Sci U S A* 2006; 103(12): 4552-4557.
- [268] Gunnarsdottir ED, Pingitore RA, Spring BJ, Konopka LM, Crayton JW, Milo T, Shirazi P. Individual differences among cocaine users. *Addict Behav* 2000; 25(5): 641-652.
- [269] Gurkovskaya O, Goeders NE. Effects of CP-154,526 on responding during extinction from cocaine self-administration in rats. *Eur J Pharmacol* 2001; 432(1): 53-56.
- [270] Haber SN. Neurotransmitters in the human and nonhuman primate basal ganglia. *Hum Neurobiol* 1986; 5(3): 159-168.
- [271] Haber SN, Fudge JL, McFarland NR. Striatonigrostriatal pathways in primates form an ascending spiral from the shell to the dorsolateral striatum. *J Neurosci* 2000; 20(6): 2369-2382.
- [272] Haile CN, GrandPre T, Kosten TA. Chronic unpredictable stress, but not chronic predictable stress, enhances the sensitivity to the behavioral effects of cocaine in rats. *Psychopharmacology (Berl)* 2001; 154(2): 213-220.
- [273] Hain HS, Crabbe JC, Bergeson SE, Belknap JK. Cocaine-induced seizure thresholds: quantitative trait loci detection and mapping in two populations derived from the C57Bl/6 and DBA/2 mouse strains. *J Pharmacol Exp Ther* 2000; 293(1): 180-187.
- [274] Hall RC, Popkin MK, Stickney SK, Gardner ER. Presentation of the steroid psychoses. *J Nerv Ment Dis* 1979; 167(4): 229-236.
- [275] Halpain S, McEwen BS. Corticosterone decreases 3H-glutamate binding in rat hippocampal formation. *Neuroendocrinology* 1988; 48(3): 235-241.
- [276] Hammond GL. Molecular properties of corticosteroid binding globulin and the sex-steroid binding proteins. *Endocr Rev* 1990; 11(1): 65-79.
- [277] Haney M, Maccari S, Le Moal M, Simon H, Piazza PV. Social stress increases the acquisition of cocaine self-administration in male and female rats. *Brain Res* 1995; 698(1-2): 46-52.
- [278] Haney M, Hart CL, Foltin RW. Effects of baclofen on cocaine self-administration: opioid- and nonopioid-dependent volunteers. *Neuropsychopharmacology* 2006; 31(8): 1814-1821.
- [279] Harfstrand A, Fuxe K, Cintra A, Agnati LF, Zini I, Wikstrom AC, Okret S, Yu ZY, Goldstein M, Steinbusch H. Glucocorticoid receptor immunoreactivity in monoaminergic neurons of rat brain. *Proc Natl Acad Sci U S A* 1986; 83(24): 9779-9783.
- [280] Harrison PA, Fulkerson JA, Beebe TJ. Multiple substance use among adolescent physical and sexual abuse victims. *Child Abuse Neglect* 1997; 21(6): 529-539.
- [281] Heesch CM, Negus BH, Keffer JH, Snyder RW, Risser RC, Eichhorn EJ. Effects of cocaine on cortisol secretion in humans. *Am J Med Sci* 1995; 310(2): 61-64.

- [282] Heim C, Owens MJ, Plotsky PM, Nemeroff CB. Persistent changes in corticotropin-releasing factor systems due to early life stress: relationship to the pathophysiology of major depression and post-traumatic stress disorder. *Psychopharmacol Bull* 1997; 33(2): 185-192.
- [283] Hemby SE, Horman B, Tang W. Differential regulation of ionotropic glutamate receptor subunits following cocaine self-administration. *Brain Res* 2005; 1064(1-2): 75-82.
- [284] Herman JP, Stinus L, Le Moal M. Repeated stress increases locomotor response to amphetamine. *Psychopharmacology (Berl)* 1984; 84(3): 431-435.
- [285] Hermann G, Tovar CA, Beck FM, Sheridan JF. Kinetics of glucocorticoid response to restraint stress and/or experimental influenza viral infection in two inbred strains of mice. *J Neuroimmunol* 1994; 49(1-2): 25-33.
- [286] Holmes A, le Guisquet AM, Vogel E, Millstein RA, Leman S, Belzung C. Early life genetic, epigenetic and environmental factors shaping emotionality in rodents. *Neurosci Biobehav Rev* 2005; 29(8): 1335-1346.
- [287] Homberg JR, van den AM, Raaso HS, Wardeh G, Binnekade R, Schoffelmeer AN, De Vries TJ. Enhanced motivation to self-administer cocaine is predicted by self-grooming behaviour and relates to dopamine release in the rat medial prefrontal cortex and amygdala. *Eur J Neurosci* 2002; 15(9): 1542-1550.
- [288] Hooks MS, Jones GH, Hemby SE, Justice JB, Jr. Environmental and pharmacological sensitization: effects of repeated administration of systemic or intra-nucleus accumbens cocaine. *Psychopharmacology (Berl)* 1993; 111(1): 109-116.
- [289] Hooks MS, Jones GH, Liem BJ, Justice J. Sensitization and individual differences to IP amphetamine, cocaine, or caffeine following repeated intracranial amphetamine infusions. *Pharmacol Biochem Behav* 1992; 43(3): 815-823.
- [290] Hooks MS, Jones GH, Smith AD, Neill DB, Justice JB, Jr. Response to novelty predicts the locomotor and nucleus accumbens dopamine response to cocaine. *Synapse* 1991; 9(2): 121-128.
- [291] Hooks MS, Jones GH, Smith AD, Neill DB, Justice JB, Jr. Individual differences in locomotor activity and sensitization. *Pharmacol Biochem Behav* 1991; 38(2): 467-470.
- [292] Hooks MS, Juncos JL, Justice JB, Jr., Meiergerd SM, Povlock SL, Schenk JO, Kalivas PW. Individual locomotor response to novelty predicts selective alterations in D1 and D2 receptors and mRNAs. *J Neurosci* 1994; 14(10): 6144-6152.
- [293] Hooks MS, Sorg BA, Kalivas PW. The relationship between mRNA levels and the locomotor response to novelty. *Brain Res* 1994; 663(2): 312-316.
- [294] Hope BT, Nye HE, Kelz MB, Self DW, Iadarola MJ, Nakabeppu Y, Duman RS, Nestler EJ. Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and other chronic treatments. *Neuron* 1994; 13(5): 1235-1244.
- [295] Horger BA, Giles MK, Schenk S. Preexposure to amphetamine and nicotine predisposes rats to self-administer a low dose of cocaine. *Psychopharmacologia* 1992; 107(2-3): 271-276.
- [296] Horger BA, Shelton K, Schenk S. Preexposure sensitizes rats to the rewarding effects of cocaine. *Pharmacol Biochem Behav* 1990; 37(4): 707-711.
- [297] Hosoi T, Okuma Y, Nomura Y. Electrical stimulation of afferent vagus nerve induces IL-1beta

- expression in the brain and activates HPA axis. *AJP - Reg Integr Comp Physiol* 2000; 279(1): R141-R147.
- [298] Hotsenpiller G, Giorgetti M, Wolf ME. Alterations in behaviour and glutamate transmission following presentation of stimuli previously associated with cocaine exposure. *Eur J Neurosci* 2001; 14(11): 1843-1855.
- [299] Hubner CB, Moreton JE. Effects of selective D1 and D2 dopamine antagonists on cocaine self-administration in the rat. *Psychopharmacologia* 1991; 105(2): 151-156.
- [300] Hummel KP. Accessory adrenal cortical nodules in the mouse. *Anat Rec* 1958; 132(3): 281-295.
- [301] Hurd YL, McGregor A, Ponten M. In vivo amygdala dopamine levels modulate cocaine self-administration behaviour in the rat: D1 dopamine receptor involvement. *Eur J Neurosci* 1997; 9(12): 2541-2548.
- [302] Ihalainen JA, Tanila H. In vivo regulation of dopamine and noradrenaline release by alpha2A-adrenoceptors in the mouse nucleus accumbens. *J Neurochem* 2004; 91(1): 49-56.
- [303] Imperato A, Angelucci L, Casolini P, Zocchi A, Puglisi-Allegra S. Repeated stressful experiences differently affect limbic dopamine release during and following stress. *Brain Res* 1992; 577(2): 194-199.
- [304] Imperato A, Puglisi-Allegra S, Grazia SM, Casolini P, Bacchi S, Angelucci L. Cortical and limbic dopamine and acetylcholine release as neurochemical correlates of emotional arousal in both aversive and non-aversive environmental changes. *Neurochem Int* 1992; 20(suppl): 265S-270S.
- [305] Imperato A, Cabib S, Puglisi-Allegra S. Repeated stressful experiences differently affect the time-dependent responses of the mesolimbic dopamine system to the stressor. *Brain Res* 1993; 601(1-2): 333-336.
- [306] Ishihara T, Akiyama K, Kashihara K. Enhanced AP-1 binding in brain induced by D1 and D2 agonists in methamphetamine-sensitized rats. *Neuroreport* 1998; 9(17): 3913-3917.
- [307] Ito R, Dalley JW, Robbins TW, Everitt BJ. Dopamine release in the dorsal striatum during cocaine-seeking behavior under the control of a drug-associated cue. *J Neurosci* 2002; 22(14): 6247-6253.
- [308] Izawa R, Jaber M, Deroche-Gamonet V, Sillaber I, Kellendonk C, Le Moal M, Tronche F, Piazza PV. Gene expression regulation following behavioral sensitization to cocaine in transgenic mice lacking the glucocorticoid receptor in the brain. *Neuroscience* 2006; 137(3): 915-924.
- [309] Jaber M, Robinson SW, Missale C, Caron MG. Dopamine receptors and brain function. *Neuropharmacology* 1997; 35(11): 1503-1519.
- [310] Jackson A, Mead AN, Rocha BA, Stephens DN. AMPA receptors and motivation for drug: effect of the selective antagonist NBQX on behavioural sensitization and on self-administration in mice. *Behav Pharmacol* 1998; 9(5-6): 457-467.
- [311] Jackson DM, Westlind-Danielsson A. Dopamine receptors: molecular biology, biochemistry and behavioural aspects. *Pharmacol Ther* 1994; 64(2): 291-370.
- [312] Jacobsen LK, Staley JK, Malison RT, Zoghbi SS, Seibyl JP, Kosten TR, Innis RB. Elevated central

- serotonin transporter binding availability in acutely abstinent cocaine-dependent patients. *Am J Psychiatry* 2000; 157(7): 1134-1140.
- [313] Jamensky NT, Gianoulakis C. Content of dynorphins and kappa-opioid receptors in distinct brain regions of C57BL/6 and DBA/2 mice. *Alcohol Clin Exp Res* 1997; 21(8): 1455-1464.
- [314] Janowsky A, Mah C, Johnson RA, Cunningham CL, Phillips TJ, Crabbe JC, Eshleman AJ, Belknap JK. Mapping genes that regulate density of dopamine transporters and correlated behaviors in recombinant inbred mice. *J Pharmacol Exp Ther* 2001; 298(2): 634-643.
- [315] Jansen ASP, Schmidt ED, Voorn P, Tilders FJH. Substance induced plasticity in noradrenergic innervation of the paraventricular hypothalamic nucleus. *Eur J Neurosci* 2003; 17(2): 298-306.
- [316] Jentsch JD, Taylor JR. Impulsivity resulting from frontostriatal dysfunction in drug abuse: implications for the control of behavior by reward-related stimuli. *Psychopharmacology (Berl)* 1999; 146(4): 373-390.
- [317] Joels M, De Kloet ER. Control of neuronal excitability by corticosteroid hormones. *Trends Neurosci* 1992; 15(1): 25-30.
- [318] Joels M, Pu Z, Wiegert O, Oitzl MS, Krugers HJ. Learning under stress: how does it work? *Trends Cogn Sci* 2006; 10(4): 152-158.
- [319] Johnson LR, Farb C, Morrison JH, McEwen BS, LeDoux JE. Localization of glucocorticoid receptors at postsynaptic membranes in the lateral amygdala. *Neuroscience* 2005; 136(1): 289-299.
- [320] Jones BC, Hou X, Cook MN. Effect of exposure to novelty on brain monoamines in C57BL/6 and DBA/2 mice. *Physiol Behav* 1996; 59(2): 361-367.
- [321] Jones BC, Sarrieau A, Reed CL, Azar MR, Mormede P. Contribution of sex and genetics to neuroendocrine adaptation to stress in mice. *Psychoneuroendocrinology* 1998; 23(5): 505-517.
- [322] Jones GH, Mittleman G, Robbins TW. Attenuation of amphetamine-stereotypy by mesostriatal dopamine depletion enhances plasma corticosterone: implications for stereotypy as a coping response. *Behav Neural Biol* 1989; 51(1): 80-91.
- [323] Kabbaj M, Norton CS, Kollack-Walker S, Watson SJ, Robinson TE, Akil H. Social defeat alters the acquisition of cocaine self-administration in rats: role of individual differences in cocaine-taking behavior. *Psychopharmacology (Berl)* 2001; 158(4): 382-387.
- [324] Kalia M, Woodward DJ, Smith WK, Fuxe K. Rat medulla oblongata. IV. Topographical distribution of catecholaminergic neurons with quantitative three-dimensional computer reconstruction. *J Comp Neurol* 1985; 233(3): 350-364.
- [325] Kalivas PW, Churchill L, Klitenick MA. GABA and enkephalin projection from the nucleus accumbens and ventral pallidum to the ventral tegmental area. *Neuroscience* 1993; 57(4): 1047-1060.
- [326] Kalivas PW, Duffy P. Repeated cocaine administration alters extracellular glutamate in the ventral tegmental area. *J Neurochem* 1998; 70(4): 1497-1502.
- [327] Kalivas PW, Duffy P. Time course of extracellular dopamine and behavioral sensitization to cocaine. I. Dopamine axon terminals. *J Neurosci* 1993; 13(1): 266-275.
- [328] Kalivas PW, Volkow ND. The neural basis of addiction: a pathology of motivation and choice. *Am J Psychiatry* 2005; 162(8): 1403-1413.

- [329] Kalivas PW. Glutamate systems in cocaine addiction. *Curr Opin Pharmacol* 2004; 4(1): 23-29.
- [330] Kantak KM, Black Y, Valencia E, Green-Jordan K, Eichenbaum HB. Dissociable effects of lidocaine inactivation of the rostral and caudal basolateral amygdala on the maintenance and reinstatement of cocaine-seeking behavior in rats. *J Neurosci* 2002; 22(3): 1126-1136.
- [331] Kapur S, Mamo D. Half a century of antipsychotics and still a central role for dopamine D2 receptors. *Prog Neuropsychopharmacol Biol Psychiatry* 2003; 27(7): 1081-1090.
- [332] Karler R, Calder LD, Bedingfield JB. Cocaine behavioral sensitization and the excitatory amino acids. *Psychopharmacology (Berl)* 1994; 115(3): 305-310.
- [333] Karler R, Chaudhry IA, Calder LD, Turkanis SA. Amphetamine behavioral sensitization and the excitatory amino acids. *Brain Res* 1990; 537(1-2): 76-82.
- [334] Karlsgodt KH, Lukas SE, Elman I. Psychosocial stress and the duration of cocaine use in non-treatment seeking individuals with cocaine dependence. *Am J Drug Alcohol Abuse* 2003; 29(3): 539-551.
- [335] Karreman M, Westerink BHC, Moghaddam B. Excitatory amino acid receptors in the ventral tegmental area regulate dopamine release in the ventral striatum. *J Neurochem* 1996; 67(2): 601-607.
- [336] Karszen AM, Meijer OC, van dS, I, Lucassen PJ, de Lange EC, de Boer AG, De Kloet ER. Multidrug resistance P-glycoprotein hampers the access of cortisol but not of corticosterone to mouse and human brain. *Endocrinology* 2001; 142(6): 2686-2694.
- [337] Karst H, Joels M. Corticosterone slowly enhances miniature excitatory postsynaptic current amplitude in mice CA1 hippocampal cells. *J Neurophysiol* 2005; 94(5): 3479-3486.
- [338] Karst H, Berger S, Turiault M, Tronche F, Schutz G, Joels M. Mineralocorticoid receptors are indispensable for nongenomic modulation of hippocampal glutamate transmission by corticosterone. *Proc Natl Acad Sci U S A* 2005; 102(52): 19204-19207.
- [339] Kavaliers M, Ossenkopp KP. Corticosterone rapidly reduces male odor preferences in female mice. *Neuroreport* 2001; 12(13): 2999-3002.
- [340] Keibian JW, Calne DB. Multiple receptors for dopamine. *Nature* 1979; 277(5692): 93-96.
- [341] Keller-Wood ME, Dallman MF. Corticosteroid inhibition of ACTH secretion. *Endocr Rev* 1984; 5(1): 1-24.
- [342] Kelley AE. Ventral striatal control of appetitive motivation: role in ingestive behavior and reward-related learning. *Neurosci Biobehav Rev* 2004; 27(8): 765-776.
- [343] Kelley AE, Domesick VB, Nauta WJ. The amygdalostriatal projection in the rat - an anatomical study by anterograde and retrograde tracing methods. *Neuroscience* 1982; 7(3): 615-630.
- [344] Kelly PH, Iversen SD. Selective 6OHDA-induced destruction of mesolimbic dopamine neurons: abolition of psychostimulant-induced locomotor activity in rats. *Eur J Pharmacol* 1976; 40(1): 45-56.
- [345] Kelly PH, Seviour PW, Iversen SD. Amphetamine and apomorphine responses in the rat following 6-OHDA lesions of the nucleus accumbens septi and corpus striatum. *Brain Res* 1975; 94(3): 507-522.
- [346] Kelz MB, Chen J, Carlezon WA, Jr., Whisler K, Gilden L, Beckmann AM, Steffen C, Zhang YJ,

- Marotti L, Self DW, Tkatch T, Baranauskas G, Surmeier DJ, Neve RL, Duman RS, Picciotto MR, Nestler EJ. Expression of the transcription factor deltaFosB in the brain controls sensitivity to cocaine. *Nature* 1999; 401(6750): 272-276.
- [347] Kendler KS, Bulik CM, Silberg J, Hettema JM, Myers J, Prescott CA. Childhood sexual abuse and adult psychiatric and substance use disorders in women: an epidemiological and cotwin control analysis. *Arch Gen Psychiatry* 2000; 57(10): 953-959.
- [348] Kendler KS, Prescott CA. Cocaine use, abuse and dependence in a population-based sample of female twins. *Br J Psychiatry* 1998; 173: 345-350.
- [349] Kendler KS, Jacobson KC, Prescott CA, Neale MC. Specificity of genetic and environmental risk factors for use and abuse/dependence of cannabis, cocaine, hallucinogens, sedatives, stimulants, and opiates in male twins. *Am J Psychiatry* 2003; 160(4): 687-695.
- [350] Kenny PJ, Boutrel B, Gasparini F, Koob GF, Markou A. Metabotropic glutamate 5 receptor blockade may attenuate cocaine self-administration by decreasing brain reward function in rats. *Psychopharmacology (Berl)* 2005; 179(1): 247-254.
- [351] Khroyan TV, Barrett-Larimore RL, Rowlett JK, Spealman RD. Dopamine D1- and D2-like receptor mechanisms in relapse to cocaine-seeking behavior: effects of selective antagonists and agonists. *J Pharmacol Exp Ther* 2000; 294(2): 680-687.
- [352] Khroyan TV, Platt DM, Rowlett JK, Spealman RD. Attenuation of relapse to cocaine seeking by dopamine D1 receptor agonists and antagonists in non-human primates. *Psychopharmacology (Berl)* 2003; 168(1-2): 124-131.
- [353] Kilts CD, Schweitzer JB, Quinn CK, Gross RE, Faber TL, Muhammad F, Ely TD, Hoffman JM, Drexler KPG. Neural activity related to drug craving in cocaine addiction. *Arch Gen Psychiatry* 2001; 58(4): 334-341.
- [354] Klebaur JE, Bevins RA, Segar TM, Bardo MT. Individual differences in behavioral responses to novelty and amphetamine self-administration in male and female rats. *Behav Pharmacol* 2001; 12(4): 267-275.
- [355] Knych ET, Eisenberg RM. Effect of amphetamine on plasma corticosterone in the conscious rat. *Neuroendocrinology* 1979; 29(2): 110-118.
- [356] Koeppe MJ, Gunn RN, Lawrence AD, Cunningham VJ, Dagher A, Jones T, Brooks DJ, Bench CJ, Grasby PM. Evidence for striatal dopamine release during a video game. *Nature* 1998; 393(6682): 266-268.
- [357] Konradi C, Cole RL, Heckers S, Hyman SE. Amphetamine regulates gene expression in rat striatum via transcription factor CREB. *J Neurosci* 1994; 14(9): 5623-5634.
- [358] Koob GF. Stress, corticotropin-releasing factor, and drug addiction. *Ann NY Acad Sci* 1999; 897: 27-45.
- [359] Koob GF, Cador M. Psychomotor stimulant sensitization: the corticotropin-releasing factor-steroid connection. Commentary on Wise and Leeb "Psychomotor-stimulant sensitization: a unitary phenomenon?". *Behav Pharmacol* 1993; 4(4): 351-354.
- [360] Koob GF, Le Moal M. Drug abuse: hedonic homeostatic dysregulation. *Science* 1997; 278(5335): 52-58.

- [361] Koob GF, Le Moal M. Drug addiction, dysregulation of reward, and allostasis. *Neuropsychopharmacology* 2001; 24(2): 97-129.
- [362] Kornet M, Goosen C, Thyssen JH, Van Ree JM. Endocrine profile during acquisition of free-choice alcohol drinking in rhesus monkeys; treatment with desglycinamide-(Arg8)-vasopressin. *Alcohol Alcoholism* 1992; 27(4): 403-410.
- [363] Kosten TA, Miserendino MJ, Kehoe P. Enhanced acquisition of cocaine self-administration in adult rats with neonatal isolation stress experience. *Brain Res* 2000; 875(1-2): 44-50.
- [364] Kosten TR, Rounsaville BJ, Kleber HD. A 2.5-year follow-up of depression, life crises, and treatment effects on abstinence among opioid addicts. *Arch Gen Psychiatry* 1986; 43(8): 733-738.
- [365] Kosten TA, Sanchez H, Zhang XY, Kehoe P. Neonatal isolation enhances acquisition of cocaine self-administration and food responding in female rats. *Behav Brain Res* 2004; 151(1-2): 137-149.
- [366] Kosten TA, Zhang XY, Kehoe P. Heightened cocaine and food self-administration in female rats with neonatal isolation experience. *Neuropsychopharmacology* 2005; 31(1): 70-76.
- [367] Kosten TR, Oliveto A, Sevarino KA, Gonsai K, Feingold A. Ketoconazole increases cocaine and opioid use in methadone maintained patients. *Drug Alcohol Depend* 2002; 66(2): 173-180.
- [368] Kovacs P, Hernadi I. Alpha2 antagonist yohimbine suppresses maintained firing of rat prefrontal neurons in vivo. *Neuroreport* 2003; 14(6): 833-836.
- [369] Krueger RJ, Lin CM, Fung YK. Effect of chronic administration of nicotine or cocaine on steroidogenesis in rat adrenocortical cells. *J Pharm Pharmacol* 1991; 43(3): 200-203.
- [370] Kruzich PJ, See RE. Differential contributions of the basolateral and central amygdala in the acquisition and expression of conditioned relapse to cocaine-seeking behavior. *J Neurosci* 2001; 21(14): RC155.
- [371] Kuhar MJ, Jaworski JN, Hubert GW, Philpot KB, Dominguez G. Cocaine- and amphetamine-regulated transcript peptides play a role in drug abuse and are potential therapeutic targets. *AAPS J* 2005; 7(1): E259-E265.
- [372] Kuhar MJ, Ritz MC, Boja JW. The dopamine hypothesis of the reinforcing properties of cocaine. *Trends Neurosci* 1991; 14(7): 299-302.
- [373] Kumer SC, Vrana KE. Intricate regulation of tyrosine hydroxylase activity and gene expression. *J Neurochem* 1996; 67(2): 443-462.
- [374] Lakshmi V, Sakai RR, McEwen BS, Monder C. Regional distribution of 11 beta-hydroxysteroid dehydrogenase in rat brain. *Endocrinology* 1991; 128(4): 1741-1748.
- [375] Le Moine C, Bloch B. D1 and D2 dopamine receptor gene expression in the rat striatum: sensitive cRNA probes demonstrate prominent segregation of D1 and D2 mRNAs in distinct neuronal populations of the dorsal and ventral striatum. *J Comp Neurol* 1995; 355(3): 418-426.
- [376] Le AD, Harding S, Juzysch W, Funk D, Shaham Y. Role of alpha-2 adrenoceptors in stress-induced reinstatement of alcohol seeking and alcohol self-administration in rats. *Psychopharmacology (Berl)* 2005; 179(2): 366-373.
- [377] Leibeltern G. Sigmund Freud and cocaine. *Wien Klin Wochenschr* 1983; 95(21): 765-769.
- [378] Lecca D, Piras G, Driscoll P, Giorgi O, Corda MG. A differential activation of dopamine output in

- the shell and core of the nucleus accumbens is associated with the motor responses to addictive drugs: a brain dialysis study in Roman high- and low-avoidance rats. *Neuropharmacology* 2004; 46(5): 688-699.
- [379] Leccese AP, Lyness WH. Lesions of dopamine neurons in the medial prefrontal cortex: effects on self-administration of amphetamine and dopamine synthesis in the brain of the rat. *Neuropharmacology* 1987; 26(9): 1303-1308.
- [380] Lee B, Platt DM, Rowlett JK, Adewale AS, Spealman RD. Attenuation of behavioral effects of cocaine by the Metabotropic Glutamate Receptor 5 Antagonist 2-Methyl-6-(phenylethynyl)-pyridine in squirrel monkeys: comparison with dizocilpine. *J Pharmacol Exp Ther* 2005; 312(3): 1232-1240.
- [381] Leibowitz SF, Eidelman D, Suh JS, Diaz S, Sladek CD. Mapping study of noradrenergic stimulation of vasopressin release. *Exp Neurol* 1990; 110(3): 298-305.
- [382] Leibowitz SF, Diaz S, Tempel D. Norepinephrine in the paraventricular nucleus stimulates corticosterone release. *Brain Res* 1989; 496(1-2): 219-227.
- [383] Lepsch LB, Gonzalo LA, Magro FJ, Delucia R, Scavone C, Planeta CS. Exposure to chronic stress increases the locomotor response to cocaine and the basal levels of corticosterone in adolescent rats. *Addict Biol* 2005; 10(3): 251-256.
- [384] Leri F, Flores J, Rodaros D, Stewart J. Blockade of stress-induced but not cocaine-induced reinstatement by infusion of noradrenergic antagonists into the bed nucleus of the stria terminalis or the central nucleus of the amygdala. *J Neurosci* 2002; 22(13): 5713-5718.
- [385] Lester J, Fink S, Aronin N, DiFiglia M. Colocalization of D1 and D2 dopamine receptor mRNAs in striatal neurons. *Brain Res* 1993; 621(1): 106-110.
- [386] Levine S. Developmental determinants of sensitivity and resistance to stress. *Psychoneuroendocrinology* 2005; 30(10): 939-946.
- [387] Levy AD, Li Q, Alvarez Sanz MC, Rittenhouse PA, Kerr JE, Van de Kar LD. Neuroendocrine responses to cocaine do not exhibit sensitization following repeated cocaine exposure. *Life Sci* 1992; 51(12): 887-897.
- [388] Lewander T. Effects of amphetamine on urinary and tissue catecholamines in rats after inhibition of its metabolism with desmethyylimipramine. *Eur J Pharmacol* 1968; 5(1): 1-9.
- [389] Leyton M, Stewart J. Preexposure to foot-shock sensitizes the locomotor response to subsequent systemic morphine and intra-nucleus accumbens amphetamine. *Pharmacol Biochem Behav* 1990; 37(2): 303-310.
- [390] Li Y, Acerbo MJ, Robinson TE. The induction of behavioural sensitization is associated with cocaine-induced structural plasticity in the core (but not shell) of the nucleus accumbens. *Eur J Neurosci* 2004; 20(6): 1647-1654.
- [391] Li Y, Hu XT, Berney TG, Vartanian AJ, Stine CD, Wolf ME, White FJ. Both glutamate receptor antagonists and prefrontal cortex lesions prevent induction of cocaine sensitization and associated neuroadaptations. *Synapse* 1999; 34(3): 169-180.
- [392] Li Y, Kolb B, Robinson TE. The location of persistent amphetamine-induced changes in the density

- of dendritic spines on medium spiny neurons in the nucleus accumbens and caudate-putamen. *Neuropsychopharmacology* 2003; 28(6): 1082-1085.
- [393] Li Y, Robinson TE, Bhatnagar S. Effects of maternal separation on behavioural sensitization produced by repeated cocaine administration in adulthood. *Brain Res* 2003; 960(1-2): 42-47.
- [394] Li Y, Wolf ME. Ibotenic acid lesions of prefrontal cortex do not prevent expression of behavioral sensitization to amphetamine. *Behav Brain Res* 1997; 84(1-2): 285-289.
- [395] Liang KC, Chen LL, Huang TE. The role of amygdala norepinephrine in memory formation: involvement in the memory enhancing effect of peripheral epinephrine. *Chin J Physiol* 1995; 38(2): 81-91.
- [396] Liang KC, Juler RG, McGaugh JL. Modulating effects of posttraining epinephrine on memory: involvement of the amygdala noradrenergic system. *Brain Res* 1986; 368(1): 125-133.
- [397] Lindley SE, Bengoechea TG, Schatzberg AF, Wong DL. Glucocorticoid effects on mesotelencephalic dopamine neurotransmission. *Neuropsychopharmacology* 1999; 21(3): 399-407.
- [398] Liposits Z, Paull WK. Association of dopaminergic fibers with corticotropin releasing hormone (CRH)-synthesizing neurons in the paraventricular nucleus of the rat hypothalamus. *Histochemistry* 1989; 93(2): 119-127.
- [399] Lorrain DS, Arnold GM, Vezina P. Previous exposure to amphetamine increases incentive to obtain the drug: long-lasting effects revealed by the progressive ratio schedule. *Behav Brain Res* 2000; 107(1-2): 9-19.
- [400] Lowy MT, Gault L, Yamamoto BK. Adrenalectomy attenuates stress-induced elevations in extracellular glutamate concentrations in the hippocampus. *J Neurochem* 1993; 61(5): 1957-1960.
- [401] Loy R, Koziell DA, Lindsey JD, Moore RY. Noradrenergic innervation of the adult rat hippocampal formation. *J Comp Neurol* 1980; 189(4): 699-710.
- [402] Lu L, Dempsey J, Shaham Y, Hope BT. Differential long-term neuroadaptations of glutamate receptors in the basolateral and central amygdala after withdrawal from cocaine self-administration in rats. *J Neurochem* 2005; 94(1): 161-168.
- [403] Lu L, Grimm JW, Shaham Y, Hope BT. Molecular neuroadaptations in the accumbens and ventral tegmental area during the first 90 days of forced abstinence from cocaine self-administration in rats. *J Neurochem* 2003; 85(6): 1604-1613.
- [404] Lu L, Liu Z, Huang M, Zhang Z. Dopamine-dependent responses to cocaine depend on corticotropin-releasing factor receptor subtypes. *J Neurochem* 2003; 84(6): 1378-1386.
- [405] Lu L, Shepard JD, Scott Hall F, Shaham Y. Effect of environmental stressors on opiate and psychostimulant reinforcement, reinstatement and discrimination in rats: a review. *Neurosci Biobehav Rev* 2003; 27(5): 457-491.
- [406] Lu W, Wolf ME. Expression of dopamine transporter and vesicular monoamine transporter 2 mRNAs in rat midbrain after repeated amphetamine administration. *Brain Res Mol Brain Res* 1997; 49(1-2): 137-148.
- [407] Lu X-Y, Behnam Ghasemzadeh M, Kalivas PW. Expression of D1 receptor, D2 receptor, substance P and enkephalin messenger RNAs in the neurons projecting from the nucleus accumbens. *Neuroscience* 1997; 82(3): 767-780.

- [408] Lucas LR, Angulo JA, Le Moal M, McEwen BS, Piazza PV. Neurochemical characterization of individual vulnerability to addictive drugs in rats. *Eur J Neurosci* 1998; 10(10): 3153-3163.
- [409] Lyness WH, Friedle NM, Moore KE. Destruction of dopaminergic nerve terminals in nucleus accumbens: effect on d-amphetamine self-administration. *Pharmacol Biochem Behav* 1979; 11(5): 553-556.
- [410] MacLennan AJ, Maier SF. Coping and the stress-induced potentiation of stimulant stereotypy in the rat. *Science* 1983; 219(4588): 1091-1093.
- [411] Maldonado R, Robledo P, Chover AJ, Caine SB, Koob GF. D1 dopamine receptors in the nucleus accumbens modulate cocaine self-administration in the rat. *Pharmacol Biochem Behav* 1993; 45(1): 239-242.
- [412] Maldonado R, Valverde O, Berrendero F. Involvement of the endocannabinoid system in drug addiction. *Trends Neurosci* 2006; 29(4): 225-232.
- [413] Malkoski SP, Dorin RI. Composite glucocorticoid regulation at a functionally defined negative glucocorticoid response element of the human corticotropin-releasing hormone gene. *Mol Endocrinol* 1999; 13(10): 1629-1644.
- [414] Mangat HS, Islam A, Heigenskind C, Mustafa A, Winblad B, Adem A. Long-term adrenalectomy decreases NMDA receptors in rat hippocampus. *Neuroreport* 1998; 9(9): 2011-2014.
- [415] Mantsch JR, Goeders NE. Ketoconazole does not block cocaine discrimination or the cocaine-induced reinstatement of cocaine-seeking behavior. *Pharmacol Biochem Behav* 1999; 64(1): 65-73.
- [416] Mantsch JR, Goeders NE. Ketoconazole blocks the stress-induced reinstatement of cocaine-seeking behavior in rats: relationship to the discriminative stimulus effects of cocaine. *Psychopharmacology (Berl)* 1999; 142(4): 399-407.
- [417] Mantsch JR, Katz ES. Elevation of glucocorticoids is necessary but not sufficient for the escalation of cocaine self-administration by chronic electric footshock stress in rats. *Neuropsychopharmacology* 2007; 32(2): 367-376.
- [418] Mantsch JR, Saphier D, Goeders NE. Corticosterone facilitates the acquisition of cocaine self-administration in rats: opposite effects of the type II glucocorticoid receptor agonist dexamethasone. *J Pharmacol Exp Ther* 1998; 287(1): 72-80.
- [419] Marek GJ, Aghajanian GK. 5-HT<sub>2A</sub> receptor or alpha<sub>1</sub>-adrenoceptor activation induces excitatory postsynaptic currents in layer V pyramidal cells of the medial prefrontal cortex. *Eur J Pharmacol* 1999; 367(2-3): 197-206.
- [420] Marinelli M, Le Moal M, Piazza PV. Acute pharmacological blockade of corticosterone secretion reverses food restriction-induced sensitization of the locomotor response to cocaine. *Brain Res* 1996; 724(2): 251-255.
- [421] Marinelli M, Piazza PV. Interaction between glucocorticoid hormones, stress and psychostimulant drugs. *Eur J Neurosci* 2002; 16(3): 387-394.
- [422] Marinelli M, Piazza PV, Deroche V, Maccari S, Le Moal M, Simon H. Corticosterone circadian secretion differentially facilitates dopamine-mediated psychomotor effect of cocaine and morphine. *J Neurosci* 1994; 14(5 Pt 1): 2724-2731.

- [423] Marinelli M, Rouge-Pont F, Deroche V, Barrot M, Jesus-Oliveira C, Le Moal M, Piazza PV. Glucocorticoids and behavioral effects of psychostimulants. I: locomotor response to cocaine depends on basal levels of glucocorticoids. *J Pharmacol Exp Ther* 1997; 281(3): 1392-1400.
- [424] Marinelli M, Rouge-Pont F, Jesus-Oliveira C, Le Moal M, Piazza PV. Acute blockade of corticosterone secretion decreases the psychomotor stimulant effects of cocaine. *Neuropsychopharmacology* 1997; 16(2): 156-161.
- [425] Martin-Iverson MT, Szostak C, Fibiger HC. 6-Hydroxydopamine lesions of the medial prefrontal cortex fail to influence intravenous self-administration of cocaine. *Psychopharmacology (Berl)* 1986; 88(3): 310-314.
- [426] Mash DC, Staley JK, Izenwasser S, Basile M, Rutenber AJ. Serotonin transporters upregulate with chronic cocaine use. *J Chem Neuroanat* 2000; 20(3-4): 271-280.
- [427] Matthews JC, Collins A. Interactions of cocaine and cocaine congeners with sodium channels. *Biochem Pharmacol* 1983; 32(3): 455-460.
- [428] Matthews K, Hall FS, Wilkinson LS, Robbins TW. Retarded acquisition and reduced expression of conditioned locomotor activity in adult rats following repeated early maternal separation: effects of prefeeding, d-amphetamine, dopamine antagonists and clonidine. *Psychopharmacologia* 1996; 126(1): 75-84.
- [429] Matthews K, Robbins TW, Everitt BJ, Caine SB. Repeated neonatal maternal separation alters intravenous cocaine self-administration in adult rats. *Psychopharmacology (Berl)* 1999; 141(2): 123-134.
- [430] Mattingly BA, Hart TC, Lim K, Perkins C. Selective antagonism of dopamine D1 and D2 receptors does not block the development of behavioral sensitization to cocaine. *Psychopharmacology (Berl)* 1994; 114(2): 239-242.
- [431] Mayer JL, Klumpers L, Maslam S, De Kloet ER, Joels M, Lucassen PJ. Brief treatment with the glucocorticoid receptor antagonist mifepristone normalises the corticosterone-induced reduction of adult hippocampal neurogenesis. *J Neuroendocrinol* 2006; 18(8): 629-631.
- [432] McClung CA, Nestler EJ. Regulation of gene expression and cocaine reward by CREB and Delta-FosB. *Nat Neurosci* 2003; 6(11): 1208-1215.
- [433] McEwen BS. Allostasis, allostatic load, and the aging nervous system: role of excitatory amino acids and excitotoxicity. *Neurochem Res* 2000; 25(9-10): 1219-1231.
- [434] McFarland K, Davidge SB, Lapish CC, Kalivas PW. Limbic and motor circuitry underlying foot-shock-induced reinstatement of cocaine-seeking behavior. *J Neurosci* 2004; 24(7): 1551-1560.
- [435] McFarland K, Kalivas PW. The circuitry mediating cocaine-induced reinstatement of drug-seeking behavior. *J Neurosci* 2001; 21(21): 8655-8663.
- [436] McFarland K, Lapish CC, Kalivas PW. Prefrontal glutamate release into the core of the nucleus accumbens mediates cocaine-induced reinstatement of drug-seeking behavior. *J Neurosci* 2003; 23(8): 3531-3537.
- [437] McGaugh JL, Cahill L, Roozendaal B. Involvement of the amygdala in memory storage: interaction with other brain systems. *Proc Natl Acad Sci U S A* 1996; 93(24): 13508-13514.
- [438] McGregor A, Baker G, Roberts DC. Effect of 6-hydroxydopamine lesions of the medial prefrontal

- cortex on intravenous cocaine self-administration under a progressive ratio schedule of reinforcement. *Pharmacol Biochem Behav* 1996; 53(1): 5-9.
- [439] McGregor A, Roberts DCS. Dopaminergic antagonism within the nucleus accumbens or the amygdala produces differential effects on intravenous cocaine self-administration under fixed and progressive ratio schedules of reinforcement. *Brain Res* 1993; 624(1-2): 245-252.
- [440] McKinzie DL, Rodd-Henricks ZA, Dagon CT, Murphy JM, McBride WJ. Cocaine is self-administered into the shell region of the nucleus accumbens in Wistar rats. *Ann N Y Acad Sci* 1999; 877: 788-791.
- [441] McLellan AT, Lewis DC, O'Brien CP, Kleber HD. Drug dependence, a chronic medical illness: implications for treatment, insurance, and outcomes evaluation. *JAMA* 2000; 284(13): 1689-1695.
- [442] Meador-Woodruff JH, Mansour A, Bunzow JR, Tol HHMV, Watson SJ, Civelli O. Distribution of D2 dopamine receptor mRNA in rat brain. *Proc Natl Acad Sci U S A* 1989; 86(19): 7625-7628.
- [443] Meador-Woodruff JH, Mansour A, Grandy DK, Damask SP, Civelli O, Watson J. Distribution of D5 dopamine receptor mRNA in rat brain. *Neurosci Lett* 1992; 145(2): 209-212.
- [444] Meaney MJ, Brake W, Gratton A. Environmental regulation of the development of mesolimbic dopamine systems: a neurobiological mechanism for vulnerability to drug abuse? *Psychoneuroendocrinology* 2002; 27(1-2): 127-138.
- [445] Meaney MJ, Diorio J, Francis D, Widdowson J, LaPlante P, Caldji C, Sharma S, Seckl JR, Plotsky PM. Early environmental regulation of forebrain glucocorticoid receptor gene expression: implications for adrenocortical responses to stress. *Dev Neurosci* 1996; 18(1-2): 49-72.
- [446] Meijer OC, van der Laan S, Lachize S, Steenbergen PJ, De Kloet ER. Steroid receptor coregulator diversity: what can it mean for the stressed brain? *Neuroscience* 2006; 138(3): 891-899.
- [447] Meliska CJ, Bartke A, McGlacken G, Jensen RA. Ethanol, nicotine, amphetamine, and aspartame consumption and preferences in C57BL/6 and DBA/2 mice. *Pharmacol Biochem Behav* 1995; 50(4): 619-626.
- [448] Mendelson JH, Teoh SK, Mello NK, Ellingboe J, Rhoades E. Acute effects of cocaine on plasma adrenocorticotrophic hormone, luteinizing hormone and prolactin levels in cocaine-dependent men. *J Pharmacol Exp Ther* 1992; 263(2): 505-509.
- [449] Mendrek A, Blaha CD, Phillips AG. Pre-exposure of rats to amphetamine sensitizes self-administration of this drug under a progressive ratio schedule. *Psychopharmacology (Berl)* 1998; 135(4): 416-422.
- [450] Mikics E, Barsy B, Barsvari B, Haller J. Behavioral specificity of non-genomic glucocorticoid effects in rats: Effects on risk assessment in the elevated plus-maze and the open-field. *Horm Behav* 2005; 48(2): 152-162.
- [451] Mikics E, Kruk MR, Haller J. Genomic and non-genomic effects of glucocorticoids on aggressive behavior in male rats. *Psychoneuroendocrinology* 2004; 29(5): 618-635.
- [452] Mink JW, Thach WT. Basal ganglia intrinsic circuits and their role in behavior. *Curr Opin Neurobiol* 1993; 3(6): 950-957.
- [453] Misra K, Pandey SC. Differences in basal levels of CREB and NPY in nucleus accumbens regions

- between C57BL/6 and DBA/2 mice differing in inborn alcohol drinking behavior. *J Neurosci Res* 2003; 74(6): 967-975.
- [454] Mittleman G, Jones GH, Robbins TW. Sensitization of amphetamine-stereotypy reduces plasma corticosterone: implications for stereotypy as a coping response. *Behav Neural Biol* 1991; 56(2): 170-182.
- [455] Mittleman G, Van Brunt CL, Matthews DB. Schedule-induced ethanol self-administration in DBA/2J and C57BL/6J mice. *Alcohol Clin Exp Res* 2003; 27(6): 918-925.
- [456] Miyashita T, Williams CL. Epinephrine administration increases neural impulses propagated along the vagus nerve: role of peripheral beta-adrenergic receptors. *Neurobiol Learn Mem* 2006; 85(2): 116-124.
- [457] Mobbs D, Greicius MD, Abdel-Azim E, Menon V, Reiss AL. Humor modulates the mesolimbic reward centers. *Neuron* 2003; 40(5): 1041-1048.
- [458] Moffett MC, Harley J, Francis D, Sanghani SP, Davis WI, Kuhar MJ. Maternal separation and handling affects cocaine self-administration in both the treated pups as adults and the dams. *J Pharmacol Exp Ther* 2006; 317(3): 1210-1218.
- [459] Moghaddam B. Stress activation of glutamate neurotransmission in the prefrontal cortex: implications for dopamine-associated psychiatric disorders. *Biol Psychiatry* 2002; 51(10): 775-787.
- [460] Moghaddam B, Bolinao ML, Stein-Behrens B, Sapolsky R. Glucocorticoids mediate the stress-induced extracellular accumulation of glutamate. *Brain Res* 1994; 655(1-2): 251-254.
- [461] Moldow RL, Fischman AJ. Cocaine induced secretion of ACTH, beta-endorphin, and corticosterone. *Peptides* 1987; 8(5): 819-822.
- [462] Monsma FJ, Jr., Mahan LC, McVittie LD, Gerfen CR, Sibley DR. Molecular cloning and expression of a D1 dopamine receptor linked to adenylyl cyclase activation. *Proc Natl Acad Sci U S A* 1990; 87(17): 6723-6727.
- [463] Monsma FJ, McVittie LD, Gerfen CR, Mahan LC, Sibley DR. Multiple D2 dopamine receptors produced by alternative RNA splicing. *Nature* 1989; 342(6252): 926-929.
- [464] Morgan D, Smith MA, Roberts DCS. Binge self-administration and deprivation produces sensitization to the reinforcing effects of cocaine in rats. *Psychopharmacologia* 2005; 178(2-3): 309-316.
- [465] Morse AC, Erwin VG, Jones BC. Strain and housing affect cocaine self-selection and open-field locomotor activity in mice. *Pharmacol Biochem Behav* 1993; 45(4): 905-912.
- [466] Mulholland PJ, Self RL, Hensley AK, Little HJ, Littleton JM, Prendergast MA. A 24 h corticosterone exposure exacerbates excitotoxic insult in rat hippocampal slice cultures independently of glucocorticoid receptor activation or protein synthesis. *Brain Res* 2006; 1082(1): 165-172.
- [467] Muller CP, Huston JP. Determining the region-specific contributions of 5-HT receptors to the psychostimulant effects of cocaine. *Trends Pharmacol Sci* 2006; 27(2): 105-112.
- [468] Najavits LM, Runkel R, Neuner C, Frank AF, Thase ME, Crits-Christoph P, Blaine J. Rates and symptoms of PTSD among cocaine-dependent patients. *J Stud Alcohol* 2003; 64(5): 601-606.
- [469] Nestler EJ. Cellular responses to chronic treatment with drugs of abuse. *Crit Rev Neurobiol* 1993; 7(1): 23-39.

- [470] Ng GY, O'Dowd BF, George SR. Genotypic differences in brain dopamine receptor function in the DBA/2J and C57BL/6J inbred mouse strains. *Eur J Pharmacol* 1994; 269(3): 349-364.
- [471] Ng GY, O'Dowd BF, George SR. Genotypic differences in mesolimbic enkephalin gene expression in DBA/2J and C57BL/6J inbred mice. *Eur J Pharmacol* 1996; 311(1): 45-52.
- [472] Nikulina EM, Covington HE, III, Ganschow L, Hammer RP, Jr., Miczek KA. Long-term behavioral and neuronal cross-sensitization to amphetamine induced by repeated brief social defeat stress: Fos in the ventral tegmental area and amygdala. *Neuroscience* 2004; 123(4): 857-865.
- [473] Noble EP. Polymorphisms of the D2 dopamine receptor gene and alcoholism and other substance use disorders. *Alcohol Alcoholism* 1994; 2(suppl): 35-43.
- [474] Noble EP. Addiction and its reward process through polymorphisms of the D2 dopamine receptor gene: a review. *Eur Psychiatry* 2000; 15(2): 79-89.
- [475] Noble EP, Blum K, Khalsa ME, Ritchie T, Montgomery A, Wood RC, Fitch RJ, Ozkaragoz T, Sheridan PJ, Anglin MD. Allelic association of the D2 dopamine receptor gene with cocaine dependence. *Drug Alcohol Depend* 1993; 33(3): 271-285.
- [476] Noble EP. The D2 dopamine receptor gene: a review of association studies in alcoholism and phenotypes. *Alcohol* 1998; 16(1): 33-45.
- [477] Nothen MM, Cichon S, Hemmer S, Hebebrand J, Remschmidt H, Lehmkuhl G, Poustka F, Schmidt M, Catalano M, Fimmers R. Human dopamine D4 receptor gene: frequent occurrence of a null allele and observation of homozygosity. *Hum Mol Genet* 1994; 3(12): 2207-2212.
- [478] O'Brien CP, Ehrman RN, Ternes JW. Classical conditioning in human opioid dependence. In: Goldberg S, Stolerman I (eds). *Behavioral analysis of drug dependence*. Academic Press, New York: 1986; 329-356.
- [479] O'Brien MS, Anthony JC. Risk of becoming cocaine dependent: epidemiological estimates for the United States, 2000-2001. *Neuropsychopharmacology* 2005; 30(5): 1006-1018.
- [480] O'Malley KL, Harmon S, Tang L, Todd RD. The rat dopamine D4 receptor: sequence, gene structure, and demonstration of expression in the cardiovascular system. *New Biol* 1992; 4(2): 137-146.
- [481] Oitzl MS, van Haarst AD, De Kloet ER. Behavioral and neuroendocrine responses controlled by the concerted action of central mineralocorticoid (MRS) and glucocorticoid receptors (GRS). *Psychoneuroendocrinology* 1997; 22(suppl 1): S87-S93.
- [482] Olson L, Fuxe K. Further mapping out of central noradrenaline neuron systems: projections of the 'subcoeruleus' area. *Brain Res* 1972; 43(1): 289-295.
- [483] Orsini C, Bonito-Oliva A, Conversi D, Cabib S. Susceptibility to conditioned place preference induced by addictive drugs in mice of the C57BL/6 and DBA/2 inbred strains. *Psychopharmacology (Berl)* 2005; 181(2): 327-336.
- [484] Orsini C, Buchini F, Piazza PV, Puglisi-Allegra S, Cabib S. Susceptibility to amphetamine-induced place preference is predicted by locomotor response to novelty and amphetamine in the mouse. *Psychopharmacology (Berl)* 2004; 172(3): 264-270.
- [485] Ortiz J, DeCaprio JL, Kosten TA, Nestler EJ. Strain-selective effects of corticosterone on locomotor

- sensitization to cocaine and on levels of tyrosine hydroxylase and glucocorticoid receptor in the ventral tegmental area. *Neuroscience* 1995; 67(2): 383-397.
- [486] Oswald LM, Wong DF, McCaul M, Zhou Y, Kuwabara H, Choi L, Brasic J, Wand GS. Relationships among ventral striatal dopamine release, cortisol secretion, and subjective responses to amphetamine. *Neuropsychopharmacology* 2005; 30(4): 821-832.
- [487] Ouimette PC, Brown PJ, Najavits LM. Course and treatment of patients with both substance use and posttraumatic stress disorders. *Addict Behav* 1998; 23(6): 785-795.
- [488] Overton PG, Tong ZY, Brain PF, Clark D. Preferential occupation of mineralocorticoid receptors by corticosterone enhances glutamate-induced burst firing in rat midbrain dopaminergic neurons. *Brain Res* 1996; 737(1-2): 146-154.
- [489] Packard MG, Knowlton BJ. Learning and memory functions of the basal ganglia. *Annu Rev Neurosci* 2002; 25(1): 563-593.
- [490] Palomo T, Archer T, Kostrzewa RM, Beninger RJ. Gene-environment interplay in schizopsychotic disorders. *Neurotox Res* 2004; 6(1): 1-9.
- [491] Pan WH, Sung JC, Fuh SM. Local application of amphetamine into the ventral tegmental area enhances dopamine release in the nucleus accumbens and the medial prefrontal cortex through noradrenergic neurotransmission. *J Pharmacol Exp Ther* 1996; 278(2): 725-731.
- [492] Park WK, Bari AA, Jey AR, Anderson SM, Spealman RD, Rowlett JK, Pierce RC. Cocaine administered into the medial prefrontal cortex reinstates cocaine-seeking behavior by increasing AMPA receptor-mediated glutamate transmission in the nucleus accumbens. *J Neurosci* 2002; 22(7): 2916-2925.
- [493] Patacchioli FR, Pontieri FE, Orzi F, Di Grezia R, Angelucci L. Increased motor response to cocaine administration following recovery from chronic corticosterone treatment in the rat. *Eur Neuropsychopharmacol* 1998; 8(1): 43-46.
- [494] Patel PD, Lopez JF, Lyons DM, Burke S, Wallace M, Schatzberg AF. Glucocorticoid and mineralocorticoid receptor mRNA expression in squirrel monkey brain. *J Psychiat Res* 2000; 34(6): 383-392.
- [495] Paterson NE, Markou A. The metabotropic glutamate receptor 5 antagonist MPEP decreased break points for nicotine, cocaine and food in rats. *Psychopharmacology (Berl)* 2005; 179(1): 255-261.
- [496] Patrick SL, Thompson TL, Walker JM, Patrick RL. Concomitant sensitization of amphetamine-induced behavioral stimulation and in vivo dopamine release from rat caudate nucleus. *Brain Res* 1991; 538(2): 343-346.
- [497] Paulson PE, Camp DM, Robinson TE. Time course of transient behavioral depression and persistent behavioral sensitization in relation to regional brain monoamine concentrations during amphetamine withdrawal in rats. *Psychopharmacology (Berl)* 1991; 103(4): 480-492.
- [498] Pauly JR, Robinson SF, Collins AC. Chronic corticosterone administration enhances behavioral sensitization to amphetamine in mice. *Brain Res* 1993; 620(2): 195-202.
- [499] Paylor R, Tracy R, Wehner J, Rudy JW. DBA/2 and C57BL/6 mice differ in contextual fear but not auditory fear conditioning. *Behav Neurosci* 1994; 108(4): 810-817.

- [500] Peltier R, Schenk S. Effects of serotonergic manipulations on cocaine self-administration in rats. *Psychopharmacology (Berl)* 1993; 110(4): 390-394.
- [501] Peters J, Kalivas PW. The group II metabotropic glutamate receptor agonist, LY379268, inhibits both cocaine- and food-seeking behavior in rats. *Psychopharmacology (Berl)* 2006; 186(2): 143-149.
- [502] Pettit HO, Ettenberg A, Bloom FE, Koob GF. Destruction of dopamine in the nucleus accumbens selectively attenuates cocaine but not heroin self-administration in rats. *Psychopharmacology (Berl)* 1984; 84(2): 167-173.
- [503] Pettit HO, Justice J. Dopamine in the nucleus accumbens during cocaine self-administration as studied by in vivo microdialysis. *Pharmacol Biochem Behav* 1989; 34(4): 899-904.
- [504] Phillips GD, Howes SR, Whitelaw RB, Robbins TW, Everitt BJ. Isolation rearing impairs the reinforcing efficacy of intravenous cocaine or intra-accumbens d-amphetamine: impaired response to intra-accumbens D1 and D2/D3 dopamine receptor antagonists. *Psychopharmacology (Berl)* 1994; 115(3): 419-429.
- [505] Phillips GD, Howes SR, Whitelaw RB, Wilkinson LS, Robbins TW, Everitt BJ. Isolation rearing enhances the locomotor response to cocaine and a novel environment, but impairs the intravenous self-administration of cocaine. *Psychopharmacology (Berl)* 1994; 115(3): 407-418.
- [506] Phillips GD, Robbins TW, Everitt BJ. Bilateral intra-accumbens self-administration of d-amphetamine: antagonism with intra-accumbens SCH-23390 and sulpiride. *Psychopharmacology (Berl)* 1994; 114(3): 477-485.
- [507] Phillips TJ, Huson MG, McKinnon CS. Localization of genes mediating acute and sensitized locomotor responses to cocaine in BXD/Ty recombinant inbred mice. *J Neurosci* 1998; 18(8): 3023-3034.
- [508] Piazza PV, Barrot M, Rouge-Pont F, Marinelli M, Maccari S, Abrous DN, Simon H, Le Moal M. Suppression of glucocorticoid secretion and antipsychotic drugs have similar effects on the mesolimbic dopaminergic transmission. *Proc Natl Acad Sci U S A* 1996; 93(26): 15445-15450.
- [509] Piazza PV, Deminiere JM, Le Moal M, Simon H. Factors that predict individual vulnerability to amphetamine self-administration. *Science* 1989; 245(4925): 1511-1513.
- [510] Piazza PV, Deminiere JM, Le Moal M, Simon H. Stress- and pharmacologically-induced behavioral sensitization increases vulnerability to acquisition of amphetamine self-administration. *Brain Res* 1990; 514(1): 22-26.
- [511] Piazza PV, Deminiere JM, Maccari S, Mormede P, Le Moal M, Simon H. Individual reactivity to novelty predicts probability of amphetamine self-administration. *Behav Pharmacol* 1990; 1(4): 339-345.
- [512] Piazza PV, Deroche V, Deminiere JM, Maccari S, Le Moal M, Simon H. Corticosterone in the range of stress-induced levels possesses reinforcing properties: implications for sensation-seeking behaviors. *Proc Natl Acad Sci U S A* 1993; 90(24): 11738-11742.
- [513] Piazza PV, Deroche-Gamonet V, Rouge-Pont F, Le Moal M. Vertical shifts in self-administration dose-response functions predict a drug-vulnerable phenotype predisposed to addiction. *J Neurosci* 2000; 20(11): 4226-4232.

- [514] Piazza PV, Le Moal M. The role of stress in drug self-administration. *Trends Pharmacol Sci* 1998; 19(2): 67-74.
- [515] Piazza PV, Le Moal ML. Pathophysiological basis of vulnerability to drug abuse: role of an interaction between stress, glucocorticoids, and dopaminergic neurons. *Annu Rev Pharmacol Toxicol* 1996; 36: 359-378.
- [516] Piazza PV, Maccari S, Deminiere JM, Le Moal M, Mormede P, Simon H. Corticosterone levels determine individual vulnerability to amphetamine self-administration. *Proc Natl Acad Sci U S A* 1991; 88(6): 2088-2092.
- [517] Piazza PV, Marinelli M, Jodogne C, Deroche V, Rouge-Pont F, Maccari S, Le Moal M, Simon H. Inhibition of corticosterone synthesis by Metyrapone decreases cocaine-induced locomotion and relapse of cocaine self-administration. *Brain Res* 1994; 658(1-2): 259-264.
- [518] Piazza PV, Rouge-Pont F, Deroche V, Maccari S, Simon H, Le Moal M. Glucocorticoids have state-dependent stimulant effects on the mesencephalic dopaminergic transmission. *Proc Natl Acad Sci U S A* 1996; 93(16): 8716-8720.
- [519] Pierce RC, Kalivas PW. Amphetamine produces sensitized increases in locomotion and extracellular dopamine preferentially in the nucleus accumbens shell of rats administered repeated cocaine. *J Pharmacol Exp Ther* 1995; 275(2): 1019-1029.
- [520] Pierce RC, Kalivas PW. A circuitry model of the expression of behavioral sensitization to amphetamine-like psychostimulants. *Brain Res Brain Res Rev* 1997; 25(2): 192-216.
- [521] Pierce RC, Reeder DC, Hicks J, Morgan ZR, Kalivas PW. Ibotenic acid lesions of the dorsal prefrontal cortex disrupt the expression of behavioral sensitization to cocaine. *Neuroscience* 1997; 82(4): 1103-1114.
- [522] Pierre PJ, Vezina P. Predisposition to self-administer amphetamine: the contribution of response to novelty and prior exposure to the drug. *Psychopharmacology (Berl)* 1997; 129(3): 277-284.
- [523] Pilla M, Perachon S, Sautel F, Garrido F, Mann A, Wermuth CG, Schwartz JC, Everitt BJ, Sokoloff P. Selective inhibition of cocaine-seeking behaviour by a partial dopamine D3 receptor agonist. *Nature* 1999; 400(6742): 371-375.
- [524] Pitts DK, Marwah J. Autonomic actions of cocaine. *Can J Physiol Pharmacol* 1989; 67(9): 1168-1176.
- [525] Plotsky PM, Meaney MJ. Early, postnatal experience alters hypothalamic corticotropin-releasing factor (CRF) mRNA, median eminence CRF content and stress-induced release in adult rats. *Brain Res Mol Brain Res* 1993; 18(3): 195-200.
- [526] Pontieri FE, Tanda G, Di Chiara G. Intravenous cocaine, morphine, and amphetamine preferentially increase extracellular dopamine in the "shell" as compared with the "core" of the rat nucleus accumbens. *Proc Natl Acad Sci U S A* 1995; 92(26): 12304-12308.
- [527] Prasad BM, Ulibarri C, Kalivas PW, Sorg BA. Effect of adrenalectomy on the initiation and expression of cocaine-induced sensitization. *Psychopharmacology (Berl)* 1996; 125(3): 265-273.
- [528] Pryce CR, Ruedi-Bettschen D, Dettling AC, Feldon J. Early life stress: long-term physiological impact in rodents and primates. *News Physiol Sci* 2002; 17(4): 150-155.

- [529] Przegalinski E, Filip M, Siwanowicz J, Nowak E. Effect of adrenalectomy and corticosterone on cocaine-induced sensitization in rats. *J Physiol Pharmacol* 2000; 51(2): 193-204.
- [530] Przegalinski E, Filip M, Frankowska M, Zaniowska M, Papla I. Effects of CP 154,526, a CRF1 receptor antagonist, on behavioral responses to cocaine in rats. *Neuropeptides* 2005; 39(5): 525-533.
- [531] Puglisi-Allegra S, Cabib S. Psychopharmacology of dopamine: the contribution of comparative studies in inbred strains of mice. *Prog Neurobiol* 1997; 51(6): 637-661.
- [532] Puglisi-Allegra S, Imperato A, Angelucci L, Cabib S. Acute stress induces time-dependent responses in dopamine mesolimbic system. *Brain Res* 1991; 554(1-2): 217-222.
- [533] Puglisi-Allegra S, Kempf E, Cabib S. Role of genotype in the adaptation of the brain dopamine system to stress. *Neurosci Biobehav Rev* 1990; 14(4): 523-528.
- [534] Pulvirenti L, Swerdlow NR, Koob GF. Microinjection of a glutamate antagonist into the nucleus accumbens reduces psychostimulant locomotion in rats. *Neurosci Lett* 1989; 103(2): 213-218.
- [535] Quirarte GL, Roozendaal B, McGaugh JL. Glucocorticoid enhancement of memory storage involves noradrenergic activation in the basolateral amygdala. *Proc Natl Acad Sci U S A* 1997; 94(25): 14048-14053.
- [536] Ramsey NF, Van Ree JM. Emotional but not physical stress enhances intravenous cocaine self-administration in drug-naive rats. *Brain Res* 1993; 608(2): 216-222.
- [537] Ranaldi R, Pocock D, Zereik R, Wise RA. Dopamine fluctuations in the nucleus accumbens during maintenance, extinction, and reinstatement of intravenous d-Amphetamine self-administration. *J Neurosci* 1999; 19(10): 4102-4109.
- [538] Ranaldi R, Wise RA. Blockade of D1 dopamine receptors in the ventral tegmental area decreases cocaine reward: possible role for dendritically released dopamine. *J Neurosci* 2001; 21(15): 5841-5846.
- [539] Reid MS, Berger SP. Evidence for sensitization of cocaine-induced nucleus accumbens glutamate release. *Neuroreport* 1996; 7(7): 1325-1329.
- [540] Reid MS, Hsu K, Jr., Berger SP. Cocaine and amphetamine preferentially stimulate glutamate release in the limbic system: studies on the involvement of dopamine. *Synapse* 1997; 27(2): 95-105.
- [541] Reul JM, De Kloet ER. Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation. *Endocrinology* 1985; 117(6): 2505-2511.
- [542] Rhoads DL. A longitudinal study of life stress and social support among drug abusers. *Int J Addict* 1983; 18(2): 195-222.
- [543] Richtand NM, Woods SC, Berger SP, Strakowski SM. D3 dopamine receptor, behavioral sensitization, and psychosis. *Neurosci Biobehav Rev* 2001; 25(5): 427-443.
- [544] Richter RM, Weiss F. In vivo CRF release in rat amygdala is increased during cocaine withdrawal in self-administering rats. *Synapse* 1999; 32(4): 254-261.
- [545] Richter RM, Pich EM, Koob GF, Weiss F. Sensitization of cocaine-stimulated increase in extracellular levels of corticotropin-releasing factor from the rat amygdala after repeated administration as determined by intracranial microdialysis. *Neurosci Lett* 1995; 187(3): 169-172.

- [546] Rifkin A. Extrapyramidal side effects: a historical perspective. *J Clin Psychiatry* 1987; 48(suppl): 3-6.
- [547] Risbrough VB, Stein MB. Role of corticotropin releasing factor in anxiety disorders: a translational research perspective. *Horm Behav* 2006; 50(4): 550-561.
- [548] Risner M, Jones BE. Role of noradrenergic and dopaminergic processes in amphetamine self-administration. *Pharmacol Biochem Behav* 1976; 5(4): 477-482.
- [549] Ritz MC, Cone EJ, Kuhar MJ. Cocaine inhibition of ligand binding at dopamine, norepinephrine and serotonin transporters: a structure-activity study. *Life Sci* 1990; 46(9): 635-645.
- [550] Ritz MC, Lamb RJ, Goldberg SR, Kuhar MJ. Cocaine receptors on dopamine transporters are related to self-administration of cocaine. *Science* 1987; 237(4819): 1219-1223.
- [551] Rivera MA, Aufderheide AC, Cartmell LW, Torres CM, Langsjoen O. Antiquity of coca-leaf chewing in the south central Andes: a 3,000 year archaeological record of coca-leaf chewing from northern Chile. *J Psychoactive Drugs* 2005; 37(4): 455-458.
- [552] Rivet JM, Stinus L, LeMoal M, Mormede P. Behavioral sensitization to amphetamine is dependent on corticosteroid receptor activation. *Brain Res* 1989; 498(1): 149-153.
- [553] Roberts AJ, Crabbe JC, Keith LD. Genetic differences in hypothalamic-pituitary-adrenal axis responsiveness to acute ethanol and acute ethanol withdrawal. *Brain Res* 1992; 579(2): 296-302.
- [554] Roberts DC. Preclinical evidence for GABAB agonists as a pharmacotherapy for cocaine addiction. *Physiol Behav* 2005; 86(1-2): 18-20.
- [555] Roberts DC, Andrews MM, Vickers GJ. Baclofen attenuates the reinforcing effects of cocaine in rats. *Neuropsychopharmacology* 1996; 15(4): 417-423.
- [556] Roberts DC, Corcoran ME, Fibiger HC. On the role of ascending catecholaminergic systems in intravenous self-administration of cocaine. *Pharmacol Biochem Behav* 1977; 6(6): 615-620.
- [557] Roberts DC, Koob GF. Disruption of cocaine self-administration following 6-hydroxydopamine lesions of the ventral tegmental area in rats. *Pharmacol Biochem Behav* 1982; 17(5): 901-904.
- [558] Roberts DC, Phelan R, Hodges LM, Hodges MM, Bennett B, Childers S, Davies H. Self-administration of cocaine analogs by rats. *Psychopharmacology (Berl)* 1999; 144(4): 389-397.
- [559] Robinson TE. Stimulant drugs and stress: factors influencing individual differences in the susceptibility to sensitization. In: Kalivas PW, Barnes C (eds). *Sensitization of the Nervous System*. Telford Press, Caldwell: 1988; 145-173.
- [560] Robinson TE, Berridge KC. The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Brain Res Rev* 1993; 18(3): 247-291.
- [561] Robinson TE, Berridge KC. Incentive-sensitization and addiction. *Addiction* 2001; 96(1): 103-114.
- [562] Robinson TE, Browman KE, Crombag HS, Badiani A. Modulation of the induction or expression of psychostimulant sensitization by the circumstances surrounding drug administration. *Neurosci Biobehav Rev* 1998; 22(2): 347-354.
- [563] Robinson TE, Gorny G, Mitton E, Kolb B. Cocaine self-administration alters the morphology of dendrites and dendritic spines in the nucleus accumbens and neocortex. *Synapse* 2001; 39(3): 257-266.

- [564] Robinson TE, Kolb B. Alterations in the morphology of dendrites and dendritic spines in the nucleus accumbens and prefrontal cortex following repeated treatment with amphetamine or cocaine. *Eur J Neurosci* 1999; 11(5): 1598-1604.
- [565] Robinson TE, Kolb B. Persistent structural modifications in nucleus accumbens and prefrontal cortex neurons produced by previous experience with amphetamine. *J Neurosci* 1997; 17(21): 8491-8497.
- [566] Robinson TG, Beart PM. Excitant amino acid projections from rat amygdala and thalamus to nucleus accumbens. *Brain Res Bull* 1988; 20(4): 467-471.
- [567] Rocha BA, Fumagalli F, Gainetdinov RR, Jones SR, Ator R, Giros B, Miller GW, Caron MG. Cocaine self-administration in dopamine-transporter knockout mice. *Nat Neurosci* 1998; 1(2): 132-137.
- [568] Rocha BA, Odom LA, Barron BA, Ator R, Wild SA, Forster MJ. Differential responsiveness to cocaine in C57BL/6J and DBA/2J mice. *Psychopharmacology (Berl)* 1998; 138(1): 82-88.
- [569] Rodd-Henricks ZA, McKinzie DL, Li TK, Murphy JM, McBride WJ. Cocaine is self-administered into the shell but not the core of the nucleus accumbens of Wistar rats. *J Pharmacol Exp Ther* 2002; 303(3): 1216-1226.
- [570] Roozendaal B. Stress and memory: opposing effects of glucocorticoids on memory consolidation and memory retrieval. *Neurobiol Learn Mem* 2002; 78(3): 578-595.
- [571] Roozendaal B, Carmi O, McGaugh JL. Adrenocortical suppression blocks the memory-enhancing effects of amphetamine and epinephrine. *Proc Natl Acad Sci U S A* 1996; 93(4): 1429-1433.
- [572] Roozendaal B, de Quervain DJF, Ferry B, Setlow B, McGaugh JL. Basolateral amygdala-nucleus accumbens interactions in mediating glucocorticoid enhancement of memory consolidation. *J Neurosci* 2001; 21(7): 2518-2525.
- [573] Roozendaal B, McGaugh JL. The memory-modulatory effects of glucocorticoids depend on an intact stria terminalis. *Brain Res* 1996; 709(2): 243-250.
- [574] Roozendaal B, Okuda S, Van der Zee EA, McGaugh JL. Glucocorticoid enhancement of memory requires arousal-induced noradrenergic activation in the basolateral amygdala. *Proc Natl Acad Sci U S A* 2006; 103(17): 6741-6746.
- [575] Rossi-Arnaud C, Ammassari-Teule M. What do comparative studies of inbred mice add to current investigations on the neural basis of spatial behaviors? *Exp Brain Res* 1998; 123(1-2): 36-44.
- [576] Rots NY, Cools AR, Berod A, Voorn P, Rostene W, De Kloet ER. Rats bred for enhanced apomorphine susceptibility have elevated tyrosine hydroxylase mRNA and dopamine D2-receptor binding sites in nigrostriatal and tuberoinfundibular dopamine systems. *Brain Res* 1996; 710(1-2): 189-196.
- [577] Rouge-Pont F, Marinelli M, Le Moal M, Simon H, Piazza PV. Stress-induced sensitization and glucocorticoids. II. Sensitization of the increase in extracellular dopamine induced by cocaine depends on stress-induced corticosterone secretion. *J Neurosci* 1995; 15(11): 7189-7195.
- [578] Rouge-Pont F, Piazza PV, Kharouby M, Le Moal M, Simon H. Higher and longer stress-induced increase in dopamine concentrations in the nucleus accumbens of animals predisposed to amphetamine self-administration. A microdialysis study. *Brain Res* 1993; 602(1): 169-174.

- [579] Rouge-Pont F, Abrous DN, Le Moal M, Piazza PV. Release of endogenous dopamine in cultured mesencephalic neurons: influence of dopaminergic agonists and glucocorticoid antagonists. *Eur J Neurosci* 1999; 11(7): 2343-2350.
- [580] Rouge-Pont F, Deroche V, Moal ML, Piazza PV. Individual differences in stress-induced dopamine release in the nucleus accumbens are influenced by corticosterone. *Eur J Neurosci* 1998; 10(12): 3903-3907.
- [581] Roullet P, Lassalle JM. Genetic variation, hippocampal mossy fibres distribution, novelty reactions and spatial representation in mice. *Behav Brain Res* 1990; 41(1): 61-70.
- [582] Saal D, Dong Y, Bonci A, Malenka RC. Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. *Neuron* 2003; 37(4): 577-582.
- [583] Sabban EL, Kvetnansky R. Stress-triggered activation of gene expression in catecholaminergic systems: dynamics of transcriptional events. *Trends Neurosci* 2001; 24(2): 91-98.
- [584] Sajadi AA, Samaei SA, Rashidy-Pour A. Intra-hippocampal microinjections of anisomycin did not block glucocorticoid-induced impairment of memory retrieval in rats: an evidence for non-genomic effects of glucocorticoids. *Behav Brain Res* 2006; 173(1): 158-162.
- [585] Saladin ME, Drobos DJ, Coffey SF, Dansky BS, Brady KT, Kilpatrick DG. PTSD symptom severity as a predictor of cue-elicited drug craving in victims of violent crime. *Addict Behav* 2003; 28(9): 1611-1629.
- [586] Saleptsi E, Bichescu D, Rockstroh B, Neuner F, Schauer M, Studer K, Hoffmann K, Elbert T. Negative and positive childhood experiences across developmental periods in psychiatric patients with different diagnoses - an explorative study. *BMC Psychiatry* 2004; 4(1): 40.
- [587] Salomon L, Lanteri C, Glowinski J, Tassin JP. Behavioral sensitization to amphetamine results from an uncoupling between noradrenergic and serotonergic neurons. *Proc Natl Acad Sci U S A* 2006; 103(19): 7476-7481.
- [588] Samaha AN, Robinson TE. Why does the rapid delivery of drugs to the brain promote addiction? *Trends Pharmacol Sci* 2005; 26(2): 82-87.
- [589] Sandi C, Venero C, Guaza C. Novelty-related rapid locomotor effects of corticosterone in rats. *Eur J Neurosci* 1996; 8(4): 794-800.
- [590] Saphier D, Welch JE, Farrar GE, Goeders NE. Effects of intracerebroventricular and intrahypothalamic cocaine administration on adrenocortical secretion. *Neuroendocrinology* 1993; 57(1): 54-62.
- [591] Sapolsky RM, Romero LM, Munck AU. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev* 2000; 21(1): 55-89.
- [592] Sarnyai Z. Neurobiology of stress and cocaine addiction. Studies on corticotropin-releasing factor in rats, monkeys, and humans. *Ann N Y Acad Sci* 1998; 851: 371-387.
- [593] Sarnyai Z, Biro E, Gardi J, Vecsernyes M, Julesz J, Telegdy G. Brain corticotropin-releasing factor mediates 'anxiety-like' behavior induced by cocaine withdrawal in rats. *Brain Res* 1995; 675(1-2): 89-97.
- [594] Sarnyai Z, Biro E, Gardi J, Vecsernyes M, Julesz J, Telegdy G. Alterations of corticotropin-releasing

- factor-like immunoreactivity in different brain regions after acute cocaine administration in rats. *Brain Res* 1993; 616(1-2): 315-319.
- [595] Sarnyai Z, Biro E, Penke B, Telegdy G. The cocaine-induced elevation of plasma corticosterone is mediated by endogenous corticotropin-releasing factor (CRF) in rats. *Brain Res* 1992; 589(1): 154-156.
- [596] Sarnyai Z, Biro E, Telegdy G. Cocaine-induced elevation of plasma corticosterone is mediated by different neurotransmitter systems in rats. *Pharmacol Biochem Behav* 1993; 45(1): 209-214.
- [597] Sarnyai Z, Hohn J, Szabo G, Penke B. Critical role of endogenous corticotropin-releasing factor (CRF) in the mediation of the behavioral action of cocaine in rats. *Life Sci* 1992; 51(26): 2019-2024.
- [598] Sarnyai Z, McKittrick CR, McEwen BS, Kreek MJ. Selective regulation of dopamine transporter binding in the shell of the nucleus accumbens by adrenalectomy and corticosterone-replacement. *Synapse* 1998; 30(3): 334-337.
- [599] Sarnyai Z, Shaham Y, Heinrichs SC. The role of corticotropin-releasing factor in drug addiction. *Pharmacol Rev* 2001; 53(2): 209-243.
- [600] Sarnyai Z, Szabo G, Kovacs GL, Telegdy G. Opposite actions of oxytocin and vasopressin in the development of cocaine-induced behavioral sensitization in mice. *Pharmacol Biochem Behav* 1992; 43(2): 491-494.
- [601] Sarnyai Z, Veldhuis JD, Mello NK, Mendelson JH, Eros-Sarnyai M, Mercer G, Gelles H, Kelly M. The concordance of pulsatile ultradian release of adrenocorticotropin and cortisol in male rhesus monkeys. *J Clin Endocrinol Metab* 1995; 80(1): 54-59.
- [602] Schank JR, Ventura R, Puglisi-Allegra S, Alcaro A, Cole CD, Liles LC, Seeman P, Weinshenker D. Dopamine beta-hydroxylase knockout mice have alterations in dopamine signaling and are hypersensitive to cocaine. *Neuropsychopharmacology* 2006; 31: 2221-2230.
- [603] Scheinman RI, Gualberto A, Jewell CM, Cidowski JA, Baldwin AS, Jr. Characterization of mechanisms involved in transrepression of NF-kappa B by activated glucocorticoid receptors. *Mol Cell Biol* 1995; 15(2): 943-953.
- [604] Schenk S, Gittings D. Effects of SCH 23390 and eticlopride on cocaine-seeking produced by cocaine and WIN 35,428 in rats. *Psychopharmacology (Berl)* 2003; 168(1-2): 118-123.
- [605] Schenk S, Horger BA, Peltier R, Shelton K. Supersensitivity to the reinforcing effects of cocaine following 6-hydroxydopamine lesions to the medial prefrontal cortex in rats. *Brain Res* 1991; 543(2): 227-235.
- [606] Schenk S, Lacelle G, Gorman K, Amit Z. Cocaine self-administration in rats influenced by environmental conditions: implications for the etiology of drug abuse. *Neurosci Lett* 1987; 81(1-2): 227-231.
- [607] Schenk S, Partridge B. Sensitization to cocaine's reinforcing effects produced by various cocaine pretreatment regimens in rats. *Pharmacol Biochem Behav* 2000; 66(4): 765-770.
- [608] Schmidt ED, Schoffelmeer ANM, De Vries TJ, Wardeh G, Dogterom G, Bol JGJM, Binnekade R, Tilders FJH. A single administration of interleukin-1 or amphetamine induces long-lasting

- increases in evoked noradrenaline release in the hypothalamus and sensitization of ACTH and corticosterone responses in rats. *Eur J Neurosci* 2001; 13(10): 1923-1930.
- [609] Schmidt ED, Tilders FJ, Binnekade R, Schoffelmeer AN, De Vries TJ. Stressor- or drug-induced sensitization of the corticosterone response is not critically involved in the long-term expression of behavioural sensitization to amphetamine. *Neuroscience* 1999; 92(1): 343-352.
- [610] Schmidt ED, Tilders FJ, Janszen AW, Binnekade R, De Vries TJ, Schoffelmeer AN. Intermittent cocaine exposure causes delayed and long-lasting sensitization of cocaine-induced ACTH secretion in rats. *Eur J Pharmacol* 1995; 285(3): 317-321.
- [611] Seale TW, Carney JM. Genetic determinants of susceptibility to the rewarding and other behavioral actions of cocaine. *J Addict Dis* 1991; 10(1-2): 141-162.
- [612] See RE, Kruzich PJ, Grimm JW. Dopamine, but not glutamate, receptor blockade in the basolateral amygdala attenuates conditioned reward in a rat model of relapse to cocaine-seeking behavior. *Psychopharmacology (Berl)* 2001; 154(3): 301-310.
- [613] Self DW, Barnhart WJ, Lehman DA, Nestler EJ. Opposite modulation of cocaine-seeking behavior by D1- and D2-like dopamine receptor agonists. *Science* 1996; 271(5255): 1586-1589.
- [614] Self DW, Genova LM, Hope BT, Barnhart WJ, Spencer JJ, Nestler EJ. Involvement of cAMP-dependent protein kinase in the nucleus accumbens in cocaine self-administration and relapse of cocaine-seeking behavior. *J Neurosci* 1998; 18(5): 1848-1859.
- [615] Selye H. A syndrome produced by diverse nocuous agents. *Nature* 1936; 138: 32.
- [616] Setlow B, Roozendaal B, McGaugh JL. Involvement of a basolateral amygdala complex-nucleus accumbens pathway in glucocorticoid-induced modulation of memory consolidation. *Eur J Neurosci* 2000; 12(1): 367-375.
- [617] Shaham Y, Erb S, Leung S, Buczek Y, Stewart J. CP-154,526, a selective, non-peptide antagonist of the corticotropin-releasing factor1 receptor attenuates stress-induced relapse to drug seeking in cocaine- and heroin-trained rats. *Psychopharmacology (Berl)* 1998; 137(2): 184-190.
- [618] Shaham Y, Erb S, Stewart J. Stress-induced relapse to heroin and cocaine seeking in rats: a review. *Brain Res Brain Res Rev* 2000; 33(1): 13-33.
- [619] Shaham Y, Funk D, Erb S, Brown TJ, Walker CD, Stewart J. Corticotropin-releasing factor, but not corticosterone, is involved in stress-induced relapse to heroin-seeking in rats. *J Neurosci* 1997; 17(7): 2605-2614.
- [620] Shaham Y, Highfield D, Delfs J, Leung S, Stewart J. Clonidine blocks stress-induced reinstatement of heroin seeking in rats: an effect independent of locus coeruleus noradrenergic neurons. *Eur J Neurosci* 2000; 12(1): 292-302.
- [621] Shalev U, Marinelli M, Baumann MH, Piazza PV, Shaham Y. The role of corticosterone in food deprivation-induced reinstatement of cocaine seeking in the rat. *Psychopharmacology (Berl)* 2003; 168(1-2): 170-176.
- [622] Shanks N, Griffiths J, Zalcman S, Zacharko RM, Anisman H. Mouse strain differences in plasma corticosterone following uncontrollable footshock. *Pharmacol Biochem Behav* 1990; 36(3): 515-519.
- [623] Shilling PD, Kelsoe JR, Segal DS. Dopamine transporter mRNA is up-regulated in the substantia

- nigra and the ventral tegmental area of amphetamine-sensitized rats. *Neurosci Lett* 1997; 236(3): 131-134.
- [624] Shippenberg TS, Heidbreder C. Sensitization to the conditioned rewarding effects of cocaine: pharmacological and temporal characteristics. *J Pharmacol Exp Ther* 1995; 273(2): 808-815.
- [625] Shoaib M, Swanner LS, Beyer CE, Goldberg SR, Schindler CW. The GABAB agonist baclofen modifies cocaine self-administration in rats. *Behav Pharmacol* 1998; 9(3): 195-206.
- [626] Simar MR, Saphier D, Goeders NE. Differential neuroendocrine and behavioral responses to cocaine in Lewis and Fischer rats. *Neuroendocrinology* 1996; 63(1): 93-100.
- [627] Sinha R, Garcia M, Paliwal P, Kreek MJ, Rounsaville BJ. Stress-induced cocaine craving and hypothalamic-pituitary-adrenal responses are predictive of cocaine relapse outcomes. *Arch Gen Psychiatry* 2006; 63(3): 324-331.
- [628] Sinha R, Lacadie C, Skudlarski P, Fulbright RK, Rounsaville BJ, Kosten TR, Wexler BE. Neural activity associated with stress-induced cocaine craving: a functional magnetic resonance imaging study. *Psychopharmacology (Berl)* 2005; 183(2): 171-180.
- [629] Smith JE, Co C, Yin X, Sizemore GM, Liguori A, Johnson III, Martin TJ. Involvement of cholinergic neuronal systems in intravenous cocaine self-administration. *Neurosci Biobehav Rev* 2004; 27(8): 841-850.
- [630] Sokoloff P, Giros B, Martres MP, Bouthenet ML, Schwartz JC. Molecular cloning and characterization of a novel dopamine receptor (D3) as a target for neuroleptics. *Nature* 1990; 347(6289): 146-151.
- [631] Sonuga-Barke EJS. Causal models of attention-deficit/hyperactivity disorder: from common simple deficits to multiple developmental pathways. *Biol Psychiatry* 2005; 57(11): 1231-1238.
- [632] Sora I, Wichems C, Takahashi N, Li XF, Zeng Z, Revay R, Lesch KP, Murphy DL, Uhl GR. Cocaine reward models: conditioned place preference can be established in dopamine- and in serotonin-transporter knockout mice. *Proc Natl Acad Sci U S A* 1998; 95(13): 7699-7704.
- [633] Sorg BA, Chen SY, Kalivas PW. Time course of tyrosine hydroxylase expression after behavioral sensitization to cocaine. *J Pharmacol Exp Ther* 1993; 266(1): 424-430.
- [634] Sorg BA, Davidson DL, Kalivas PW, Prasad BM. Repeated daily cocaine alters subsequent cocaine-induced increase of extracellular dopamine in the medial prefrontal cortex. *J Pharmacol Exp Ther* 1997; 281(1): 54-61.
- [635] Sorg BA, Kalivas PW. Effects of cocaine and footshock stress on extracellular dopamine levels in the ventral striatum. *Brain Res* 1991; 559(1): 29-36.
- [636] Spencer RL, Young EA, Choo PH, McEwen BS. Adrenal steroid type I and type II receptor binding: estimates of in vivo receptor number, occupancy, and activation with varying level of steroid. *Brain Res* 1990; 514(1): 37-48.
- [637] Stachowiak MK, Rigual RJ, Lee PH, Viveros OH, Hong JS. Regulation of tyrosine hydroxylase and phenylethanolamine N-methyltransferase mRNA levels in the sympathoadrenal system by the pituitary-adrenocortical axis. *Brain Res* 1988; 427(3): 275-286.
- [638] Stein-Behrens BA, Lin WJ, Sapolsky RM. Physiological elevations of glucocorticoids potentiate glutamate accumulation in the hippocampus. *J Neurochem* 1994; 63(2): 596-602.

- [639] Steketee JD. Neurotransmitter systems of the medial prefrontal cortex: potential role in sensitization to psychostimulants. *Brain Res Brain Res Rev* 2003; 41(2-3): 203-228.
- [640] Stern E, Silbersweig DA, Chee KY, Holmes A, Robertson MM, Trimble M, Frith CD, Frackowiak RS, Dolan RJ. A functional neuroanatomy of tics in Tourette syndrome. *Arch Gen Psychiatry* 2000; 57(8): 741-748.
- [641] Sternberg DB, Isaacs KR, Gold PE, McGaugh JL. Epinephrine facilitation of appetitive learning: attenuation with adrenergic receptor antagonists. *Behav Neural Biol* 1985; 44(3): 447-453.
- [642] Sulzer D, Sonders MS, Poulsen NW, Galli A. Mechanisms of neurotransmitter release by amphetamines: a review. *Prog Neurobiol* 2005; 75(6): 406-433.
- [643] Sun W, Akins CK, Mattingly AE, Rebec GV. Ionotropic glutamate receptors in the ventral tegmental area regulate cocaine-seeking behavior in rats. *Neuropsychopharmacology* 2005; 30(11): 2073-2081.
- [644] Sun W, Rebec GV. The role of prefrontal cortex D1-like and D2-like receptors in cocaine-seeking behavior in rats. *Psychopharmacology (Berl)* 2005; 177(3): 315-323.
- [645] Sunahara RK, Guan HC, O'Dowd BF, Seeman P, Laurier LG, Ng G, George SR, Torchia J, Van Tol HH, Niznik HB. Cloning of the gene for a human dopamine D5 receptor with higher affinity for dopamine than D1. *Nature* 1991; 350(6319): 614-619.
- [646] Sunahara RK, Niznik HB, Weiner DM, Stormann TM, Brann MR, Kennedy JL, Gelernter JE, Rozmahel R, Yang YL, Israel Y. Human dopamine D1 receptor encoded by an intronless gene on chromosome 5. *Nature* 1990; 347(6288): 80-83.
- [647] Suto N, Austin JD, Tanabe LM, Kramer MK, Wright DA, Vezina P. Previous exposure to VTA amphetamine enhances cocaine self-administration under a progressive ratio schedule in a D1 dopamine receptor dependent manner. *Neuropsychopharmacology* 2002; 27(6): 970-979.
- [648] Svenningsson P, Nairn AC, Greengard P. DARPP-32 mediates the actions of multiple drugs of abuse. *AAPS J* 2005; 7(2): E353-E360.
- [649] Svingos AL, Periasamy S, Pickel VM. Presynaptic dopamine D(4) receptor localization in the rat nucleus accumbens shell. *Synapse* 2000; 36(3): 222-232.
- [650] Swanson CJ, Kalivas PW. Regulation of locomotor activity by metabotropic glutamate receptors in the nucleus accumbens and ventral tegmental area. *J Pharmacol Exp Ther* 2000; 292(1): 406-414.
- [651] Szantai E, Szmola R, Sasvari-Szekely M, Guttman A, Ronai Z. The polymorphic nature of the human dopamine D4 receptor gene: a comparative analysis of known variants and a novel 27 bp deletion in the promoter region. *BMC Genet* 2005; 6(1): 39.
- [652] Szilagyi A, Boor K, Szekely A, Gaszner P, Kalasz H, Sasvari-Szekely M, Barta C. Combined effect of promoter polymorphisms in the dopamine D4 receptor and the serotonin transporter genes in heroin dependence. *Neuropsychopharmacol Hung* 2005; 7(1): 28-33.
- [653] Szumlinski KK, Dehoff MH, Kang SH, Frys KA, Lominac KD, Klugmann M, Rohrer J, Griffin III, Toda S. Homer proteins regulate sensitivity to cocaine. *Neuron* 2004; 43(3): 401-413.
- [654] Taber MT, Das S, Fibiger HC. Cortical regulation of subcortical dopamine release: mediation via the ventral tegmental area. *J Neurochem* 1995; 65(3): 1407-1410.

- [655] Takahashi T, Kimoto T, Tanabe N, Hattori TA, Yasumatsu N, Kawato S. Corticosterone acutely prolonged N-methyl-d-aspartate receptor-mediated Ca<sup>2+</sup> elevation in cultured rat hippocampal neurons. *J Neurochem* 2002; 83(6): 1441-1451.
- [656] Tanaka SC, Doya K, Okada G, Ueda K, Okamoto Y, Yamawaki S. Prediction of immediate and future rewards differentially recruits cortico-basal ganglia loops. *Nat Neurosci* 2004; 7(8): 887-893.
- [657] Tang WX, Fasulo WH, Mash DC, Hemby SE. Molecular profiling of midbrain dopamine regions in cocaine overdose victims. *J Neurochem* 2003; 85(4): 911-924.
- [658] Tessari M, Pilla M, Andreoli M, Hutcheson DM, Heidbreder CA. Antagonism at metabotropic glutamate 5 receptors inhibits nicotine- and cocaine-taking behaviours and prevents nicotine-triggered relapse to nicotine-seeking. *Eur J Pharmacol* 2004; 499(1-2): 121-133.
- [659] Tiberi M, Jarvie KR, Silvia C, Falardeau P, Gingrich JA, Godinot N, Bertrand L, Yang-Feng TL, Fremeau RT, Jr., Caron MG. Cloning, molecular characterization, and chromosomal assignment of a gene encoding a second D1 dopamine receptor subtype: differential expression pattern in rat brain compared with the D1A receptor. *Proc Natl Acad Sci U S A* 1991; 88(17): 7491-7495.
- [660] Tidey JW, Miczek KA. Acquisition of cocaine self-administration after social stress: role of accumbens dopamine. *Psychopharmacology (Berl)* 1997; 130(3): 203-212.
- [661] Tiffany ST. A cognitive model of drug urges and drug-use behavior: role of automatic and nonautomatic processes. *Psychol Rev* 1990; 97(2): 147-168.
- [662] Toda S, McGinty JF, Kalivas PW. Repeated cocaine administration alters the expression of genes in corticolimbic circuitry after a 3-week withdrawal: a DNA macroarray study. *J Neurochem* 2002; 82(5): 1290-1299.
- [663] Todtenkopf MS, Mihalakopoulos A, Stellar JR. Withdrawal duration differentially affects c-fos expression in the medial prefrontal cortex and discrete subregions of the nucleus accumbens in cocaine-sensitized rats. *Neuroscience* 2002; 114(4): 1061-1069.
- [664] Tolliver BK, Belknap JK, Woods WE, Carney JM. Genetic analysis of sensitization and tolerance to cocaine. *J Pharmacol Exp Ther* 1994; 270(3): 1230-1238.
- [665] Tolliver BK, Carney JM. Locomotor stimulant effects of cocaine and novel cocaine analogs in DBA/2J and C57BL/6J inbred mice. *Pharmacol Biochem Behav* 1995; 50(2): 163-169.
- [666] Tolliver BK, Carney JM. Sensitization to stereotypy in DBA/2J but not C57BL/6J mice with repeated cocaine. *Pharmacol Biochem Behav* 1994; 48(1): 169-173.
- [667] Tracy JL, de Leon J, Qureshi G, McCann EM, McGrory A, Josiassen RC. Repetitive behaviors in schizophrenia: a single disturbance or discrete symptoms? *Schizophr Res* 1996; 20(1-2): 221-229.
- [668] Trapp T, Holsboer F. Heterodimerization between mineralocorticoid and glucocorticoid receptors increases the functional diversity of corticosteroid action. *Trends Pharmacol Sci* 1996; 17(4): 145-149.
- [669] Tricot JP. Cocaine: half a century of therapeutic use (1880-1930). *Verh K Acad Geneesk Belg* 1991; 53(5): 487-496.
- [670] Tsuang MT, Lyons MJ, Eisen SA, Goldberg J, True W, Lin N, Meyer JM, Toomey R, Faraone SV,

- Eaves L. Genetic influences on DSM-III-R drug abuse and dependence: a study of 3,372 twin pairs. *Am J Med Genet* 1996; 67(5): 473-477.
- [671] Tsujimoto S, Sawaguchi T. Neuronal activity representing temporal prediction of reward in the primate prefrontal cortex. *J Neurophysiol* 2005; 93(6): 3687-3692.
- [672] U'Prichard DC, Bylund DB, Snyder SH. (+/-)-[3H]Epinephrine and (-)[3H]dihydroalprenolol binding to beta1- and beta2-noradrenergic receptors in brain, heart, and lung membranes. *J Biol Chem* 1978; 253(14): 5090-5102.
- [673] U'Prichard DC, Snyder SH. Differential labelling of alpha and beta-noradrenergic receptors in calf cerebellum membranes with 3H-adrenaline. *Nature* 1977; 270(5634): 261-263.
- [674] U'Prichard DC, Greenberg DA, Sheehan P, Snyder SH. Regional distribution of alpha-noradrenergic receptor binding in calf brain. *Brain Res* 1977; 138(1): 151-158.
- [675] Ujike H, Akiyama K, Otsuki S. D-2 but not D-1 dopamine agonists produce augmented behavioral response in rats after subchronic treatment with methamphetamine or cocaine. *Psychopharmacology (Berl)* 1990; 102(4): 459-464.
- [676] Ungless MA, Whistler JL, Malenka RC, Bonci A. Single cocaine exposure in vivo induces long-term potentiation in dopamine neurons. *Nature* 2001; 411(6837): 583-587.
- [677] Upchurch M, Wehner JM. Differences between inbred strains of mice in Morris water maze performance. *Behav Genet* 1988; 18(1): 55-68.
- [678] Valadez A, Schenk S. Persistence of the ability of amphetamine preexposure to facilitate acquisition of cocaine self-administration. *Pharmacol Biochem Behav* 1994; 47(1): 203-205.
- [679] Valjent E, Corvol JC, Pages C, Besson MJ, Maldonado R, Caboche J. Involvement of the extracellular signal-regulated kinase cascade for cocaine-rewarding properties. *J Neurosci* 2000; 20(23): 8701-8709.
- [680] Van Bockstaele EJ, Pickel VM. GABA-containing neurons in the ventral tegmental area project to the nucleus accumbens in rat brain. *Brain Res* 1995; 682(1-2): 215-221.
- [681] van den Buuse M, Morris M, Chavez C, Martin S, Wang J. Effect of adrenalectomy and corticosterone replacement on prepulse inhibition and locomotor activity in mice. *Br J Pharmacol* 2004; 142(3): 543-550.
- [682] van der Elst MC, Roubos EW, Ellenbroek BA, Veening JG, Cools AR. Apomorphine-susceptible rats and apomorphine-unsusceptible rats differ in the tyrosine hydroxylase-immunoreactive network in the nucleus accumbens core and shell. *Exp Brain Res* 2005; 160(4): 418-423.
- [683] van der Elst MCJ, Verheij MMM, Roubos EW, Ellenbroek BA, Veening JG, Cools AR. A single exposure to novelty differentially affects the accumbal dopaminergic system of apomorphine-susceptible and apomorphine-unsusceptible rats. *Life Sci* 2005; 76(12): 1391-1406.
- [684] van der Kam EL, Ellenbroek BA, Cools AR. Gene - environment interactions determine the individual variability in cocaine self-administration. *Neuropharmacology* 2005; 48(5): 685-695.
- [685] Van Eekelen JA, De Kloet ER. Co-localization of brain corticosteroid receptors in the rat hippocampus. *Prog Histochem Cytochem* 1992; 26(1-4): 250-258.
- [686] Van Eekelen JA, Jiang W, De Kloet ER, Bohn MC. Distribution of the mineralocorticoid and the glucocorticoid receptor mRNAs in the rat hippocampus. *J Neurosci Res* 1988; 21(1): 88-94.

- [687] Van Ree JM, Burbach-Bloemarts EM, Wallace M. Vasopressin neuropeptides and acquisition of heroin and cocaine self-administration in rats. *Life Sci* 1988; 42(10): 1091-1099.
- [688] Van Ree JM, Niesink RJM, Van Wolfswinkel L, Ramsey NF, Kornet M, Van Furth WR, Vanderschuren LJM, Gerrits MAFM, Van den Berg CL. Endogenous opioids and reward. *Eur J Pharmacol* 2000; 405(1-3): 89-101.
- [689] Van Tol HH, Bunzow JR, Guan HC, Sunahara RK, Seeman P, Niznik HB, Civelli O. Cloning of the gene for a human dopamine D4 receptor with high affinity for the antipsychotic clozapine. *Nature* 1991; 350(6319): 610-614.
- [690] Van Tol HH, Wu CM, Guan HC, Ohara K, Bunzow JR, Civelli O, Kennedy J, Seeman P, Niznik HB, Jovanovic V. Multiple dopamine D4 receptor variants in the human population. *Nature* 1992; 358(6382): 149-152.
- [691] Vanderschuren LJ, Beemster P, Schoffelmeer AN. On the role of noradrenaline in psychostimulant-induced psychomotor activity and sensitization. *Psychopharmacology (Berl)* 2003; 169(2): 176-185.
- [692] Vanderschuren LJ, Everitt BJ. Drug seeking becomes compulsive after prolonged cocaine self-administration. *Science* 2004; 305(5686): 1017-1019.
- [693] Vanderschuren LJ, Kalivas PW. Alterations in dopaminergic and glutamatergic transmission in the induction and expression of behavioral sensitization: a critical review of preclinical studies. *Psychopharmacology (Berl)* 2000; 151(2-3): 99-120.
- [694] Vanderschuren LJ, Schmidt ED, De Vries TJ, Van Moorsel CA, Tilders FJ, Schoffelmeer AN. A single exposure to amphetamine is sufficient to induce long-term behavioral, neuroendocrine, and neurochemical sensitization in rats. *J Neurosci* 1999; 19(21): 9579-9586.
- [695] Vanderschuren LJ, Schoffelmeer AN, Mulder AH, De Vries TJ. Dopaminergic mechanisms mediating the long-term expression of locomotor sensitization following pre-exposure to morphine or amphetamine. *Psychopharmacology (Berl)* 1999; 143(3): 244-253.
- [696] Vanderschuren LJ, Schoffelmeer AN, Van Leeuwen SD, Hof L, Jonker AJ, Voorn P. Compartment-specific changes in striatal neuronal activity during expression of amphetamine sensitization are the result of drug hypersensitivity. *Eur J Neurosci* 2002; 16(12): 2462-2468.
- [697] Vanderschuren LJM, Di Ciano P, Everitt BJ. Involvement of the dorsal striatum in cue-controlled cocaine seeking. *J Neurosci* 2005; 25(38): 8665-8670.
- [698] Vanderschuren LJM, Everitt BJ. Behavioral and neural mechanisms of compulsive drug seeking. *Eur J Pharmacol* 2005; 526(1-3): 77-88.
- [699] Venero C, Borrell J. Rapid glucocorticoid effects on excitatory amino acid levels in the hippocampus: a microdialysis study in freely moving rats. *Eur J Neurosci* 1999; 11(7): 2465-2473.
- [700] Ventura R, Alcaro A, Cabib S, Conversi D, Mandolesi L, Puglisi-Allegra S. Dopamine in the medial prefrontal cortex controls genotype-dependent effects of amphetamine on mesoaccumbens dopamine release and locomotion. *Neuropsychopharmacology* 2004; 29(1): 72-80.
- [701] Ventura R, Alcaro A, Mandolesi L, Puglisi-Allegra S. In vivo evidence that genetic background controls impulse-dependent dopamine release induced by amphetamine in the nucleus accumbens. *J Neurochem* 2004; 89(2): 494-502.

- [702] Ventura R, Cabib S, Alcaro A, Orsini C, Puglisi-Allegra S. Norepinephrine in the prefrontal cortex is critical for amphetamine-induced reward and mesoaccumbens dopamine release. *J Neurosci* 2003; 23(5): 1879-1885.
- [703] Ventura R, Cabib S, Puglisi-Allegra S. Genetic susceptibility of mesocortical dopamine to stress determines liability to inhibition of mesoaccumbens dopamine and to behavioral 'despair' in a mouse model of depression. *Neuroscience* 2002; 115(4): 999-1007.
- [704] Ventura R, Cabib S, Puglisi-Allegra S. Opposite genotype-dependent mesocorticolimbic dopamine response to stress. *Neuroscience* 2001; 104(3): 627-631.
- [705] Ventura R, Puglisi-Allegra S. Environment makes amphetamine-induced dopamine release in the nucleus accumbens totally impulse-dependent. *Synapse* 2005; 58(3): 211-214.
- [706] Vezina P, Stewart J. The effect of dopamine receptor blockade on the development of sensitization to the locomotor activating effects of amphetamine and morphine. *Brain Res* 1989; 499(1): 108-120.
- [707] Vezina P, Stewart J. Amphetamine administered to the ventral tegmental area but not to the nucleus accumbens sensitizes rats to systemic morphine: lack of conditioned effects. *Brain Res* 1990; 516(1): 99-106.
- [708] Vezina P. Sensitization of midbrain dopamine neuron reactivity and the self-administration of psychomotor stimulant drugs. *Neurosci Biobehav Rev* 2004; 27(8): 827-839.
- [709] Vezina P, Lorrain DS, Arnold GM, Austin JD, Suto N. Sensitization of midbrain dopamine neuron reactivity promotes the pursuit of amphetamine. *J Neurosci* 2002; 22(11): 4654-4662.
- [710] Volkow N, Li TK. The neuroscience of addiction. *Nat Neurosci* 2005; 8(11): 1429-1430.
- [711] Volkow ND, Wang GJ, Telang F, Fowler JS, Logan J, Childress AR, Jayne M, Ma Y, Wong C. Cocaine cues and dopamine in dorsal striatum: mechanism of craving in cocaine addiction. *J Neurosci* 2006; 26(24): 6583-6588.
- [712] Vorel SR, Ashby CR, Jr., Paul M, Liu X, Hayes R, Hagan JJ, Middlemiss DN, Stemp G, Gardner EL. Dopamine D3 receptor antagonism inhibits cocaine-seeking and cocaine-enhanced brain reward in rats. *J Neurosci* 2002; 22(21): 9595-9603.
- [713] Vrana SL, Vrana KE, Koves TR, Smith JE, Dworkin SI. Chronic cocaine administration increases CNS tyrosine hydroxylase enzyme activity and mRNA levels and tryptophan hydroxylase enzyme activity levels. *J Neurochem* 1993; 61(6): 2262-2268.
- [714] Wachtel SR, Charnot A, De Wit H. Acute hydrocortisone administration does not affect subjective responses to d-amphetamine in humans. *Psychopharmacology (Berl)* 2001; 153(3): 380-388.
- [715] Wagner FA, Anthony JC. From first drug use to drug dependence; developmental periods of risk for dependence upon marijuana, cocaine, and alcohol. *Neuropsychopharmacology* 2002; 26(4): 479-488.
- [716] Walker CD, Rivest RW, Meaney MJ, Aubert ML. Differential activation of the pituitary-adrenocortical axis after stress in the rat: use of two genetically selected lines (Roman low- and high-avoidance rats) as a model. *J Endocrinol* 1989; 123(3): 477-485.
- [717] Walters CL, Blendy JA. Different requirements for cAMP response element binding protein in

- positive and negative reinforcing properties of drugs of abuse. *J Neurosci* 2001; 21(23): 9438-9444.
- [718] Wang JB, Liu ZF, Kofuji P, Burt DR. The GABA(A) receptor gamma1-subunit in seizure prone (DBA/2) and resistant (C57BL/6) mice. *Brain Res Bull* 1998; 45(4): 421-425.
- [719] Ward AS, Collins ED, Haney M, Foltin RW, Fischman MW. Ketoconazole attenuates the cortisol response but not the subjective effects of smoked cocaine in humans. *Behav Pharmacol* 1998; 9(7): 577-586.
- [720] Watanabe Y, Weiland NG, McEwen BS. Effects of adrenal steroid manipulations and repeated restraint stress on dynorphin mRNA levels and excitatory amino acid receptor binding in hippocampus. *Brain Res* 1995; 680(1-2): 217-225.
- [721] Wee S, Anderson KG, Baumann MH, Rothman RB, Blough BE, Woolverton WL. Relationship between the serotonergic activity and reinforcing effects of a series of amphetamine analogs. *J Pharmacol Exp Ther* 2005; 313(2): 848-854.
- [722] Wee S, Wang Z, He R, Zhou J, Kozikowski AP, Woolverton WL. Role of the increased noradrenergic neurotransmission in drug self-administration. *Drug Alcohol Depend* 2006; 82(2): 151-157.
- [723] Wei P, Inamdar N, Vedeckis WV. Transrepression of c-jun gene expression by the glucocorticoid receptor requires both AP-1 sites in the c-jun promoter. *Mol Endocrinol* 1998; 12(9): 1322-1333.
- [724] Weidenfeld J, Newman ME, Itzik A, Feldman S. Adrenocortical axis responses to adrenergic and glutamate stimulation are regulated by the amygdala. *Neuroreport* 2005; 16(11): 1245-1249.
- [725] Weil-Malherbe H, Axelrod J, Tomchick R. Blood-brain barrier for adrenaline. *Science* 1959; 129(3357): 1226-1227.
- [726] Weiner DM, Levey AI, Sunahara RK, Niznik HB, O'Dowd BF, Seeman P, Brann MP. D1 and D2 Dopamine Receptor mRNA in Rat Brain. *Proc Natl Acad Sci U S A* 1991; 88(5): 1859-1863.
- [727] Weinshank RL, Adham N, Macchi M, Olsen MA, Branchek TA, Hartig PR. Molecular cloning and characterization of a high affinity dopamine receptor (D1 beta) and its pseudogene. *J Biol Chem* 1991; 266(33): 22427-22435.
- [728] Weiss IC, Domeney AM, Heidbreder CA, Moreau JL, Feldon J. Early social isolation, but not maternal separation, affects behavioral sensitization to amphetamine in male and female adult rats. *Pharmacol Biochem Behav* 2001; 70(2-3): 397-409.
- [729] Wiener HL, Reith MEA. Correlation between cocaine-induced locomotion and cocaine disposition in the brain among four inbred strains of mice. *Pharmacol Biochem Behav* 1990; 36(3): 699-701.
- [730] Williams CL, McGaugh JL. Reversible lesions of the nucleus of the solitary tract attenuate the memory-modulating effects of posttraining epinephrine. *Behav Neurosci* 1993; 107(6): 955-962.
- [731] Wills TA, Vaccaro D, McNamara G. Novelty seeking, risk taking, and related constructs as predictors of adolescent substance use: an application of Cloninger's theory. *J Subst Abuse* 1994; 6(1): 1-20.
- [732] Windle RJ, Wood SA, Shanks N, Lightman SL, Ingram CD. Ultradian rhythm of basal corticosterone

- release in the female rat: dynamic interaction with the response to acute stress. *Endocrinology* 1998; 139(2): 443-450.
- [733] Wise RA, Bozarth MA. A psychomotor stimulant theory of addiction. *Psychol Rev* 1987; 94(4): 469-492.
- [734] Wise RA, Newton P, Leeb K, Burnette B, Pocock D, Justice JB, Jr. Fluctuations in nucleus accumbens dopamine concentration during intravenous cocaine self-administration in rats. *Psychopharmacologia* 1995; 120(1): 10-20.
- [735] Wise RA, Rompre PP. Brain dopamine and reward. *Annu Rev Psychol* 1989; 40: 191-225.
- [736] Wolf ME, White FJ, Nassar R, Brooderson RJ, Khansa MR. Differential development of autoreceptor subsensitivity and enhanced dopamine release during amphetamine sensitization. *J Pharmacol Exp Ther* 1993; 264(1): 249-255.
- [737] Womer DE, Jones BC, Erwin VG. Characterization of dopamine transporter and locomotor effects of cocaine, GBR 12909, epidepride, and SCH 23390 in C57BL and DBA mice. *Pharmacol Biochem Behav* 1994; 48(2): 327-335.
- [738] Wong DL. Epinephrine biosynthesis: hormonal and neural control during stress. *Cell Mol Neurobiol* 2006; 26(4-6): 889-898.
- [739] Woolverton WL. Evaluation of the role of norepinephrine in the reinforcing effects of psychomotor stimulants in rhesus monkeys. *Pharmacol Biochem Behav* 1987; 26(4): 835-839.
- [740] Xi ZX, Gilbert J, Campos AC, Kline N, Ashby CR, Jr., Hagan JJ, Heidbreder CA, Gardner EL. Blockade of mesolimbic dopamine D3 receptors inhibits stress-induced reinstatement of cocaine-seeking in rats. *Psychopharmacology (Berl)* 2004; 176(1): 57-65.
- [741] Xi ZX, Newman AH, Gilbert JG, Pak AC, Peng XQ, Ashby CR, Jr., Gitajn L, Gardner EL. The novel dopamine D3 receptor antagonist NGB 2904 inhibits cocaine's rewarding effects and cocaine-induced reinstatement of drug-seeking behavior in rats. *Neuropsychopharmacology* 2006; 31(7): 1393-1405.
- [742] Xi ZX, Ramamoorthy S, Baker DA, Shen H, Samuvel DJ, Kalivas PW. Modulation of group II metabotropic glutamate receptor signaling by chronic cocaine. *J Pharmacol Exp Ther* 2002; 303(2): 608-615.
- [743] Yajie D, Lin K, Baoming L, Lan M. Enhanced cocaine self-administration in adult rats with adolescent isolation experience. *Pharmacol Biochem Behav* 2005; 82(4): 673-677.
- [744] Yamada S, Yokoo H, Nishi S. Changes in sensitivity of dopamine autoreceptors in rat striatum after subchronic treatment with methamphetamine. *Eur J Pharmacol* 1991; 205(1): 43-47.
- [745] Yamaguchi K, Hama H, Watanabe K, Adachi C. Effect of dopamine injection into the anteroventral third ventricular region and the paraventricular nucleus on vasopressin secretion in conscious rats. *Acta Endocrinol (Copenh)* 1992; 127(5): 420-424.
- [746] Yamaguchi M, Suzuki T, Abe S, Hori T, Kurita H, Asada T, Okado N, Arai H. Repeated cocaine administration differentially affects NMDA receptor subunit (NR1, NR2A-C) mRNAs in rat brain. *Synapse* 2002; 46(3): 157-169.
- [747] Yentis SM, Vlassakov KV. Vassily von Anrep, forgotten pioneer of regional anesthesia. *Anesthesiology* 1999; 90(3): 890-895.

- [748] Yi SJ, Johnson KM. Chronic cocaine treatment impairs the regulation of synaptosomal 3H-DA release by D2 autoreceptors. *Pharmacol Biochem Behav* 1990; 36(3): 457-461.
- [749] Yin HH, Knowlton BJ, Balleine BW. Lesions of dorsolateral striatum preserve outcome expectancy but disrupt habit formation in instrumental learning. *Eur J Neurosci* 2004; 19(1): 181-189.
- [750] Yokel RA, Wise RA. Attenuation of intravenous amphetamine reinforcement by central dopamine blockade in rats. *Psychopharmacology (Berl)* 1976; 48(3): 311-318.
- [751] Young EA, Abelson J, Lightman SL. Cortisol pulsatility and its role in stress regulation and health. *Front Neuroendocrinol* 2004; 25(2): 69-76.
- [752] Zaborszky L, Alheid GF, Beinfeld MC, Eiden LE, Heimer L, Palkovits M. Cholecystokinin innervation of the ventral striatum: a morphological and radioimmunological study. *Neuroscience* 1985; 14(2): 427-453.
- [753] Zaborszky L, Vadasz C. The midbrain dopaminergic system: anatomy and genetic variation in dopamine neuron number of inbred mouse strains. *Behav Genet* 2001; 31(1): 47-59.
- [754] Zahm DS, Brog JS. On the significance of subterritories in the "accumbens" part of the rat ventral striatum. *Neuroscience* 1992; 50(4): 751-767.
- [755] Zhang L, Zhou R, Li X, Ursano RJ, Li H. Stress-induced change of mitochondria membrane potential regulated by genomic and non-genomic GR signaling: a possible mechanism for hippocampus atrophy in PTSD. *Med Hypotheses* 2006; 66(6): 1205-1208.
- [756] Zhang XY, Sanchez H, Kehoe P, Kosten TA. Neonatal isolation enhances maintenance but not reinstatement of cocaine self-administration in adult male rats. *Psychopharmacology (Berl)* 2005; 177(4): 391-399.
- [757] Zhou QY, Grandy DK, Thambi L, Kushner JA, Tol HHMV, Cone R, Pribnow D, Salon J, Bunzow JR, Civelli O. Cloning and expression of human and rat D1 dopamine receptors. *Nature* 1990; 347(6288): 76-80.
- [758] Zhou Y, Spangler R, Ho A, Jeanne Kreek M. Increased CRH mRNA levels in the rat amygdala during short-term withdrawal from chronic 'binge' cocaine. *Mol Brain Res* 2003; 114(1): 73-79.
- [759] Zhu BG, Zhu DH, Chen YZ. Rapid enhancement of high affinity glutamate uptake by glucocorticoids in rat cerebral cortex synaptosomes and human neuroblastoma clone SK-N-SH: possible involvement of G-protein. *Biochem Biophys Res Commun* 1998; 247(2): 261-265.
- [760] Zocchi A, Orsini C, Cabib S, Puglisi-Allegra S. Parallel strain-dependent effect of amphetamine on locomotor activity and dopamine release in the nucleus accumbens: an in vivo study in mice. *Neuroscience* 1998; 82(2): 521-528.
- [761] Zubieta JK, Gorelick DA, Stauffer R, Ravert HT, Dannals RF, Frost JJ. Increased mu opioid receptor binding detected by PET in cocaine-dependent men is associated with cocaine craving. *Nat Med* 1996; 2(11): 1225-1229.

