



Universiteit
Leiden

The Netherlands

Interference effects with surface plasmons

Kuzmin, N.V.

Citation

Kuzmin, N. V. (2008, January 10). *Interference effects with surface plasmons*. *Casimir PhD Series*. LION, Quantum Optics Group, Faculty of Science, Leiden University. Retrieved from <https://hdl.handle.net/1887/12551>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12551>

Note: To cite this publication please use the final published version (if applicable).

STELLINGEN

behorende bij het proefschrift
Interference effects with surface plasmons

I

The magnetic field is the most convenient tool for describing effects of surface plasmons on plane and structured metallo-dielectric interfaces.

This thesis, Chapter 1.

II

In plasmon-assisted two-slit experiments interference takes place both on the observation screen and in the slits.

This thesis, Chapter 2.

III

The effect of plasmonic cross-talk can easily be missed by choosing appropriate illumination conditions.

This thesis, Chapter 2.

IV

Ohmic damping in the metal is responsible for the retardation-plate like behavior of a single subwavelength-wide slit in a metal film.

This thesis, Chapter 6.

V

The ten orders of magnitude size difference between a melting iceberg and a nanoparticle changing its phase has welcome implications for nanophotonics: in the nanoworld spring and autumn may take only a few billionths of a second.

All change, please, N.I. Zheludev, Nature Photonics, 1, p.551 (2007).

VI

The ratio of the real and imaginary parts of the surface-plasmon propagation constant provides a useful figure-of-merit for comparing plasmonic materials for use at short wavelengths.

VII

A *hyperlens* is a more convenient device than a *superlens* (perfect lens) for imaging objects with subwavelength resolution.

Negative refraction makes a perfect lens,

*J.B. Pendry, PRL **85**, p.3966 (2000).*

Optical hyperlens: far-field imaging beyond the diffraction limit,

*Z. Jacob, L.V. Alekseyev and E. Narimanov, Opt. Exp. **14**, p.8247 (2006).*

VIII

It is deceptive to ignore the damping of a surface plasmon when describing its properties.

Plasmonics: optics at the nanoscale,

*A. Polman and H.A. Atwater, Materials Today, **8**, p.56 (2005).*

IX

Myriads of fractals are daily produced and thrown out into trash bins.

Mechanically deformed crumpled surfaces,

M.A.F. Gomes, T.I. Ren, I.M. Rodrigues and C.B.S. Furtado,

*J.Phys.D **22**, p.1217 (1989).*

Geometry of Crumpled Paper, D.L. Blair and A. Kudrolli,

*PRL **94**, p.166107 (2005).*

X

Glasses are nowadays made from plastic.

XI

Parties and optical experiments thrive when the lights are dimmed, or out.

Nikolay Kuzmin
Leiden, 10 januari 2008