



Universiteit
Leiden
The Netherlands

From NSD1 to Sotos syndrome : a genetic and functional analysis

Visser, R.

Citation

Visser, R. (2011, May 26). *From NSD1 to Sotos syndrome : a genetic and functional analysis*. Retrieved from <https://hdl.handle.net/1887/17673>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/17673>

Note: To cite this publication please use the final published version (if applicable).

Chapter 8

Genome-wide SNP array analysis in patients with features of Sotos syndrome

Remco Visser¹, Antoinet Gijsbers², Claudia Ruivenkamp², Marcel Karperien³, H. Maarten Reeser⁴, Martijn H. Breuning², Sarina G. Kant² and Jan M. Wit¹

Horm Res Paediatr 2010; 73: 265-274

1. Department of Pediatrics, Leiden University Medical Center, Leiden, The Netherlands
2. Center for Human and Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands
3. Department of Tissue Regeneration, University of Twente, Enschede, The Netherlands
4. Department of Pediatrics, Juliana Children's Hospital, The Hague, the Netherlands



Abstract

Background

Sotos syndrome is characterized by overgrowth, facial dysmorphism and learning impairment. Haploinsufficiency of *NSD1* accounts for approximately 60%-90% of the patients. Consequently, a considerable number of patients with features of Sotos syndrome remain without a molecular diagnosis. To date, target-gene approaches in these patients have not been successful.

Methods

Twenty-six Sotos syndrome-like patients were analyzed with a high-resolution whole-genome SNP array and segregation was studied in the parents.

Results

Four possible pathogenic copy-number variants including deletions of 10p12.32-p12.31, 14q13.1, Xq21.1-q21.31 and a duplication of 15q11.2-q13.1 were detected. They varied in size from 155 kb to 13.36 Mb. The 10p12.32-p12.31 deletion revealed a candidate gene (*PLXDC2*) for overgrowth. The 14q13.1 deletion affected only the *NPAS3* gene and the patient carrying this deletion displayed mental retardation as the main feature. The Xq21.1-q21.31 deletion and the 15q11.2-q13.1 duplication encompassed multiple genes of which several could be associated with phenotypic expression.

Conclusion

The high-resolution genome-wide SNP array approach resulted in a detection rate of 15% of novel abnormalities and is therefore a powerful method to attain a molecular diagnosis in Sotos syndrome-like patients. Identified candidate genes provide directions for future screening of larger patient cohorts.

Introduction

Sotos syndrome (SoS; OMIM 117550) is an autosomal dominant overgrowth disorder characterized by three cardinal features: height and/or head circumference more than +2.0 SDS, facial dysmorphism and learning disability (1). The typical craniofacial features include macrodolichocephaly, a broad forehead with a receding hairline, a prominent chin and downslanting of the palpebral fissures (2). SoS is caused by haploinsufficiency of the *NSD1* gene at 5q35.2-35.3. The detection percentage varies but overall *NSD1* abnormalities are detected in ~60% - 90% of the cases (3). Therefore, there are a considerable number of patients suspected of SoS but without a molecular explanation. The phenotypic spectrum of these patients is usually broad, varying from a classical SoS phenotype to patients exhibiting only a few SoS features. The latter group is sometimes referred to as “Sotos-like” (4,5).

In order to identify genetic alterations in SoS patients without *NSD1* abnormalities, several gene-targeted approaches have been performed (6-8). In 78 overgrowth syndrome patients in whom *NSD1* abnormalities were excluded, the *NSD* gene family members *NSD2* and *NSD3* were screened but no aberrations were detected (6). Furthermore, screening of the *NSD1* interacting protein 1 gene (*NIZP1*) did not reveal any sequence abnormalities in 97 patients referred for *NSD1* analysis [Visser et al., unpublished data] and no *NIZP1* deletions were detected in 12 typical non-*NSD1* SoS patients (7). In 38 patients with SoS features, a duplication of *IGF1R* was found in a single patient (8). Recently, *RNF135* (OMIM 611358) abnormalities were identified as the cause of a new overgrowth syndrome in combination with learning disability (9). However, analysis of *RNF135* in 160 patients referred for *NSD1* screening did not reveal any alterations (10). These results render the gene-targeted approach rather unsuccessful in detecting a molecular diagnosis in these patients. Therefore, in this study we have chosen a high-resolution genome-wide approach using a Single Nucleotide Polymorphism (SNP) array with the objective of detecting pathogenic copy-number variants (CNVs) in 26 patients with certain features of SoS.

Patients and methods

Patients

The present study includes 26 patients described previously by de Boer et al. (11) in whom *NSD1* mutations or microdeletions were excluded. From family B of the original study, only the proband (no. 41) was included. From 3 patients (no. 21, 29 and 52), DNA was no longer available. In patient 17 a duplication of *IGF1R* had been previously found and this patient was therefore excluded from the present analysis (8). According to the clinical scoring system previously used (11), this study included 3 'typical' SoS patients, 15 from the 'dubious' SoS group, and 8 from the 'atypical' SoS group. Phenotypic details and results of the clinical scoring system are shown in Table 1. Approval was obtained from the Medical Ethical Committee of the Leiden University Medical Center and consent was given by the patients and/or their parents or legal guardians. Standard deviation scores (SDS) for growth are expressed in reference to the Dutch population (12) using the Growth Analyser version 3.5 software (<http://www.growthanalyser.org/>). Weight SDS refers to the SDS for weight corrected for height. Target height was calculated with a correction for the secular trend as described previously (13). Growth parameters at birth were corrected for gestational age using Swedish references (14).

SNP arrays

Genome-wide high-density SNP array screening was performed with HumanCNV370-Duo Genotyping BeadChips (Illumina Inc., Eindhoven, The Netherlands). These chips contain ~318,000 SNP probes and ~52,000 non-polymorphic probes per chip, which results in a median spacing of 1 probe per 5 kb. Procedures were performed according to manufacturer's protocol (<http://www.illumina.com>). Fluorescence intensities were read with the BeadArray Reader (Illumina) and data files were analyzed with the Beadstudio Data Analysis Software Version 3.2.

CNV validation

Copy-number polymorphisms were excluded with the Database of Genomic Variants (DGV; <http://projects.tcag.ca/variation/>). Detected CNVs were mapped to the human genome assembly (NCBI build 36.1) using the UCSC genome browser (<http://genome.ucsc.edu/>). They were confirmed using a SNP array from a different manufacturer, i.e. the GeneChip Human Mapping 500k array Set (Affymetrix Inc., Santa Clara, CA). These chips contain

Table 1. Clinical scores for 26 patients diagnosed as having Sotos syndrome and the results of the molecular studies¹

	Patient ²	Sex	Age (years)	Fac	1	2	3	4	5	6	Gr ³	Ba ³	Hc ³	Dev ³	Sum ³	MSD1	SNP array
Group 1	6	M	4.6	5	+	+	-	+	+	+	1	2	1	1	10	-	-
	7	M	5.8	5	+	+	+	+	+	+	1	2	1	1	10	-	-
	15	M	15.3	3	-	-	+	+	-	-	2	2	1	1	9	-	-
Group 2	24	M	3.8	3	+	-	+	-	+	+	0	2	1	1	7	-	-
	25	M	4.1	3	+	-	+	-	+	+	1	2	0	1	7	-	-
	30	M	5.6	3	-	+	-	+	+	-	0	2	1	1	7	-	-
	31	F	7.2	3	-	+	+	-	+	+	0	2	1	1	7	-	-
	32	F	9.2	3	-	+	-	+	-	-	0	2	1	1	7	-	-
	33	M	20.6	3	+	+	-	-	+	+	0	2	1	1	7	-	-
	37	M	1.8	3	-	-	-	+	+	+	0	2	0	1	6	-	-
	39	F	4.8	3	-	+	-	+	+	+	1	0	1	1	6	-	del(X)(q21.1q21.31)
	40	F	6.3	0	-	-	-	-	-	-	2	2	1	1	6	-	-
	41	M	10.7	3	-	+	-	+	-	-	0	2	0	1	6	-	-
	42	F	14.0	3	-	+	+	-	+	+	0	1	1	1	6	-	-
	43	M	36.3	3	-	+	-	-	+	+	0	1	1	1	6	-	-
	45	M	6.2	3	-	+	-	-	+	+	1	0	0	1	5	-	del(10)(p12.32p12.31)
	46	F	10.2	0	-	-	-	-	+	-	1	2	1	1	5	-	-
	49	M	48.4	3	-	+	+	-	-	-	0	1	0	1	5	-	dup(15)(q11.2q13.1)
Group 3	50	M	7.6	3	-	-	+	-	+	-	0	0	1	0	4	-	-
	51	M	10.1	0	-	-	-	-	-	-	2	1	0	1	4	-	-

Table 1. (continued)

	Patient ²	Sex	Age (years)	Fac	1	2	3	4	5	6	Gr ³	Ba ³	Hc ³	Dev ³	Sum ³	NSD1	SNP array
Group 3	53	M	12.8	0	-	+	-	-	-	-	2	0	1	0	3	-	-
	54	F	5.7	0	-	+	-	-	-	-	2	0	0	0	2	-	-
	56	M	8.6	0	-	-	-	-	-	-	0	0	1	1	2	-	-
	57	M	8.8	0	-	-	-	-	-	-	0	0	1	1	2	-	-
	58	M	7.3	0	-	-	-	-	-	-	0	0	0	1	1	-	del(14)(q13.1)
	59	M	9.6	0	-	-	-	-	-	-	0	0	0	1	1	-	-

Group 1 = Typical Sotos syndrome; Group 2 = Dubious Sotos syndrome; Group 3 = Atypical Sotos syndrome;

Fac = Facial characteristics (1 = down-slant palpebral fissures, 2 = high arched palate, 3 = prominent jaw, 4 = dolichocephaly, 5 = frontal bossing, 6 = high hairline); Gr = growth; Ba = Bone age; Hc = Head circumference; Dev = Development; Mut = mutation; Del = Deletion; Dup = Duplication .

¹ Table is adapted from De Boer et al. (11) with permission from S. Karger AG, Basel.

² Numbering of patients is according the original article of De Boer et al. (11)

³ For the details about the marks used in the clinical scoring system we refer to De Boer et al. (11)

another set of SNP probes, either using the restriction enzyme *NspI* (~262,000 SNPs) or *StyI* (~238,000 SNPs), and are prepared and analyzed following different practical procedures and analysis methods. If available, parental DNA was analyzed as well using the Affymetrix platform. Practical procedures were conducted following manufacturer's guidelines (<http://www.affymetrix.com>). SNP copy number was assessed using CNAG v2 software (15).

X-inactivation study


X-inactivation was investigated at the androgen receptor locus as described previously (16).

Results

Genome-wide SNP array was performed in 26 patients with SoS features. In total we identified five deletions (patients 6, 39, 45, 54, 58) and one duplication (patient 49) which were not found in the DGV. The CNV plots are shown in Figure 1 and are summarized in Table 2. Table 3 lists the genes involved per CNV. Based on the size, the *de novo* character and/or the genes involved, the chromosomal alterations identified in patients 39, 45, 49 and 58 were categorized as possibly pathogenic.

The deletion in patient 6 was inherited from a phenotypically normal mother suggesting absence of a causal correlation between the deletion and the phenotype in the child. Furthermore, in patient 54 the father (possibly affected) and two of his siblings (normal phenotype) were carrier of the deletion detected. Therefore, the chromosomal imbalances detected in patients 6 and 54 were considered to be likely non-pathogenic and these two patients will not be described here.

In patient 39 a 13.36-Mb deletion was found encompassing Xq21.1-q21.31. The deletion involved 25 Refseq genes and occurred in the paternally derived chromosome. The X-inactivation study showed bi-allelic activity, with the paternal chromosome being more active (71% versus 29%). Although chromosome analysis at the age of three had been described as a normal 46,XX karyotype, with the current techniques the deletion was visible in the repeated karyogram (data not shown).






Figure 1. Copy-number analysis plots of the CNVs detected

The log R ratios (Y-axis) of the SNP probes for the detected CNVs and their normal flanking regions are shown. A log R ratio of > 0.3 or < -0.3 indicates a duplication or deletion respectively. Corresponding chromosomes, chromosomal bands and positions (in base pairs) are depicted below each CNV and are according to the NCBI Build 36.1 genome assembly in the UCSC genome browser.

In patient 45 a 550-kb deletion was detected at 10p12.32-p12.31 deleting the last 12 exons of a total of 14 exons, of the *Plexin Domain Containing 2* (*PLXDC2*) gene. *PLXDC2* encodes a transmembrane protein and is expressed in mouse brain, limbs, spinal cord, dorsal root ganglia, lung buds and heart (17). Unfortunately parental DNA was not available to test inheritance. In the DGV, variants were reported in four individuals, including loss of exon 1 (variant 9140), loss of exon 2 (variant 2855), loss of exon 2 and 3 (variant 9141) and gain of exon 2 (variant 2855) (18,19).

A 6.53-Mb-sized duplication was found in patient 49 encompassing 15q11.2-q13.1 and harboring 20 Refseq genes. The proximal and distal breakpoints were mapped within two known low copy repeats (LCRs) (20). Non-allelic homologous recombination between

Table 2. Identified CNVs by genome-wide SNP array screening

Patient	Chromosome	SNP start	SNP end	Start position	End position	Size	Type	Genes	Inheritance
Possible pathogenic CNVs									
39	Xq21.1-q21.31	rs5912576	rs3126874	77400259	90759007	13.36Mb	Deletion	25	<i>de novo</i>
45	10p12.32-p12.31	rs10827906	rs11012086	20262098	20812000	550kb	Deletion	1	N.D. ¹
49	15q11.2-q13.1	rs34742996	rs8040604	20315855	26849239	6.53Mb	Duplication	20	N.D. ¹
58	14q13.1	rs10872871	rs6571577	32456510	32611701	155kb	Deletion	1	mother carrier ²
Likely non-pathogenic CNVs									
6	11p14.2	rs12275693	rs1442927	26971983	27197527	225kb	Deletion	2	maternal
54	1p22.1-p21.3	rs2747038	rs1041282	94007455	94705022	698kb	Deletion	5	father, paternal uncle and aunt

¹ N.D. Not Determined because parental DNA was not available

² Only maternal DNA available

Table 3. List of Refseq genes involved in the deleted or duplicated regions

Possible pathogenic CNVs	
Patient	Refseq Genes
39	<i>CYSLTR1, ZCCHC5, LPAR4, P2RY10, GPR174, ITM2A, TBX22, FAM46D, BRWD3, NSBP1, SH3BGRL, POU3F4, CYLC1, RPS6KA6, HDX, APOOL, SATL1, ZNF711, POF1B, CHM, DACH2, KLHL4, CPXCR1, TGIF2LX, PABPC5</i>
45	<i>PLXDC2</i>
49	<i>TUBGCP5, CYFIP1, NIPA2, NIPA1, GOLGA8E, MKRN3, MAGEL2, NDN, C15orf2, SNRPN, SNURF, UBE3A, ATP10A, GABRB3, GABRA5, GABRG3, OCA2, HERC2, GOLGA8G, FLJ32679</i>
58	<i>NPAS3</i>
Likely non-pathogenic CNVs	
Patient	Refseq Genes
6	<i>FIBIN, BBOX1</i>
54	<i>DNTTIP2, GCLM, ABCA4, ARHGAP29, ABCD3</i>

N.B. Genes are ordered according to their genomic position.

LCRs at 15q11-q13 cause interstitial deletions which account for approximately 70% of the patients with Prader-Willi syndrome or Angelman syndrome (21). Reciprocal duplications of 15q11-q13 occur less frequently and breakpoints are clustered in the same LCRs as Prader-Willi and Angelman syndromes: BP1 or BP2 for proximal and BP3 for distal breakpoints (22). The same breakpoints, BP1 and BP3, were found in our patient

In patient 58 a 155-kb deletion of 14q13.1 was identified comprising the first exon of the *Neural Pas Domain Protein 3 (NPAS3)* gene, which was also detected in his mother. Two of his brothers were tested and both were confirmed to be non-carriers. To exclude compound heterozygosity, sequence analysis of *NPAS3* was performed in our patient but no pathogenic changes were detected (data not shown). *NPAS3* has two transcript isoforms (NM_022123 and NM_173159) which both include exon 1 and encode for a neuronal transcription factor (23).

Discussion

Studies examining *NSD1* abnormalities in SoS patients have refined the clinical diagnostic criteria, resulting in higher detection rates of *NSD1* abnormalities in classical SoS patients. The diagnostic challenge nowadays lies therefore in the heterogeneous group of patients with a few phenotypic features of SoS, but without *NSD1* abnormalities. With a genome-wide approach four possible pathogenic CNVs were detected in patients previously diagnosed as 'dubious' SoS (patients 39, 45, 49) and 'atypical' SoS (patient 58) (11). Phenotypic details and possible phenotypic expression of the genes involved are discussed per patient below.

Patient 39

This female patient was born as the first child of healthy, non-consanguineous white Dutch parents at 35⁺² weeks of gestation after premature rupture of membranes and gestational diabetes. At birth, weight was 3,240 grams (+1.9 SDS) and length 50 cm (+1.5 SDS). Parental heights were 192.2 cm (+1.2 SDS) and 168.3 cm (-0.4 SDS) for the father and mother, respectively, which results in a target height of 178.2 cm (+1.2 SDS). Her motor milestones were within the normal range. Audiologic examination at 2.2 years of age was normal. At the age of 2.9 years, her height was 107 cm (+3.2 SDS) and occipital frontal circumference (OFC) was 54.4 cm (+3.4 SDS). She showed dysmorphic features including dolichocephaly, frontal bossing, parietal balding, a high, narrow palate (Figure 2A-C) and there was an advanced osseous maturation of 6 months. Two IQ tests at around 3 years of age showed an IQ of 57 and 74. Concerning her family history, her father's growth pattern and head circumference were reportedly similar, although his OFC measured 60 cm (+1.4 SDS). Two younger siblings of the proband showed normal development. Our patient was last seen at the age of 9.4 years, with a height of 152.4 cm (+2.0 SDS), weight 49.5 kg (+1.2 SDS) and OFC 57.5 cm (+3.0 SDS). She receives special education. During a recent routine checkup for amblyopia, abnormalities of the retina were detected and retinitis pigmentosa was suspected.

An Xq21.1-q21.31 deletion with a size of 13.36 Mb was detected encompassing 25 genes. In the literature several genes in this region have been associated with phenotypic expression. Firstly, deletions of Xq21 in male patients were reported to cause a contiguous gene syndrome including choroideremia (*CHM* gene), mental retardation (possibly *RPS6KA6*) and X-linked deafness type 3 (*POU3F4* gene) (24). In our patient audiologic testing at the age of 2 was normal and presently she has no apparent hearing loss. However, recently

Figure 2. Facial features of the patients carrying a possible pathogenic CNV


Written permission for publication of the photographs was obtained. For patient 45 photographs were not available.

A-C: Patient 39 at the ages of 2.9, 4 and 9.4 years.

D: Patient 49 at the age of 51 years.

E: Patient 58 at the age of 15.8 years.

F: Hands and fingers of patient 58.



retinal abnormalities were found which might be the first signs of choroideremia. The variable phenotypic expression of deleted genes in our patient could be explained by the incompletely skewed X-inactivation.

Secondly, *BRWD3* is another gene possibly related to our patient's phenotype, because loss of function mutations were identified in 4 male patients with X-linked mental retardation, macrocephaly and dysmorphic features (25). Interestingly, one of them also showed tall stature ($> +1.9$ SDS) and his affected uncle was tall with a final height of $+1.3$ SDS. Thirdly, a gene which might also be associated with tall stature, is *ITM2A*. This gene encodes a transmembrane protein which has been shown to be involved in chondrogenesis and a significant association with height was recently found in a genome-wide association study (26,27).

Thus, although the implication of multiple genes complicates linking individual genes to the phenotypic features in our patient, the deleted region contains likely candidate genes explaining the retinal alterations, overgrowth and mental retardation.

Patient 45

This male patient was born as the first child of healthy, non-consanguineous white Dutch parents at a gestational age of 35⁺³ weeks with a birth weight of 3,500 grams (+2.2 SDS). The mother was hospitalized for preeclampsia, but the delivery was uneventful. Parental heights were 185.0 cm (+0.2 SDS) and 179.7 cm (+1.4 SDS) for the father and mother, which results in a target height of 193.4 cm (+1.3 SDS). Although his motor development milestones were normal, his gross and fine motor skills were assessed as delayed during childhood. He spoke his first words at the age of 30 months and talked complete sentences at the age of 60 months. For this language and speech development delay he attended special education, after which he proceeded to a mainstream elementary school. During childhood he had an increased statural growth near the +2.5 SDS curve. Bone age at the age of 2.5 conformed with calendar age. At the age of 6.2 years his height was 133.1 cm (+2.4 SDS), weight 29.9 kg (+0.6 SDS), OFC 55 cm (+1.9 SDS) and arm span 131.5 cm (+2.0 SDS). His facial features included frontal bossing, a receding hairline and a high arched palate. A simian crease was found in his right hand. His IQ was tested to be 84. Chromosome analysis showed a normal male 46,XY karyotype.

A small deletion was found, affecting the last 12 of 14 exons of *PLXDC2*. A patient (patient KK) with a 4.3-Mb interstitial deletion of 10p12.1-p12.31 was described, including amongst others *PLXDC2* in the deleted region (28). This patient also showed overgrowth with height, weight and OCF more than +1.6 SDS, mild dysmorphic features and a delay in language development (28). Patients with larger interstitial deletions or 10pter deletions have been reported but they showed postnatal growth retardation (29). In addition to the much larger size of the deletion, involvement of *PLXDC2* in these deletions could not always be confirmed (Figure 3).

Due to the missing parental DNA for inheritance studies and the detection of variants within *PLXDC2* in 4 control individuals (18,19), the contribution of haploinsufficiency of *PLXDC2* to a clinical phenotype remains equivocal. However, all the more since growth and height data of the 4 control individuals are lacking, phenotypic expression should be considered as well. The findings in our patient suggest then that *PLXDC2* is a candidate gene to be involved in growth regulation and/or intellectual development.




Figure 3. Schematic representation of reported patients with deletions of the short arm of chromosome 10

On top, the chromosomal bands of 10p are shown in scale with the physical distance in Mb according the NCBI Build 36.1 genome assembly in the UCSC genome browser (below). The deletions of patients 45, KK, GM3470 and MAR comprise *PLXDC2* (horizontal black line), while the deletions of patients MEG, TAT, KAN and ROB might include *PLXDC2*. For each patient, horizontal black bars depict non-deleted chromosomal regions. White bars represent the undetermined breakpoint regions and the blank spaces in between the deleted region. Mapping data for patient KK was obtained from Yatsenko et al. (28) and for patients GM3470, MAR, MEG, TAT, KAN and ROB from Schuffenhauer et al. (29) and references therein.

Patient 49

This male patient of Indonesian descent was first seen at the age of 45.2 years. His height was 1.78 cm (-0.8 SDS), weight 78.5 kg (+0.3 SDS) and OFC 60 cm (+1.4 SDS). The patient presented with a long face, a prominent jaw, a high arched palate and strabismus divergens (Figure 2D). He had thoracic kyphosis and scoliosis. He was institutionalized for his moderate to severe mental retardation. In his 40s he developed periods of depression followed by severe behavioural problems such as screaming, restlessness and aggression. His medical history showed hypertension and Parkinsonism but his family history was normal. Chromosome analysis revealed a normal 46,XY karyotype. The patient died at the age of 52, due to complications after surgery for a femoral fracture.

A duplication of 15q11-q13 was identified, which is a region containing imprinted and non-imprinted genes. The clinical phenotype of the 15q11-q13 duplication is variable even within families and is associated with varying levels of mental retardation and problems in motor coordination (30). On physical examination, hypotonia, decreased deep tendon reflexes and joint laxity are frequently found. Distinct dysmorphic features are not present, although an anti-mongoloid slant of the eyes and thick or pouting lips have been described (30). An association with autism spectrum disorders has been postulated, but this is not a common feature (30,31). Paternal inherited duplications have been reported to be associated with a normal or less severe phenotype (30,32), but affected patients have been reported as well (22,33,34). Considering the phenotypic features associated with 15q11-q13 duplications, it is likely that the chromosomal imbalance in our patient is the cause of his mental retardation and behavioural problems.

Patient 58

This male patient was born as the 7th child of healthy non-consanguineous white Dutch parents at 42 weeks of gestational age after an uneventful pregnancy. At birth, weight was 3550 grams (-0.5 SDS) and length 54 cm (+1.2 SDS). His target height was 186 cm (+0.3 SDS). Although he was hypotonic for which he received physiotherapy from 7 to 14 months of age, his motor development milestones were within the normal range. He followed 2 years of mainstream elementary school, but continued with special education. His IQ was tested to be 76 at the age of 7.3 years. He was last seen at the age of 15.8 with a height of 180.3 cm (+0.3 SDS), weight 84 kg (+1.9 SDS) and OFC 57.4 cm (+0.6 SDS). He did not show any major dysmorphic features, although he had short fingers and a simian crease in his left hand. His father died at the age of 40 years, due to a ruptured abdominal aneurysm. Both his father and mother attended normal elementary schools and followed lower secondary education. He had 9 siblings who were healthy, but learning problems were prevalent. There was no history of psychiatric problems.

A small deletion of the first exon of *NPAS3* was identified and his mother was confirmed to be a carrier. Previously, a mother and daughter have been described who carried a balanced reciprocal translocation t(9;14)(q34;q13) disrupting *NPAS3* on chromosome 14 without disrupting genes on chromosome 9 (35). The mother showed mild learning disability and was diagnosed with schizophrenia (36). The daughter had severe learning impairment and suffered from schizophreniform psychosis (35). Additionally, a recent study revealed 4 *NPAS3*

related haplotypes which were associated with bipolar disorder and schizophrenia (37). The association of *NPAS3* with these psychiatric disorders remains subject for discussion since our patient and mother did not show signs of psychiatric illness. Yet, there is still a possibility that our patient will develop schizophrenia since the onset of schizophrenia in males is usually in their late teens and early 20s (38). Another possibility is that due to the translocation, the disruption of *NPAS3* resulted in a dominant negative effect and therefore caused a more severely affected phenotype (36). More pronounced behavioural abnormalities were also seen in homozygous knockout mice, while only a nonsignificant trend was observed in the heterozygous mutants (39). Compound heterozygosity in our patient was excluded.

The findings in our patient support the postulated association of *NPAS3* with intellectual development, although admittedly the intelligence of the mother of our patient was borderline normal. In contrast, they are not in support of a causative relation of *NPAS3* with psychiatric disorders and more patients with *NPAS3* abnormalities are necessary to elucidate this association.

Conclusion

A genome-wide SNP array analysis detected 4 possible pathogenic CNVs in 26 (15%) patients with features of SoS. This approach is therefore a powerful method to provide a molecular explanation in patients with features of SoS but without *NSD1* alterations, although molecular validation is still required to establish a definite causal relationship. A candidate gene for overgrowth (*PLXDC2*) was identified and a deletion of *NPAS3* in 2 carriers without psychiatric illness warrants caution of linking this gene to schizophrenia. Screening of these genes in larger patient populations is necessary in order to further delineate the associated phenotype.

Acknowledgements

We are very grateful to the patients and parents for their participation in this study, as well as to the referring physicians with special thanks to dr. Veenstra-Knol, Department of Clinical

Genetics, University Medical Centre Groningen, the Netherlands. We express our gratitude to dr. R.J. Sinke, Department of Medical Genetics, University Medical Center Utrecht, the Netherlands, for performing the X-inactivation study. R.Visser was supported by grant No. 920-03-325 from The Netherlands Organisation for Health Research and Development and this research was supported by Stinafo (Dutch fund for disabled children).

References

1. Tatton-Brown K, Douglas J, Coleman K, Baujat G, Cole TR, Das S, et al. Genotype-phenotype associations in Sotos syndrome: an analysis of 266 individuals with NSD1 aberrations. *Am J Hum Genet* 2005; 77: 193-204
2. Tatton-Brown K, Rahman N. Clinical features of NSD1-positive Sotos syndrome. *Clin Dysmorphol* 2004; 13: 199-204
3. Visser R, Matsumoto N. NSD1 and Sotos Syndrome. In: Epstein CJ, Erickson RP, Wynshaw-Boris A, eds. *Inborn errors of development* (2nd edition). New York, Oxford University Press, 2008; Chapter 113: 1032-1037
4. Rio M, Clech L, Amiel J, Faivre L, Lyonnet S, Le Merrer M, et al. Spectrum of NSD1 mutations in Sotos and Weaver syndromes. *J Med Genet* 2003; 40: 436-440
5. Douglas J, Hanks S, Temple IK, Davies S, Murray A, Upadhyaya M, et al. NSD1 mutations are the major cause of Sotos syndrome and occur in some cases of Weaver syndrome but are rare in other overgrowth phenotypes. *Am J Hum Genet* 2003; 72: 132-143
6. Douglas J, Coleman K, Tatton-Brown K, Hughes HE, Temple IK, Cole TR, et al. Evaluation of NSD2 and NSD3 in overgrowth syndromes. *Eur J Hum Genet* 2005; 13: 150-153
7. Saugier-veber P, Bonnet C, Afenjar A, Drouin-Garraud V, Coubes C, Fehrenbach S, et al. Heterogeneity of NSD1 alterations in 116 patients with Sotos syndrome. *Hum Mutat* 2007; 28: 1098-1107
8. Kant SG, Kriek M, Walenkamp MJ, Hansson KB, van Rhijn A, Clayton-Smith J, et al. Tall stature and duplication of the insulin-like growth factor I receptor gene. *Eur J Med Genet* 2007; 50: 1-10
9. Douglas J, Cilliers D, Coleman K, Tatton-Brown K, Barker K, Bernhard B, et al. Mutations in RNF135, a gene within the NF1 microdeletion region, cause phenotypic abnormalities including overgrowth. *Nat Genet* 2007; 39: 963-965
10. Visser R, Koelma N, Vijfhuizen L, van der Wielen MJ, Kant SG, Breuning MH, et al. RNF135 mutations are not present in patients with Sotos syndrome-like features. *Am J Med Genet A* 2009; 149A: 806-808
11. de Boer L, Kant SG, Karperien M, van Beers L, Tjon J, Vink GR, et al. Genotype-phenotype correlation in patients suspected of having sotos syndrome. *Horm Res* 2004; 62: 197-207
12. Fredriks AM, van Buuren S, Burgmeijer RJ, Meulmeester JF, Beuker RJ, Brugman E, et al. Continuing positive secular growth change in The Netherlands 1955-1997. *Pediatr Res* 2000; 47: 316-323
13. de Boer L, Hoogerbrugge CM, van Doorn J, Buul-Offers SC, Karperien M, Wit JM. Plasma insulin-like growth factors (IGFs), IGF-Binding proteins (IGFBPs), acid-labile subunit (ALS) and IGFBP-3 proteolysis in individuals with clinical characteristics of Sotos syndrome. *J Pediatr Endocrinol Metab* 2004; 17: 615-627

14. Niklasson A, Ericson A, Fryer JG, Karlberg J, Lawrence C, Karlberg P. An update of the Swedish reference standards for weight, length and head circumference at birth for given gestational age (1977-1981). *Acta Paediatr Scand* 1991; 80: 756-762
15. Nannya Y, Sanada M, Nakazaki K, Hosoya N, Wang L, Hangaishi A, et al. A robust algorithm for copy number detection using high-density oligonucleotide single nucleotide polymorphism genotyping arrays. *Cancer Res* 2005; 65: 6071-6079
16. Allen RC, Zoghbi HY, Moseley AB, Rosenblatt HM, Belmont JW. Methylation of HpaII and HhaI sites near the polymorphic CAG repeat in the human androgen-receptor gene correlates with X chromosome inactivation. *Am J Hum Genet* 1992; 51: 1229-1239
17. Miller SF, Summerhurst K, Runker AE, Kerjan G, Friedel RH, Chedotal A, et al. Expression of Plxdc2/TEM7R in the developing nervous system of the mouse. *Gene Expr Patterns* 2007; 7: 635-644
18. Pinto D, Marshall C, Feuk L, Scherer SW. Copy-number variation in control population cohorts. *Hum Mol Genet* 2007; 16 Spec No. 2: R168-R173
19. Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, Andrews TD, et al. Global variation in copy number in the human genome. *Nature* 2006; 444: 444-454
20. Christian SL, Fantes JA, Mewborn SK, Huang B, Ledbetter DH. Large genomic duplicons map to sites of instability in the Prader-Willi/Angelman syndrome chromosome region (15q11-q13). *Hum Mol Genet* 1999; 8: 1025-1037
21. Nicholls RD, Knepper JL. Genome organization, function, and imprinting in Prader-Willi and Angelman syndromes. *Annu Rev Genomics Hum Genet* 2001; 2: 153-175
22. Roberts SE, Dennis NR, Browne CE, Willatt L, Woods G, Cross I, et al. Characterisation of interstitial duplications and triplications of chromosome 15q11-q13. *Hum Genet* 2002; 110: 227-234
23. Brunskill EW, Witte DP, Shreiner AB, Potter SS. Characterization of npas3, a novel basic helix-loop-helix PAS gene expressed in the developing mouse nervous system. *Mech Dev* 1999; 88: 237-241
24. Yntema HG, van den HB, Kissing J, van Duijnhoven G, Poppelaars F, Chelly J, et al. A novel ribosomal S6-kinase (RSK4; RPS6KA6) is commonly deleted in patients with complex X-linked mental retardation. *Genomics* 1999; 62: 332-343
25. Field M, Tarpey PS, Smith R, Edkins S, O'Meara S, Stevens C, et al. Mutations in the BRWD3 gene cause X-linked mental retardation associated with macrocephaly. *Am J Hum Genet* 2007; 81: 367-374
26. Van den Plas D, Merregaert J. In vitro studies on Itm2a reveal its involvement in early stages of the chondrogenic differentiation pathway. *Biol Cell* 2004; 96: 463-470
27. Gudbjartsson DF, Walters GB, Thorleifsson G, Stefansson H, Halldorsson BV, Zusmanovich P, et al. Many sequence variants affecting diversity of adult human height. *Nat Genet* 2008; 40: 609-615
28. Yatsenko SA, Yatsenko AN, Szigeti K, Craigen WJ, Stankiewicz P, Cheung SW, et al. Interstitial deletion of 10p and atrial septal defect in DiGeorge 2 syndrome. *Clin Genet* 2004; 66: 128-136

29. Schuffenhauer S, Lichtner P, Peykar-Derakhshandeh P, Murken J, Haas OA, Back E, et al. Deletion mapping on chromosome 10p and definition of a critical region for the second DiGeorge syndrome locus (DGS2). *Eur J Hum Genet* 1998; 6: 213-225
30. Bolton PF, Dennis NR, Browne CE, Thomas NS, Veltman MW, Thompson RJ, et al. The phenotypic manifestations of interstitial duplications of proximal 15q with special reference to the autistic spectrum disorders. *Am J Med Genet* 2001; 105: 675-685
31. Bolton PF, Veltman MW, Weisblatt E, Holmes JR, Thomas NS, Youings SA, et al. Chromosome 15q11-13 abnormalities and other medical conditions in individuals with autism spectrum disorders. *Psychiatr Genet* 2004; 14: 131-137
32. Cook EH, Jr., Lindgren V, Leventhal BL, Courchesne R, Lincoln A, Shulman C, et al. Autism or atypical autism in maternally but not paternally derived proximal 15q duplication. *Am J Hum Genet* 1997; 60: 928-934
33. Mohandas TK, Park JP, Spellman RA, Filiano JJ, Mamourian AC, Hawk AB, et al. Paternally derived de novo interstitial duplication of proximal 15q in a patient with developmental delay. *Am J Med Genet* 1999; 82: 294-300
34. Veltman MW, Thompson RJ, Craig EE, Dennis NR, Roberts SE, Moore V, et al. A paternally inherited duplication in the Prader-Willi/Angelman syndrome critical region: a case and family study. *J Autism Dev Disord* 2005; 35: 117-127
35. Kamnasaran D, Muir WJ, Ferguson-Smith MA, Cox DW. Disruption of the neuronal PAS3 gene in a family affected with schizophrenia. *J Med Genet* 2003; 40: 325-332
36. Pickard BS, Malloy MP, Porteous DJ, Blackwood DH, Muir WJ. Disruption of a brain transcription factor, NPAS3, is associated with schizophrenia and learning disability. *Am J Med Genet B Neuropsychiatr Genet* 2005; 136: 26-32
37. Pickard BS, Christoforou A, Thomson PA, Fawkes A, Evans KL, Morris SW, et al. Interacting haplotypes at the NPAS3 locus alter risk of schizophrenia and bipolar disorder. *Mol Psychiatry* 2009; 14: 874-884
38. Leung A, Chue P. Sex differences in schizophrenia, a review of the literature. *Acta Psychiatr Scand Suppl* 2000; 401: 3-38
39. Erbel-Sieler C, Dudley C, Zhou Y, Wu X, Estill SJ, Han T, et al. Behavioral and regulatory abnormalities in mice deficient in the NPAS1 and NPAS3 transcription factors. *Proc Natl Acad Sci USA* 2004; 101: 13648-13653

