Cover Page



Universiteit Leiden



The handle <u>http://hdl.handle.net/1887/35931</u> holds various files of this Leiden University dissertation

Author: Deventer, Sjoerd van Title: Tracking the big ones : novel dynamics of organelles and macromolecular complexes during cell division and aging Issue Date: 2015-10-21

Chapter 4:

N-terminal acetylation and replicative age affect proteasome localization and cell fitness during aging

Sjoerd J. van Deventer¹, Victoria Menendez-Benito¹, Fred van Leeuwen², Jacques Neefjes¹

Journal of Cell Science, Jan 2015, 128 (1): 109-117

¹ Division of Cell Biology 2 and ² Division of Gene Regulation, Netherlands Cancer Institute, Amsterdam, The Netherlands

Abstract

Specific degradation of proteins is essential for virtually all cellular processes and is carried out predominantly by the proteasome. The proteasome is important for clearance of damaged cellular proteins. Damaged proteins accumulate over time and excess damaged proteins can aggregate and induce the death of old cells. In yeast, the localization of the proteasome changes dramatically during aging, possibly in response to altered proteasome activity requirements. We followed two key parameters of this process: the distribution of proteasomes in nuclear and cytosolic compartments, and the formation of cytoplasmic aggregate-like structures called proteasome storage granules (PSGs). Whereas replicative young cells efficiently relocalized proteasomes from the nucleus to the cytoplasm and formed PSGs, replicative old cells were less efficient in relocalizing the proteasome and had less PSGs. By using a microscopy-based genome-wide screen, we identified genetic factors involved in these processes. Both relocalization of the proteasome and PSG formation were affected by two of the three N-acetylation complexes. These N-acetylation complexes also had different effects on the longevity of cells, indicating that each N-acetylation complex has different roles in proteasome location and aging.

Introduction

The proteasome is a major intracellular protease and controls many processes, including protein quality control. Protein quality control is required to prevent accumulation of damaged proteins during the lifespan of a cell (Amm *et al.*, 2014; Koga *et al.*, 2011). Insufficient recognition and clearance of damaged proteins can yield harmful protein aggregates (Powers *et al.*, 2009; Schmidt and Finley, 2014). A proper functioning ubiquitin-proteasome system (UPS) might prevent protein aggregation and counteract cellular aging.

Several studies report an age-dependent decline in UPS activity in various model systems (Carrard *et al.*, 2002; Dasuri *et al.*, 2009; Lee *et al.*, 1999; Vernace *et al.*, 2007a; Vernace *et al.*, 2007b). Other studies suggest a causative relation between UPS activity and aging. Enhancing proteasome activity by overexpression of the proteasome assembly chaperone Ump1 improves budding yeast longevity under starvation conditions (Chen *et al.*, 2006). Increasing proteasome levels by overexpressing Rpn4, a protein which drives the transcription of the proteasome subunits, also increases the replicative lifespan in *S. cerevisiae* (Kruegel *et al.*, 2011). These studies suggest that the UPS system decays with age and limits the lifespan of cells and organisms. Manipulating UPS therefore might have dramatic effects on the aging process.

For several reasons, *S. cerevisiae* is an important model organism to elucidate the molecular basis of processes related to aging. First, cell division is asymmetrical with a distinguishable mother and daughter cell. This allows tracking of a single cell over time, even during division. Second, the number of cell divisions can be quantified by counting the bud scars left on the mother cell after budding of a new generation. The asymmetrical cell division defines two forms of aging; chronological aging and replicative aging (Kaeberlein, 2010; Jazwinski *et al.*, 1989). Chronological aging is defined as the time between the budding from the mother, the birth, until the daughter cell dies. This aging is usually addressed on a

population level by measuring the viability of a liquid culture upon starvation (Kaeberlein, 2010). Replicative aging is aging as a result of cell division and defined by the number of daughter cells produced by an individual mother cell. Replicative aging in yeast is used to model aging of mitotically active mammalian cells (Kaeberlein, 2010; Mortimer and Johnston, 1959). Chronological and replicative aging are overlapping processes (Delaney *et al.*, 2013; Kennedy *et al.*, 1994; Murakami *et al.*, 2012), exemplified by the observation that, during starvation of a liquid yeast culture, the replicative age of a cell at the start of starvation highly affects the chronological age that will be reached (Allen *et al.*, 2006; Aragon *et al.*, 2008). The studies in yeast have revealed many insights into the various molecular processes such as neurodegenerative disorders (Clay and Barral, 2013; Tenreiro and Outeiro, 2010).

Here, we followed two proteasome-related processes that occur during chronological aging in yeast: nuclear-cytoplasmic relocalization of proteasomes, and the formation of cytoplasmic proteasome storage granules (PSGs). PSGs are aggregate-like structures that contain the proteasome and form early during yeast starvation (Laporte *et al.*, 2008). The replicative age of cells had a major effect on these processes. Replicative young cells efficiently relocalized the proteasome from the nucleus and formed PSGs, unlike replicative old cells. A genome-wide knockout screen revealed that proteasome relocalization and PSG formation involves two of the three N-acetylation complexes, each having a particular effect on proteasome localization. The N-acetylation complexes were found to affect cell fitness in different ways. One N-acetylation complex, NatC, both affected proteasome location and fitness of old cells.

Results

Proteasome localization during starvation correlates with replicative age

Proteasomes equally distribute over the nucleus and cytoplasm in mammalian cells (Reits *et al.*, 1997). In the budding yeast *Saccharomyces cerevisiae*, proteasomes accumulate in the nucleus when cells have sufficient nutrients (Russell *et al.*, 1999). This changes when yeast cells exhaust glucose in the growth medium, a process that leads to starvation. During starvation, cells relocalize their proteasome from the nucleus into the cytosol and form cytoplasmic PSGs (Laporte *et al.*, 2008). Starvation is apparently a factor controlling the intracellular distribution of proteasomes.

We visualized proteasomes in live yeast cells by tagging the catalytically active β 1 subunit (Pre3) of the proteasome with GFP. Efficient and quantitative introduction of the β 1–GFP in 20S proteasomes was confirmed by native gel electrophoresis (supplementary material Fig. S1). The GFP-labeled proteasomes had a similar distribution during starvation as reported for non-modified proteasomes previously (Laporte *et al.*, 2008). We observed that cells in starvation show a wide heterogeneity in proteasome localization (Fig. 1A). Based on proteasome localization, we defined four localization phenotypes: (1) cells with proteasome accumulation in the nucleus (Nuclear); (2) cells displaying dots of cytoplasmic proteasomes (Nuclear + PSG); and (4) cells without any of these phenotypes, where proteasomes are approximately equally distributed between the cytoplasmic and nuclear compartments (Equal) (Fig. 1B). In a typical 5-day starvation experiment the majority of the cells are either PSG or Equal, whereas a small portion of the cells is Nuclear or Nuclear

+ PSG.

Given that these cells are genetically identical and grow under identical conditions, it is expected that other factors should be responsible for this heterogeneity. These could include replicative aging. Replicative aging results from asymmetrical cell division of budding yeasts in which damaged cell components are typically retained in the mother cell (Kaeberlein, 2010). After each cell division, chitinous scar tissue is left on the cell wall of the mother, which is called a bud scar. The number of bud scars can be visualized with Calcofluor White (CFW), which marks the replicative age (Pringle, 1991). CFW staining distinguishes three age groups: (1) virgin cells without bud scars; (2) young mother cells with one or two bud scars, and (3) old mothers with more than two bud scars (Fig. 1C). We quantified the respective proteasome localization phenotypes per age group. PSG formation inversely related to age as ~30% of the old mothers, ~60% of the young mothers



Fig. 1: Proteasome localization in nutrient-starved cells correlates with replicative age

(A) Live cell microscopy of yeast cells in starvation shows various 20S proteasome localizations as is visualized by endogenous expression of β 1(Pre3)-GFP. Cells were stained with CalcoFluor White to assess the replicative age of individual cells. (B) β 1-GFP localization and Hoechst staining define four different phenotypes: cells with cytosolic Proteasome Storage Granules (PSG), cells with nuclear enrichment of proteasomes (Nuclear), cells that display both (Nuclear + PSG) and cells without a clear enrichment of proteasomes in PSGs or nuclei (Equal). (C) Based on Calcofluor White staining of bud scars, three different replicative age groups were defined; Virgin daughter cells without bud scars (V), Young Mothers with 1-2 bud scars (YM) and Old Mothers with more than two bud scars (OM). (D) The prevalence of the different proteasome phenotypes in living cells from each age group was calculated by dividing the number of cells with a certain phenotype in a particular age group over the total number of cells in this age group. Values and standard deviations are based on three independent experiments and significance was calculated with a paired, two-tailed t-test (* = p<0.05, ** = p<0.01). (Scale bars, 5 µm)

and ~65% of the virgin cells displayed this phenotype (Fig. 1D). Nuclear accumulation of proteasomes correlated with replicative age in the small population of yeast where this phenotype was observed. This suggests that impairment of proteasome relocalization and/or PSG formation can be associated with replicative age. The prevalence of the other two phenotypes did not differ significantly between the different age groups. Similar results were found in yeasts expressing proteasomes labeled through another 20S subunit [α 2–GFP (Pre8)] or a 19S subunit (Rpn1–GFP) (supplementary material Fig. S2).

Identification of genes affecting proteasome localization during starvation

To identify genetic factors controlling proteasome localization during starvation, we performed a microscopy-based yeast knockout screen. We considered two explanations for the maintenance of nuclear enrichment of the proteasome during starvation: altered proteasome biogenesis in the nucleus or altered nuclear retention of the proteasome. Therefore, we tagged the β 1 subunit (Pre3) of the proteasome with a fluorescent Recombination-Induced Tag Exchange (RITE) cassette (Verzijlbergen *et al.*, 2010), to differentially label new and old proteasomes. Integration of the RITE cassette behind the β 1 gene results in a GFP-tagged proteasome produced before tag exchange, whereas new proteasomes (produced after tag exchange due to translocation of an estrogen receptor (ER)-coupled Cre-recombinase to the nucleus after addition of β -estradiol (Verzijlbergen *et al.*, 2010)) will be labeled with mRFP. The genetic GFP-for-mRFP swapping is permanent and induced after two days of starvation. When recombination was induced at this time point little or no synthesis of proteasomes was detected in wild-type (WT) cells. Similar results have been obtained by Menendez-Benito *et al.*, 2013).

To obtain a screening library, the β 1–RITE strain was crossed with the MATa haploid knockout (KO) collection (Thermo Scientific) using SGA technology (Tong et al., 2001) (Fig. 2A1). This high-throughput crossing yielded 4263 knockout strains containing a RITEtagged proteasome. These strains were subjected to a 5-day starvation protocol, including the induction of tag exchange (switch) at day 2 (Fig. 2A2). To efficiently analyze thousands of samples by microscopy, cells were fixed, stained with Hoechst 33342 and spotted on an object glass using a DNA microarray printer (Narayanaswamy et al., 2006) (Fig. 2A3). Each spot, typically consisting of ~2000 cells, was imaged by confocal microscopy (Fig. 2A4). A CellProfiler image analysis pipeline was designed for guantification of the proteasome phenotypes of interest (Carpenter et al., 2006) (Fig. 2A5). This pipeline assessed the nuclear cytosolic distribution of the proteasome by dividing the mean GFP fluorescence in the nucleus over the mean GFP fluorescence in the cytoplasm. Three successive rounds of screening identified three hits with a nuclear retention phenotype of the proteasome: $hul5\Delta$, $uba3\Delta$ and $mak10\Delta$ (Fig. 2A6; Fig. 2B). These results were verified by repeating the experiment with independently made knockout strains. Loss of HUL5, UBA3 or MAK10 increased the population of cells with nuclear accumulation of proteasomes (Fig. 2C). Little or no synthesis of new (mRFP tagged) β 1 was detected in either WT or KO cells, thus implying that the nuclear enrichment is not due to de novo synthesis. A plating assay before and after recombination confirmed the successful genetic recombination (GFP to mRFP) in these cells (supplementary material Fig. S3A). When recombination is induced at an earlier time point in starvation (after 1 day), synthesis of new (mRFP tagged) proteasomes could be observed in both WT cells and the three screen hits (supplementary material Fig. S3B). The RITE technology was only used for identifying the hits.

Fig. 2



Fig. 2: Genome-wide screen identifies genes affecting nuclear proteasome localization during starvation

(A) Schematic overview of the screening. (1) A yeast knock-out library was crossed with a β 1-GFP \rightarrow mRFP RITE strain. (2) Tag recombination (switch) was performed after two days during a five day starvation experiment. (3) Samples were fixed, stained with Hoechst and printed on yeast arrays. (4) Microscopic imaging of GFP (old proteasomes), RFP (new proteasomes) and Hoechst (nuclei) was performed. (5) Images were analyzed by CellProfiler. (6) *mak10*Δ was one of the hits for a nuclear proteasome enrichment. (B) Confocal microscopy images of three hits showing nuclear enrichment of GFP labeled proteasome in the nucleus; *hul5*Δ, *mak10*Δ and *uba3*Δ. Only background signal is observed for the mRFP proteasome. (C) Quantification of nuclear/cytosolic ratios of GFP in WT and nuclear retention hits. Values and standard deviations are based on 5 independent experiments and significance was calculated with a paired, two tailed t-test (* = p<0.05, ** = p<0.01). (Scale bar, 5 µm)

Loss of N-acetylation by NatC causes nuclear enrichment of the proteasome without affecting PSG formation, and both phenotypes are affected by replicative age

The Mak10 protein is a subunit of the N-acetyltransferase C (NatC) complex. NatC associates to the ribosome for co-translational N-terminal acetylation of a subset of proteins (Starheim et al., 2012). The NatC complex further consists of Mak31 and the catalytic subunit Mak3 (Polevoda and Sherman, 2000). In our screening, Mak3 and Mak31 were just below the threshold, but independently generated knockouts of all three individual NatC subunits showed increased nuclear retention of proteasomes during starvation, whereas the number of cells displaying PSGs was not significantly altered (Fig. 3A,C; supplementary material Fig. S4A,B), indicating a specific role for the NatC complex in the nuclear enrichment of the proteasome. The presence of cells with both nuclear enrichment of proteasomes and cytoplasmic PSGs means that PSG formation has been uncoupled from the nuclear-to-cytosolic relocalization of proteasomes (Fig. 3A,B). A catalytically inactive NatC mutant (Mak3 N123A and Y130A) (Polevoda and Sherman, 2000) showed the same phenotype (Fig. 3A,B). NatC activity is apparently involved in the nuclear-to-cytosolic relocalization of proteasomes under starvation conditions. CFW staining was used to assess a potential correlation of proteasome localization with replicative age. Nuclear enrichment of proteasomes correlated with replicative age, whereas PSG formation correlated negatively with replicative age (Fig. 3C,D). Cells displaying both nuclear enrichment and PSG formation also showed a weak correlation, but 'Equal' cells did not. Similar results with respect to the correlation between replicative age and proteasome localization and the effect of NatC deficiency were found for $\alpha 2$ -GFP- (Pre8) and Rpn1–GFP-expressing cells (supplementary material Fig. S2). These results



Fig. 3: Loss of N-acetylation by NatC causes nuclear retention of the proteasome without affecting PSG formation; both phenotypes are affected by replicative age

(A) Fixed cell microscopy of Hoechst stained mak3A cells or cells expressing a catalytically inactive Mak3 (MAK3-CD) shows an increased population of cells displaying nuclear retention of proteasomes after a five day starvation period. (B) The prevalence of the different phenotypes in the total population was scored in three independent experiments. (C) Live cell microscopy of mak3∆ cells stained with Calcofluor White after a five day starvation period. (D) The prevalence of each proteasome phenotype in living cells was scored in three different age groups. Quantifications are based on three independent experiments and significance was calculated with paired two-tailed t-tests (* = p<0.05, ** = p<0.01). (Scale bars, 5 µm)

are similar to those observed for WT cells (Fig. 1D), indicating that NatC does not affect aging-related relocalization of proteasomes.

Nuclear-to-cytosolic relocalization of the proteasome during starvation requires NatB and NatC, PSG formation requires only NatB

The main N-terminal acetyltransferases in yeast are NatA, NatB and NatC, contributing to respectively ~50%, ~20% and ~20% of the N-terminal acetylome. Each of these complexes recognizes specific substrates depending on their N-terminal sequences (Starheim *et al.*, 2012). To test whether nuclear enrichment of proteasomes only depends on NatC-mediated N-acetylation, the subunits of the NatA (Ard1, Nat1 and Nat5) and the NatB (Nat3 and Mdm20) complex were knocked out. Deficiency of NatA subunits did not alter proteasome distribution following starvation (Fig. 4A,B;supplementary material Fig. S4C,D). However, deletion of the various NatB subunits increased the population with nuclear proteasomes and reduced the cells displaying PSGs (Fig. 4A,B; supplementary material Fig. S4E,F). Although both NatB and NatC knockouts induced nuclear proteasome



Fig. 4: Nuclear-cytosolic relocalization of the proteasome during starvation requires N-acetylation by NatB and NatC

(A) Fixed cell microscopy of starved $ard1\Delta$ (NatA deficient), $nat3\Delta$ (NatB deficient) and $mak3\Delta$ (NatC deficient) cells. Nuclei were visualized with a Hoechst staining. (B) Prevalence of the different proteasome localization phenotypes was scored in the total population. Values and standard deviations are based on a biological triplicate. (C) Live cell imaging of starved $ard1\Delta$ and $nat3\Delta$ cells. Cells were stained with CalcoFluor White to assess their replicative age. (D) Prevalence of the different proteasome phenotypes in the three age groups in living cells was quantified in three independent experiments and significance was calculated with a paired two-tailed T-test. (* = p<0.05, ** = p<0.01) (Scale bars, 5 µm)

enrichment, they had different effects on the cytoplasmic proteasome pool. NatC inactivation still allowed PSG formation, whereas NatB inactivation prevented formation of PSGs (Fig. 4A,B; supplementary material Fig. S4C–F). This suggests a specific role of the NatB complex in PSG formation. Cells with both PSGs and nuclear retention were hardly detected among NatB-knockout cells (Fig. 4B). As for WT and NatC-deficient cells, nuclear enrichment induced by NatB inactivation correlated with replicative age, whereas the prevalence of 'Equal' cells decreased with replicative age. (Fig. 4C,D; compare to Fig. 3D). Surprisingly, NatA-deficient cells did show a WT-like prevalence of the different proteasome phenotypes in the total population, but there was no correlation of these phenotypes with replicative age.

These results would suggest that the mechanism underlying the different proteasome localizations in cells involves selective N-acetylation and can be (at least partially) uncoupled from aging effects that also require N-acetylation. NatB and NatC, unlike NatA, are involved in the effects on proteasome distribution. Their combined inactivation might further accelerate these effects, and a NatB+NatC double knockout strain was made. This double knockout (unlike the single knockouts) had severe growth defects, preventing a fair comparison with the single knockout strains. Given that the effects on nuclear enrichment of the proteasome were specific to NatB and NatC, N-acetylation of one or more NatB and NatC substrates must be involved in nuclear-to-cytosolic proteasome distribution. Based on the N-terminal sequence requirements of each Nat complex a list of potential substrates were defined in the yeast proteome (Arnesen et al., 2009; Polevoda and Sherman, 2003). The role of N-acetylation of selected candidate substrates was tested by making an N-terminal MX- to MP- (X2P) mutation, resulting in an N-terminus that cannot be N-acetylated (Polevoda and Sherman, 2003). Preventing N-acetylation of α5 (Pup2), a6 (Pre5), Rpn9, Fub1, Avo2, Hul5 or Nup100 failed to phenotypically mimic cells lacking NatB or NatC (supplementary material Fig. S4G). Whether NatB and NatC act on proteasome distribution by modifying a single target or many, is as yet unclear.

NatA and NatB control general cell fitness during starvation and NatC fitness of old cells only

As proteasome composition and activity was found to influence longevity in starvation, we wondered whether proteasome localization would also correlate with cell fitness. Cellular fitness in starvation can be determined by assaying the ability of cells to restart their cell cycle when nutrients are added. This is determined by plating equal numbers of cells and quantifying the number of colony-forming units (CFUs). When grown in the presence of sufficient nutrients, the reproductive capacity of NatA- and NatC-deficient cells is similar to WT cells, whereas NatB-deficient cells show a lower reproductive capacity (Fig. 5A) (Polevoda *et al.*, 1999). After a 5-day starvation period, both NatA- and NatB-deficient cells showed lower CFUs than wild-type cells, whereas the reproductive capacity in NatC-deficient cells seemed to be unaffected (Fig. 5B). Proteasome localization in NatA-deficient cells was similar to WT, unlike that in NatB-deficient cells. Given that NatA and NatB deficiency both decrease the number of CFUs measured, proteasome localization cannot be directly related to reproductive capacity in starvation.

Given that NatC-deficient cells show a strong correlation of proteasome localization with replicative age, we wondered whether NatC also affected reproductive capacity in an age-dependent manner. We determined the fitness of old versus young cells in a starved population of the various mutant yeast strains by staining the yeast cells with CFW and

then separating young and old yeast cells by FACS sorting (Pringle, 1991). Microscopy on the sorted populations verified separation of virgin and old mother cells (Fig. 5C). Equal numbers of cells from the different populations were subsequently plated and the number of CFUs determined. About 75% of the young WT cells and ~30% of the old WT cells were able to form colonies upon plating (Fig. 5D). This correlation between replicative age and cell fitness is similar to results previously reported by Allen *et al.* (Allen *et al.*, 2006). The low number of CFUs measured for old as well as young NatA- and NatB-deficient cells was expected, based on their general effects on cell fitness (Fig. 5B) and is in agreement



Fig. 5: In starvation, loss of NatA and NatB has a general effect on reproductive capacity, whereas loss of NatC specifically affects old cells

The reproductive capacity of the different strains in log phase (A) and starvation (B) was assessed by a plating assay. Loss of NatA and NatB compromised reproductive capacity in starved cells, whereas loss of NatC did not. Representative plating assay from three independent experiments. (C) After Calcofluor White (CFW) staining of a starved culture, the cells with the lowest (negative) and highest (positive) CFW signal were sorted to obtain populations of virgin cells and old mothers, respectively. The CFW image is the maximum projection of a 5 μ m Z-stack, the DIC image is a single scan in the middle of the Z-stacks. (D) CFUs were counted for old and young cells. Loss of NatC affects the reproductive capacity of old cells, while loss of NatA and NatB reduces the reproductive capacity of both young and old cells. Values and standard deviations are based on three independent experiments. Significance was calculated with a paired two-tailed T-test. (* = p<0.05, ** = p<0.01)

with findings for NatA by Aragon *et al.* (Aragon *et al.*, 2008). Surprisingly, WT and NatC-knockout young cells were equally fit, whereas the fitness of the old NatC-knockout cells was reduced to only 30% of old WT cells. NatC deficiency not only affected localization of proteasomes in replicative old cells, but it also had a selective effect on the fitness of old mothers.

Discussion

The proteasome is located in both the cytosol and nucleosol. Its subunits are made in the cytosol where proteasomes are assembled in precursor complexes that can be imported into the nucleus for full formation of the complex (Lehmann *et al.*, 2002). Nuclear import and export of mature proteasomes is a very slow process in mammalian cells and the details of this process are poorly understood (Reits *et al.*, 1997). Given that the nuclear

envelope disintegrates during mitosis in mammalian cells, the boundary between the two pools of proteasomes is lost and nuclear proteasomes mix with cytosolic proteasomes. Nuclear proteasomes could thus have a cytosolic origin and vice versa. This should be different in budding yeast, where the nuclear envelope is maintained during cell division and proteasomes tend to accumulate in the nucleus. However, nuclear-to-cytosolic relocalization of proteasomes is observed upon glucose exhaustion, which is followed by rapid nuclear import of mature proteasomes upon re-addition of nutrients. It would therefore be expected that proteasome distribution is under some kind of control (Laporte *et al.*, 2008).

We report that nuclear-to-cytosolic relocalization of the proteasome upon starvation correlates with the replicative age of yeast cells. Young cells are better capable of relocalizing the proteasome from the nucleus and of forming cytosolic PSGs, whereas old mother cells usually fail to do so. This correlation might be due to an age-dependent defect in the proteasome relocalization machinery, but could also be due to asymmetrical division of damaged cell components upon cell division. In general, damaged protein accumulates in mother cells leaving daughter cells with a fresh set of proteins (Henderson and Gottschling, 2008; Kaeberlein, 2010; Nyström and Liu, 2014). Perhaps a low amount of damaged proteins allows young cells to store their proteasomes in PSGs, whereas old cells might need to maintain a larger pool of active proteasomes to handle accumulated protein damage at the cost of PSGs. A nuclear enrichment of the proteasome might help replicative old cells to handle protein stress in the nucleus (Gardner *et al.*, 2005) but it might also contribute to protein quality control in the cytosol (Heck *et al.*, 2010; Prasad *et al.*, 2010).

The reported association between the capacity of the ubiquitin-proteasome system and viability during aging (Carrard et al., 2002; Kruegel et al., 2011; Tonoki et al., 2009; Vernace et al., 2007b) suggests that the localization of the proteasome also affects viability. This can only be tested in a system where proteasome localization can be manipulated during the aging process. We performed a microscopy-based screen to identify proteins that control proteasome localization in yeast cells. We identified two of the three complexes involved in N-terminal acetylation of proteins as controlling nuclear-to-cytosolic relocalization of the proteasome. About half of the yeast proteome can be N-acetylated by these complexes (Starheim et al., 2012). N-acetylation can both stabilize and de-stabilize proteins as well as affect the intracellular localization and activity of proteins (Arnesen, 2011; Hwang et al., 2010; Scott et al., 2011). Given that the acetylation complexes NatB and NatC but not NatA control the nuclear-to-cytosolic relocalization of proteasomes, a selective set of substrates rather than the general N-acetylation process seems to be responsible. The different phenotypes for the NatB and NatC knockouts allowed testing of the link between nuclear-to-cytosolic relocalization of proteasomes and the formation of PSGs. Although NatC-deficient old mother cells accumulated significantly more nuclear proteasomes, the prevalence of cytosolic PSGs was unaffected. In addition, a population of cells with both nuclear retention and PSG formation was observed. Both observations suggest that nuclear accumulation of proteasomes in older mothers does not necessarily prevent formation of cytosolic PSGs. This suggests that the mere enrichment of proteasomes in the cytosolic compartment is not a prerequisite for PSG formation.

As proteasome localization during aging can be manipulated by inactivation of either NatB or NatC, the effect on cell fitness during aging can be determined. Although NatA and NatB deficiency strongly affected cell fitness in all age groups tested, NatC deficiency

selectively affected fitness of old mothers. As for the localization of the proteasome, the different Nat complexes affected cell fitness in a Nat-complex-specific manner. Whether this is the result of modification of part of the proteome or of one defined substrate, is at present unclear. We excluded some proteins that were potentially involved, like Hul5 (supplementary material Fig. S4G), another hit in our screening, as single candidates but that does not exclude other proteins.

Acetylation of lysine side chains of various proteins has been connected to the aging process. This is exemplified by the yeast deacetylase Sir2. Sir2 reduces lifespan upon deletion and prolongs it upon overexpression (Wierman and Smith, 2014). Homologs of Sir2 in several other organisms, including the SIRT proteins in mammals, have been linked to aging and age-related diseases (Donmez and Guarente, 2010). Furthermore, calorific restriction can increase age and has been associated with altered acetylation status of many mitochondrial proteins (Hebert *et al.*, 2013), which has been extensively studied in neurodegeneration diseases (Guedes-Dias and Oliveira, 2013). These examples indicate a role of acetyl modifications of lysine side chains in processes associated with aging. Here, we have shown that N-terminal acetylation, a different, stable and often co-translational modification with the same chemical group, is associated with proteasome distribution and fitness during aging. The mechanisms controlling proteasome localization and fitness at old age involve specific N-acetylation complexes and result in a further expansion of the role of the small acetyl modification.

Materials and Methods

Yeast strains and plasmids

With the exception of the strains used for screening, all strains were derived from NKI4103 (Verzijlbergen *et al.*, 2010). Gene knockouts were made by PCR-mediated gene disruption based on pRS plasmids (Baker Brachmann *et al.*, 1998). N-terminal mutations were made by PCR amplification of the pYM-N10 and pYM-N11 plasmids using the S1 and S4 primers extended with 40 bp of sequence homologous to the endogenous sequence (Janke *et al.*, 2004). The N-terminal mutation was introduced in the 40 bp endogenous sequence. The two mutations for the catalytically inactive N123A-Y130A-Mak3 were generated by Delitto Perfetto technology (Storici and Resnick, 2006). All strains are described in supplementary material Table S1.

Growth Conditions

Yeast cells were grown in liquid YEPD cultures of 5 ml at 30°C. To prevent recombination of the RITE cassette, cells were grown in presence of Hygromycin (200 μ g/ml, Invitrogen). Liquid cultures were starved by inoculating 5 ml of YEPD with 0.5 ml of an overnight culture followed by a 5-day starvation period.

Library construction

NKI4103 (Verzijlbergen *et al.*, 2010) was crossed with the MATa haploid knockout collection (ThermoScientific) by Synthetic Genetic Array analysis (Tong and Boone, 2006) using a RoToR HAD (Singer Instruments) with the following modifications. After mating, diploids were selected and kept on Hygromycin, G418 and CloNat triple selection on rich medium for 1 night. After 2 weeks on sporulation medium, MATa haploid clones containing both a gene knockout and the RITE tagging system were selected. The first

two rounds of selection generated haploid MATa cells (YC-His+Can+SAEC) (van Leeuwen and Gottschling, 2002). The next two selection rounds selected the knockout and the RITE system (YC-His+Can+SAEC+MSG+ Hygromycin, G418 and CloNat).

Microscopy and image analysis

Fixed microscopy samples were prepared by fixing ~10⁸ cells in 4% formaldehyde and mounted in Vectashield (Vector Laboratories) on ConA-coated coverslips. Live-cell samples were prepared by resuspending ~10⁸ cells in 100 µl 40°C 1% UltraPure[™] LMP Agarose (Invitrogen) in PBS which is squeezed between a cover glass and an object glass. Imaging was performed at room temperature within 1 hour after mounting. Hoechst 33342 (Invitrogen, 1 µg/ml) or Calcofluor White (CFW, Sigma-Aldrich, 2 µg/ml) staining was performed before mounting the sample. Images were made on a Leica SP5 (Leica Microsystems), using a $63\times$ objective and a 405-nm laser to excite Hoechst 33342 and CFW, a 488 nm laser for GFP and 561 nm for mRFP. 5 µm thick Z-stacks were made with 15 slides. Image analysis was performed on maximum Z-projections by scoring different phenotypes, counting 200–500 cells per biological replicate.

Screening

The screening of the β 1-RITE + KO library was performed in batches of 384 strains. A RoToR HAD (Singer Instruments) was used to transfer the strains from 384-well glycerol stocks to a YEPD+Hygromycin agar plate. 120 µl start cultures in 96-well plates were inoculated from the YEPD plate, grown overnight and used to start a 4 ml culture. A 2-day starvation period was followed with a switch assay as described by Verzijlbergen *et al.* (Verzijlbergen *et al.*, 2010) and another 3-day starvation period. Samples were then fixed in 4% formaldehyde, stained with Hoechst and spotted on a yeast array (Narayanaswamy *et al.*, 2006). Microscopic analysis was performed on a Leica AOBS LSCM (Leica Microsystems) using 405-, 488- and 561-nm laser light to excite Hoechst 33342, GFP and mRFP respectively. High-throughput image analysis was performed using CellProfiler software (Carpenter *et al.*, 2006). The screening results of selected candidates were validated by two additional rounds of analysis and independent generation of the knockout yeast strains.

Flow-based sorting of old and young cells

To isolate replicative old and young cells, $\sim 2 \times 10^7$ cells of a starved culture were stained with CFW and FACS sorted on a MoFlo-Astrios (Beckman Coulter) using 405 nm excitation and collecting fluorescence emission with a 450 nm (30-nm bandpass) filter. The 2.5% of the cells with the highest and lowest CFW signal were isolated and ~250 yeasts were subsequently plated on YEPD plates (Allen *et al.*, 2006). CFUs were counted after 3 days culture at 30°C.

Protein extraction and native gel analysis

Native protein samples were made by washing a cell pellet of ~10⁸ cells in PBS plus protease inhibitors (1 mM PMSF, 5 mM benzamidine, 1 µg/ml pepstatin, 1 µg/ml leupepting) and resuspended in buffer A (20 mM Tris-HCl pH 7.4, 5 mM MgCl2, 1 mM DTT, 1 mM ATP) plus protease inhibitors. Cells were lysed in buffer A by using glass beads, which were removed before addition of a blue loading buffer (5×; 50% glycerol and Bromophenol Blue). Samples were loaded on a NativePAGE[™] 3–12% Bis-Tris gel (Life Technologies) and ran in NativePAGE[™] running buffer (Life Technologies). GFP fluorescence was visualized on 4

a ProXPRESS (Perkin Elmer) machine with 480 nm (30-nm bandpass) excitation and 550 nm (40-nm bandpass) emission filters. To visualize untagged proteasomes, the gel was incubated with 100 μ M suc-LLVY-AMC (Enzo Life Sciences) in the presence of 1 mM ATP, 1 mM DTT and 0.02% SDS (Elsasser *et al.*, 2005). Gel scans were made with 390 nm (70-nm bandpass) excitation and 450 nm (20-nm bandpass) emission filters.

Acknowledgements

We thank L. Oomen and L. Brocks (Digital Microscopy Facility, Netherlands Cancer Institute) for technical microscopy assistance, A. Pfauth (Flow Cytometry Facility, Netherlands Cancer Institute) for yeast cell sorts and W. Brugman (Central Microarray and Deep Sequencing Core Facility, Netherlands Cancer Institute) for printing yeast arrays.

References

Allen, C., Buttner, S., Aragon, A., Thomas, J., Meirelles, O., Jaetao, J., Benn, D., Ruby, S., Veenhuis, M., Madeo, F. *et al.* (2006). Isolation of quiescent and nonquiescent cells from yeast stationary-phase cultures. J Cell Biol 174, 89-100.

Amm, I., Sommer, T. and Wolf, D. (2014). Protein quality control and elimination of protein waste: The role of the ubiquitin proteasome system. Biochim Biophys Acta 1843, 182-196.

Aragon, A. D., Rodriguez, A. L., Meirelles, O., Roy, S., Davidson, G. S., Tapia, P. H., Allen, C., Joe, R., Benn, D. and Werner-Washburne, M. (2008). Characterization of Differentiated Quiescent and Nonquiescent Cells in Yeast Stationary-Phase Cultures. Molecular Biology of the Cell 19, 1271-1280.

Arnesen, T. (2011). Towards a Functional Understanding of Protein N-Terminal Acetylation. PLoS Biol 9, e1001074.

Arnesen, T., Van Damme, P., Polevoda, B., Helsens, K., Evjenth, R., Colaert, N., Varhaug, J., Vandekerckhove, J., Lillehaug, J., Sherman, F. *et al.* (2009). Proteomics analyses reveal the evolutionary conservation and divergence of N-terminal acetyltransferases from yeast and humans. PNAS 106, 8157-8162.

Baker Brachmann, C., Davies, A., Cost, G., Caputo, E., Li, J., Hieter, P. and Boeke, J. (1998). Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: A useful set of strains and plasmids for PCR-mediated gene disruption and other applications. Yeast 14, 115-132.

Carpenter, A., Jones, T., Lamprecht, M., Clarke, C., Kang, I., Friman, O., Guertin, D., Chang, J., Lindquist, R., Moffat, J. *et al.* (2006). CellProfiler: image analysis software for identifying and quantifying cell phenotypes. Genome Biol 7, R100.

Carrard, G., Bulteau, A., Petropoulos, I. and Friguet, B. (2002). Impairment of proteasome structure and function in aging. Int J Biochem Cell Biol 34, 1461-1474.

Chen, Q., Thorpe, J., Dohmen, J., Li, F. and

Keller, J. (2006). Ump1 extends yeast lifespan and enhances viability during oxidative stress: Central role for the proteasome? Free Radic Biol Med 40, 120-126.

Clay, L. and Barral, Y. (2013). New approaches to an age-old problem. Current Opinion in Biotechnology 24, 784-789.

Dasuri, K., Zhang, L., Ebenezer, P., Liu, Y., Fernandez-Kim, S. and Keller, J. (2009). Aging and dietary restriction alter proteasome biogenesis and composition in the brain and liver. Mech Ageing Dev 130, 777-783.

Delaney, J., Murakami, C., Chou, A., Carr, D., Schleit, J., Sutphin, G., An, E., Castanza, A., Fletcher, M., Goswami, S. *et al.* (2013). Dietary restriction and mitochondrial function link replicative and chronological aging in *Saccharomyces cerevisiae*. Exp Geront 48, 1006-1013.

Donmez, G. and Guarente, L. (2010). Aging and disease: connections to sirtuins. Aging Cell 9, 285-290.

Elsasser, S., Schmidt, M. and Finley, D. (2005). Characterization of the Proteasome Using Native Gel Electrophoresis. In Methods in Enzymology, vol. Volume 398 (ed. J. D. Raymond), pp. 353-363: Academic Press.

Gardner, R., Nelson, Z. and Gottschling, D. (2005). Degradation-Mediated Protein Quality Control in the Nucleus. Cell 120, 803-815.

Guedes-Dias, P. and Oliveira, J. (2013). Lysine deacetylases and mitochondrial dynamics in neurodegeneration. Biochim Biophys Acta 1832, 1345-1359.

Hebert, A., Dittenhafer-Reed, K., Yu, W., Bailey, D., Selen, E., Boersma, M., Carson, J., Tonelli, M., Balloon, A., Higbee, A. *et al.* (2013). Calorie Restriction and SIRT3 Trigger Global Reprogramming of the Mitochondrial Protein Acetylome. Mol Cell 49, 186-199.

Heck, J., Cheung, S. and Hampton, R. (2010). Cytoplasmic protein quality control degradation mediated by parallel actions of the E3 ubiquitin ligases Ubr1 and San1. PNAS 107, 1106-1111.

Henderson, K. and Gottschling, D. (2008). A mothers sacrifice: what is she keeping for herself? Current Opinion in Cell Biology 20, 723-728.

Hwang, C., Shemorry, A. and Varshavsky, A. (2010). N-Terminal Acetylation of Cellular Proteins Creates Specific Degradation Signals. Science 327, 973-977.

Janke, C., Magiera, M., Rathfelder, N., Taxis, C., Reber, S., Maekawa, H., Moreno-Borchart, A., Doenges, G., Schwob, E., Schiebel, E. *et al.* (2004). A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. Yeast 21, 947-962.

Kaeberlein, M. (2010). Lessons on longevity from budding yeast. Nature 464, 513-519.

Kennedy, B. K., Austriaco, N. R. and Guarente, L. (1994). Daughter cells of *Saccharomyces cerevisiae* from old mothers display a reduced life span. J Cell Biol 127, 1985-1993.

Koga, H., Kaushik, S. and Cuervo, A. (2011). Protein homeostasis and aging: The importance of exquisite quality control. Ageing Research Reviews 10, 205-215.

Kruegel, U., Robison, B., Dange, T., Kahlert, G. and Delaney, J. (2011). Elevated Proteasome Capacity Extends Replicative Lifespan in *Saccharomyces cerevisiae*. PLoS Genet, 1371.

Laporte, D., Salin, B., Daignan-Fornier, B. and Sagot, I. (2008). Reversible cytoplasmic localization of the proteasome in quiescent yeast cells. J Cell Biol 181, 737-745.

Lee, C., Klopp, R., Weindruch, R. and Prolla, T. (1999). Gene Expression Profile of Aging and Its Retardation by Caloric Restriction. Science 285, 1390-1393.

Lehmann, A., Janek, K., Braun, B., Kloetzel, P. and Enenkel, C. (2002). 20 S proteasomes are imported as precursor complexes into the nucleus of yeast. J Mol Biol 317, 401-413.

Menendez-Benito, V., van Deventer, S. J., Jimenez-Garcia, V., Roy-Luzarraga, M., van Leeuwen, F. and Neefjes, J. (2013). Spatiotemporal analysis of organelle and macromolecular complex inheritance. Proceedings of the National Academy of Sciences 110, 175-180.

Michal Jazwinski, S., Egilmez, N. K. and Chen, J. B. (1989). Replication control and cellular life span. Experimental Gerontology 24, 423-436.

Mortimer, R. K. and Johnston, J. R. (1959). Life Span of Individual Yeast Cells. Nature 183, 1751-1752.

Murakami, C., Delaney, J., Chou, A., Carr, D., Schleit, J., Sutphin, G., An, E., Castanza, A., Fletcher, M., Goswami, S. *et al.* (2012). pH neutralization protects against reduction in replicative lifespan following chronological aging in yeast. Cell Cycle 11, 3087-3096.

Narayanaswamy, R., Niu, W., Scouras,

A., Hart, G. T., Davies, J., Ellington, A., Iyer, V. and Marcotte, E. (2006). Systematic profiling of cellular phenotypes with spotted cell microarrays reveals mating-pheromone response genes. Genome Biol 7, R6.

Nystrom, T. and Liu, B. (2014). The mystery of aging and rejuvenation; a budding topic. Current Opinion in Microbiology 18, 61-67.

Polevoda, B., Norbeck, J., Takakura, H., Blomberg, A. and Sherman, F. (1999). Identification and specificities of N-terminal acetyltransferases from *Saccharomyces cerevisiae*. The EMBO Journal 18, 6155-6168.

Polevoda, B. and Sherman, F. (2000). N-terminal Acetylation of Eukaryotic Proteins. J Biol Chem 275, 36479-36482.

Polevoda, B. and Sherman, F. (2003). N-terminal Acetyltransferases and Sequence Requirements for N-terminal Acetylation of Eukaryotic Proteins. J Mol Biol 325, 595-622.

Powers, E., Morimoto, R., Dillin, A., Kelly, J. and Balch, W. (2009). Biological and Chemical Approaches to Diseases of Proteostasis Deficiency. Annu Rev Biochem 78, 959-991.

Prasad, R., Kawaguchi, S. and Ng, D. (2010). A Nucleus-based Quality Control Mechanism for Cytosolic Proteins. Mol Biol Cell 21, 2117-2127.

Pringle, J. (1991). [52] Staining of bud scars and other cell wall chitin with Calcofluor. In Methods in Enzymology

Guide to Yeast Genetics and Molecular Biology, (ed. G. Christine Guthrie), pp. 732-735: Academic Press.

Reits, E., Benham, A., Plougastel, B., Neefjes, J. and Trowsdale, J. (1997). Dynamics of proteasome distribution in living cells. EMBO J 16, 6087-6094.

Russell, S., Steger, K. and Johnston, S. (1999). Subcellular Localization, Stoichiometry, and Protein Levels of 26 S Proteasome Subunits in Yeast. J Biol Chem 274, 21943-21952.

Schmidt, M. and Finley, D. (2014). Regulation of proteasome activity in health and disease. Biochim Biophys Acta 1843, 13-25.

Scott, D., Monda, J., Bennett, E., Harper, J. W. and Schulman, B. (2011). N-Terminal Acetylation Acts as an Avidity Enhancer Within an Interconnected Multiprotein Complex. Science 334, 674-678.

Starheim, K., Gevaert, K. and Arnesen, T. (2012). Protein N-terminal acetyltransferases: when the start matters. Trends Biochem Sci 37, 152-161.

Storici, F. and Resnick, M. (2006). The Delitto Perfetto Approach to *In Vivo* Site Directed Mutagenesis and Chromosome Rearrangements with Synthetic Oligonucleotides in Yeast. In Methods in Enzymology

DNA Repair, Part B, (ed. L. C. Judith), pp. 329-345: Academic Press.

Tenreiro, S. and Outeiro, T. (2010). Simple is good: yeast models of neurodegeneration. FEMS Yeast Research 10, 970-979.

Tong, A. and Boone, C. (2006). Synthetic genetic array analysis in Saccharomyces cerevisiae. Methods Mol Biol 313:171-92.

Tong, A., Evangelista, M., Parsons, A., Xu, H., Bader, G., Pag, N., Robinson, M., Raghibizadeh, S., Hogue, C., Bussey, H. et al. (2001). Systematic Genetic Analysis with Ordered Arrays of Yeast Deletion Mutants. Science 294, 2364-2368.

Tonoki, A., Kuranaga, E., Tomioka, T., Hamazaki, J., Murata, S., Tanaka, K. and Miura, M. (2009). Genetic Evidence Linking Age-Dependent Attenuation of the 26S Proteasome with the Aging Process. Mol Cel Biol 29, 1095-1106.

van Leeuwen, F. and Gottschling, D. (2002). Assays for gene silencing in yeast. In Methods in Enzymology

Guide to Yeast Genetics and Molecular and Cell Biology - Part B, (ed. C. G. a. Gerald), pp. 165-186: Academic Press.

Vernace, V., Arnaud, L., Schmidt-Glenewinkel, T. and Figueiredo-Pereira, M. (2007a). Aging perturbs 26S proteasome assembly in Drosophila melanogaster. FASEB J 21, 2672-2682.

Vernace, V., Schmidt-Glenewinkel, T. and Figueiredo-Pereira, M. (2007b). Aging and regulated protein degradation: who has the UPPer hand? Aging Cell 6, 599-606.

Verzijlbergen, K., Menendez-Benito, V., van Welsem, T., van Deventer, S., Lindstrom, D., Ovaa, H., Neefjes, J., Gottschling, D. and van Leeuwen, F. (2010). Recombination-induced tag exchange to track old and new proteins. PNAS 107, 64-68.

Wierman, M. and Smith, J. (2014). Yeast sirtuins and the regulation of aging. FEMS Yeast Research 14, 73-88.



Supplemental Figures:

Fluorescent gel scanner: GFP

Fig S1: β1-GFP is efficiently incorporated in the 20S proteasome in WT and NatA, B or C deficient cells A fluorescent gel scan of a native gel shows one GFP fluorescent band around 720 kDa for both β 1-GFP and NatA, B or C deficient starved cells. This band runs at the same height as a purified 20S control (Mouse 20S proteasome, Boston Biochem) and is also present in a WT sample without GFP as was visualized by incubating the gel with a suc-LLVY-AMC (Enzo Life Sciences) proteasome activity probe. The marker (M) was visualized by Coomassie staining of the same gel. We conclude that the presence of the GFP tag on $\beta 1$ doesn't induce the presence of proteasome assembly intermediates or a pool of unincorporated substrates in β1-GFP or NatA, B or C deficient cells.

Fluorescent gel scanner: LLVY-AMC

Proteasome localization and aging



Fig S2: GFP tagged a2 (Pre8) and Rpn1 show the same proteasome localization as β 1-GFP in WT and NatA,B or C deficient cells

(A) Live cell microscopy of starved WT and NatA ($ard1\Delta$), NatB ($nat3\Delta$) or NatC ($mak3\Delta$) deficient cells endogenously expressing a2(Pre8)-GFP. Cells were stained with CalcoFluor White to asses replicative age. (B) The prevalence of the different phenotypes in the different age groups was scored in two independent experiments (~200 live cells per replicate). (C) The same experiment as in (A), but now with cells endogenously expressing RPN1-GFP, a component of the 19S base particle. (D) Prevalence of the different proteasome phenotypes in the three age groups of living cells. (Scale bars, 5 μ m) 4

Fig. S3



Fig S3: Plating assay confirms efficient recombination in hul5 Δ , mak10 Δ and uba3 Δ cells

(A) Genetic recombination of the RITE cassette results in the loss of Hygromycin resistance, which can be used to assess the percentage of cells that underwent a recombination event as described by (Verzijlbergen *et al.*, 2010). Samples were taken just before (Pre) and ~16h after (Post) recombination was induced. In both WT and mutant cells the recombination without induction is low and the induced recombination is high. (B) Live cell microscopy of WT cells and the three nuclear retention hits. Images were taken after 5 days starvation, but the recombination was induced after one day starvation instead of two days. Under those conditions synthesis of new (mRFP-tagged) proteasome can be detected, thereby validating the RITE tool in starvation conditions. (Scale bar, 5 μ m)

Proteasome localization and aging



Fig S4: Loss of the non-catalytic subunits of the different N-acetylation complexes results in the same phenotype as loss of the catalytic subunit and N-acetylation status of defined proteins did not affect the localization of the proteasome in starved cells

(A) Loss of NatC activity by knockout of one of its subunits (Mak3, Mak10 and Mak31) or exchange for a catalytic inactive Mak3 leads to an altered proteasome localization after a five day starvation period. (B) Cells showing nuclear retention, PSGs or a combination of both were scored in three independent experiments. Approximately 200 cells were scored in two independent samples per condition. Significance was calculated with a non-paired, two-tailed t-test (* = p<0.05, ** = p<0.01). (C) Loss of any single NatA subunit (Nat1, Nat5 and Ard1) does not lead to different proteasome localization after a five day starvation period. (D) Quantification as in (B). (E) Loss of any single NatB subunit (Nat3 and Mdm20) alters proteasome localization after a five day starvation period. (F) Quantification as in (B). (G) Several predicted NatC substrate proteins were N-terminally mutated to test whether their N-acetylation status was important for proteasome localization in starvation. An X2P mutation was made of $\alpha 6$ (Pre5), $\alpha 5$ (Pup2), Nup100, Avo2, Hul5, FUB1 and RPN9 and these strains were subjected to a five day starvation period. All X2P mutats showed a proteasome localization similar to the WT and can thus be excluded as a NatC substrate influencing proteasome localization. (Scale bars, 5 µm)

4

Supplementary Table 1:

List of used strains:

Strain:	Genotype:
NKI4103*	MAT@ can1d::STE2pr-Sp_his5 lyp1d his3d1 leu2d0 ura3d0 met15d0 LYS2+ pre3::PRE3-V5-loxP-HA-yEGFP-HYG-loxP-T7-mRFP lyp1d::NATMX-GPD_CRE_EBD78
NKI5537	NKI4103 + mak3Δ::URA3
NKI5538	NKI4103 + mak31Δ::LEU2
NKI5539	NKI4103 + mak10Δ::KanMX4
NKI5540	NKI4103 + nat3∆::URA3
NKI5541	NKI4103 + mdm20∆::URA3
NKI5542	NKI4103 + ard1Δ::URA3
NKI5543	NKI4103 + nat1Δ::URA3
NKI5544	NKI4103 + nat5Δ::URA3
NKI5545	NKI4103 + mak3Δ::KanMX4, nat3Δ::URA3
NKI5546	NKI4103 + mak3::MAK3(N123A,Y130A)
NKI4101*	MAT@ can1d::STE2pr-Sp_his5 lyp1d his3d1 leu2d0 ura3d0 met15d0 LYS2+ pre3::PRE3-V5-loxP-HA-yEGFP-HYG-loxP-T7-mRFP
NKI5547	NKI4101 + pre5::NatNT2-PCYC1-PRE5(F2P)
NKI5548	NKI4101 + pup2::NatNT2-PCYC1-PUP2 (F2P)
NKI5549	NKI4101 + pre5::NatNT2-PCYC1-PRE5(F2P) pup2::KanMX4-PCYC1-PUP2(F2P)
NKI5550	NKI4101 + nup100::KanMX4-PCYC1-NUP100(F2P)
NKI5551	NKI4101 + avo2::NatNT2-PCYC1-AVO2(L2P)-3xFlag-KanMX4
NKI5552	NKI4101 + hul5::NatNT2-PCYC1-HUL5(L2P)
NKI5553	NKI4101 + fub1::NatNT2-PCYC1-FUB1(I2P)
NKI5554	NKI4101 + rpn9::NatNT2-PCYC1-RPN9(L2P)
NKI4105	MAT@ can1d::STE2pr-Sp_his5 lyp1d his3d1 leu2d0 ura3d0 met15d0 LYS2+ pre8::PRE8-V5-loxP-HA-yEGFP-HYG-loxP-T7-mRFP lyp1d::NATMX-GPD_CRE_EBD78
NKI5555	NKI4105 + ard1Δ::URA3
NKI5556	NKI4105 + nat3∆::URA3
NKI5557	NKI4105 + mak3Δ::URA3
NKI4121	MAT@ can1d::STE2pr-Sp_his5 lyp1d his3d1 leu2d0 ura3d0 met15d0 LYS2+ rpn1::RPN1-V5-loxP-HA-yEGFP-HYG-loxP-T7-mRFP lyp1d::NATMX-GPD_CRE_EBD78
NKI5558	NKI4121 + ard1Δ::URA3
NKI5559	NKI4121 + nat3Δ::URA3
NKI5560	NKI4121 + mak3Δ::URA3

* As described in Verzijlbergen et al 2009

Proteasome localization and aging

#