

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/29572> holds various files of this Leiden University dissertation.

Author: Berg, Rosaline van den


Title: Spondyloarthritis : recognition, imaging, treatment

Issue Date: 2014-10-29

First update of the current evidence for the management of ankylosing spondylitis with non-pharmacological treatment and non-biological drugs: a systematic literature review for the ASAS/EULAR management recommendations in ankylosing spondylitis

Rheumatology (Oxford) 2012;51:1388-96

R. van den Berg
X. Baraliakos
J. Braun
D. van der Heijde



ABSTRACT

Objective

To perform a systematic literature review as a basis for the update of the Assessment in SpondyloArthritis International Society and European League Against Rheumatism (ASAS/EULAR) recommendations for the management of AS with non-pharmacological interventions and non-biologic drugs.

Methods

The search was performed in PubMed, EMBASE, PEDro and Cochrane between 1 January 2005 and 1 December 2009, and in abstracts of EULAR and ACR meetings (2007-09). Effect sizes for outcomes on pain, disease activity, spinal mobility and physical function and level of evidence were presented.

Results

Of 2383 papers, 35 with complete data were included. Physical therapy exercises in various modalities have positive effects on BASFI, BASDAI, pain and mobility function. Various NSAIDs including coxibs improve BASDAI, disease activity and BASFI. No effect of SSZ and MTX on any variable was found. Surgical interventions of the spine and the hip can give excellent results by restoring function.

Conclusion

This concise summary of current evidence for non-pharmacological interventions and non-biologic drugs formed the basis for the update of the ASAS/EULAR recommendations for the management of AS.

INTRODUCTION

AS is a chronic, inflammatory rheumatic disease, generally starting early in life¹⁻⁴. Inflammatory back pain due to sacroiliitis and spondylitis, and formation of syndesmophytes leading to ankylosis of the spine, characterize AS^{4,5}. Although AS is difficult to treat, the treatment armamentarium of AS has been broadened since the discovery of anti-TNF- α agents as effective treatments⁶⁻⁸.

Clinicians need to be aware of the relative benefits and risks of the available treatments, and need to have evidence-based information about the most efficacious strategies in particular patient settings⁴.

In 2005 Zochling *et al.*⁴ performed a systematic literature search for evidence based recommendations by the Assessment in SpondyloArthritis International Society and European League Against Rheumatism (ASAS/EULAR) for the management of AS. In 2010 an update of this systematic literature search was performed to serve as a base for the development of an update of the ASAS/EULAR recommendations⁹. The details and results of the performed systematic review on non-pharmacological interventions and non-biologic drugs are presented in this article. The results on biologics are presented in the article by Baraliakos elsewhere in this journal.

METHODS

Participants and outcome measures

Participants were defined as patients with a diagnosis of AS or axial spondyloarthritis. The required treatments were non-pharmacological interventions and non-biologic drugs. There were no restrictions with regard to type of non-pharmacological intervention, or to dose, duration or route of administration of non-biologic drugs.

The primary outcomes of interest include pain, disease activity (including BASDAI), spinal mobility (including BASMI) and physical function (including BASFI).

Inclusion/exclusion criteria

Randomized controlled trials (RCTs) and controlled trials are the ideal study designs for inclusion in this review.

However, the aim of this review is to provide evidence of all types of non-pharmacological interventions and non-biologic drugs. Since not all types of treatment can be studied within RCTs alone, the main focus of interest was also on systematic reviews, uncontrolled trials/cohort studies, case-control studies and cross-sectional studies. Studies about non-axial spondyloarthritides and other inflammatory joint conditions, animal studies, non-clinical outcome studies and non-treatment studies, narrative review articles, commentaries, guidelines, case reports, letters and editorials and studies in other languages than English, Dutch and German were excluded.

Studies about biologic drugs were also excluded because those studies will be included in a search performed by Baraliakos.

Systematic literature search

A search strategy was built in collaboration with an experienced librarian, based on the previous search of Zochling *et al.*⁴. The systematic literature search for published papers was performed in the electronic databases PubMed, EMBASE, PEDro and Cochrane between 1 January 2005, which is the end date of the literature search by Zochling *et al.*⁴, and 1 December 2009. The complete search strategies for the databases are provided in supplementary appendix S1, available as supplementary data at Rheumatology Online.

Abstracts of rheumatology scientific meetings (EULAR, ACR) from the years 2007, 2008 and 2009 were searched by hand to ensure that all potential studies were identified for this review. Furthermore, references of relevant reviews and included papers were hand searched for information on any other relevant studies.

Selection of studies

One reviewer (RvdB) assessed each title and abstract on suitability for inclusion in the review, according to the inclusion and exclusion criteria described above. Papers not addressing the topic of interest were excluded and reasons recorded. The full-text paper was assessed when further information was required to determine if the inclusion criteria were met.

Data extraction and categorizing evidence

The included papers were assessed using the full-text paper by one reviewer (RvdB) to extract relevant data, including patient characteristics and details of treatment.

If necessary, authors were contacted to provide any required additional information. The results were reported to the ASAS/EULAR expert committee at the beginning of the recommendation development process. All included papers were categorized according to their level of evidence (see legend in table 1)³. The assigned levels are shown in table 1.

Data analysis

Since different types of studies are included about various types of treatments, the results are very heterogeneous and therefore the results cannot be pooled. Yet, the results are analysed and presented per type of treatment.

Estimation of effectiveness

Per treatment group, the Cohen's effect size (Cohen's ES; mean change in score divided by the baseline SD) was calculated, and the standardized response mean (SRM; mean change divided by the SD of the change) was calculated where possible³⁶. To compare the effect between treatment groups, treatment ES was calculated (mean change in the index group minus the mean change in the comparator group divided by a pooled baseline SD). For each ES, the corresponding 95% CI was constructed. An ES of 0.2 or 0.3 is considered a small change, around 0.5 as moderate and >0.8 as a large change, and a negative ES indicates worse.

RESULTS

Treatment modalities and types of research evidence

The general search revealed 3179 papers; 1638 in PubMed, 1486 in EMBASE, 14 in PEDro, 34 in Cochrane and 7 abstracts. After eliminating duplicates, 2383 papers remained. Of those, 2347 papers were excluded (supplementary appendix S2, available as supplementary data at Rheumatology Online) and 35 papers were included, of which 3 are Cochrane reviews and 1 abstract (supplementary appendix S3, available as supplementary data at Rheumatology Online). An overview of the included papers is shown in the supplementary appendix S4, available as supplementary data at Rheumatology Online.

Non-pharmacological treatment

No studies on treatments about diet, education, self-help groups or lifestyle modification were present within this search.

Table 1: Cohen's effect size (Cohen's ES) with 95% confidence interval (95% CI) for various outcome measures and different management modalities.

Intervention	Assessment point	No. of patients	Level of evidence	Cohen's ES BASFI (95% CI)	Cohen's ES BASDAI (95% CI)	Cohen's ES Pain (95% CI)	Cohen's ES Disease act. (95% CI)	Cohen's ES BASMI (95% CI)
Exercise therapy								
Group exercise ¹⁵	6 weeks	22	2b	0.27 (-0.33, 0.86)	-	0.50 (-0.10, 1.10)	-	0.26 (-0.34, 0.86)
Home exercise		16		0 (-0.69, 0.69)	-	0.20 (-0.50, 0.89)	-	0.05 (-0.64, 0.69)
Exercise group ¹³	6 weeks	22	3	0.10 (-0.49, 0.69)	-	-	-	-
Exercise Group ¹⁴	8 weeks	16	1b	0.41 (-0.29, 1.11)	0.38 (-0.31, 1.08)	-	-	-
Control group		16		0.22 (-0.47, 0.92)	0.24 (-0.46, 0.93)	-	-	0.10 (-0.60, 0.79)
Hospital exercise ¹⁷	12 weeks	23		0.94 (0.34, 1.55)	1.00 (0.39, 1.61)	0.76 (0.16, 1.36)	-	-
Home exercise		23		0.22 (-0.36, 0.80)	0.88 (0.28, 1.49)	0.11 (-0.47, 0.69)	-	-
Hospital exercise	6 months	23	1b	0.78 (0.18, 1.38)	0.84 (0.24, 1.45)	0.48 (-0.10, 1.07)	-	-
Home exercise		23		0.48 (-0.11, 1.06)	1.12 (0.50, 1.74)	0.48 (-0.10, 1.07)	-	-
Home exercise ¹⁸	12 weeks	25		0.63 (0.07, 1.20)	0.94 (0.35, 1.52)	-	-	-
Control group		18	2b	0.18 (-0.48, 0.83)	0.19 (-0.48, 0.84)	-	-	-

Table 1: continued

Exercise GPR method ¹⁹	20	0.57 (-0.06, 1.21)	1.07 (0.41, 1.73)	1.28 (0.60, 1.96)	-	-
Conventional exercise	21	0.66 (0.04, 1.28)	1.20 (0.54, 1.86)	1.03 (0.38, 1.67)	-	-
Control group	15	0.13 (-0.59, 0.85)	0.32 (-0.40, 1.04)	0.19 (-0.53, 0.91)	-	-
Balneotherapy						
Exercise + Stangerbath ²⁰	29	1.05 (-0.08, 2.17)	1.71 (0.33, 3.09)	-	-	0.30 (-0.44, 1.05)
Exercise	28	0.20 (-0.48, 0.88)	0.36 (-0.43, 1.15)	-	-	0.06 (-0.52, 0.63)
Balneotherapy ²²	20	-	-	0.81 (-0.26, 1.88)	-	-
Balneotherapy + NSAIDs	21	-	-	0.83 (-0.26, 1.92)	-	-
NSAID	20	-	-	0.58 (-0.38, 1.54)	-	-
Balneotherapy	20	-	-	0.97 (-0.17, 2.11)	-	-
Balneotherapy + NSAIDs	21	-	-	1.10 (-0.10, 2.10)	-	-
NSAID	20	-	-	0.96 (-0.18, 2.09)	-	-
Rehabilitation ²⁴	52	2.23 (1.74, 2.72)	-	5.10 (4.31, 5.89)	-	-
	3	1.70 (1.25, 2.14)	-	3.62 (3.00, 4.24)	-	-
	12 weeks	1.16 (0.75, 1.58)	-	2.06 (1.58, 2.54)	-	-

Table 1: continued

NSAIDs							
Celecoxib (200 mg) ²⁷	153		0.35 (0.12, 0.57)	0.60 (0.37, 0.83)	1.96 (1.69, 2.23)	1.11 (0.87, 1.35)	0.52 (0.29, 0.75)
Celecoxib (400 mg)	150	1b	0.41 (0.18, 0.64)	0.80 (0.56, 1.03)	1.79 (1.52, 2.06)	1.29 (1.05, 1.54)	0.13 (-0.10, 0.36)
Diclofenac (150 mg)	155		0.35 (0.12, 0.57)	0.83 (0.60, 1.06)	1.84 (1.57, 2.10)	1.28 (1.03, 1.52)	0.18 (-0.04, 0.40)
Etoricoxib (90 mg) ²⁵	22	3	1.08 (0.45, 1.71)	1.08 (0.45, 1.71)	1.44 (0.77, 2.10)	0.93 (0.31, 1.56)	-
DMARDs							
Sulfasalazine (2g) ³²	120	1b	0.19 (-0.06, 0.44)	1.11 (0.84, 1.38)	-	-	-
Sulfasalazine (3g) ³³	187	1b	0.29 (0.08, 0.49)	1.24 (1.02, 1.46)	0.95 (0.74, 1.17)	-	0.08 (-0.13, 0.28)
Etanercept (50 mg)	379		0.48 (0.33, 0.62)	1.96 (1.79, 2.14)	1.68 (1.51, 1.84)	-	0.34 (0.20, 0.48)
Sulfasalazine (2g) ³⁴	16	3	-	0.43 (-0.28, 1.13)	-	-	-
MTX (15mg up to 20mg) ³¹	20	3	0.01 (-0.61, 0.63)	0.00 (-0.62, 0.62)	0.00 (-0.62, 0.62)	-	0.23 (-0.12, 1.14)
Other							
Radium-224 ³⁶	278	3	0.68 (0.39, 0.97)	0.86 (0.57, 1.15)	1.14 (0.85, 1.44)	1.22 (0.92, 1.52)	-

Table 1: continued

Surgery	No. of patients	Results
Hip resurfacing ³⁷	23 (38 hips)	Both groups ROM ↑: resurfacing group significantly better than THR group
Total hip replacement (THR)	25 (41 hips)	
Posterior correction and fixation without anterior fusion (posterior opening-wedge osteotomy) ³⁹	30	Neurologic deficit ↑ good correction of kyphosis
Open wedge osteotomy (OWO) ³⁸	51	Both techniques have good clinical outcome, patient quality of life ↑, high patient satisfaction: no difference between the two techniques
Closed wedge osteotomy (CWO)	66	
Cervicothoracic extension osteotomy ⁴⁰	26	Neck pain and swallowing problems ↑ Both groups restored horizontal gaze
Cervical extension osteotomy: Conventional technique ⁴¹	114	High patient satisfaction, function ↑, psycho-social body image ↑: no difference in outcome between the two groups
Cervical extension osteotomy: Current technique	17	
Smith-Peterson osteotomy ⁴²	12	Pain and neurologic deficits ↑, high patient satisfaction
Lumbar closing wedge osteotomy ⁴³	11	Horizontal gaze restored in all patients
Cervical osteotomy: sitting position ⁴⁵	11	Good correction of kyphosis, no loss of correction
Cervical osteotomy: prone position	5	
Cervical decancellation closing wedge osteotomy ⁴⁶	8	Horizontal gaze ↑, good subjective outcome
Closing wedge osteotomy ⁴⁴	21	Quality of life ↑, good functional outcome

Significant ES in **bold italics**. Category of evidence: Ia = meta-analysis of randomized controlled trials (RCT); Ib = RCT; IIa = controlled study without randomization; IIb = quasi-experimental study; III = non-experimental descriptive studies (comparative, correlation and case-control studies); IV = expert committee reports or opinion or clinical experience of respected authorities, or both.

Exercise therapy

The effect of physiotherapy has been reviewed in a Cochrane review in 2008³⁷. The results of this review show that individual home-based or supervised exercise programmes are better than no intervention at all on pain, physical function, spinal mobility and patient global assessment, and that supervised group physiotherapy is better than home exercise³⁷. Besides the Cochrane review, nine papers were identified^{10-15, 38-40} of which three were already included in the Cochrane review³⁸⁻⁴⁰. In the six additional papers, the effects of various exercises in AS patients are compared (supplementary appendix S4, available as supplementary data at Rheumatology Online). The results of these six studies confirm the results of the Cochrane review. Various types of exercise [supervised group, home and Global Posture Reeducation (GPR) method exercise] have moderate to good effects on BASFI, BASDAI, pain and mobility, as shown by the calculated Cohen's ES and SRM (table 1 and figure 1). The calculated treatment ES showed that supervised group physiotherapy is better than home exercise on BASFI, pain and mobility, and slightly better on BASDAI. Home exercise is better than no exercise at all on BASFI and BASDAI (table 2).

Although most papers had level 1b evidence, the studies investigated various exercises with variable durations and had small patient samples. Therefore, many ES are not statistically significant, showing only a trend (table 1).

Balneotherapy, spa therapy and rehabilitation

The same Cochrane review also revealed that combined inpatient spa exercise therapy followed by group physiotherapy is better than group physiotherapy alone³⁷. In addition to the Cochrane review, four RCTs^{16, 17, 42, 43} about various types of balneotherapy and spa therapy in AS patients were identified (level 1b evidence), of which two were already presented in the Cochrane review^{42, 43}.

As in the exercise therapy studies, the studies about balneotherapy included only small patient numbers in various therapies, resulting in not statistically significant ES.

However, the trend shows that balneotherapy in all its modalities is (moderate) effective on BASFI, BASDAI and pain, as shown by the calculated Cohen's ES and treatment ES (tables 1 and 2). The effect of balneotherapy on pain is equal to the effect of NSAIDs (either mono or combined)¹⁷. Stangerbath therapy combined with exercises is effective on BASFI and BASDAI, only directly after therapy¹⁶ (figure 1). One level 3 evidence study about the effect of inpatient rehabilitation was identified that showed a strong effect on BASFI, pain and OWD (table 1)¹⁸.

NSAIDs

Three studies about the effects of different NSAIDs in AS patients were identified^{19, 20, 41}. The effect of celecoxib (200 and 400mg daily) in comparison with diclofenac (150mg daily) (level 1b evidence)¹⁹, the effect of etoricoxib (90mg daily) (level 3 evidence)²⁰ and the effect of NSAIDs in continuous usage in comparison with NSAID usage on demand (level 1b evidence)⁴¹ were investigated.

The latter study is a follow-up study of a double-blind RCT about the effect of celecoxib 200mg versus ketoprofen 200mg versus placebo after 6 weeks⁴⁴. This study was already included in the review of Zochling *et al.*⁴, showing a significant improvement in pain and function after 6 weeks of use of both NSAIDs in comparison with placebo⁴⁴. The follow-up study showed that measures of disease activity, including pain and BASDAI, were stable over a time period of 24 months in both the continuous and on-demand groups and not statistically significant between the groups⁴¹. Although the clinical effects of both treatment strategies are similar, inhibition of structural damage progression in the spine is better with continuous use than with on-demand use⁴¹.




Figure 1: SRM of different outcome parameters.

The calculated Cohen's ES and SRM of the other two studies showed that all NSAIDs have statistically significantly moderate to good effect on BASFI, BASDAI, disease activity and pain (table 1 and figure 1). Various NSAIDs have a similar effect as assessed by treatment ES (table 2). Furthermore, no new signs of toxicity were discovered.

DMARDs

In 2005 and 2006, two Cochrane reviews summarized the effects of MTX and SSZ, respectively^{45,46}. The MTX review showed that there is no evidence to support any benefit of MTX in the treatment of AS. One additional open-label study about the effect of MTX was found besides the Cochrane review. The calculated Cohen's ES did not show any improvement on BASFI, BASDAI, pain or mobility (table 1 and figure 1)²⁴.

The SSZ Cochrane review showed some benefit of SSZ in reducing ESR and easing morning stiffness, yet no benefit in physical function, pain, spinal mobility and disease activity⁴⁶. These results are confirmed by three additional identified SSZ studies (level 1b and 3 evidences) not included in the SSZ Cochrane review²¹⁻²³ (supplementary appendix S4, available as supplementary data at Rheumatology Online). Although the calculated Cohen's ES revealed moderate to good effect on BASDAI and pain (table 1, figure 1), the calculated treatment ES showed that the effect of SSZ on these outcome parameters is not better than the effect of placebo, as shown in one study, and that the effect of SSZ was statistically significantly worse than the effect of etanercept (ETN) (table 2). No new signs of toxicity for SSZ and MTX were found.

Other therapies

Two studies about other types of therapy were identified. One study investigated the effect of probiotics compared with placebo (level 1b evidence)⁴⁷ (supplementary appendix S4, available as supplementary data at Rheumatology Online). The calculated ES showed that probiotics do no better on global well-being and functional index than placebo (tables 1 and 2). The other study investigated the effect of radium chloride on BASFI, BASDAI, pain and disease activity in an uncontrolled design (level 3 evidence) (table 1). The calculated ES demonstrated a moderate effect on BASFI and good effect on BASDAI, pain and disease activity (table 1 and figure 1)²⁵.

Surgical interventions

Total hip replacement

Total hip replacement (THR) is a frequently used procedure in AS patients with hip involvement. This search revealed one study about hip surgery and resurfacing of the hip (Zimmer, Wintherthur, Switzerland). The authors proposed that hip resurfacing might be an option instead of THR for young AS patients with hip involvement. They compared the effects of resurfacing with THR on pain relief, function and mobility in 38 resurfaced hips (23 AS patients) and 41 THRs (25 AS patients) over a mean follow-up time of 34.5 months. Both groups showed significant pain relief and good restoration of function and mobility²⁶ (table 1).

Spine

Although spinal surgery to resolve fixed kyphotic deformity is accompanied by severe risks, it can give excellent functional results by restoring balance and horizontal vision, as shown by all nine included papers in this search²⁷⁻³⁵. These papers review the different available techniques. All included papers are case series, and therefore low-quality studies (level of evidence 3) (supplementary Appendices S4 and S5, available as supplementary data at Rheumatology Online).

One study compared open wedge osteotomy (OWO) of the cervical spine with closed wedge osteotomy (CWO).

No difference in correction of kyphosis between the two techniques was found²⁸.

Table 2: Treatment ES with 95% confidence interval (95% CI) for the different interventions. The interventions are grouped together. The level of evidence, duration and point of measurement are described in this table.

Intervention	Comparator	Assessment point	Treatment ES BASFI (95% CI)	Treatment ES BASDAI (95% CI)	Treatment ES Pain (95% CI)	Treatment ES Disease act. (95% CI)	Treatment ES BASMI (95% CI)
Exercise therapy							
Group exercise ¹⁵	Home exercise	6 weeks	0.30 (-0.35, 0.95)	-0.06 (-0.71, 0.58)	0.31 (-0.34, 0.96)	-	0.20 (-0.44, 0.85)
Exercise Group ¹⁴	Control group	8 weeks	0.17 (-0.52, 0.87)	0.00 (-0.69, 0.69)	-	-	0.53 (-0.17, 1.24)
Hospital exercise ¹⁷	Home exercise	12 weeks	0.59 (0.01, 1.16)	0.22 (-0.36, 0.80)	0.68 (0.10, 1.26)	-	-
Hospital exercise	Home exercise	6 months	0.15 (-0.43, 0.72)	-0.17 (-0.74, 0.41)	0.04 (-0.54, 0.61)	-	-
Home exercise ¹⁸	Control group	12 weeks	0.49 (-0.12, 1.12)	0.84 (0.21, 1.47)	-	-	-
Exercise (GPR method) ¹⁹	Conventional exercise		0.01 (-0.60, 0.62)	-0.04 (-0.65, 0.57)	0.17 (-0.44, 0.79)	-	-
Conventional exercise	Control group	12 weeks	0.44 (-0.23, 1.11)	0.82 (0.13, 1.51)	0.93 (0.24, 1.63)	-	-
Exercise (GPR method)	Control group		0.41 (-0.26, 1.09)	0.75 (0.06, 1.44)	1.15 (0.43, 1.87)	-	-
Balneotherapy							
Exercise + stangerbath ²⁰	Exercise	3 weeks	0.65 (0.12, 1.18)	1.49 (0.91, 2.08)	-	-	0.21 (-0.31, 0.73)

Table 2: continued

Balneootherapy ²²	Balneootherapy + NSAIDs	-	-	0.08 (-0.53, 0.70)	-	-
Balneootherapy + NSAIDs	NSAIDs	3 weeks	-	0.36 (-0.26, 0.98)	-	-
Balneootherapy	NSAIDs		-	0.43 (-0.18, 1.04)	-	-
Balneootherapy	Balneootherapy + NSAIDs		-	0.00 (-0.61, 0.61)	-	-
Balneootherapy + NSAIDs	NSAIDs	6 months	-	0.31 (-0.31, 0.92)	-	-
Balneootherapy	NSAIDs		-	0.28 (-0.32, 0.89)	-	-
NSAIDs						
Continuous NSAID use (200-400 mg) ²⁶	On-demand NSAID use (200-400 mg)	104 weeks	-0.15 (-0.42, 0.12)	-0.053 (-0.32, 0.21)	0.1 (-0.17, 0.37)	-
Celecoxib (200 mg) ²⁷	Diclofenac (150 mg)	12 weeks	0 (-0.22, 0.22)	-0.30 (-0.52, 0.07)	-0.17 (-0.39, 0.06)	-
Celecoxib (400 mg)	Diclofenac (150 mg)		0.04 (-0.18, 0.27)	-0.10 (-0.32, 0.13)	-0.06 (-0.28, 0.17)	-
DMARDs						
Sulfasalazine (2 g) ³²	Placebo	24 weeks	0.08 (-0.17, 0.33)	0.15 (-0.10, 0.40)	0.05 (-0.20, 0.31)	-
Sulfasalazine (3 g) ³³	Etanercept (50 mg)	16 weeks	-0.19 (-0.37, -0.01)	-0.76 (-0.94, -0.58)	-0.70 (-0.88, -0.52)	-0.23 (-0.41, -0.06)

The interventions are grouped together. The level of confidence, duration and point of measurement are described in this table. Significant ES in **italics**.

Another study compared the conventional technique of cervical extension osteotomy with a new technique in which the patients have a modified larger lateral resection area than with the conventional technique. Again, no differences between the two techniques were found concerning functional improvement, satisfaction or complications³⁰. Similarly, a prone or a sitting position during the procedure demonstrated no difference in correction³³.

For thoracolumbar deformities, polysegmental wedge osteotomy might be associated with lower risks. However, the correction is often insufficient in the case of calcified intervertebral discs. Theoretically, CWO is superior to OWO in terms of efficiency and minimal loss of correction and lower accompanied risks, although technically difficult^{28,34}. For pseudoarthrosis, posterior correction is an effective treatment (posterior opening wedge osteotomy), as well as fixation without anterior fusion²⁷. The data from the included papers do not show whether a specific technique gives better results for any specific indication.

DISCUSSION

This systematic review is an update of the review by Zochling *et al.*⁴ and identified available nonpharmacological and non-biologic pharmacological treatments effective for symptomatic control of AS. The results of this search confirm the 2005 findings for physiotherapy⁴; exercises in various modalities, individually at home or in a group and under supervision, land or water based, have positive effects on BASFI, BASDAI, pain and mobility function. However, the small numbers of participants, the heterogeneity of the interventions and outcome measures, and deficiency in reporting data result in wide intervals and lack of strong evidence.

Zochling *et al.*⁴ revealed that different kinds of NSAIDs and coxibs improve spinal and peripheral joint pain and function. The current search confirmed these results by showing that various NSAIDs including coxibs improve BASDAI, disease activity and BASFI.

In 2005 no effect of SSZ or MTX on back pain and function was demonstrated⁴, which is confirmed by new research. The current search revealed no effect of SSZ and MTX on pain, nor on BASFI and BASDAI.

THR is still the standard procedure in AS patients with hip involvement. Although a small study showed positive effects of hip resurfacing techniques²⁶, it must be carefully considered whether resurfacing techniques are indeed a good alternative for THR given the recent developments and accompanying problems with the resurfacing techniques from another brand. The articular surface replacement hip prosthesis from the manufacturer DePuy (Warsaw, IN, USA) has been recalled from the market because of failing of the prosthesis. Metal debris from wear of the implant led to a reaction that destroyed the soft tissues surrounding the joint, causing long-term disability and a high revision rate of 12% over 5 years⁴⁸.

Surgical interventions of the spine give excellent results by restoring horizontal gaze and function yet are considered with high risks. Furthermore, it is still unclear which procedure of spine surgery is the best for any specific indication.

CONCLUSION

This review presents a concise summary of the current evidence available for therapeutic interventions for the management of AS, both non-pharmacological and pharmacological, excluding biologics. This overview formed the basis for the update of the ASAS/EULAR recommendations for the management of AS.

SUPPLEMENTARY DATA

Supplementary data are available at Rheumatology Online.

REFERENCES

1. Braun J, Pham T, Sieper J, *et al.* International ASAS consensus statement for the use of anti-tumour necrosis factor agents in patients with ankylosing spondylitis. *Ann Rheum Dis* 2003;62:817-24.
2. van der Heijde D, Landewé R, Baraliakos X, *et al.* Radiographic findings following two years of infliximab therapy in patients with ankylosing spondylitis. *Arthritis Rheum* 2008;58:3063-70.
3. Zochling J, van der Heijde D, Burgos-Vargas R, *et al.* ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis* 2006;65:442-52.
4. Zochling J, van der Heijde D, Dougados M, *et al.* Current evidence for the management of ankylosing spondylitis: a systematic literature review for the ASAS/EULAR management recommendations in ankylosing spondylitis. *Ann Rheum Dis* 2006;65:423-32.
5. Baraliakos X, Listing J, Rudwaleit M, *et al.* Radiographic progression in patients with ankylosing spondylitis after 2 years of treatment with the tumour necrosis factor alpha antibody infliximab. *Ann Rheum Dis* 2005;64:1462-6.
6. Braun J, Brandt J, Listing J, *et al.* Treatment of active ankylosing spondylitis with infliximab: a randomised controlled multicentre trial. *Lancet* 2002;359:1187-93.
7. Davis J, van der Heijde D, Braun J, *et al.* Recombinant human tumor necrosis factor receptor, (etanercept) for treating ankylosing spondylitis—a randomized, controlled trial. *Arthritis Rheum* 2003;48:3230-6.
8. van der Heijde D, Kivitz A, Schiff M, *et al.* Efficacy and safety of adalimumab in patients with ankylosing spondylitis: results of a multicenter, randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2006;54:2136-46.
9. Braun J, van den Berg R, Baraliakos X, *et al.* update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis* 2011;70:896-904.
10. Karapolat H, Akkoc Y, Sari I, *et al.* Comparison of group-based exercise versus home-based exercise in patients with ankylosing spondylitis: effects on Bath Ankylosing Spondylitis Indices, quality of life and depression. *Clin Rheumatol* 2008;27:695-700.
11. Ortancil O, Sarikaya S, Sapmaz P, *et al.* The effect(s) of a six-week home-based exercise program on the respiratory muscle and functional status in ankylosing spondylitis. *J Clin Rheumatol* 2009;15:68-70.
12. Widberg K, Hossein K, Hafstrom I. Self- and manual mobilization improves spine mobility in men with ankylosing spondylitis—a randomized study. *Clin Rehabil* 2009;23: 599-608.
13. Cagliyan A, Kotevoglou N, Onal T, *et al.* Does group exercise program add anything more to patients with ankylosing spondylitis? *J Back Musculoskelet Rehabil* 2007;20:79-85.
14. Durmus D, Alayli G, Cil E, *et al.* Effects of a home-based exercise program on quality of life, fatigue, and depression in patients with ankylosing spondylitis. *Rheumatol Int* 2009;29:673-7.
15. Durmus D, Alayli G, Uzun O, *et al.* Effects of two exercise interventions on pulmonary functions in the patients with ankylosing spondylitis. *Joint Bone Spine* 2009;76:150-5.
16. Gurcay E, Yuzer S, Eksioglu E, *et al.* Stanger bath therapy for ankylosing spondylitis: illusion or reality? *Clin Rheumatol* 2008;27:913-7.
17. Yurtkuran M, Ay A, Karakoc Y. Improvement of the clinical outcome in ankylosing spondylitis by balneotherapy. *Joint Bone Spine* 2005;72:303-8.
18. Lubrano E, D'Angelo S, Parsons W, *et al.* Effectiveness of rehabilitation in active ankylosing spondylitis assessed by the ASAS response criteria. *Rheumatology* 2007;46: 1672-5.
19. Sieper J, Klopsch T, Richter M, *et al.* Comparison of two different dosages of celecoxib with diclofenac for the treatment of active ankylosing spondylitis: results of a 12-week randomised, double-blind, controlled study. *Ann Rheum Dis* 2008;67:323-9.
20. Jarrett SJ, Sivera F, Cawkwell L, *et al.* MRI and clinical findings in patients with ankylosing spondylitis eligible for anti-tumour necrosis factor therapy after a short course of etoricoxib. *Ann Rheum Dis* 2009;68:1466-9.
21. Braun J, Zochling J, Baraliakos X, *et al.* Efficacy of sulfasalazine in patients with inflammatory back pain due to undifferentiated spondyloarthritis and early ankylosing spondylitis: a multicentre randomised controlled trial. *Ann Rheum Dis* 2006;65:1147-53.

22. Braun J, van der Horst-Bruinsma I, Huang F, *et al.* Clinical efficacy and safety of etanercept versus sulfasalazine in patients with ankylosing spondylitis: a randomized, double-blind trial. *Arthritis Rheum* 2011;63:1543-51.
23. Genc H, Duyur CB, Nacir B, *et al.* The effects of sulfasalazine treatment on enthesal abnormalities of inflammatory rheumatic diseases. *Clin Rheumatol* 2007; 26:1104-10.
24. Haibel H, Brandt H, Song I, *et al.* No efficacy of subcutaneous methotrexate in active ankylosing spondylitis: a 16-week open-label trial. *Ann Rheum Dis* 2007;66:419-21.
25. Alberding A, Stierle H, Brandt J, *et al.* Effectiveness and safety of radium chloride in the treatment of ankylosing spondylitis. Results of an observational study. *Z Rheumatol* 2006;65:245-51.
26. Li J, Xu W, Xu L, *et al.* Hip resurfacing arthroplasty for ankylosing spondylitis. *J Arthroplasty* 2009;24:1285-91.
27. Chang K, Tu M, Huang H, *et al.* Posterior correction and fixation without anterior fusion for pseudoarthrosis with kyphotic deformity in ankylosing spondylitis. *Spine* 2006;31:E408-13.
28. Chang K, Chen Y, Lin C, *et al.* Closing wedge osteotomy versus opening wedge osteotomy in ankylosing spondylitis with thoracolumbar kyphotic deformity. *Spine* 2005;30:1584-93.
29. Belanger T, Milam R, Roh J, *et al.* Cervicothoracic extension osteotomy for chin-on-chest deformity in ankylosing spondylitis. *J Bone Joint Surg Am* 2005;87:1732-8.
30. Simmons E, DiStefano R, Zheng Y, *et al.* Thirty-six years experience of cervical extension osteotomy in ankylosing spondylitis: techniques and outcomes. *Spine* 2006;31:3006-12.
31. Kim K, Lee S, Suk K, *et al.* Spinal pseudarthrosis in advanced ankylosing spondylitis with sagittal plane deformity: clinical characteristics and outcome analysis. *Spine* 2007;32:1641-7.
32. Min K, Hahn F, Leonardi M. Lumbar spinal osteotomy for kyphosis in ankylosing spondylitis: the significance of the whole body kyphosis angle. *J Spinal Disord Tech* 2007;20:149-53.
33. Langeloo D, Journee H, Pavlov PW, *et al.* Cervical osteotomy in ankylosing spondylitis: evaluation of new developments. *Eur Spine J* 2006;15:493-500.
34. Tokala D, Lam K, Freeman B, *et al.* C7 decancellation closing wedge osteotomy for the correction of fixed cervico-thoracic kyphosis. *Eur Spine J* 2007;16:1471-8.
35. Brox J, Helle A, Sorensen R, *et al.* Functional outcome after lumbar closing wedge osteotomy in ankylosing spondylitis. *Int Orthop* 2009;33:1049-53.
36. Wu J, Jiang G, Wei W. Confidence intervals of effect size in randomized comparative parallel-group studies. *Stat Med* 2006;25:639-51.
37. Dagfinrud H, Kvien T, Hagen K. Physiotherapy interventions for ankylosing spondylitis. *Cochrane Database Syst Rev* 2008; CD002822.
38. Fernandez-de-Las-Penas C, Alonso-Blanco C, Morales-Cabezas M, *et al.* Two exercise interventions for the management of patients with ankylosing spondylitis: a randomized controlled trial. *Am J Phys Med Rehabil* 2005;84:407-19.
39. Fernandez-de-Las-Penas C, Alonso-Blanco C, Alguacil-Diego IM, *et al.* One-year follow-up of two exercise interventions for the management of patients with ankylosing spondylitis: a randomized controlled trial. *Am J Phys Med Rehabil* 2006; 85:559-67.
40. Ince G, Sarpel T, Durgun B, *et al.* Effects of a multimodal exercise program for people with ankylosing spondylitis. *Phys Ther* 2006;86:924-35.
41. Wanders A, van der Heijde D, Landewé R, *et al.* Nonsteroidal antiinflammatory drugs reduce radiographic progression in patients with ankylosing spondylitis: a randomized clinical trial. *Arthritis Rheum* 2005;52:1756-65.
42. Codish S, Dobrovinsky S, Abu S, *et al.* Spa therapy for ankylosing spondylitis at the Dead Sea. *Isr Med Assoc J* 2005;7:443-6.
43. Altan L, Bingol U, Aslan M, *et al.* The effect of balneotherapy on patients with ankylosing spondylitis. *Scand J Rheumatol* 2006;35:283-9.
44. Dougados M, Behier J, Jolchine I, *et al.* Efficacy of celecoxib, acyclooxigenase 2-specific inhibitor, in the treatment of ankylosing spondylitis—a six-week controlled study with comparison against placebo and against a conventional nonsteroidal antiinflammatory drug. *Arthritis Rheum* 2001;44:180-5.
45. Chen J, Liu C, Lin J. Methotrexate for ankylosing spondylitis. *Cochrane Database Syst Rev* 2006;CD004524.

46. Chen J, Liu C. Sulfasalazine for ankylosing spondylitis. *Cochrane Database Syst Rev* 2005;CD004800.
47. Brophy S, Burrows CL, Brooks C, *et al.* Internet-based randomised controlled trials for the evaluation of complementary and alternative medicines: probiotics in spondyloarthropathy. *BMC Musculoskelet Disord* 2008;9:4.
48. Cohen D. Out of joint: the story of the ASR. *BMJ* 2011;342: d2905.

